



## RESEARCH ARTICLE

# Identifying ways to maximise cervical screening uptake: a qualitative study of GPs' and practice nurses' cervical cancer screening-related behaviours [version 1; peer review: 2 approved]

Mairead O'Connor <sup>1\*</sup>, Lisa A. McSherry<sup>2\*</sup>, Stephan U. Dombrowski<sup>3</sup>, Jill J. Francis<sup>4</sup>, Cara M. Martin<sup>5</sup>, John J. O'Leary <sup>5</sup>, Linda Sharp<sup>6</sup>

<sup>1</sup>School of Public Health, Univesity College Cork, Cork, Ireland

<sup>2</sup>National Cancer Registry of Ireland, Kinsale Road, Cork, Ireland, Ireland

<sup>3</sup>Faculty of Kinesiology, University of New Brunswick, Fredericton, NB, Canada

<sup>4</sup>School of Health Sciences, City University London, London,, UK

<sup>5</sup>Trinity College Dublin, Dublin, Ireland

<sup>6</sup>Institute of Health and Society, Newcastle University, Newcastle, UK

\* Equal contributors

**V1** First published: 05 May 2021, 4:44  
<https://doi.org/10.12688/hrbopenres.13246.1>  
 Latest published: 05 May 2021, 4:44  
<https://doi.org/10.12688/hrbopenres.13246.1>

## Abstract

**Background:** Cervical screening uptake is declining in several countries. Primary care practitioners could play a greater role in maximising uptake, but better understanding is needed of practitioners' cervical screening-related behaviours. Among general practitioners (GPs) and practice nurses, we aimed to identify cervical screening-related clinical behaviours; clarify practitioners' roles/responsibilities; and determine factors likely to influence clinical behaviours.

**Methods:** Telephone interviews were conducted with GPs and practice nurses in Ireland. Interview transcripts were analysed using the Theoretical Domains Framework (TDF), a comprehensive psychological framework of factors influencing clinical behaviour.

**Results:** 14 GPs and 19 practice nurses participated. Key clinical behaviours identified were offering smears and encouraging women to attend for smears. Smeartaking responsibility was considered a predominantly female role. Of 12 possible theoretical domains, 11 were identified in relation to these behaviours. Those judged to be the most important were beliefs about capabilities; environmental context and resources; social influences; and behavioural regulation. Difficulties in obtaining smears from certain subgroups of women and inexperience of some GPs in smeartaking arose in relation to beliefs about capabilities. The need for public health education and

## Open Peer Review

Reviewer Status

Invited Reviewers

1

2

version 1

05 May 2021



report



report

1. **John Campbell**, University of Exeter, Exeter, UK

2. **Suhailah M Ali** , College of Nursing, University of Kirkuk, Kirkuk, Iraq

Any reports and responses or comments on the article can be found at the end of the article.

reluctance of male practitioners to discuss cervical screening with female patients emerged in relation to social influences.

**Conclusions:** We identified - for the first time - primary care practitioners' cervical-screening related clinical behaviours, their perceived roles and responsibilities, and factors likely to influence behaviours. The results could inform initiatives to enable practitioners to encourage women to have smear tests which in turn, may help increase cervical screening uptake.

### Keywords

cervical screening, women, primary care practitioners, clinical behaviours, the Theoretical Domains Framework

**Corresponding author:** Mairead O'Connor ([maireadoconnor@ucc.ie](mailto:maireadoconnor@ucc.ie))

**Author roles:** **O'Connor M:** Visualization, Writing – Review & Editing; **McSherry LA:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Dombrowski SU:** Methodology, Writing – Review & Editing; **Francis JJ:** Conceptualization, Methodology, Writing – Review & Editing; **Martin CM:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **O'Leary JJ:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **Sharp L:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Methodology, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** CMM and JJOL have received funding from Hologic (<https://www.hologic.com/>) for evaluating the CERVISTA HPV test.

**Grant information:** This research was funded by the Friends of the Coombe Hospital charity (no grant number available), based at the Coombe Women and Infants University Hospital, Dublin, Ireland. This study was nested within the ATHENS trial, under the umbrella of the CERVIVA research consortium ([www.cerviva.ie](http://www.cerviva.ie)). MO'C has worked on the following HRB grants since 1 January 2017: ICE-2015-1037 [ICE Research Fellow] and APA 2016-1874 [co-applicant]. CMM has held the following HRB grants since 1 January 2017: ILP-PHR-2017-103 [Lead applicant], APA 2016-1874 [co-applicant] and ICE-2015-1037 [co-applicant]. JJ'OL has held the following HRB grants since 1 January 2017: HRB-ARPP-A-2018-018 [co-applicant], ILP-2017-022 [co-applicant]. APA 2016-1874 [Co-Lead applicant] and ICE-2015-1037 [Lead applicant]. LS has held the following HRB grants since 1 January 2017: ILP-PHR-2017-013 [co-applicant], APA 2016-1874 [co-applicant] and ICE-2015-1037 [co-applicant].

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2021 O'Connor M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** O'Connor M, McSherry LA, Dombrowski SU *et al.* **Identifying ways to maximise cervical screening uptake: a qualitative study of GPs' and practice nurses' cervical cancer screening-related behaviours [version 1; peer review: 2 approved]** HRB Open Research 2021, 4:44 <https://doi.org/10.12688/hrbopenres.13246.1>

**First published:** 05 May 2021, 4:44 <https://doi.org/10.12688/hrbopenres.13246.1>

## Introduction

Well-organised cervical screening is effective in reducing cervical cancer incidence and mortality<sup>1,2</sup>. One of the keys to screening success is maximising uptake<sup>3</sup>. Thus, it is concerning that screening uptake is declining in several developed countries, including England, Sweden, and Australia<sup>4</sup>. For example, in England, the proportion of eligible women being screened has declined from 76% in 2011 to 72% in 2017<sup>5</sup>. In the U.S, a downward trend in smear test use has been shown from 2000 to 2015<sup>6</sup>. There is currently considerable interest in developing initiatives and strategies to reverse these concerning trends. For example, the Cervical Screening Programme (CSP) in England has affirmed that halting this trend is a key programme objective and are supporting research in this area<sup>7</sup>.

Cervical screening is a significant component of the primary care workload. Evidence is emerging that the screening-related attitudes and clinical behaviours of primary care practitioners (e.g. general practitioners (GPs), practice nurses) influence women's cervical screening behaviours. For example, a previous bad experience in attending for a smear deters women from attending for another smear<sup>8,9</sup>. In contrast, a considerate smear taker who takes care to minimise pain and discomfort may positively influence women's future participation. Moreover, women report that GP prompting, and a positive GP attitude, are important motivators for having smears<sup>10</sup>, while a negative or dismissive GP attitude is a significant barrier<sup>11</sup>, findings replicated in various settings<sup>12,13</sup>.

To date, the evidence of practitioners' influences on screening uptake has concentrated on doctors. Practice nurses are increasingly involved in delivering cervical screening in several settings, including the UK, Ireland and Australia<sup>14-16</sup>. It seems likely, therefore, that practice nurses may influence women's screening-related behaviours in similar ways to GPs<sup>17-19</sup>.

The influence of primary care practitioners on women's behaviours raises the possibility that there may be opportunities to engage practitioners more actively and/or directly in maximising cervical screening uptake. As a first step, there is a need to better understand what practitioners consider to be their responsibilities with regard to screening and what factors may influence their screening-related clinical behaviours. We undertook a qualitative study with general practitioners (GPs) and practice nurses to explore different clinical behaviours around cervical cancer prevention - HPV testing, HPV vaccination and smear tests/cytological screening. The results on HPV-related clinical behaviours (e.g. initiating a discussion about HPV with female patients, recommending HPV vaccination) have been previously published<sup>20</sup>. The current paper focuses on results related to cervical screening related clinical behaviours with the following aims: to: (1) identify cervical screening-related clinical behaviours; (2) clarify practitioners' roles and what they consider to be their responsibilities in relation to cervical screening; and (3) determine factors likely to influence their cervical-screening related clinical behaviours.

## Methods

### Setting

The study setting was Ireland which has a mixed public/private healthcare system. At the time the study was conducted, Ireland's health service (the Health Service Executive (HSE)) was divided up into four geographical health board areas covering the Republic of Ireland (HSE Mid-Eastern, HSE North-Eastern, HSE Southern and HSE Western). GPs are effectively private practitioners but may provide services for certain patients (typically those with low incomes) under the public system for which they are reimbursed by the state<sup>21</sup>. The national screening programme, CervicalCheck, began in 2008 and offers free cervical screening for women aged 25–65 years in primary care. Women are invited, through a centralised call-recall system. A woman may choose to have a screening test (at her general practice or with any other registered screening test taker). CervicalCheck reimburses practitioners for screening tests taken within guidelines; additional/opportunistic screening tests are not reimbursed. There are currently no screening uptake targets for individual practices, but nationally, the programme achieved 75% coverage of the eligible population in its first five years of operation<sup>22</sup>. Until very recently, the programme used smear tests as its primary screening tool – women were invited for smear tests every 3 years (for women aged 25–44) or 5 years (for older women). Traditionally, GPs were the primary smeaftakers, but now, practice nurses carry out the vast majority of cervical screening tests within CervicalCheck<sup>23</sup>, although GPs retain clinical responsibility for the provision of the service in their practice. In March 2020, the programme began replacing traditional cervical cytology (smear) with primary HPV testing in line with best international recommendations – now women aged 25–29 years are screened every 3 years and those aged 30–65 are screened every 5 years. Irrespective of the recent changes to the screening programme, the overall screening process from the point of view of both the woman and the test taker (e.g. registering as a screening test taker, a woman choosing where to have her screening test) remains largely unchanged.

### Participants

GPs and practice nurses working in Ireland were eligible to participate. GPs were recruited via postal invitation from a group of 145 GPs who had: been randomly sampled from a national GP database (comprising approximately 2,000 registered GPs in Ireland), participated in a cervical screening survey in 2007, and indicated they were potentially willing to assist with further research<sup>24</sup>. The group was diverse in terms of personal and practice characteristics (in 2007). A purposive sample was recruited from this group of 145 GPs. Sampling strata was defined in terms of variables that had been found in the 2007 survey to be strongly associated with attitudes towards smear taking. These variables were: GP gender, years since graduation, area of practice location (HSE health board area). Since there is no national practice nurse database two routes were used to recruit nurses to ensure maximum sample variation. Attendees at a national Irish Practice Nurses Association (IPNA)

conference were invited to participate and postal invitations were dispatched to randomly selected nurses via area-based practice nurse professional development coordinators (PDCs) across Ireland. All practitioners who returned a reply slip were contacted for interview. Ethical approval was obtained from the Irish College of General Practitioners.

## Procedures

Semi-structured interviews took place between November 2010 and February 2011. Participants received an information sheet about the research and provided written informed consent before the interview commenced. Telephone interviews (lasting 16 to 50 minutes in duration) were conducted (by LAMcS, ATHENS research investigator. ATHENS - an intervention trial of HPV education and support in primary care) and guided by a topic guide (see extended data<sup>25</sup>). The guide included open “core” questions and clinical scenarios designed to elicit information about cervical cancer screening-related clinical behaviours, roles/responsibilities, and drivers of clinical behaviours. The same core questions were asked of both GPs and practice nurses. The order in which the core questions were asked, and the content and order of follow-up questions and prompts, varied between practitioners.

Recruitment continued until new issues ceased to emerge for GPs and practice nurses separately (i.e. data saturation was reached in each practitioner group (practitioner group – (1) GPs or (2) practice nurses<sup>26</sup>). All except three interviews were audio-recorded and transcribed verbatim (the three interviews were not audio-recorded as study participants declined consent for their interviews to be recorded). When interviews were not recorded, the interviewer took detailed notes contemporaneously. The study adhered to the Standards for reporting qualitative research: a synthesis of recommendations (SRQR)<sup>27</sup> (see extended data for completed SRQR checklist<sup>25</sup>).

## Analysis

GP and practice nurse interviews were analysed collectively. No qualitative data software was used for analysis. Analysis was conducted following the Framework Analysis approach<sup>28,29</sup>, to identify key cervical screening-related clinical behaviours and roles/responsibilities of practitioners. We used the Theoretical Domains Framework (TDF) as the coding framework<sup>30</sup> to determine the factors likely to influence the behaviours. Developed through an expert consensus process, the TDF summarises multiple psychological and organisational theories regarding influences on clinical behaviour in 12 theoretical domains: *knowledge; skill, social/professional role and identity; beliefs about capabilities; beliefs about consequences; motivation and goals; memory, attention and decision processes; environmental context and resources; social influences; emotion; behavioural regulation; and nature of the behaviour*. It is a leading theoretical framework for developing interventions aimed at changing clinical behaviour. Two investigators (LAMcS, LS) read and reread all transcripts, independently coded these, combined codes into subthemes and allocated these, and participants’ direct quotes, to the TDF domains. For analytical rigour, a second iteration of this process was performed with uncertainties

resolved in discussion with co-investigators with expertise in behaviour change (JJF, SUD). The domains (themes) considered strongest/dominant were those: (1) mentioned by most practitioners; (2) where most subthemes were identified; and (3) which were discussed at greatest length<sup>31</sup>. Whether subthemes arose solely among male/female GPs, practice nurses or both was noted. Illustrative quotes are provided to supplement narrative descriptions. Study participants were not invited to provide feedback on the interview data but were sent a final report on study findings.

## Results

Of the 145 GPs contacted, 19 responded (i.e. returned a reply slip). All 19 GPs who responded were interviewed. Of the 30 practice nurses approached through PDCs, ten were interviewed; four nurses were recruited from the annual conference. [Table 1](#) summarises participants’ characteristics.

### Clinical behaviours

Two cervical screening-related clinical behaviours were identified among both GPs and practice nurses: (1) offering smears to women and (2) encouraging women to attend for smears.

### Roles and responsibilities

Taking smears was considered a predominantly female role with responsibility falling to female GPs and practice nurses (all of whom are female). Female practitioners frequently assumed that women prefer to have the option of a female smearer, making comments like “I do.....99% of them because the other two GPs are male (PN5020)” and “it’s a male doctor and a lot of the patients don’t seem to.....[want to]....avail of a male (PN5040).” Male GPs performed smears less often and made comments like “I do an occasional one when a patient requests it (GP0133).”

### Factors influencing Clinical behaviours

[Table 2](#) displays the 12 theoretical domains, subthemes identified which relate to the individual domains, and illustrative quotes. All but one of the 12 theoretical domains, *motivation and goals*, influenced both offering smears and encouraging women to attend ([Table 2](#)). The dominant domains were: *beliefs about capabilities, environmental context and resources, behavioural regulation and social influences*. These are described in more detail below.

In terms of *beliefs about capabilities*, both practitioner groups spoke about difficulties in getting women to attend for smears. Once patients attend, practitioners reported difficulties in dealing with certain subgroups (e.g., women who are extremely anxious or have a history of sexual abuse). This was mentioned particularly frequently by female GPs. Some practice nurses described referring any cases expected to be “difficult” to GPs. One practice nurse noted that GPs may be reluctant to admit inexperience in smear-taking, which may have implications for quality assurance.

A lack of time to take smears, inadequate facilities and equipment, limited appointment times and limited practice

**Table 1. Characteristics of practitioners interviewed.**

Characteristic		GPs	Practice Nurses
Sex	Female	13	14
	Male	6	0
Health board area	HSE Mid-Eastern	4	3
	HSE North-Eastern	3	4
	HSE Southern	7	0
	HSE Western	5	7
Location of practice	City	6	6
	Other	13	8
Solo GP practice	Yes	6	7
	No	13	7
Practice nurse(s) employed in practice	Yes	16	-
	No	3	-
Years since graduation <sup>1</sup>	<10 years	2	-
	10–19 years	2	-
	20+ years	14	-

<sup>1</sup>from responses to 2007 GP survey

support staff emerged in relation to *environmental context and resources*. Practitioners also noted that some women have accessibility issues making it difficult for them to attend the practice. Some practice nurses mentioned that difficulties arise because CervicalCheck does not hold women's full smear and gynaecological histories (the programme databases only hold information from 2008 onwards). For example, CervicalCheck may be unaware that a woman has had a hysterectomy and may send her a smear invitation. GPs expressed a desire for clearer CervicalCheck guidelines on patient follow up after an abnormal smear.

In terms of behavioural regulation, some practitioners recognised opportunities to increase uptake by offering dedicated smear clinics and extended practice hours. Practice nurses suggested practical tools, such as flipcharts to use during consultations or clinic posters advertising the service, as potential means of improving the service. Many practitioners wanted to have the option of offering free smears to women outside the screening age range, if the practitioner felt this was warranted (e.g., for some women under 25). Furthermore, some criticised

the policy of discouraging opportunistic smears at the practitioners' discretion, and expressed concern that the recommended recall period was too long.

As regards *social influences*, both GPs and practice nurses believed there is a need for more public education around smears; they considered that many women do not know much about screening or smear tests, in part because the programme is relatively new. Practice nurses recognised that some women are not keen, or may be afraid, to attend for smears. GPs considered that male GPs may be reluctant to broach the topic of smears with female patients. They also noted that women are not yet familiar with nurse-led services and this may discourage women from attending.

## Discussion

### Summary of main findings

The major cervical screening-related clinical behaviours identified among primary care practitioners were offering smears and encouraging women to attend for smears. Responsibility for taking smears was considered a predominantly female role,

**Table 2. Factors influencing behaviours related to cervical screening.**

Theme / construct domain	Subtheme / specific belief	Practitioner*		Sample quotes
		GPs	Practice Nurses	
<b>1. Knowledge</b>	Unaware that they can still take a smear if the patient is not already registered	ü	-	<i>"It also depends on whether she is actually registered with CervicalCheck, because we are being told they won't process smears unless they are registered" - GP 0058</i>
	Unaware that they can post 1-2 smear samples back at a time	ü	-	<i>"The cardboard boxes which are packed in sixes or fours I can't remember so there's kind of a posting issue that you kind of accumulate... get a few together for the post.... A small issue" - GP 0133</i>
	Unaware that they can take a smear between days 10-20 of woman's cycle	-	ü	<i>"Sometimes the women arrive and they're not in the required time, they're not between day 10 and 20" - Practice nurse 5017</i>
	Difficulty taking a smear from some patients e.g., obese, learning difficulties	ü	ü	<i>"Sometimes there are technical issues e.g. if you have a very obese patient or a patient with an arthritic hip that can be physically challenging" - GP 0136</i>
<b>2. Skill</b>	Difficulty visualising the cervix	ü	ü	<i>"Occasionally it can be difficult to visualise the cervix" - GP 0090</i>
	Difficulty assessing whether a patient has been sexually active	-	ü	<i>"It's very hard sometimes to I suppose in one sense assess whether this person had been sexually active or not.... It sometimes can be a sensitive issue on bringing up whether they've been sexually active or not" - Practice nurse 5034</i>
	Ability to put patients at ease	ü	-	
	Ability to explain to patient what smear test is for & differentiate it from STI testing	ü	-	<i>"Often people come for smears and what they're really looking for is STI testing" - GP 0003</i>
<b>3. Social / professional role &amp; identity</b>	GPs becoming deskilled	ü	-	<i>"I wouldn't be particularly upskilled at the moment" - GP 0026</i>
	Role of smearer is shifting:			
	Male GPs less likely to take smears	ü	ü	<i>"Harder for male GPs as most women want to go to a female doctor" - GP 0016</i>
	GP role more auxiliary i.e., nurse may refer to GP if complications arise	ü	ü	<i>"The nurse does the smears ur and we supervise" - GP 0026</i>
	Feel responsible for recall	ü	-	<i>"It looks like it's down to us to make sure they're recalled for their second smear" - GP 0141</i>
	CervicalCheck training not mandatory for GPs	-	ü	



Theme / construct domain	Subtheme / specific belief	Practitioner*		Sample quotes
		GPs	Practice Nurses	
<b>4. Beliefs about capabilities</b>	Difficulty dealing with awkward or sensitive situations e.g. anxious woman	ü	ü	"What can make it difficult? Well I guess mmm a difficult exam, just ur either patient anxiety or 'fretfulness" -GP 0026
	Need for good local contacts for advice & to refer technically challenging smears	ü	ü	
	Difficulty getting women to attend for smears	ü	ü	"The only problem is getting them to come in" - GP 0034
	Difficulty dealing with women who have never been sexually active	ü	-	"The only time I'd have difficulty is with is with somebody if they haven't been sexually active. Now it's not a barrier really I suppose but you have to tease that out obviously with the person" -GP 0072
<b>5. Beliefs about consequences</b>	GP reluctance to admit inexperience	-	ü	"We've had people who've never taken a smear test before and you mightn't realise it. And sometimes people won't tell you that mmmm I think sometimes medical people are not as quick to say I've never done that before or I'm not 100% sure" - Practice nurse 5001
	Practitioners are no longer in control of the recall system	-	ü	"What I found hard to come to terms with at the start was I felt I had control of the actual system" - Practice nurse 5040
	CervicalCheck not aware of patient history - caused some problems	ü	ü	"We're told under 25 is not of much significance.... one would wonder about that with you know promiscuity at the rate it's at, at the moment with these young girls..." - GP 0129
	Belief that it is an unpleasant / intrusive test for women	ü	ü	"I suppose the ideal thing would that mmm it wouldn't be a test you know the test itself is physically mmm unpleasant.... Ideally if you could swab from the mouth.... Or take a urine sample or something like that. But the actual having to get the sample from the place you have to get it is going to make it difficult" - GP 0058
<b>6. Motivation and goals</b>	Belief that women who don't have a smear 'there and then' will not come back at another time	ü	ü	
	Belief that payment for taking smears is too low	ü	-	"The payment (for smear taking) is derisory, it's ridiculous, I mean the amount of work and time that goes into it" - GP 0058
<b>7. Memory, attention and decision processes</b>	Having a reminder system for follow-ups	ü	ü	"I actually keep the manual register as well so ur it means that if any smear result doesn't come back I can see it" - Practice nurse 5002

Theme / construct domain	Subtheme / specific belief	Practitioner*		Sample quotes
		GPs	Practice Nurses	
<b>8. Environmental context and resources</b>	Don't have enough time to take smears	ü	ü	"And I suppose time..... Its not difficult its just time really" - GP 0058
	Inadequate facilities / equipment	ü	ü	
	Limited practice staff support	ü	ü	"I think its having nursing time really. I think having more support" - GP 0060
	Limited appointment times	ü	ü	"Maybe times are a problem - their surgery is 9-5 so it might be difficult for women to get there if they are working" - Practice nurse 5026
	Transport / accessibility issues for some patients	ü	ü	"Transport here is a big thing for some people..... They live quite rural and they don't have a lift in at the last minute" - Practice nurse 5040
	Lack of foreign language leaflets / language barriers	-	ü	"Sometimes explaining to immigrant population is difficult" - Practice nurse 5036
	Need better advertising of service	-	ü	
	Difficulty getting records for women with history of hysterectomy	-	ü	"The big problem we're having a lot now is em women with histories of hysterectomies.....but the problem is getting back to the hospitals. Like some of these might have had their hysterectomy like 20 - 20 odd years ago. Em. And we don't know the full extent" - Practice nurse 5004
	Women not used to CervicalCheck programme e.g. recall system	ü	ü	"Some women find it difficult to accept the recall system that's now in place" - GP 0129
	Need for more public education	ü	ü	"I think the general public need to be made more aware of this service" - GP 0129
<b>9. Social influences</b>	Women don't know a lot about smears	ü	ü	"I'm surprised at people's lack of knowledge" - GP 0003
	Women choose who to go to for smears	ü	-	"Patients just make an appointment for a smear with the doctor or nurse of their choice" - GP 0051
	Women not used to nurse led service	ü	-	"A lot of people aren't that use to the whole nurse thing you know" - GP 0060
	Male GPs reluctant to broach topic with female patients	ü	-	"I think generally speaking ur there is a hole there where single-handed GPs may not broach the subject with female patients and the female patients may not feel comfortable going to the men"- GP 0072
	Some women don't want to have smears taken then and there	-	ü	
	Women are not keen / afraid to come in for smears	-	ü	"A lot of patients are actually afraid to come for a smear" - Practice nurse 5023
<b>10. Emotion</b>	Frustration because of difficulty in getting smear from some patients	-	ü	"There's nothing more frustrating you have, twenty minutes to do a smear and you can't find the cervix" - Practice nurse 5023
	Awkwardness of practitioner (if woman outside of programme age range wants a smear)	ü	-	"It's just awkward" -GP 0003



Theme / construct domain	Subtheme / specific belief	Practitioner*		Sample quotes
		GPs	Practice Nurses	
11. Behavioural regulation	Belief that patients under 25 years should have access to the programme	ü	ü	
	Belief that not allowing opportunistic smearing was a mistake	ü	ü	"I do think there is a bit more of a place for somebody just doing mmm and opportunistic smear. That we're no longer able to do" - GP 0046
	Concern about recall period	ü	ü	
	Offering appointment system / dedicated smear clinics	ü	ü	"I facilitate working ladies maybe on Saturday evening if I'm working duty for the weekend... we do everything to facilitate the appointment" - GP 0129
	Having a reminder system for follow-ups	ü	ü	"Send her a text to remind her to go (for follow-up test)" - Practice nurse 5017
	Offering extended clinic opening hours	ü	-	
	Having leaflets available to give to patients	ü	ü	"I give the ladies a leaflet explaining the results before when they come and do their smear" - Practice nurse 5034
	Having posters and signs advertising the service	-	ü	"We have posters up in the, still up in the clinic" - GP 0060
	Having "Flipcharts" to aid patient consultations	-	ü	"It's a cardboard chart with like a ring thing at the top and it just explains what a smear test is...but it's quite simplified in fairness for the patients" - Practice nurse 5003
	Having good secretarial staff to book clinics and liaise with women	ü	-	
12. Nature of the behaviour	Having clearer CervicalCheck guidelines	ü	-	"If there were much clearer guidelines.... What was set out as protocol and has actually come in are quite different" - GP 0141
	Having a standard set of letters to cover every eventuality	ü	-	
	Having access to records for women with a history of hysterectomy	-	ü	
	More consistency in what staff record on smear forms (can affect follow-up recommendations)	ü	-	Some people mightn't put down as much information on the form as others say in the surgery and it could come back like a three year recall for by rights when it should have been on an annual recall" - GP 0141
	Team meetings involving nurses and GPs	ü	-	
	Take a lot of smears	ü	-	
	Girls starting sexual behaviour at an earlier age	-	ü	"In my, my bigger surgery mmm the, the women, well the girls, they start having sex a lot younger" - Practice nurse 5017

\* ü= mentioned by at least one practitioner

- = not mentioned by any practitioners

with male GPs taking less responsibility for this task. Of the 12 theoretical domains, 11 were judged to influence clinical practice. This perhaps indicates that practitioners consider cervical screening to be a complex issue, influenced by multiple factors. The domains judged to be the most important behavioural influences were *beliefs about capabilities, environmental context and resources, social influences* and *behavioural regulation*.

### Comparison with existing literature

It was possible that practitioners, especially practice nurses, might have seen their role in relation to cervical screening as being limited to providing a service (*i.e.*, taking smears) for CervicalCheck. In fact, nurses and female GPs considered that encouraging women to attend for smears also formed part of their role. This suggests that female practitioners may be willing to engage with initiatives to enhance screening uptake. It might also have been expected *a priori* that practitioners would consider providing patients with information about screening or smears to be part of their role, but this did not emerge from the interviews. It would be interesting to see if this also applies in other settings.

The move by male GPs away from responsibility for smearing – which emerged in several interviews with both GPs and practice nurses – was striking. In addition, it was particularly difficult to get male GPs to participate in the study, in part because (as they told us when they declined to take part) they viewed smearing as outside their remit. Others have shown that women have strong preferences for female smear takers and are more likely to attend for screening with female GPs and practice nurses<sup>32,33</sup>. This is consistent with the assumptions of practitioners in this study that women prefer a female smear taker. As regards maximising screening uptake, it is important that women receive consistent messages and encouragement from GPs; this implies that male GPs should be involved in strategies aimed at encouraging women to have smears, even if they do not take smears themselves. However, the findings of this study suggest that engaging male GPs with such strategies may prove difficult.

Previous research has shown that women with a history of sexual abuse are less likely to attend for cervical screening<sup>34</sup>. Our findings indicate that smear consultations with such women (and other “challenging” groups, such as women with learning difficulties) are also perceived as difficult by practitioners. While this may be unique to Ireland because of the relative infancy of CervicalCheck, it is also possible that the increasingly dominant role of practice nurses in smearing means that GPs’ belief in their own capabilities in these more challenging situations is declining. Since women who have a bad experience when having a smear may not reengage with the service<sup>8,9</sup>, it is important to ensure that practitioners have the skills and confidence to manage these types of consultations; support, training and/or interventions for practitioners in this area are, therefore, warranted.

Practical issues, such as limited appointment times and inadequate facilities, impacted on practitioners’ clinical behaviours.

These, and similar issues, have also emerged as reasons why women fail to attend for smear tests or follow-up of abnormal smear results. For example, women cite difficulties in getting an appointment that fits with work/childcare commitments as a barrier to cervical screening attendance<sup>9,33</sup>. Providing convenient appointment times – perhaps outwith usual hours – may be a simple way in which screening uptake could be enhanced.

In terms of the screening protocol in Ireland, there is little evidence that screening women under 25 years is effective<sup>35</sup>. In our study, practitioners seemed to understand this but some – GPs especially – felt that they should have discretion to take a smear in a younger woman if they considered it warranted. In the UK the degree to which practitioners encourage women to attend for smears is related to their attitudes to the national cervical screening programme<sup>36</sup>. In Ireland it remains to be seen whether practitioners who do not fully agree with CervicalCheck guidelines will actively encourage women to have smears and/or comply with strategies to enhance uptake.

For women, emotional issues (such as embarrassment) impact on screening participation<sup>9</sup>, but these did not arise as a major issue for most practitioners. This is unsurprising since practitioners have been trained not to talk about sexual behaviour in relation to smears, which probably removes much of the potential for embarrassment/awkwardness from the consultation. However, the increasing integration of HPV testing into screening may present difficulties. Some practitioners are uncomfortable discussing the sexual behaviour aspects of HPV in relation to cervical screening<sup>24</sup> and this may hinder future strategies designed to maximise uptake.

Around the time CervicalCheck was established, in 2008, there was some controversy over the level of reimbursement that would be provided by the programme to practitioners for taking smears: it was perceived as too low by some GPs who made their views well known. In light of this, it is noteworthy that payments did not emerge as an issue in relation to the domain of *motivation and goals*. This may reflect the fact that the level of reimbursement is now a *fait accompli* and the economic situation has changed dramatically. Alternatively, it is possible that any GPs who remain disgruntled with the payment simply declined to participate in this study. No other aspects of *motivation and goals* emerged as important influences on practitioners’ screening-related clinical behaviours. The most probable explanation for this is that, in contrast to the time before CervicalCheck, when smears were mainly taken opportunistically, a standard national screening protocol is now in place and practitioners are not responsible for call/recall.

### Implications for practice

Most empirical research around increasing cervical screening uptake has focused on understanding women’s screening-related views. This study suggests that practitioners may be receptive to playing a more active role in encouraging women to have smears, and our findings in relation to the influences on practitioners’ behaviours could inform development of strategies or

interventions to enhance these behaviours and, in turn, positively impact on uptake. Ensuring practitioners are comfortable communicating with patients about smears and that they have skills and capabilities to manage “difficult” smear related consultations could also help enhance uptake. Education and training in this area should target all practitioners, so that women who have a male GP are not disadvantaged. Cervical screening programmes are changing and are increasingly based on screening using HPV testing rather than cytology. The overall screening process will remain similar, despite the switch to HPV screening. For example, from the perspective of the woman undergoing an HPV test will be physically the same as having a smear test. However, GPs and practice nurses consider HPV a complex and challenging topic<sup>24</sup>. In addition women are attached to and have confidence in smears, and concerns about programmes changing to HPV testing<sup>32,37</sup>. Practitioners need to be adequately supported and prepared for these changes, and the associated patient communication issues, so that screening uptake is not negatively affected.

### Strengths and limitations

The qualitative design allowed us to gain an in-depth understanding of the issues around cervical screening from practitioners’ perspectives. Data of such richness and depth would not have been available using a quantitative approach. The data may somewhat overemphasise experiences of practitioners with an interest in women’s health and views of practitioners with more positive attitudes towards screening than others since we interviewed GPs who had previously expressed an interest in this area. In addition, almost three quarters of the GPs interviewed had been in general practice for more than 20 years, but it is entirely unknown how this may influence their clinical practice and beliefs in relation to cervical screening. Overall, the interviewees’ characteristics were diverse and the process’ credibility and findings are evidenced in the diversity of themes and opinions which emerged. Despite practice nurses having responsibility for smearing in several healthcare systems, as far as we are aware, this is the first study to directly explore their roles and behaviours. Interviews for the study were conducted in late 2010 and early 2011 when the CervicalCheck programme was in its infancy. A lot has changed over the last 10 years regarding cervical cancer prevention in Ireland e.g. increased knowledge and awareness among women and the general population of cervical screening, HPV infection and the HPV vaccines. From the perspective of GPs and practice nurses, influences on their cervical-screening related clinical behaviours may also have changed. Using the TDF for analysis meant that we did not have to select, *a priori*, a single psychological theory of behaviour, thus minimising the likelihood of missing important influences on clinical behaviour. However, the TDF does not specify relationships between the domains<sup>38,39</sup>. Finally, while we were only able to determine in a qualitative way which of the domains were likely to be the most important drivers of clinical behaviour, the study was intended to be hypothesis-generating rather than hypothesis-testing.

### Conclusions

We identified - for the first time - primary care practitioners’ cervical-screening related clinical behaviours, their perceived

roles and responsibilities, and the factors likely to influence behaviours. In addition to taking smears, practitioners considered it their responsibility to encourage women to have smears, but smearing was viewed as a predominantly female role. The results could inform development of strategies to: motivate and enable practitioners to encourage women to have smear/screening tests; and/or improve women’s access and experiences; these in turn, may help increase cervical screening uptake. However, engaging male practitioners with these initiatives may prove difficult.

### Data availability

#### Underlying data

There are no quantitative data associated with this article. The audio files and transcripts generated during the current study are confidential. In the consent document, participants were not asked to consent to sharing of data beyond the research team and their collaborators. A comprehensive set of quotes reflecting the transcripts are available in [Table 2](#). Researchers seeking to access the underlying data (i.e. audio files and transcripts) will need to apply directly to the Irish College of General Practitioners Research Ethics Committee for approval. The Committee can be contacted at [research@icgp.ie](mailto:research@icgp.ie). Should approval be granted, the authors are happy to facilitate access.

#### Extended data

Figshare: Extended Data: Identifying ways to maximise cervical screening uptake: a qualitative study of GPs’ and practice nurses’ cervical cancer screening-related behaviours <https://doi.org/10.6084/m9.figshare.14132105.v1><sup>25</sup>.

- Interview topic guide
- SRQR checklist

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

### Acknowledgements

We thank the GPs and practice nurses who participated in the telephone interviews. We are grateful for the assistance of the Professional Development Coordinators for Practice Nurses in recruiting practice nurses; members of the Steering Group and Intervention Development Group for their input; Dr Judith Murphy for her input into the initial stages of the study; Claire O’Callaghan and Kate Higney for transcribing the interviews; Damien Foley for designing the ATHENS logo; and staff at the CervicalCheck training unit for helpful discussions.

The ATHENS group comprised:

Investigators: LAMcS, SUD, JJJ, CMM, JJO’L and LS

Steering Group: Dr Brenda Corcoran, Ms Ann McGill, Ms Eileen O’Donovan, Dr Eamonn Shanahan, Dr Alan Smith, and investigators as listed above

Intervention Development Group: Ms Carol McNamara, Ms Louise McKee, Dr Genevieve McGuire, LAMcS, SUD and LS.

## References

- International Agency for Research on Cancer Working Group: **IARC handbook of cervical cancer prevention, Cervix cancer screening**. Lyon: IARC, 2005; 10. [Reference Source](#)
- Day NE: **Screening for cancer of the cervix**. *J Epidemiol Community Health*. 1989; **43**(2): 103–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chang HC, Hsiung HY, Chen SI, et al.: **Comparison of a community outreach service with opportunity screening for cervical cancer using Pap smears**. *J Public Health (Oxf)*. 2007; **29**(2): 165–72. [PubMed Abstract](#) | [Publisher Full Text](#)
- Lancucki L, Fender M, Koukari A, et al.: **A fall-off in cervical screening coverage of younger women in developed countries**. *J Med Screen*. 2010; **17**(2): 91–6. [PubMed Abstract](#) | [Publisher Full Text](#)
- NHS Screening & Immunisations Team: **Cervical Screening Programme, England, 2016-17**. 2017; (accessed 21 January 2021). [Reference Source](#)
- Hall IJ, Tangka FKL, Sabatino SA, et al.: **Patterns and Trends in Cancer Screening in the United States**. *Prev Chronic Dis*. 2018; **15**: E97. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Public Health England (PHE) screening: **NHS Screening Programmes in England: 2017 to 2018**. 2019; (accessed 12 February 2020). [Reference Source](#)
- Blomberg K, Ternstedt BM, Törnberg S, et al.: **How do women who choose not to participate in population-based cervical cancer screening reason about their decision?** *Psychooncology*. 2008; **17**(6): 561–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Waller J, Bartoszek M, Marlow L, et al.: **Barriers to cervical cancer screening attendance in England: a population-based survey**. *J Med Screen*. 2009; **16**(4): 199–204. [PubMed Abstract](#) | [Publisher Full Text](#)
- Twinn S, Cheng F: **Increasing uptake rates of cervical cancer screening amongst Hong Kong Chinese women: the role of the practitioner**. *J Adv Nurs*. 2000; **32**(2): 335–42. [PubMed Abstract](#) | [Publisher Full Text](#)
- O'Connor M, Murphy J, Martin C, et al.: **Motivators for women to attend cervical screening: the influential role of GPs**. *Fam Pract*. 2014; **31**(4): 475–82. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ackerson K, Pohl J, Low LK: **Personal influencing factors associated with Pap smear testing and cervical cancer**. *Policy Polit Nurs Pract*. 2008; **9**(1): 50–60. [PubMed Abstract](#) | [Publisher Full Text](#)
- McAlearney AS, Oliveri JM, Post DM, et al.: **Trust and distrust among Appalachian women regarding cervical cancer screening: a qualitative study**. *Patient Educ Couns*. 2012; **86**(1): 120–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peate I: **Cervical cancer. 1: Role of nurses in the primary healthcare setting**. *Br J Nurs*. 1999; **8**(11): 730–4. [PubMed Abstract](#) | [Publisher Full Text](#)
- Herity B, McDonald P, Johnson Z, et al.: **A pilot study of cervical screening in an inner city area—lessons for a national programme**. *Cytopathology*. 1997; **8**(3): 161–70. [PubMed Abstract](#) | [Publisher Full Text](#)
- Mills J, Chamberlain-Salaun J, Christie L, et al.: **Australian nurses in general practice, enabling the provision of cervical screening and well women's health care services: a qualitative study**. *BMC Nurs*. 2012; **11**: 23. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rennie D, Boxsell J, Pedretti K: **A team care model of cervical screening in a general practice**. *Aust Fam Physician*. 2015; **44**(7): 515–8. [PubMed Abstract](#)
- Gannon M, Dowling M: **Increasing the uptake of cervical screening programmes**. *Br J Nurs*. 2008; **17**(20): 1280–4. [PubMed Abstract](#) | [Publisher Full Text](#)
- Cooper M: **Reasons for non-attendance at cervical screening**. *Nurs Stand*. 2011; **25**(26): 41–6. [PubMed Abstract](#)
- McSherry LA, Dombrowski SU, Francis JJ, et al.: **'It's a can of worms': understanding primary care practitioners' behaviours in relation to HPV using the Theoretical Domains Framework**. *Implement Sci*. 2012; **7**: 73. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Smith S, Normand C: **Equity in health care: the Irish perspective**. *Health Econ Policy Law*. 2011; **6**(2): 205–17. [PubMed Abstract](#) | [Publisher Full Text](#)
- CervicalCheck programme report 2012-2013**. 2015; (accessed 22 January 2020). [Reference Source](#)
- The National Screening Service: **Standards for Quality Assurance in Cervical Screening. Quality Assurance in Primary Care and Other Cervical Screening Settings**. 2020; (accessed 31 March 2021). [Reference Source](#)
- Murphy J, Sharp L, Carsin AE: **Attitudes and knowledge of GP's towards Human Papillomavirus infection, testing and vaccination: results from a national survey in Ireland**. *J Epidemiol Comm Health*. 2008; **62**: A21–2. [Reference Source](#)
- O'Connor M: **Extended data study materials maximising screening uptake**. *docx.figshare*. Online resource. 2021. <http://www.doi.org/10.6084/m9.figshare.14132105.v1>
- Strauss A, Corbin J: **Basics of Qualitative Research: Techniques and Procedures for Developing Grounded Theory**, 2nd edn. New York: Sage Publications Inc; 1998. [Reference Source](#)
- O'Brien BC, Harris IB, Beckman TJ, et al.: **Standards for reporting qualitative research: a synthesis of recommendations**. *Acad Med*. 2014; **89**(9): 1245–1251. [PubMed Abstract](#) | [Publisher Full Text](#)
- Pope C, Ziebland S, Mays N: **Qualitative research in health care. Analysing qualitative data**. *BMJ*. 2000; **320**(7227): 114–116. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ritchie J, Lewis J: **Qualitative Research Practice - A Guide for Social Science Students and Researchers**. London, Thousand Oaks, CA: Sage Publications Ltd; 2003. [Reference Source](#)
- Michie S, Johnston M, Abraham C, et al.: **Making psychological theory useful for implementing evidence based practice: a consensus approach**. *Qual Saf Health Care*. 2005; **14**(1): 26–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Islam R, Timmouth AT, Francis JJ, et al.: **A cross-country comparison of intensive care physicians' beliefs about their transfusion behaviour: A qualitative study using the Theoretical Domains Framework**. *Implement Sci*. 2012; **7**: 93. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McRae J, Martin C, O'Leary J, et al.: **"If you can't treat HPV, why test for it?" Women's attitudes to the changing face of cervical cancer prevention: a focus group study**. *BMC Womens Health*. 2014; **14**: 64. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chorley AJ, Marlow LAV, Forster AS, et al.: **Experiences of cervical screening and barriers to participation in the context of an organised programme: a systematic review and thematic synthesis**. *Psychooncology*. 2017; **26**(2): 161–172. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cadman L, Waller J, Ashdown-Barr L, et al.: **Barriers to cervical screening in women who have experienced sexual abuse: an exploratory study**. *J Fam Plann Reprod Health Care*. 2012; **38**(4): 214–20. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sasieni P, Castanon A, Cuzick J: **Effectiveness of cervical screening with age: population based case-control study of prospectively recorded data**. *BMJ*. 2009; **339**: b2968. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chew-Graham C, Mole E, Evans LJ, et al.: **Informed consent? How do primary care professionals prepare women for cervical smears: a qualitative study**. *Patient Educ Couns*. 2006; **61**(3): 381–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Obermair HM, Dodd RH, Bonner C, et al.: **'It has saved thousands of lives, so why change it?' Content analysis of objections to cervical screening programme changes in Australia**. *BMJ Open*. 2018; **8**(2): e019171. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Francis JJ, Stockton C, Eccles MP, et al.: **Evidence-based selection of theories for designing behaviour change interventions: using methods based on theoretical construct domains to understand clinicians' blood transfusion behaviour**. *Br J Health Psychol*. 2009; **14**(Pt 4): 625–46. [PubMed Abstract](#) | [Publisher Full Text](#)
- Atkins L, Francis J, Islam R, et al.: **A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems**. *Implement Sci*. 2017; **12**(1): 77. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

## Open Peer Review

Current Peer Review Status:  

---

### Version 1

Reviewer Report 16 August 2021

<https://doi.org/10.21956/hrbopenres.14409.r29993>

© 2021 Ali S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Suhailah M Ali** 

College of Nursing, University of Kirkuk, Kirkuk, Iraq

This paper presents a very important issue concerning primary care practitioners' and practice nurses' perceived roles and responsibilities, in terms of cervical-screening related clinical behaviours. The study aimed to determine factors likely to influence clinical behaviours. 19 GPs and 14 practice nurses in Ireland participated. Authors utilized a well-established Theoretical Domains Framework (TDF) to analyse the interview transcripts. The table describing the included 12 possible theoretical domains was useful to gain an in-depth understanding of the issues around cervical screening from practitioners' perspectives. The findings of this study could inform development of interventions to enhance practitioners' clinical behaviours to promote cervical screening uptake.

**Introduction:** The research question clearly identifies the importance of factors influencing primary care practitioners' and practice nurses' cervical screening-related clinical behaviours to encourage cervical screening uptake. The introduction is well written and provides clear justification.

**Methods:** The setting of the study is well described. It is evident to the reader what current practices are regarding cervical screening in Ireland with regards to switching to the primary HPV testing in line with recent international recommendations. Authors described well how the participants were recruited and how it evolved as the study progressed. Recruitment continued until the decision was reached that there was sufficient depth of information to meet the purpose of the study.

**Result:** The results of this qualitative study are well presented to address the research questions. Table 2 enabled us to get an in depth snapshot of the challenging contexts in which the practitioners' and practice nurses' worked.

However, there are few minor points I noticed that ought to be rectified:

- In the result section, page 5: Table 1 shows 19 GPs and 14 practice nurses participated in the study. Whereas in the abstract page 1, result section: authors stated 14 GPs and 19 practice nurses participated? This need to be consistent.



**Discussion:** Authors clearly explained how this study has attempted to identify factors likely to influence practitioner's clinical behaviours and they made explicit statements from their study to compare with the existing literature. The implications for practice seemed appropriate and key messages are accurate indicating that practitioners are willing to consider a more active role in encouraging women to uptake cervical smears. Limitations of the paper are well presented. However, I can not see suggestions for further investigation in the area of practice which leaves the reader with the implication of the phenomenon being researched and no onward plan for future research.

In summary, this is a professional written paper with high level of in-depth analysis and a well-structured presentation of arguments. The findings make significant contribution to the field of cervical screening uptake.

### References

1. Atkins L, Francis J, Islam R, O'Connor D, et al.: A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implementation Science*. 2017; **12** (1). [Publisher Full Text](#)

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Not applicable

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** cervical cancer screening , qualitative studies, mixed methods studies, systematic reviews

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 01 June 2021

<https://doi.org/10.21956/hrbopenres.14409.r29408>



© 2021 Campbell J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



### John Campbell

Collaboration for Academic Primary Care (APEX), University of Exeter, Exeter, UK

Thank you for the opportunity to review this interesting article setting out to address three clear aims in respect of the observed decline in cervical screening. A qualitative study of 14 GPs and 19 practice nurses is reported.

**Abstract** - The abstract is broadly acceptable although does lack some precision and clarity in respect of the presentation of results. Exactly what behaviours are being referred to line 3? In addition, a very substantial claim is made round the originality of this study. Although this is not my principal area of expertise or interest I'd be very surprised if this was the first such study to explore primary care practitioners' behaviours in respect of cervical screening

**Introduction** - the introduction presents a compelling and well-articulated case for the paper, noting the decline in cervical screening uptake, some potential issues impacting on the likelihood that women will respond to screening requests and the previous experience on doctors (although I would want to be reassured that this was comprehensive).

**Methods** - the setting of the study is well described, including a short summary of the Irish health Service. The situation is described where the responsibility for taking smears has moved from GPs to practice nurses. Certainly for some GPs this is a frustration, and may be undermining of the whole cervical cytology programme.

The authors clearly outline their approach to sampling for the qualitative research, but it seems likely that participants would be subject to selection bias having taken part in previous related research (for GPs). The approach to practice nurses seems more comprehensive, with recruitment via a conference or to nurse development coordinators.

The interview procedures are standard, and well described. Recruitment continued until saturation was evident. Analysis involved a framework approach with theoretical domains framework as the coding framework.

**Results** - recruitment appears to have covered a range of participants as defined by gender, geographical distribution, practice setting, clinical experience etc.

A range of quotes are provided in respect of roles and responsibilities in respect of smear taking.

A useful table (2) identifies factors influencing behaviours relating to cervical screening, accompanied by useful quotes. The results focus on belief as well as organisation factors relating to time, clinical behaviours, and external social influences.

**Discussion** - the findings are neatly summarised and presented in the context of a reasonable comparison with relevant international literature. Key issues identified in the results are followed

through in the discussion and a comprehensive contextualisation of the findings is presented and discussed. A range of implications for practice are presented although I was surprised that some of the anticipated views of male GPs is not reflected in the implications for practice – for many experienced GPs, taking smears was a normal part of routine practice and it is possibly to the detriment of overall clinical care, including to the experience of women, that this clinical procedure is now seen to be within the domain of nurses or female GPs.

A range of strengths and limitations of the work are presented and reasonable conclusions are drawn (although, again, I would question whether this is the first such study in this area).

Overall however, this is a carefully conducted piece of important research which is well presented, well written, and provides an interesting range of significant findings. I have no hesitation in supporting its publication with only minor changes.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Not applicable

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Primary care; access; quality

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---