

# Cell Line Specific Modulation of Extracellular A $\beta$ <sub>42</sub> by Hsp40

Anna Carnini<sup>1</sup>, Lucas O. M. Scott<sup>1</sup>, Eva Ahrendt<sup>1</sup>, Juliane Proft<sup>1</sup>, Robert J. Winkfein<sup>1</sup>, Sung-Woo Kim<sup>2</sup>, Michael A. Colicos<sup>1</sup>, Janice E. A. Braun<sup>1\*</sup>

**1** Department of Physiology and Pharmacology, Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada, **2** Department of Biochemistry and Molecular Biology, Southern Alberta Cancer Research Institute, University of Calgary, Calgary, Alberta, Canada

## Abstract

Heat shock proteins (Hsps) are a set of molecular chaperones involved in cellular repair. They provide protective mechanisms that allow cells to survive potentially lethal insults. In response to a conditioning stress their expression is increased. Here we examined the connection between Hsps and A $\beta$ <sub>42</sub>, the amyloid peptide involved in the pathological sequence of Alzheimer's disease (AD). Extracellular A $\beta$ <sub>42</sub> associates with neuronal cells and is a major constituent of senile plaques, one of the hallmarks of AD. Although Hsps are generally thought to prevent accumulation of misfolded proteins, there is a lack of mechanistic evidence that heat shock chaperones directly modulate A $\beta$ <sub>42</sub> toxicity. In this study we show that neither extracellular A $\beta$ <sub>42</sub> nor A $\beta$ <sub>42</sub>/PrP<sup>C</sup> trigger the heat shock response in neurons. To address the influence of the neuroprotective heat shock response on cellular A $\beta$ <sub>42</sub>, Western analysis of A $\beta$ <sub>42</sub> was performed following external A $\beta$ <sub>42</sub> application. Five hours after a conditioning heat shock, A $\beta$ <sub>42</sub> association with CAD cells was increased compared to control neurons. However, at forty-eight hours following heat shock A $\beta$ <sub>42</sub> levels were reduced compared to that found for control cells. Moreover, transient transfection of the stress induced Hsp40, decreased CAD levels of A $\beta$ <sub>42</sub>. In contrast to CAD cells, hippocampal neurons transfected with Hsp40 retained A $\beta$ <sub>42</sub> indicating that Hsp40 modulation of A $\beta$ <sub>42</sub> proteostasis is cell specific. Mutation of the conserved HPD motif within Hsp40 significantly reduced the Hsp40-mediated A $\beta$ <sub>42</sub> increase in hippocampal cultures indicating the importance of this motif in regulating cellular A $\beta$ <sub>42</sub>. Our data reveal a biochemical link between Hsp40 expression and A $\beta$ <sub>42</sub> proteostasis that is cell specific. Therefore, increasing Hsp40 therapeutically with the intention of interfering with the pathogenic cascade leading to neurodegeneration in AD should be pursued with caution.

**Citation:** Carnini A, Scott LOM, Ahrendt E, Proft J, Winkfein RJ, et al. (2012) Cell Line Specific Modulation of Extracellular A $\beta$ <sub>42</sub> by Hsp40. PLoS ONE 7(5): e37755. doi:10.1371/journal.pone.0037755

**Editor:** Mark R. Cookson, National Institutes of Health, United States of America

**Received:** January 19, 2012; **Accepted:** April 24, 2012; **Published:** May 29, 2012

**Copyright:** © 2012 Carnini et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by funding from the Canadian Institute of Health Research MOP#74713 (Dr. Braun) and MOP#7944 (Dr. Colicos) and the Alberta Prion Research Institute. Dr. Braun is a Senior Scholar of the Alberta Heritage Foundation for Medical Research. Dr. Colicos is a Scholar of the Alberta Heritage Foundation for Medical Research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: braunj@ucalgary.ca

## Introduction

Alzheimer's disease (AD), an age-dependent neurodegenerative disease that is estimated to affect 35 million people worldwide is characterized by amyloid deposits, neurofibrillary tangles, selective neuronal loss, cognitive decline and memory loss [1,2]. Multiple lines of evidence suggest an imbalance between the production and clearance of A $\beta$ <sub>1–42</sub>, a 42 residue long  $\beta$  amyloid protein that spontaneously self aggregates into dimers, oligomers, protofibrils, and fibrils and initiates a toxic sequence of events leading to synaptic dysfunction and dementia [3]. A $\beta$ <sub>42</sub> as well as A $\beta$ <sub>40</sub> are derived from the amyloid precursor protein (APP) by the sequential proteolytic processing of  $\alpha$ ,  $\beta$  and  $\gamma$  secretases (reviewed: [4,5]. Following proteolysis, the peptides can be secreted or transferred to the endosomal/lysosomal system. Intraneuronal A $\beta$ <sub>42</sub> is comprised of both uptake of A $\beta$ <sub>42</sub> from the extracellular space as well as intracellular cleavage of APP [6]. Synaptic activity increases levels of secreted, extracellular A $\beta$  peptides while reducing intracellular levels [7].

Why do the physiological mechanisms that under normal circumstances tightly regulate A $\beta$ <sub>42</sub> production, cell association

and clearance fail? Deficiencies in cellular chaperone systems are one possibility. Molecular chaperones maintain protein homeostasis by assisting nascent polypeptides to fold, protecting mature proteins from stresses and eliminating misfolded proteins. Protein quality control mechanisms are critical to neural function and defects in proteolytic pathways are widely held to lead to neurodegeneration [8]. The cellular level of chaperones might affect the toxicity of A $\beta$ <sub>42</sub>. In fact, enhancement of the cellular quality control machinery, has been proposed to prevent or delay the cascade of misfolding in conformational diseases [9,10]. In addition to maintenance of protein homeostasis (proteostasis) by constitutive chaperones, in response to a number of stressful stimuli, there is an induction of stress-induced chaperones (eg Hsp40, Hsp90, Hsp70 and Hsp27). Understanding the biochemical sequence of events that underlies A $\beta$ <sub>42</sub>-mediated neurodegeneration requires a clear understanding of the role(s) that chaperones play in the AD pathogenic cascade.

A number of chaperones are implicated in A $\beta$ <sub>42</sub> proteostasis [11]. Several chaperones have been found both in association with senile plaques [12–14] as well as endogenous A $\beta$ <sub>42</sub> [15]. These reports have given rise to the notion that molecular chaperones are

suppressors of toxic A $\beta$ <sub>42</sub> conformations leading to AD. This idea is consistent with the observations that heat shock genes appear to be induced poorly late in life and that the principal risk factor for AD is age [9]. Further support for this view has come from reports demonstrating that in experimental models, Hsp70 [16,17], Hsp27 [18] and Hsp90 [16] protect against the toxic effects of A $\beta$ <sub>42</sub>. Also, Hsp70 is reported to suppress cognitive deficits and pathological phenotypes in AD mice [19]. *In vitro* Hsp70/40 and Hsp90 suppressed early stages of A $\beta$ <sub>42</sub> assembly into aggregates but had no effect on fibrils [16]. Still, many questions remain unanswered regarding the chaperone folding paths for A $\beta$ <sub>42</sub>. For example, Mearow and colleagues have shown that heat shock of neonatal rat cortical cultures increases the detrimental effects of A $\beta$ <sub>42</sub> on cell survival while overexpressing Hsp27 protects against A $\beta$ <sub>42</sub> [18]. In mice models of Alzheimer's disease overall content of the chaperone  $\alpha$  $\beta$  crystallin is reduced [20]. However, in contrast to the concept of therapeutic rescue by chaperones, several molecular chaperones actually support the formation of the toxic A $\beta$ <sub>42</sub> oligomeric species [21–23]. This promotion of A $\beta$ <sub>42</sub> oligomerization by select chaperones has similar features to that observed in response to general anesthesia [24,25].

In addition to cellular chaperones, cellular prion protein (PrP<sup>C</sup>)/A $\beta$ <sub>42</sub> association could influence A $\beta$ <sub>42</sub> quality control. A $\beta$ <sub>42</sub> in contrast to A $\beta$ <sub>40</sub>, associates rapidly with neuronal cells [26]. Two distinct A $\beta$ <sub>42</sub> oligomeric conformations accumulate on the surface of living cells [27]. Exposure of A $\beta$ <sub>42</sub> to pH = 6 for 24 hours to mimic endosomal conditions increases A $\beta$ <sub>42</sub> binding to PC12 cells [28]. The cellular prion protein has been shown to act as a functional high affinity receptor for A $\beta$ <sub>42</sub> [29–31]. Strittmatter and colleagues report that association of PrP<sup>C</sup> with A $\beta$ <sub>42</sub> mediates downstream A $\beta$ <sub>42</sub>-impairment of hippocampal long term potentiation and that AD transgenic mice lacking PrP<sup>C</sup> accumulate A $\beta$ <sub>42</sub> but have normal survival and test normal for learning and memory [29,31]. Furthermore, transgenic overexpression of PrP<sup>C</sup> is shown to enhance amyloid plaque formation in an AD mouse model [32]. Along these lines, pathological levels of A $\beta$ <sub>42</sub> have been shown to disrupt PrP<sup>C</sup> modulation of NMDA (N-Methyl-d-aspartate) receptor activity [33]. However, in contrast, Balducci et al report that A $\beta$ <sub>42</sub> impairs consolidation of long-term recognition memory in mice independent of PrP<sup>C</sup>, raising questions regarding the role of A $\beta$ <sub>42</sub>/PrP<sup>C</sup> association in AD progression [30]. Additionally, other molecules (eg STI1) are recognized to bind PrP<sup>C</sup>, but whether these agents compete with A $\beta$ <sub>42</sub> for binding is unknown [34]. Following cell association, insoluble A $\beta$ <sub>42</sub> aggregates localize to endosome/lysosome compartments [35,36]. Curiously, one report reveals that PrP<sup>C</sup> inhibits  $\beta$ -secretase cleavage of amyloid precursor protein and reduces A $\beta$ <sub>42</sub> levels [37].

That said, which molecular chaperones directly regulate A $\beta$ <sub>42</sub> proteostasis and/or toxicity remain to be established. In this study we have monitored the association of A $\beta$ <sub>42</sub> with cultured neural cells following induction of the heat shock. Our findings demonstrate that heat shock initially increases A $\beta$ <sub>42</sub> association with CAD neuroblastoma cells but is followed by a decline in cellular A $\beta$ <sub>42</sub> levels at 48 hours. Transient transfection experiments revealed that Hsp40 decreased cellular levels of A $\beta$ <sub>42</sub> in CAD cell but increased cellular levels of A $\beta$ <sub>42</sub> in hippocampal cultures. We evaluated the influence of exogenously applied soluble PrP<sup>C</sup> on cellular uptake/processing of A $\beta$ <sub>42</sub> to test the hypothesis that soluble PrP<sup>C</sup> would bind to A $\beta$ <sub>42</sub> and reduce association with cell anchored PrP<sup>C</sup>, thereby blocking an early event in the A $\beta$ <sub>42</sub> pathogenic cascade. Here we document that PrP<sup>C</sup> failed to decrease cellular association of A $\beta$ <sub>42</sub>. Our data reveal a biochemical link between cellular levels of Hsp40 and

A $\beta$ <sub>42</sub> that is cell line specific. This raises the possibility that Hsp40 is involved in the pathogenic cascade leading to dementia and neurodegeneration in AD.

## Results and Discussion

### Extracellular A $\beta$ <sub>42</sub> does not Trigger the Heat Shock Response


Heat shock chaperones are induced in response to a number of cell stressors such as temperature, ischemia and heavy metals. During aging when heat shock genes are thought to be induced poorly, humans are susceptible to AD. To gain insight into the involvement of the heat shock response in A $\beta$ <sub>42</sub> pathogenic cascades, we carried out biochemical studies to establish whether the treatment of neuroblastoma cells with A $\beta$ <sub>42</sub> triggers the expression of the stress-induced chaperones. Mouse CAD neuroblastoma cells were incubated with a high concentration (25  $\mu$ M) of A $\beta$ <sub>42</sub> for 48 hours, rinsed in PBS and solubilized. 30  $\mu$ g of supernatant (1% TX-100/0.1% SDS soluble proteins) and 10  $\mu$ l of total pellet (1% TX-100/0.1% SDS insoluble proteins) were subjected to Western analysis. **Figure 1** demonstrates that A $\beta$ <sub>42</sub> was clearly found in both soluble and insoluble CAD cell fractions. The expression of cellular Hsp70 (heat shock protein of 70 kDa), Hsp25 (Heat shock protein of 25 kDa) and Hsp40 (Heat shock protein of 40 kDa) in CAD cells did not change following treatment with A $\beta$ <sub>42</sub> for 48 hours. Hsp70 and Hsp25 were not detectable in either the presence or absence of A $\beta$ <sub>42</sub>. Moreover, Hsp40 showed modest expression in control CAD cells as previously described [38], and no change was observed in response to A $\beta$ <sub>42</sub> treatment. The expression levels of the constitutive chaperone Hsc70 (Heat shock cognate protein of 70 kDa) also did not change in response to A $\beta$ <sub>42</sub>. Actin is shown as a loading control.

### A $\beta$ <sub>42</sub> does not Block the Heat Shock Response

The heat shock response is a highly conserved cell survival program that enhances cell survival to subsequent insults [9]. Since interference with the heat shock response would be expected to reduce protein surveillance and triage mechanisms and downstream cell survival, we next tested whether A $\beta$ <sub>42</sub> altered induction of the heat shock chaperones. CAD cells were incubated with 3  $\mu$ M A $\beta$ <sub>42</sub> then heat shocked at 43°C for 40 min and allowed to recover for 5 hours prior to lysis and Western analysis. **Figure 2** clearly shows that Hsp70 is induced by heat shock and that A $\beta$ <sub>42</sub> does not alter the induction of this stress inducible chaperone. As expected, A $\beta$ <sub>42</sub> treatment of CAD cells triggers apoptotic pathways as shown by activation of caspase 3, a marker of programmed cell death. The A $\beta$ <sub>42</sub> activation of caspase 3 was not altered by heat shock. Heat shock treatment increased cellular levels of soluble A $\beta$ <sub>42</sub> ~3.5 fold, suggesting that following heat shock neurons are more prone to the cellular toxicity of A $\beta$ <sub>42</sub> (**Figure 2**). This observation is in line with a previous study demonstrating that A $\beta$ <sub>42</sub> decreases cortical neuron cell survival and that heat shock renders neurons more vulnerable to A $\beta$ <sub>42</sub> treatment [18]. Taken together, our data demonstrate that, although a large number of stressors activate the neuroprotective heat shock response, A $\beta$ <sub>42</sub> failed to increase the expression of the heat shock chaperones. Moreover, removal/disruption of the protective effects of a conditioning heat shock against cell death is not part of the A $\beta$ <sub>42</sub> pathogenic cascade.

### Soluble PrP<sup>C</sup> does not Block Cell Association of A $\beta$ <sub>42</sub>

PrP<sup>C</sup> has been proposed to act as a functional high affinity receptor for A $\beta$ <sub>42</sub> [29–31]. Based on these reports, we wanted to



**Figure 1. A $\beta$ <sub>42</sub> does not trigger the heat shock response in CAD neural cells.** CAD cells were incubated with 25  $\mu$ M A $\beta$ <sub>42</sub> for 48 hours, washed in PBS and lysed. 30  $\mu$ g of soluble protein or 10  $\mu$ l of the insoluble fraction were resolved by SDS-PAGE and the indicated proteins were evaluated by Western blot analysis.  $\beta$ -actin is shown as a loading control. Data are representative of 3 separate experiments. doi:10.1371/journal.pone.0037755.g001

determine how PrP<sup>C</sup> may impact the cellular association of A $\beta$ <sub>42</sub>. Therefore, we examined the possibility that soluble recombinant PrP<sup>C</sup> would competitively displace A $\beta$ <sub>42</sub> from CAD cells thereby reducing the association of A $\beta$ <sub>42</sub> with the neuronal cell culture. Treatment with 0.5  $\mu$ M recombinant PrP<sup>C</sup> did not induce Hsp70 or activation of caspase 3. Also, PrP<sup>C</sup> failed to alter the heat shock induction of Hsp70 or the A $\beta$ <sub>42</sub> induced activation of caspase 3. No difference in cellular A $\beta$ <sub>42</sub> association was observed between A $\beta$ <sub>42</sub> and PrP<sup>C</sup>/A $\beta$ <sub>42</sub> treated control (no-heat shock) cells. In fact, rather than inhibit cellular association of A $\beta$ <sub>42</sub>, PrP<sup>C</sup>/A $\beta$ <sub>42</sub> co-treatment followed by heat shock revealed that PrP<sup>C</sup> increased


A $\beta$ <sub>42</sub> levels in the soluble CAD cell fractions (**Figure 2**). CAD cells express endogenous PrP<sup>C</sup>, a glycosylphosphatidyl (GPI) anchored plasma membrane protein [39,40] (**Figure 3**) that is subject to N-linked glycosylation and non-, mono- and di-glycosylated versions of PrP<sup>C</sup> simultaneously exist [41]. Recombinant bovine PrP<sup>C25–232</sup> migrated further on SDS-PAGE than native unglycosylated PrP<sup>C</sup> therefore rendering cell association of exogenous recombinant PrP<sup>C</sup> distinguishable from endogenous PrP<sup>C</sup>. Exogenous recombinant PrP<sup>C</sup> was observed to associate with cells (**Figure 3**). In the absence of CAD cells recombinant PrP<sup>C</sup> was observed to undergo partial breakdown following heat shock for 40 min at 43°C. Taken together our results show that exogenous PrP<sup>C</sup> does not block cellular A $\beta$ <sub>42</sub> association, in fact, following heat shock PrP<sup>C</sup>/A $\beta$ <sub>42</sub> co-treatment increased cell associated A $\beta$ <sub>42</sub>.

#### Heat Shock Promotes Time Dependent Clearance of A $\beta$ <sub>42</sub>

Although heat shock facilitated the pathogenicity of A $\beta$ <sub>42</sub> as measured by its increased cellular association, we speculated that this may be due to the physical effects of heat shock on membrane permeability rather than the conformational processing of PrP<sup>C</sup> by stress induced chaperones. To gain further insight into the relationship between cellular uptake/clearance of A $\beta$ <sub>42</sub> and the heat shock response, we carried out immunoblot analysis on CAD cells in which the heat shock was given at an earlier time point thereby increasing the time A $\beta$ <sub>42</sub> is exposed to the stress-induced chaperones. **Figure 4** shows that when a 40 min heat shock was given starting at the time that A $\beta$ <sub>42</sub> was applied to CAD cells, the cellular levels of soluble A $\beta$ <sub>42</sub> at the 42 hour time point was reduced. A $\beta$ <sub>42</sub> does not always resolve as a discrete band by SDS-PAGE depending on abundance and the characteristic wide A $\beta$ <sub>42</sub> band is shown in Figure 4. Cellular levels of Hsp70 and Hsp40, which are elevated 3 hours following heat shock [38], remained elevated 48 hours following heat shock and at the 48 time point translocation of Hsp70 (but not Hsp40) to the detergent insoluble fraction was observed. Our observations demonstrate that while soluble A $\beta$ <sub>42</sub> is increased 5 hours following heat shock (**Figure 2**), soluble A $\beta$ <sub>42</sub> is decreased 48 hours following heat shock (**Figure 4**) indicating that with time heat shock chaperones increase cellular A $\beta$ <sub>42</sub> clearance. Geldanamycin-treatment of CAD cells which induces Hsp40 but not Hsp70 [38] was also observed to reduce cellular levels of A $\beta$ <sub>42</sub> (data not shown). **Figure 4 (lower panel)** clearly shows that CAD cell levels of A $\beta$ <sub>42</sub> increase in response to increasing A $\beta$ <sub>42</sub> concentrations. Again, PrP<sup>C</sup>, (either bovine upper panel or mouse lower panel **Figure 4**) did not reduce cell association of A $\beta$ <sub>42</sub> and did not alter the heat shock response.

**Figure 5** shows that in the absence of cells, heat shock *per se* does not cause degradation or oligomerization of A $\beta$ <sub>42</sub>. Also, PrP<sup>C</sup> does not initiate any shifts in the molecular weight of A $\beta$ <sub>42</sub> indicative of proteolysis or SDS-resistant oligomerization. In contrast a 15 kDa breakdown of PrP<sup>C</sup> was observed following heat shock. **Figure 6** shows that neither PrP<sup>C</sup> nor A $\beta$ <sub>42</sub> were found to alter cellular levels of the constitutive chaperones DnaJA1/Hdj2, DnaJA2/Rdj2, DnaJA3/Tid1, DnaJA4, Hsc70 or the stress induced chaperones Hsp70, Hsp40 and Hsp25, indicating that a generalized reduction in these molecular chaperone levels is not an underlying mechanism of A $\beta$ <sub>42</sub> induced neurodegeneration.

Taken together, these results indicate that heat shock chaperones have a role in clearance of cellular A $\beta$ <sub>42</sub>, and may perhaps be involved in reducing A $\beta$ <sub>42</sub> pathogenesis.



**Figure 2. Heat shock initially increases A $\beta$ <sub>42</sub> association with CAD neural cells.** In four independent experiments CAD cells were incubated with 3  $\mu$ M A $\beta$ <sub>42</sub> and/or 0.5  $\mu$ M bovine rPrP<sup>C</sup>, after ~42 hours cells were subjected to heat-shock at 43°C for 40 minutes and then allowed to recover for 5 hours. The cells were rinsed in PBS and lysed. 30  $\mu$ g of the soluble protein or 10  $\mu$ l of the insoluble fraction was resolved by SDS-PAGE and subjected to Western blot analysis. Data are representative of 4 separate experiments.  
doi:10.1371/journal.pone.0037755.g002


### Modulation of A $\beta$ <sub>42</sub> by Hsp40 is Cell Line Specific

To further investigate the role that specific inducible chaperones play in heat shock induced reduction of A $\beta$ <sub>42</sub>, CAD cells were transfected with the stress induced J protein Hsp40 and then challenged with the toxic A $\beta$ <sub>42</sub> (Figure 7). Both heat shock and Hsp40 transfection reduced soluble and insoluble A $\beta$ <sub>42</sub> (monomer) at 48 hours. Quantitative immunoblotting uncovered a 50%  $\pm$  3 Hsp40-mediated reduction compared to a smaller heat shock-mediated decreases 88%  $\pm$  6 in insoluble monomeric A $\beta$ <sub>42</sub>. Soluble A $\beta$ <sub>42</sub> monomer was found to decrease to 67%  $\pm$  12 following Hsp40 transfection and 48%  $\pm$  15 following heat shock. Hsp40 and heat shock both caused changes in A $\beta$ <sub>42</sub> oligomerization, initially increasing the A $\beta$ <sub>42</sub> 72 kDa oligomer followed by a decrease at 48 hrs (Figure 7). These experiments clearly establish Hsp40 as a chaperone that influences cellular clearance of A $\beta$ <sub>42</sub>. Transfection does not induce the stress response (Figure 7C). Likewise, residues encoding amino acids 106–126 of PrP<sup>C</sup> as well as a scrambled control do not induce Hsp70 or increase Hsp40 levels.

We then asked the question whether Hsp40 also regulates A $\beta$ <sub>42</sub> levels in primary hippocampal co-cultures of neurons and glia. Rat hippocampal neurons were isolated on postnatal day 0, transfected by electroporation with cDNA for myc-tagged Hsp40 or pCMV vector. Extensive neuritic outgrowth was found in both control

and transfected cultures (Figure 8). Bassoon (presynaptic-red) and neurofilament (green) expressing cells are apparent. DAPI staining is shown in blue. Healthy neurons are visible in both control and A $\beta$ <sub>42</sub>-treated cultures. 3 days post-transfection cultures were treated with 10  $\mu$ M A $\beta$ <sub>42</sub> and 4 days later cultures were washed in PBS and cellular (total) A $\beta$ <sub>42</sub> was determined by quantitative immunoblotting. Total lysates were prepared by direct cell lysis in TX100 lysis buffer followed by SDS sample buffer to ensure that all A $\beta$ <sub>42</sub> present was evaluated by Western blot analysis. Exposure of primary hippocampal cultures to extracellular A $\beta$ <sub>42</sub> resulted in rapid A $\beta$ <sub>42</sub> clearance. Figure 9 shows that 4 days following 10  $\mu$ M A $\beta$ <sub>42</sub> exposure primary rat hippocampal cultures, which have high endogenous levels of Hsp40, reduce A $\beta$ <sub>42</sub> as does vector transfected neurons. To our surprise, in contrast to CAD neural cultures, transfection of hippocampal neurons with Hsp40 increased (59.3 fold) A $\beta$ <sub>42</sub> monomer levels over nontransfected cultures. These results reveal stark differences in neuronal processing of A $\beta$ <sub>42</sub> following Hsp40 transfection between CAD cell cultures and hippocampal cultures.

To begin to address the mechanism of the Hsp40-mediated increase in A $\beta$ <sub>42</sub> hippocampal co-cultures were transfected with a mutated form of Hsp40 predicted to bind to client proteins but not have chaperone activity. Hsp40 interacts with and activates the ATPase Hsp70/Hsc70 via its J domain. An HPD (histidine-



**Figure 3. PrP<sup>C</sup> associates with CAD cells.** (A) CAD cells were incubated with 3  $\mu$ M A $\beta_{42}$  and/or 0.5  $\mu$ M bovine rPrP<sup>C</sup>, after ~42 hours cells were subjected to heat-shock at 43°C for 40 minutes and then allowed to recover for 5 hours. The cells were rinsed in PBS and lysed. 30  $\mu$ g of the soluble protein or 10  $\mu$ l of the insoluble fraction was resolved by SDS-PAGE and cellular PrP<sup>C</sup> was determined by Western analysis. (B) Purified rPrP<sup>C</sup> was added to DMEM/F12 tissue culture media, subjected to heat-shock (43°C for 40 minutes) and then incubated at 37°C for 48 hours in the absence of cell lines and dissolved directly in sample buffer.  
doi:10.1371/journal.pone.0037755.g003

proline-aspartic acid) motif located within the J domain of Hsp40 is required for harnessing Hsc70/Hsp70 for conformational work. By mutating the histidine-proline-aspartic acid motif of Hsp40 to alanines (Hsp40<sub>HDP-AAA</sub>) an Hsp40 that binds client protein but does not activate Hsc70/Hsp70 ATPase is generated. Transfection of hippocampal neurons with Hsp40<sub>HDP-AAA</sub> increased A $\beta_{42}$  monomer (15.3 fold) compared to vector control (7.9 fold) (Figure 9A&B). Although Hsp40<sub>HDP-AAA</sub> increased cellular A $\beta_{42}$  levels over untreated control cells (no transfection, no extracellular A $\beta_{42}$ ) and vector control cultures (transfection with vector alone followed by application of extracellular A $\beta_{42}$ ), the Hsp40<sub>HDP-AAA</sub>-mediated increase (15.3 fold) was smaller compared to Hsp40 (59.3 fold) indicating that the HPD motif within the J domain of Hsp40 impacts directly on cell processing of A $\beta_{42}$ . Figure 9C & D show that transfection efficiency is lower in co cultures compared to CAD cells. Like that seen for CAD cells, A $\beta_{42}$  did not induce the heat shock response in hippocampal neurons. PrP<sup>C</sup> did not block the association/accumulation of A $\beta_{42}$


in hippocampal neurons (data not shown). Figure 9D compares total cellular hippocampal cultures exposed to 10, 20, 25  $\mu$ M A $\beta_{42}$  at the time of transfection (left panel) and three days following transfection (right panel) to CAD cells treated for 24 hours with 25  $\mu$ M A $\beta_{42}$ .

Taken together, these results demonstrate that modulation of A $\beta_{42}$  by Hsp40 is cell line specific.


In summary we have found that Hsp40 is able to influence cellular levels of A $\beta_{42}$ . Neural processing of extracellular A $\beta_{42}$  is dynamic. Overproduction and impaired clearance of A $\beta_{42}$ , are implicated in AD [8]. Accumulation of A $\beta$  oligomers can lead to synaptic dysfunction and disruption in neural plasticity [42–45], however, details of the molecular cascade(s) that underlie A $\beta_{42}$  neuronal toxicity remain unclear. Understanding how cells regulate cellular A $\beta_{42}$  levels requires identification of the cellular chaperone machinery involved in processing the neural amyloid pool. In this study we provide mechanistic evidence that Hsp40 regulates association/accumulation of extracellular A $\beta_{42}$  with neurons and that Hsp40-mediated regulation is cell specific.

Hsp40 is an evolutionarily ancient and widely expressed chaperone that almost certainly targets multiple client proteins. In neurons, Hsp40 is found to be concentrated in postsynaptic densities [46], in lipid rafts [47] and in association with presynaptic chaperones [38]. Furthermore, Hsp40 is linked to neurite outgrowth [48]. That said, its precise role in synaptic transmission and neurodegeneration is not yet known. Hsp40 is a member of the J protein family [49]. All J proteins have a tetrahelical Hsp70/Hsc70-interacting domain called a J domain. Via their J domain, J proteins target a wide array of cellular proteins to the ATPases Hsc70/Hsp70 for conformational work. Although our observation that Hsp40 alters A $\beta_{42}$  cellular turnover in a cell culture specific manner is in contrast to the notion that heat shock chaperones rid the cell of toxic proteins in all cells, it is possible that in disease conditions Hsp40 may protect a toxic protein (eg A $\beta_{42}$ ) from triage resulting in acceleration of disease progression. In fact, the J protein family determines which chaperone pathway is pursued by Hsc70/Hsp70 [50]. For example, DnaJB2(HSJ1) stimulates ubiquitination and sorting of substrates to the proteasome [51], while DnaJC6(auxilin) stimulates recycling of clathrin from clathrin coated vesicles [52] and DnaJB6 (Mrj) regulates keratin turnover [53]. Further experimentation is required to understand how chaperones like Hsp40 either aid in folding and maintenance or lead to degradation.

Cellular levels of Hsp40 routinely rise and fall as part of the heat shock program. Levels of constitutive Hsp40 vary drastically among neural cell lines [38] and classes of neurons in the adult rat brain [54] but how this correlates with A $\beta_{42}$  levels remains to be established. Although the heat shock response is a highly conserved cellular program that confers transient cyto-protection via the induction and translocation of stress induced chaperones, differences in the threshold for the induction of the heat shock response, the complement of heat shock proteins induced as well as the constitutive expression of heat shock proteins are observed among neurons. For example, Hsp27 is induced by heat shock in primary hippocampal cultures but not primary cortical cultures [18]. Also, some neurons (eg motor neurons) have a high threshold for inducing the heat shock response [55]. The cellular variations in either constitutive expression or heat shock expression of Hsp40 may well lead to selective vulnerability of neurons. Other heat shock proteins have been shown to protect against A $\beta_{42}$  neuronal toxicity. Expression of Hsp27 in cortical neuronal cultures prepared from postnatal day 1 rats protects against A $\beta_{42}$ -associated toxicity [18]. Also, Hsp70 overexpressing mice have reduced A $\beta$  [19]. Our data show that extracellular A $\beta_{42}$  does not



**Figure 4. A $\beta_{42}$  levels are reduced 48 hours following induction of the cellular heat shock response.** CAD cells were incubated with 3  $\mu$ M A $\beta_{42}$  and/or 0.5  $\mu$ M bovine rPrP<sup>C</sup>. Immediately following addition of A $\beta_{42}$  or rPrP<sup>C</sup>, the cells were subjected to heat-shock at 43°C for 40 minutes, allowed to recover for 48 hours, and washed in PBS prior to lysis. 30  $\mu$ g of soluble cellular protein or 10  $\mu$ l of the insoluble fraction was resolved by SDS-PAGE and subjected to Western blot analysis. Lower panel: CAD cells were incubated with the indicated increasing concentrations of A $\beta_{42}$  and/or 0.5  $\mu$ M mouse (L42) rPrP<sup>C</sup>. Data are representative of 3 separate experiments.  
doi:10.1371/journal.pone.0037755.g004



**Figure 5. A $\beta_{42}$  is not directly altered by heat shock.** (A) Purified A $\beta_{42}$  was added to DMEM/F12 tissue culture media, subjected to heat-shock (43°C for 40 minutes) and then incubated at 37°C for 48 hours in the absence of cell lines and dissolved directly in sample buffer. doi:10.1371/journal.pone.0037755.g005


elicit a heat shock response in neurons nor does it alter induction of the heat shock response. Moreover we demonstrate that while heat shock initially increases neural A $\beta_{42}$ , with time in culture A $\beta_{42}$  levels are reduced in neurons following heat shock compared to control cultures.

The basis of the cell specific responses to Hsp40 is likely due to differences in the chaperone networks between the cultured cells. The elaborate chaperone machinery that is present in cells rids the cell of toxic proteins often via assembly of chaperones into active chaperone complexes. Levels of chaperones and chaperone complexes may differ between CAD neuroblastoma cells and primary hippocampal cultures. Further experimentation is required to develop a detailed and comprehensive overview of differences in chaperones among neurons. In conclusion we have shown that (1) Hsp40 modulates processing of extracellular A $\beta_{42}$ . (2) Hsp40 modulation of A $\beta_{42}$  is cell specific and dependent on the conserved HPD motif within Hsp40's J domain. (3) A $\beta_{42}$  does not trigger the heat shock response or alter the threshold for induction of the heat shock response. (4) Soluble PrP<sup>C</sup> does not block association of extracellular A $\beta_{42}$ . Overall these results elucidate an important link between Hsp40 and cellular levels of A $\beta_{42}$ , which has not been illustrated previously.

## Materials and Methods

### Reagents and Chemicals

Amyloid- $\beta_{42}$  (A $\beta_{42}$ ) was from rPeptide. Recombinant bovine PrP<sup>C</sup> (rPrP<sup>C</sup>) was from Prionics. Anti-A $\beta$  6E10 monoclonal antibody and DnaJA4 monoclonal antibody were from Cedarlane Laboratories. Anti-Hsp40 rabbit polyclonal, anti-Hsp70 mouse monoclonal and anti-Hsp25 mouse monoclonal were from Stressgen. Anti-Hsc70 mouse monoclonal and anti- $\beta$ -actin mouse monoclonal were from Sigma. Geldanamycin was from Calbiochem. Anti-Rdj2 mouse monoclonal and anti-DnaJA3 (Tid1)




**Figure 6. A $\beta_{42}$  or bovine rPrP<sup>C</sup> do not alter expression of the J proteins DnaJA1(Hdj2), DnaJA2(Rdj2), DnaJA3 (Tid1) or DnaJA4 in neural cells.** CAD cells were incubated with 3  $\mu$ M A $\beta_{42}$  or 0.5  $\mu$ M bovine rPrP<sup>C</sup> for 48 hours and washed in PBS prior to lysis. (A) Western analysis of heat shock proteins (B) The J proteins; DnaJA1(Hdj2), DnaJA2(Rdj2), DnaJA3 (Tid1) or DnaJA4 were detected by Western blot analysis.  $\beta$ -actin is shown as a loading control. Data are representative of 4 separate experiments. doi:10.1371/journal.pone.0037755.g006

polyclonal antibody were from Abnova. Caspase-3 (8G10) Rabbit monoclonal antibody was from Cell Signaling Technology. DnaJ1 (Hdj2) polyclonal antibody was from USBiological. Peroxidase-conjugates AffiniPure Goat Anti-Mouse IgG (H+L) was from Cedarlane.


### CAD (CNS Catecholaminergic Derived) Mouse Neuroblastoma Cells [38,56,57]

CAD (CNS catecholaminergic derived) mouse neuroblastoma cells were seeded into 6 well plates and grown in DMEM/F12 medium supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin as previously described. Cells were lysed in




**Figure 7. Transient transfection of Hsp40 reduces cellular A $\beta$ <sub>42</sub>.** (A) CAD cells were transiently transfected with Hsp40 as indicated for 24 hours prior to the addition of 3  $\mu$ M A $\beta$ <sub>42</sub>. Immediately following addition of A $\beta$ <sub>42</sub>, indicated cells were subjected to heat shock at 43°C for 40 minutes and allowed to recover. At the indicated times cells were washed in PBS, lysed separated into soluble and insoluble fractions and cellular levels of A $\beta$ <sub>42</sub> were determined by Western analysis. Actin is shown as a loading control (B) Quantification of three independent experiments. \*p<0.05. (C) Western analysis of CAD cells transfected with Hsp40 as indicated prior to incubation with 3  $\mu$ M A $\beta$ <sub>42</sub> or 0.5  $\mu$ M PrP<sup>106-126</sup> or scrambled control. Data are representative of 4 separate experiments. doi:10.1371/journal.pone.0037755.g007





**Figure 8. Effect of A $\beta$ <sub>42</sub> treatment on Hsp40 and Hsp40<sub>HPD-AAA</sub>-transfected neurons.** Rat hippocampal neurons were transfected by electroporation with 4  $\mu$ g myc-tagged Hsp40, Hsp40<sub>HPD-AAA</sub> or vector alone and plated on silicon wafers (Silicon wafers, Silicon Quest, CA). Three days after transfection, neurons were incubated with 10  $\mu$ M A $\beta$ <sub>42</sub> (panels B,D,F,H) and 7 days post transfection hippocampal cultures were fixed with 15% Picric Acid (Sigma, P6744), 4% PFA (Sigma, P6148), blocked in donkey serum/BSA and permeabilized with 0.1% Triton-X100 (Biorad, 161-0407) for 2 hours. Cultures were immunostained with Bassoon (red) (Enzo, ADI-VAM-PS003), and Neurofilament (green) (Millipore, A1991) and DAPI (blue) (Invitrogen, D1306) and examined by microscopy via an Olympus inverted scope (BX61WI) and water submersible 10x objective (UMplanF1). The panels are as follows A,B = controls; C,D transfected with pcDNA; E,F transfected with Hsp40 and G,H transfected with Hsp40<sub>HPD-AAA</sub>; A,C,E,G = no A $\beta$ <sub>42</sub>; B,D,F,H = 10  $\mu$ M of A $\beta$ <sub>42</sub>. Neurons are healthy with axon and dendritic elaboration forming networks.  
doi:10.1371/journal.pone.0037755.g008



**Figure 9. Transfection of hippocampal co-cultures with Hsp40 or Hsp40<sub>HSPD-AAA</sub> elevates cellular A $\beta_{42}$ .** (A) Rat hippocampal neurons were transfected by electroporation with 4  $\mu$ g myc-tagged Hsp40, Hsp40<sub>HSPD-AAA</sub> or vector alone. Three days after transfection, neurons were incubated with 10  $\mu$ M A $\beta_{42}$  and 7 days post transfection hippocampal cultures were washed in PBS and cellular (total) A $\beta_{42}$  was determined by

quantitative immunoblotting. Actin is shown for reference. **(B)** Quantification of three independent experiments. \* $p < 0.01$  **(C)** CAD cells were transfected with 1, 2, 4  $\mu$ g myc-tagged Hsp40 as indicated. Hsp40 and actin were determined 3 days following transfection. **(D)** Western analysis of hippocampal primary cultures (3 or 4 days following external A $\beta$ <sub>42</sub> application) or CAD cells treated with A $\beta$ <sub>42</sub> for 24 hours. Data are representative of 3 separate experiments.  
doi:10.1371/journal.pone.0037755.g009

40 mM Tris (pH 7.4), 150 mM NaCl, 2 mM EDTA, 1 mM EGTA, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 0.1% SDS, 1% TX-100, 0.5 mM PMSF and protease inhibitor (Sigma) at 4°C for 1 hour. Lysates were centrifuged at 15000 $\times$ g for 5 minutes at 4°C and the supernatant (soluble fraction) and pellet (insoluble fraction) was collected and stored at -70°C. For transient transfection, CAD cells were washed in PBS and transiently transfected with c-myc tagged rat Hsp40 DNA using Lipofectamine-2000 (Invitrogen) in Opti-MEM. For heat shock experiments cells were incubated at 43°C for 40 minutes and then returned to 37°C. Amyloid- $\beta$ <sub>1-42</sub>-TFA was dissolved in 1% NH<sub>4</sub>OH as recommended by the supplier (rPeptide) to a concentration of 1 mg/mL. The resulting solution was sonicated for 1 minute, aliquoted and stored at -70°C. With this preparation, A $\beta$ <sub>42</sub> will be mainly in the monomeric form (~95%) [58]. Recombinant bovine Prion Protein (Prionics, amino acids PrP<sup>C25-242</sup>) was suspended in H<sub>2</sub>O, aliquoted and stored at -70°C. PrP<sup>C106-126</sup> and the scrambled PrP<sup>C106-126</sup> (Anaspec) were resuspended in DMSO to a final stock concentration of 1 mM and stored at -20°C [59]. Recombinant mouse His-tagged L42 prion protein was prepared as previously described [60,61]. Aliquots were diluted in culture media immediately prior to treatment of cells. Protein concentration of the soluble CAD cell fraction was determined by Bradford assay (BioRad). The 1% TX-100/0.1% SDS insoluble cell fraction was dissolved directly in sample buffer.

### Primary Hippocampal Cell Culture [62,63]

Dissociated primary co-cultures of neurons and glia were isolated by dissection from Sprague Dawley rats (Charles River) at postnatal day 0 as previously described [62]. Animals were anesthetized on ice and sacrificed by decapitation. Hippocampi were removed and incubated in 40  $\mu$ l/ml of papain. After 30 minutes cells were washed three times in fresh Eagle's basal media (GIBCO-Invitrogen) supplemented with B-27, penicillin, streptomycin, L-glutamine and 4% feta bovine serum. Cells were triturated using three decreasing calibers of pipettes and the cell solution was then transfected by electroporation (Bio Rad Gene Pulsor Xcell: settings 150 volts, pulse length 25, 5 pulses, pulse

interval 0.1 and cuvette size 4 mm) with cDNA of either myc-tagged Hsp40 or Hsp40<sub>HPD-AAA</sub> or vector control (pCDNA3.1) and then plated. For immunostaining, transfected cells were plated on silicon wafers (Silicon wafers, Silicon Quest, CA). Co-cultures were grown for 7 days in a 5% CO<sub>2</sub> incubator. After transfection, A $\beta$ <sub>42</sub> was added to the culture and 3 days later the cells were washed in PBS, lysed in 40 mM Tris (pH 7.4), 150 mM NaCl, 2 mM EDTA, 1 mM EGTA, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 0.1% SDS, 1% TX-100, 0.5 mM PMSF and protease inhibitor (Sigma) at 4°C for 1 hour. Total cell lysates were solubilized directly in sample buffer and evaluated by Western blot analysis. The University of Calgary Conjoint Faculties Research Ethics Board specifically approved this study (protocol number M09008).

### Immunoblotting

Proteins were electrotransferred from polyacrylamide gels to 0.2  $\mu$ m nitrocellulose membrane in 20 mM Tris, 150 mM glycine and 12% methanol. Membranes were blocked with in PBS with 0.1% Tween 20, 4% milk and incubated with primary antibody overnight at 4°C. The membranes were washed and incubated with horseradish peroxidase-coupled secondary antibody. The signal was developed using West Pico reagent (Pierce Biotechnology Inc.) and exposed to Kodak film. Bound antisera were quantitated by Biorad Fluor-S MultiImager Max and QuantityOne 4.2.1 software.

### Acknowledgments

We are grateful to Dr. Frank Jirik and Dr. Pankaj Tailor for supplying PrP<sup>C106-126</sup> and PrP<sup>Cscrambled</sup> peptides and for scientific discussions during the course of these experiments.

### Author Contributions

Conceived and designed the experiments: AC JEAB MC. Performed the experiments: AC JEAB LS. Analyzed the data: AC JEAB MC SWK. Contributed reagents/materials/analysis tools: AC JEAB LS MC EA RW JP. Wrote the paper: JEAB.

### References

1. Querfurth HW, LaFerla FM (2010) Alzheimer's disease. *N Engl J Med* 362: 329–344.
2. Karran E, Mercken M, De Strooper B (2011) The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of therapeutics. *Nat Rev Drug Discov* 10: 698–712.
3. Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353–356.
4. Demuro A, Parker I, Stutzmann GE (2010) Calcium signaling and amyloid toxicity in Alzheimer disease. *J Biol Chem* 285: 12463–12468.
5. Sakono M, Zako T (2010) Amyloid oligomers: formation and toxicity of Abeta oligomers. *FEBS J* 277: 1348–1358.
6. Mohamed A, Posse dC (2011) Abeta internalization by neurons and glia. *Int J Alzheimers Dis* 2011: 127984. 127984 p.
7. Tampellini D, Rahman N, Lin MT, Capetillo-Zarate E, Gouras GK (2011) Impaired beta-Amyloid Secretion in Alzheimer's Disease Pathogenesis. *J Neurosci* 31: 15384–15390.
8. Bingol B, Sheng M (2011) Deconstruction for reconstruction: the role of proteolysis in neural plasticity and disease. *Neuron* 69: 22–32.
9. Westerheide SD, Morimoto RI (2005) Heat shock response modulators as therapeutic tools for diseases of protein conformation. *J Biol Chem* 280: 33097–33100.
10. Muchowski PJ, Wacker JL (2005) Modulation of neurodegeneration by molecular chaperones. *Nat Rev Neurosci* 6: 11–22.
11. Magrane J, Querfurth HW (2008) Heat Shock Proteins, Unfolded Protein Response Chaperones and Alzheimer's Disease. In: Asea AA, Brown IR, eds. *Heat Shock Proteins and the Brain: Implications for Neurodegenerative Diseases and Neuroprotection* Springer. pp 25–50.
12. Perez N, Sugar J, Charya S, Johnson G, Merrill C, et al. (1991) Increased synthesis and accumulation of heat shock 70 proteins in Alzheimer's disease. *Brain Res Mol Brain Res* 11: 249–254.
13. Hamos JE, Oblas B, Pulaski-Salo D, Welch WJ, Bole DG, et al. (1991) Expression of heat shock proteins in Alzheimer's disease. *Neurology* 41: 345–350.
14. Shinohara H, Inaguma Y, Goto S, Inagaki T, Kato K (1993) Alpha B crystallin and HSP28 are enhanced in the cerebral cortex of patients with Alzheimer's disease. *J Neurol Sci* 119: 203–208.
15. Fonte V, Kapulkin V, Taft A, Fluet A, Friedman D, et al. (2002) Interaction of intracellular beta amyloid peptide with chaperone proteins. *Proc Natl Acad Sci U S A* 99: 9439–9444.
16. Evans CG, Wisen S, Gestwicki JE (2006) Heat shock proteins 70 and 90 inhibit early stages of amyloid beta-(1–42) aggregation in vitro. *J Biol Chem* 281: 33182–33191.
17. Magrane J, Rosen KM, Smith RC, Walsh K, Gouras GK, et al. (2005) Intraneuronal beta-amyloid expression downregulates the Akt survival pathway and blunts the stress response. *J Neurosci* 25: 10960–10969.

18. King M, Nafar F, Clarke J, Mearow K (2009) The small heat shock protein Hsp27 protects cortical neurons against the toxic effects of beta-amyloid peptide. *J Neurosci Res* 87: 3161–3175.
19. Hoshino T, Murao N, Namba T, Takehara M, Adachi H, et al. (2011) Suppression of Alzheimer's disease-related phenotypes by expression of heat shock protein 70 in mice. *J Neurosci* 31: 5225–5234.
20. Ojha J, Karmegam RV, Masilamoni JG, Terry AV, Cashikar AG (2011) Behavioral defects in chaperone-deficient Alzheimer's disease model mice. *PLoS ONE* 6: e16550.
21. Stege GJ, Renkawek K, Overkamp PS, Verschuure P, van Rijk AF, et al. (1999) The molecular chaperone alphaB-crystallin enhances amyloid beta neurotoxicity. *Biochem Biophys Res Commun* 262: 152–156.
22. Sakono M, Zako T, Ueda H, Yohda M, Maeda M (2008) Formation of highly toxic soluble amyloid beta oligomers by the molecular chaperone prefoldin. *FEBS J* 275: 5982–5993.
23. Oda T, Wals P, Osterburg HH, Johnson SA, Pasinetti GM, et al. (1995) Clusterin (apoJ) alters the aggregation of amyloid beta-peptide (A beta 1-42) and forms slowly sedimenting A beta complexes that cause oxidative stress. *Exp Neurol* 136: 22–31.
24. Carnini A, Lear JD, Eckenhoff RG (2007) Inhaled anesthetic modulation of amyloid beta(1-40) assembly and growth. *Curr Alzheimer Res* 4: 233–241.
25. Eckenhoff RG, Johansson JS, Wei H, Carnini A, Kang B, et al. (2004) Inhaled anesthetic enhancement of amyloid-beta oligomerization and cytotoxicity. *Anesthesiology* 101: 703–709.
26. Bateman DA, McLaurin J, Chakrabarty A (2007) Requirement of aggregation propensity of Alzheimer amyloid peptides for neuronal cell surface binding. *BMC Neurosci* 8: 29.
27. Bateman DA, Chakrabarty A (2009) Two distinct conformations of A beta aggregates on the surface of living PC12 cells. *Biophys J* 96: 4260–4267.
28. Bateman DA, Chakrabarty A (2011) Cell surface binding and internalization of abeta modulated by degree of aggregation. *Int J Alzheimers Dis* 2011: 962352. 962352 p.
29. Lauren J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM (2009) Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* 457: 1128–1132.
30. Balducci C, Beeg M, Stravalaci M, Bastone A, Scip A, et al. (2010) Synthetic amyloid-beta oligomers impair long-term memory independently of cellular prion protein. *Proc Natl Acad Sci U S A* 107: 2295–2300.
31. Gimbel DA, Nygaard HB, Coffey EE, Gunther EC, Lauren J, et al. (2010) Memory impairment in transgenic Alzheimer mice requires cellular prion protein. *J Neurosci* 30: 6367–6374.
32. Schwarze-Eicker K, Keyvani K, Gortz N, Westaway D, Sachser N, et al. (2005) Prion protein (PrP<sup>c</sup>) promotes beta-amyloid plaque formation. *Neurobiol Aging* 26: 1177–1182.
33. You H, Tsutsui S, Hameed S, Kannanayakal TJ, Chen L, et al. (2012) Abeta neurotoxicity depends on interactions between copper ions, prion protein, and N-methyl-D-aspartate receptors. *Proc Natl Acad Sci U S A* 109: 1737–1742.
34. Caetano FA, Lopes MH, Hajj GN, Machado CF, Pinto AC, et al. (2008) Endocytosis of prion protein is required for ERK1/2 signaling induced by stress-inducible protein 1. *J Neurosci* 28: 6691–6702.
35. Yang AJ, Chandswangbhuvana D, Margol L, Glabe CG (1998) Loss of endosomal/lysosomal membrane impermeability is an early event in amyloid Abeta1-42 pathogenesis. *J Neurosci Res* 52: 691–698.
36. Yang AJ, Chandswangbhuvana D, Shu T, Henschen A, Glabe CG (1999) Intracellular accumulation of insoluble, newly synthesized abeta1-42 in amyloid precursor protein-transfected cells that have been treated with Abeta1-42. *J Biol Chem* 274: 20650–20656.
37. Parkin ET, Watt NT, Hussain I, Eckman EA, Eckman CB, et al. (2007) Cellular prion protein regulates beta-secretase cleavage of the Alzheimer's amyloid precursor protein. *Proc Natl Acad Sci U S A* 104: 11062–11067.
38. Gibbs SJ, Barren B, Beck KE, Proft J, Zhao X, et al. (2009) Hsp40 couples with the CSPalpha chaperone complex upon induction of the heat shock response. *PLoS ONE* 4: e4595.
39. Stahl N, Borchelt DR, Hsiao K, Prusiner SB (1987) Scrapie prion protein contains a phosphatidylinositol glycolipid. *Cell* 51: 229–240.
40. Naslavsky N, Stein R, Yanai A, Friedlander G, Taraboulos A (1997) Characterization of detergent-insoluble complexes containing the cellular prion protein and its scrapie isoform. *J Biol Chem* 272: 6324–6331.
41. Haraguchi T, Fisher S, Olofsson S, Endo T, Groth D, et al. (1989) Asparagine-linked glycosylation of the scrapie and cellular prion proteins. *Arch Biochem Biophys* 274: 1–13.
42. Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, et al. (2009) Soluble oligomers of amyloid beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 62: 788–801.
43. Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, et al. (2008) Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 14: 837–842.
44. Tyszkiewicz JP, Yan Z (2005) beta-Amyloid peptides impair PKC-dependent functions of metabotropic glutamate receptors in prefrontal cortical neurons. *J Neurophysiol* 93: 3102–3111.
45. Wang Q, Walsh DM, Rowan MJ, Selkoe DJ, Anwyl R (2004) Block of long-term potentiation by naturally secreted and synthetic amyloid beta-peptide in hippocampal slices is mediated via activation of the kinases c-Jun N-terminal kinase, cyclin-dependent kinase 5, and p38 mitogen-activated protein kinase as well as metabotropic glutamate receptor type 5. *J Neurosci* 24: 3370–3378.
46. Suzuki T, Usuda N, Murata S, Nakazawa A, Ohtsuka K, et al. (1999) Presence of molecular chaperones, heat shock cognate (Hsc) 70 and heat shock proteins (Hsp) 40, in the postsynaptic structures of rat brain. *Brain Res* 816: 99–110.
47. Chen S, Bawa D, Beshoh S, Gurd JW, Brown IR (2005) Association of heat shock proteins and neuronal membrane components with lipid rafts from the rat brain. *J Neurosci Res* 81: 522–529.
48. Takeuchi H, Kobayashi Y, Yoshihara T, Niwa J, Doyu M, et al. (2002) Hsp70 and Hsp40 improve neurite outgrowth and suppress intracytoplasmic aggregate formation in cultured neuronal cells expressing mutant SOD1. *Brain Res* 949: 11–22.
49. Zhao X, Braun AP, Braun JE (2008) Biological Roles of Neural J Proteins. *Cell Mol Life Sci* 65: 2385–2396.
50. Kampinga HH, Craig EA (2010) The HSP70 chaperone machinery: J proteins as drivers of functional specificity. *Nat Rev Mol Cell Biol* 11: 579–592.
51. Westhoff B, Chapple JP, van der Spuy J, Hohfeld J, Cheetham ME (2005) HSP70 is a neuronal shuttling factor for the sorting of chaperone clients to the proteasome. *Curr Biol* 15: 1058–1064.
52. Ungewickell E, Ungewickell H, Holstein SEH, Linder R, Prasad K, et al. (1995) Role of auxilin in uncoating clathrin-coated vesicles. *Nature* 378: 632–635.
53. Watson ED, Geary-Joo C, Hughes M, Cross JC (2007) The Mrj co-chaperone mediates keratin turnover and prevents the formation of toxic inclusion bodies in trophoblast cells of the placenta. *Development* 134: 1809–1817.
54. Chen S, Brown IR (2007) Neuronal expression of constitutive heat shock proteins: implications for neurodegenerative diseases. *Cell Stress Chaperones* 12: 51–58.
55. Batulan Z, Shinder GA, Minotti S, He BP, Doroudchi MM, et al. (2003) High threshold for induction of the stress response in motor neurons is associated with failure to activate HSF1. *J Neurosci* 23: 5789–5798.
56. Rosales-Hernandez A, Beck KE, Zhao X, Braun AP, Braun JE (2008) RDJ2 (DNAJA2) chaperones neural G protein signaling pathways. *Cell Stress Chaperones* 14: 71–82.
57. Xu F, Proft J, Gibbs S, Winkfein B, Johnson JN, et al. (2010) Quercetin targets cysteine string protein (CSPalpha) and impairs synaptic transmission. *PLoS ONE* 5: e11045.
58. Barghorn S, Nimmrich V, Striebinger A, Krantz C, Keller P, et al. (2005) Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease. *J Neurochem* 95: 834–847.
59. Florio T, Thellung S, Amico C, Robello M, Salmona M, et al. (1998) Prion protein fragment 106-126 induces apoptotic cell death and impairment of L-type voltage-sensitive calcium channel activity in the GH3 cell line. *J Neurosci Res* 54: 341–352.
60. Eiden M, Buschmann A, Kupfer L, Groschup MH (2006) Synthetic prions. *J Vet Med B Infect Dis Vet Public Health* 53: 251–256.
61. Zahn R, von Schroetter C, Wuthrich K (1997) Human prion proteins expressed in *Escherichia coli* and purified by high-affinity column refolding. *FEBS Lett* 417: 400–404.
62. Morales M, Colicos MA, Goda Y (2000) Actin-dependent regulation of neurotransmitter release at central synapses. *Neuron* 27: 539–550.
63. Gutierrez RC, Hung J, Zhang Y, Kertesz AC, Espina FJ, et al. (2009) Altered synchrony and connectivity in neuronal networks expressing an autism-related mutation of neuroligin 3. *Neuroscience* 162: 208–221.