

Effect of Hydrogen Peroxide on Immersion Challenge of Rainbow Trout Fry with *Flavobacterium psychrophilum*

Maya Maria Mihályi Henriksen*, Lone Madsen, Inger Dalsgaard

Technical University of Denmark, National Veterinary Institute, Bülowsvej 27, Frederiksberg C, Denmark

Abstract

An experimental model for immersion challenge of rainbow trout fry (*Oncorhynchus mykiss*) with *Flavobacterium psychrophilum*, the causative agent of rainbow trout fry syndrome and bacterial cold water disease was established in the present study. Although injection-based infection models are reliable and produce high levels of mortality attempts to establish a reproducible immersion model have been less successful. Various concentrations of hydrogen peroxide (H₂O₂) were evaluated before being used as a pre-treatment stressor prior to immersion exposure to *F. psychrophilum*. H₂O₂ accelerated the onset of mortality and increased mortality approximately two-fold; from 9.1% to 19.2% and from 14.7% to 30.3% in two separate experiments. Clinical signs observed in the infected fish corresponded to symptoms characteristically seen during natural outbreaks. These findings indicate that pre-treatment with H₂O₂ can increase the level of mortality in rainbow trout fry after exposure to *F. psychrophilum*.

Citation: Henriksen MMM, Madsen L, Dalsgaard I (2013) Effect of Hydrogen Peroxide on Immersion Challenge of Rainbow Trout Fry with *Flavobacterium psychrophilum*. PLoS ONE 8(4): e62590. doi:10.1371/journal.pone.0062590

Editor: Dongsheng Zhou, Beijing Institute of Microbiology and Epidemiology, China

Received: December 14, 2012; **Accepted:** March 22, 2013; **Published:** April 30, 2013

Copyright: © 2013 Henriksen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was performed as a part of the Danish Fish Immunology Research Centre and Network (DAFINET; www.dafinet.dk) supported by the Danish Council of Strategic Research (grant 2101-080017). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: mmah@vet.dtu.dk

Introduction

Flavobacterium psychrophilum is a Gram-negative fish pathogen and the causative agent of bacterial cold water disease (BCWD), also called rainbow trout fry syndrome (RTFS). Since first being discovered in USA in the 1940s, the disease has spread worldwide and was identified in Western Europe in the mid 1980s. Today it has a significant impact on rainbow trout (*Oncorhynchus mykiss*) and other salmonid fish in aquaculture worldwide [1]. Clinical signs of the disease include a dark coloration of the skin, pale gills and organs (kidney, intestine and liver) due to anemia. A loss of appetite and lethargic behavior is also observed.

Injection-based experimental challenges with *F. psychrophilum* have been standardized and the resulting mortality depends on several factors, including the number of colony forming units (CFU) injected, size of the fish, batch differences and the number of fish in each tank [2]. However, infection by injection is not a suitable approach for all investigations since the first line of defense, consisting of non-specific barriers, are bypassed. It has proven difficult to produce high and consistent mortality rates using cohabitation and immersion models unless stress or scarification has been applied [2–7], while immersion exposure to *F. psychrophilum* in the logarithmic phase of growth reportedly results in significant mortality rates [8].

Bath-treatment with various non-medical compounds, such as copper sulphate, chloramine-T, sodium carbonates, sodium chloride, formalin and hydrogen peroxide (H₂O₂), are routinely used against pathogens in aquaculture [9]. One previous immersion model for *F. psychrophilum* used formalin as stressor, which raised the mortality rate significantly [2]. Since the use of formalin is to be phased out by Danish fish farmers due to human

health considerations, another less harmful stressor is needed. H₂O₂ is a useful and environmentally friendly alternative to formalin and has been used against skin parasites, bacterial gill infections and mold on eggs in Denmark for over ten years [10,11]. The use of H₂O₂ has been shown to accelerate *Tenacibaculum maritimum* infections in turbot [12] and *Flavobacterium columnaris* infections in channel catfish [13]. Accordingly, H₂O₂ was an obvious candidate for a stressor in a *F. psychrophilum* immersion model.

The aim of this study was to establish a reproducible method for increasing mortality of RTFS in immersion challenge of rainbow trout fry. A reliable model is needed for studies on pathogen invasion, host immune response and efficacy of treatments. First, various concentrations of H₂O₂ were tested in experiment 1 before being combined with immersion exposure to a virulent *F. psychrophilum* strain at two different phases of bacterial growth in experiment 2. The most favorable H₂O₂ concentration and bacterial growth phase were combined in experiment 3, after which the temperature and number of animals were adjusted in experiment 4 to increase the power of the model.

Materials and Methods

Fish and Rearing Conditions

Three batches of eggs from the same stock and family of fish were used in all four experiments (see Table 1). The eggs originated from Fousing Trout Farm (Denmark) and disinfection and hatching was carried out at AquaBaltic (Bornholm, Denmark). The fish were acclimatized for at least 3 weeks in 200 L tanks containing 15°C recirculated tap water (non-chlorinated), while nitrate, nitrite and ammonia levels were monitored continuously.

The fish were fed dry commercial feed daily (INICIO Plus, BioMar A/S, Denmark). All tanks, except the 1 and 2 L tanks used during challenge and pre-treatment, were opaque in order to minimize external stress due to movement in the experimental facility. Upon arrival, at least 10 fish from each batch were sacrificed by an overdose of MS-222 (Ethyl 3-aminobenzoate methanesulfonate, Sigma), examined for *F. psychrophilum* and proven to be free of the pathogen. Samples from brain and kidney were streaked on tryptone yeast extract salts (TYES) and blood agar plates, while spleen samples were placed in TYES broth (110 rpm). Samples were incubated for 7 d at 15°C for TYES broth/plates and at 20°C for blood plates. Yellow colonies were further examined by species specific PCR. First, a pure culture was obtained by streaking and after 5–7 d of growth, a single colony was suspended in 50 µL distilled water and boiled for 5 min to lyse the bacterial cells and placed on ice. Then, species specific primers, referred to as PSY1 and PSY2, amplifying the 16S rRNA genes of *F. psychrophilum* were used [14] for PCR amplifications, which were carried out according to a previously described method [15]. In brief, Ready to Go PCR beads (cat. no. 27955901, Amersham Pharmacia Biotech, Millwaukee, USA), including all reagents, were used. Ten pmol of each primer and 1 µL of the lysed bacteria were added to each reaction. Distilled water was added to a total volume of 25 µL. Amplifications were carried out using a Biometra T- 3000 thermocycler amplifier (Biometra, Göttingen, Germany): 95°C for 5 min followed by 35 cycles consisting of 30 s at 95°C, 30 s at 57°C and 60 s at 72°C.

Before randomly selecting the experimental groups, the largest (approx. >1.5 g) and smallest (approx. <0.6 g) fish were removed in order to minimize size variation. The fish were inspected at least once every 24 h and dead fish were examined for *F. psychrophilum* as previously described. Fish surviving until the completion of each experiment were euthanized using an overdose of MS-222.

The described work was carried out in accordance with the internationally accepted guidelines for care and use of laboratory animals in research and procedures were approved by the Committee for Animal Experimentation, Ministry of Justice, Copenhagen, Denmark (J.nr. 2006/561-1204 and 2011/561-51). During experiments, the fish were inspected several times a day and precautions were taken to minimize stress by keeping the fish in opaque tanks, minimizing handling and monitoring water chemistry.

Bacterial Strain and Challenge Dose

A Danish *F. psychrophilum* strain, 950106-1/1, was used for all challenges (serotype Fd and elastin degrading). It was isolated from a clinical outbreak of RTFS in a freshwater farm in 1995, and has previously been used for several i.p. and immersion challenges [2]. The bacterial strain was stored at -80°C in TYES broth with 15–20% glycerol and pre-cultivated in 10 mL TYES broth at 15°C

for 3 d (110 rpm). Then, 100 mL TYES broth was inoculated using 0.5 mL of the pre-culture and optical density (OD) was measured at 525 nm using a spectrophotometer (Shimadzu UV-1201) at regular intervals to establish a growth curve. Logarithmic and stationary phases were determined after 24 or 48 h of growth, respectively. Before being used for challenge, each culture was examined microscopically to assert purity and CFU counts were carried out by spreading 0.1 mL of 10-fold dilutions (10⁻⁴ to 10⁻⁷) on TYES agar in duplicates.

Experiment 1: H₂O₂ as Pre-treatment Stressor

Three groups of 6 fish in duplicates (n = 36) were immersed for 60 min in aerated 1 L tanks containing final concentrations of 50, 150 or 300 mg H₂O₂ L⁻¹ (prod. no. 1072090250, Merck KGaA, Germany). H₂O₂ was added directly into the tanks already containing the fish. Subsequently, the fish were removed and placed in 20 L aquaria and observed for 14 d. The temperature during the experiment was 15±0.5°C, while H₂O₂ treatment was carried out at approximately 12°C. Based on the results, a concentration of 200 mg H₂O₂ L⁻¹ was chosen as stressor for experiment 2.

Experiment 2: H₂O₂ Combined with Bacterial Growth Phases (24 and 48 h)

Four groups of 10 fish in duplicates (n = 80) were immersed for 60 min in aerated 1 L tanks containing 200 mg H₂O₂ L⁻¹ or kept under similar conditions in water. H₂O₂ was added directly into the aquaria already containing the fish. Afterwards, the fish were removed and placed for 30 min in a tank containing a 1:9 dilution of cultures grown for either 24 h or 48 h. This resulted in bacterial concentrations of 10⁵ and 10⁷ CFU mL⁻¹ water, respectively. Finally, the fish were placed in 20 L aquaria containing tap water. The fish were observed for 30 d. The temperature during the experiment was 14.5±0.5°C, while H₂O₂ treatment and bacterial challenge were carried out at a temperature below 14°C. The animals dying after exposure to the pathogen were sampled as previously described for the bacteriological examination. Since no significant difference regarding either mortality or reisolation of the pathogen was observed, the 48 h culture was chosen for the following experiments, since it has been used successfully in previous investigations [2].

Experiment 3: 150 mg L⁻¹ H₂O₂ as Experimental Stressor

Four groups of 50 fish in duplicates (n = 400) were immersed for 60 min in aerated 2 L tanks containing 150 mg H₂O₂ L⁻¹ or kept under similar conditions in tap water. H₂O₂ was diluted in 300 mL tap water, before being added to the aquaria already containing the fish. Subsequently, the fish were removed and immersed in a 1:9 diluted broth culture (48 h) containing 10⁷ CFU mL⁻¹ for 30 min, while controls were immersed in

Table 1. Experimental parameters.

Exp.	Batch	Weight	Length	H ₂ O ₂ mg L ⁻¹	F _p CFU mL ⁻¹	°C	n (pr. replicate)
1	1	0.69 g (±0.21)	4.4 cm (±0.3)	50/150/300	–	15.5±1.0	36 (6)
2	1	0.77 g (±0.18)	4.4 cm (±0.3)	–/200	10 ⁵ (24 h)/10 ⁷ (48 h)	16.0±0.5	80 (10)
3	2	1.2 g (±0.50)	5.0 cm (±0.8)	–/150	–/10 ⁷	16.8±0.5	400 (50)
4	3	1.1 g (±0.34)	4.7 cm (±0.5)	–/200	–/10 ⁷	14.0±0.5	500 (50)

Fish batches and mean values for weight, length and temperature are stated along with standard deviation (SD). The applied H₂O₂ concentrations are given in mg L⁻¹, while the final bacterial concentrations in broth used for immersion challenge are stated in CFU mL⁻¹. The symbol ‘–’ designates the absence of F_p and H₂O₂. doi:10.1371/journal.pone.0062590.t001

sterile broth. Finally, the fish were placed in 30 L aquaria and mortality observed for 50 d. The temperature during the experiment was 16.8±0.5°C, while H₂O₂ treatment and bacterial challenge were carried out at a temperature below 14°C. To increase mortality in experiment 4, the temperature was lowered and the concentration of H₂O₂ increased.

Experiment 4:200 mg L⁻¹ H₂O₂ as Experimental Stressor

Four groups of 50 fish in duplicates for uninfected or triplicates for infected (n=500) were immersed for 60 min in aerated 2 L tanks containing 200 mg H₂O₂ L⁻¹ or kept under similar conditions in tap water. H₂O₂ was diluted in 300 mL tap water, before being added to the treatment tanks already containing the fish. Subsequently, the fish were removed and immersed in a 1:9 diluted broth culture (48 h) containing 10⁷ CFU mL⁻¹ for 30 min, while controls were immersed in sterile broth. Finally, the fish were placed in 30 L aquaria and observed for 40 d. The temperature during the experiment was 14±0.5°C, while H₂O₂ treatment and bacterial challenge were carried out at a temperature below 14°C. Conditions during treatments of the various groups were kept as identical as possible with the same number of nettings and amount of retention time in the 2 L tanks during treatments.

Statistical Analysis

Mortality data was analyzed using the generalized linear model (GLM) on the R software platform [16] and figures have been produced using GraphPad Prism 5. The relative standard deviation (RSD = (standard deviation/mean)*100) was also calculated for mortality rates of infected groups in experiment 3 and 4. Weight and length of the fish was compared between groups (where a sufficient n was available) by using 1-way ANOVA combined with Tukey’s test.

In tables and figures, significance levels are denoted by * for p=0.05, ** for p=0.01 and *** for p=0.001.

Results

Hydrogen peroxide pre-treatment was found to elevate the mortality of subsequent immersion exposure to *F. psychrophilum* in the stationary phase of growth.

Experiment 1: H₂O₂ as Pre-treatment Stressor

An apparent escape response was observed in all groups immediately after exposure to H₂O₂; the majority of fish clumped together and swam towards the tank wall. This was most prominent in tanks treated with medium and high concentrations, where the escape response was followed by fish laying on bottom of the tank. Furthermore, a few fish from the high dose group showed erratic swimming behavior for shorter periods of time. All visual changes in behavior ceased after a few min at the lowest concentration, while persisting longer in groups treated with higher concentrations of H₂O₂. All fish survived the treatment, although one death occurred in the high dosage group within the first 24 h.

Experiment 2:200 mg L⁻¹ H₂O₂ Combined with Bacterial Growth Phases (24 and 48 h)

The experiment included four groups in duplicates: (1) 24 h culture, (2) 24 h culture with H₂O₂ pre-treatment, (3) 48 h culture and (4) 48 h culture with H₂O₂ pre-treatment (Table 2).

Nine of the 40 fish pre-treated with H₂O₂ died during treatment and were accounted for as acute mortality. It was only

Table 2. H₂O₂ combined with bacterial growth phases.

Group	H ₂ O ₂	F.p (CFU mL ⁻¹)	n	Re-isolation/ dead fish	% mortality
1	-	24 h	20	0/3	15%
2	+	(10 ⁵)	12	3/3	25%
3	-	48 h	20	0/1	5%
4	+	(10 ⁷)	19	3/4	21%

Experiment 2. The symbols ‘+’ and ‘-’ designate addition or absence of the two factors, H₂O₂ (200 mg L⁻¹ for 60 min) and *F.p.* (10⁵/10⁷ CFU mL⁻¹ *F. psychrophilum* for 30 min). Successful reisolations of *F. psychrophilum* from dead fish are given as a fraction of the total number of dead fish. Finally, cumulative mortality is stated in percent. doi:10.1371/journal.pone.0062590.t002

possible to re-isolate *F. psychrophilum* from dead fish belonging to groups pre-treated with H₂O₂.

Experiment 3:150 mg L⁻¹ H₂O₂ as Experimental Stressor

The experiment consisted of four groups in duplicates: (I) untreated control, (II) H₂O₂, (III) *F. psychrophilum* and (IV) H₂O₂ and *F. psychrophilum* (Table 3).

Seven of the 200 fish pre-treated with 150 mg H₂O₂ L⁻¹ died during the treatment or subsequent transfer to tap water and were accounted for as acute mortality. Seven days after challenge, technical problems lead to an increased temperature in the room and flooding of tank III.B. This led to the loss of 23 fish, which were excluded from the experiment, since the event took place before onset of mortality.

The cumulative mortality (Table 3) was higher in both infected groups compared to their respective controls (p=0.05 for III and p=0.001 for IV). The cumulative mortality for infected groups was 9.1% for *F. psychrophilum* challenge alone and 19.2% in combination with H₂O₂ (p=0.001). Relative standard deviation for the two groups was 23% and 39%, respectively. Furthermore, H₂O₂ accelerated the onset of mortality post challenge from 9 to 3 d (Fig. 1).

Experiment 4:200 mg L⁻¹ H₂O₂ as Experimental Stressor

The experiment consisted of four groups, which were set up in either duplicates for uninfected or triplicates for infected groups: (A) untreated control and (B) H₂O₂, (C) *F. psychrophilum* and (D) H₂O₂ and *F. psychrophilum* (Table 4).

Five of the 250 fish pre-treated with 200 mg H₂O₂ L⁻¹ died during the treatment and subsequent transfer to tap water and were accounted for as acute mortality.

The cumulative mortality was higher in both infected groups compared to their respective controls (p=0.001). The cumulative mortality (Table 4) of infected groups was 14.7% (n=22) for *F. psychrophilum* challenge alone and rose to 30.3% (n=54) in combination with H₂O₂ (p=0.001). RSD for the two groups was 27% and 38%, respectively. Furthermore, H₂O₂ accelerated the onset of mortality post challenge from 10 to 4 d (Fig. 2). A statistically significant difference in mortality was found between replicate tanks D1 (20.0%) and D2 (42.9%) from group D (p=0.05), which was taken into consideration in data processing.

Discussion

Several studies have investigated the potential adverse effects of H₂O₂ on salmonids to establish recommendations for safe treatment dosages [10,11,17–20]. Possible damage caused by

Table 3. Challenge after 150 mg L⁻¹ H₂O₂.

Group	Replicate	H ₂ O ₂	<i>F.p.</i>	n	% mortality (n)	Cumulative % mortality	SD
I	A	-	-	45	2.0% (1)	2.0%	
	B	-	-	47	2.0% (1)	(a*)	
II	A	+	-	49	4.1% (2)	4.0%	
	B	+	-	44	4.4% (2)	(b***)	
III	A	-	+	49	8.0% (4)	9.1%	2.19
	B	-	+	27	11.1% (3)	(a*)(c**)	(23%)
IV	A	+	+	49	14.0% (7)	19.2%	7.42
	B	+	+	48	24.5% (12)	(b***)(c**)	(39%)

Experiment 3. The fish weighed 1.2 g and the temperature was 16.8±0.5°C. The symbols '+' and '-' designate addition or absence of the two factors, H₂O₂ (150 mg L⁻¹ for 60 min) and *F.p.* (10⁷ CFU mL⁻¹ *F. psychrophilum* for 30 min). The % cumulative mortality is stated for both replicates and groups along with SD and RSD in brackets. Statistical differences are denoted by a letter and * for p=0.05, ** for p=0.01 and *** for p=0.001. doi:10.1371/journal.pone.0062590.t003

H₂O₂ depends on several factors, including the applied concentration, exposure time, frequency of treatment, life stage of the fish and temperature during treatment. Damage generally consists of injury to the gills and larger fish are more susceptible [19]. Pathological changes may include an increase in epithelial cell granularity, edemas, lamellar fusion, epithelial hyperplasia as well as swelling and lifting of the gill epithelium. The gills are a frequent target in stress responses and of the mentioned lesions can, with the existing knowledge, be induced by many types of environmental stressor; not just H₂O₂ [23]. Treatment is generally discouraged at temperatures above 14–15°C [11,21,22], but temperatures approaching 14°C during H₂O₂ treatment did not result in increased mortality under the conditions applied in the present experiments.

Mortality due to H₂O₂ has been shown to occur within the first days after treatment; predominantly during or within hours of exposure [17,18]. It is difficult to directly compare the published

studies regarding H₂O₂, and the performance of pilot studies on small groups of fish are recommended before treating an entire population [18,19].

H₂O₂ as Pre-treatment Stressor

Previous studies have demonstrated that H₂O₂ has a fast short-term stress effect in both Atlantic salmon and sea bass [24,25]. In the present study, an escape response was observed in all fish treated with H₂O₂, although most prominent in the high dose group, in which the only mortality occurred within 24 h after exposure. Based on these results, a concentration of 200 mg H₂O₂ L⁻¹ was chosen as stressor for experiment 2, where 9 of the 40 treated fish died from the pre-treatment alone. The increased mortality is likely due to the direct administration of undiluted 35% H₂O₂ into the water. Since the density of fish was higher in experiment 2, the probability of contact with high local concentrations of H₂O₂ before dispersal in the water was




Figure 1. Challenge after 150 mg L⁻¹ H₂O₂. Cumulative mortality of the merged replicates is shown in the figure. The fish weighed 1.2 g and the temperature was 16.8±0.5°C. The experiment ran for 50 days. ◇: control, △: H₂O₂, □: *F. psychrophilum*, ○: H₂O₂+ *F. psychrophilum*. doi:10.1371/journal.pone.0062590.g001

Table 4. Challenge after 200 mg L⁻¹ H₂O₂.

Group	Replicate	H ₂ O ₂	F.p.	n	% mortality (n)	Cumulative % mortality	SD	
A	1	-	-	50	0.0% (0)	3%		
	2	-	-	50	6.0% (3)			(a ^{***})
B	1	+	-	50	8.0% (4)	5%		
	2	+	-	50	2.0% (1)			(b ^{***})
C	1	-	+	50	12.0% (5)	14.7%	4.19	
	2	-	+	50	14.3% (7)			(a ^{***})(c ^{***})
	3	-	+	50	20.0% (10)			
D	1	+	+	50	20.0% (10)	30.3%	11.57	
	2	+	+	49	42.9% (21)			(b ^{***})(c ^{***})
	3	+	+	46	28.3% (13)			

Experiment 4. The fish weighed 1.1 g and the temperature was 14±0.5°C. The symbols '+' and '-' designate addition or absence of the two factors, F.p. (10⁷ CFU mL⁻¹ *F. psychrophilum* for 30 min) and H₂O₂ (200 mg L⁻¹ for 60 min). The % cumulative mortality is stated for both replicates and groups along with SD and RSD. Statistical differences are denoted by a letter and * for p=0.05, ** for p=0.01 and *** for p=0.001. doi:10.1371/journal.pone.0062590.t004

increased. In experiment 3 and 4, H₂O₂ was diluted at in least 300 mL water before being added to the treatment tanks. This approach resulted in 2–3.5% acute mortality in addition to 4–5% mortality during the experiments, compared to a total of 2–3% in the untreated control groups. In a previous test, healthy juvenile rainbow trout were treated twice a week for seven weeks (200 mg L⁻¹ for 60 min) reported lower feed conversion ratios in the first weeks of treatment but no noteworthy mortality [26], while the same dosage given only once resulted in high mortalities within a few hours under slightly different conditions [17]. The experimental design was not intended to investigate the potential long-term consequences of H₂O₂ treatment, but the results indicate the possibility of excess mortality due to as little as a single treatment but the subject should be investigated further.

H₂O₂ Combined with Bacterial Growth Phases (24 and 48 h)

Experiment 2 was a pilot test used to indicate, whether there was a difference in mortality for fish infected with bacteria in either the logarithmic or stationary phase of growth. Furthermore, it was relevant to conduct a small scale test of H₂O₂ in combination with *F. psychrophilum*. Based on previous results, an increased mortality was expected for the 24 h culture [8], but no differences were seen. Reisolation of *F. psychrophilum* from spleen, kidney and brain of dead fish was only successful in H₂O₂ treated groups. Although the portals of entry have not been determined, *F. psychrophilum* has been found in mucus, fins, gills and stomach of infected fish and it has been speculated, that sub-optimal environmental conditions may allow the bacterium to get across skin and gills [27]. Besides




Figure 2. Challenge after 200 mg L⁻¹ H₂O₂. Cumulative mortality of the merged replicates is shown in the figure. The fish weighed 1.1 g and the temperature was 14±0.5°C. The experiment ran for 40 days. ◇: control, △: H₂O₂, □: *F. psychrophilum*, ○: H₂O₂+ *F. psychrophilum*. doi:10.1371/journal.pone.0062590.g002

stressing the host, treatment with H₂O₂ may have resulted in better access to the blood stream, possibly via the gills or by damaging mucus on either skin or gastrointestinal tract, hence allowing for a rapid spread inside the host.

Based on these findings, it was decided to proceed using H₂O₂ as a stressor and the 48 h culture. The 48 h culture has been used successfully in previous investigations [2] and resulted in a higher number of CFU. Cumulative mortality was relatively low for all treatments in experiment 2. The small size of the experimental groups could have affected the outcome in several ways and the results of this experiment cannot be considered to be conclusive. Firstly, the dynamics between healthy, infected and dead fish were influenced by the low density. Secondly, the low number of fish also decreases the statistical robustness, especially since reproducibility is a known problem regarding immersion challenge with *F. psychrophilum*. Choice of strain might have played a role, although 950106-1/1, which was used in the present study, is known to cause mortality in rainbow trout fry [2].

H₂O₂ as Experimental Stressor in Immersion Challenge

The cumulative mortality was significantly increased by pre-treatment with H₂O₂ in both experiment 3 and 4. In experiment 3, pre-treatment with H₂O₂ increased mortality due to *F. psychrophilum* from 9.1% to 19.2%. In experiment 4, the temperature was lowered and the pre-treatment dosage increased slightly, resulting in mortality rates of 14.7% for *F. psychrophilum* alone and 30.3% in combination with H₂O₂. The increased dosage of H₂O₂ may have stressed the fish more or resulted in more damage to gills or other tissues in contact with the surrounding water, such as gastrointestinal tract or skin. The lowered temperature is also likely to have played a role, since even smaller changes have been shown to have significant consequences on survival of infections [27]. At lower temperatures, the immune response of the fish is delayed [28,29], while the psychrophilic bacteria's physiological functions are impaired to a lesser degree [1]. Another explanation is a suppression of the immune system due to cortisol, which is a corticosteroid associated with stress [23]. Finally, batch differences can also have contributed to the difference in mortality seen between the two experiments.

Variation was consistent around 40% for the pre-treated and infected group, and 25% for the infected group for both experiment 3 and 4. Thus, variation did not seem to be influenced by smaller differences in handling through the experiments, since strictly uniformed management was applied in experiment 4. Both variation and proportional change in mortality induced by H₂O₂ was comparable between the two experiments and to the results obtained in a similar model using formalin [2]. Despite the variation, H₂O₂ pre-treatment consistently elevated the mortality after subsequent immersion exposure to *F. psychrophilum*. Conversely, the results underline the importance of maximizing the statistical power of the experimental model by using larger experimental groups or more replicates.

References

1. Nematollahi A, Decostere A, Pasmans F, Haesebrouck F (2003) *Flavobacterium psychrophilum* infections in salmonid fish. *Journal of Fish Diseases* 26: 563–574.
2. Madsen L, Dalsgaard I (1999) Reproducible methods for experimental infection with *Flavobacterium psychrophilum* in rainbow trout *Oncorhynchus mykiss*. *Diseases of Aquatic Organisms* 36: 169–176.
3. Borg AF (1960) Studies on myxobacteria associated with diseases in salmonid fishes. *American Association for the Advancement of Science* 8: 1–85.
4. Madetoja J, Nyman P, Wiklund T (2000) *Flavobacterium psychrophilum*, invasion into and shedding by rainbow trout *Oncorhynchus mykiss*. *Diseases of Aquatic Organisms* 43: 27–38.
5. Busch S, Dalsgaard I, Buchmann K (2003) Concomitant exposure of rainbow trout fry to *Gyrodactylus dejavini* and *Flavobacterium psychrophilum*: Effects on infection and mortality of host. *Veterinary Parasitology* 117: 117–122.
6. Holt RA (1988) *Cytophaga psychrophila*, the causative agent of bacterial cold-water disease in salmonid fish. PhD thesis, Oregon State University, Corvallis.
7. Rangdale RE (1995) Studies on rainbow trout fry syndrome (RTFS). PhD thesis, University of Stirling, Stirling.
8. Aoki M, Kondo M, Kawai K, Oshima S (2005) Experimental bath infection with *Flavobacterium psychrophilum*, inducing typical signs of rainbow trout *Oncorhynchus mykiss* fry syndrome. *Diseases of Aquatic Organisms* 67: 73–79.

Onset of mortality was accelerated, a pattern which was also observed for H₂O₂ treatment in combination with *T. maritimum* and *F. columnaris* [12,13]. The clinical signs in both experiment 3 and 4 corresponded to the symptoms seen in natural outbreaks and included anemia (resulting in pale gills and organs), splenomegaly (which could in some cases be seen through the skin as a red coloration of the abdomen) and dark pigmentation of the skin [1,30,31].

A number of previous studies have also focused on developing a reliable model for infection, which mimics natural transmission, since it is essential to gain more knowledge regarding transmission and host-pathogen interactions. Previous studies have highlighted the limited pathogenicity of *F. psychrophilum* without using various forms of stress or scarification. A number of studies have not resulted in mortality [4,5,32,33], while infections have been successfully established in other studies [2,7,8,34]. Even when mortality occurred, reproducibility has been a problem [35].

It is hard to directly compare the various studies due to differences in setup; including origin, size and health status of the fish, differences in bacterial strains and growth conditions, method of exposure and diversity in the experimental setup, including the number replicates and animals in each group. Since it is difficult to induce *F. psychrophilum* mortality without using stressors or compromising the outer barriers, it seems unlikely that immersion challenge will be standardized on the same level as seen with injections. Although mortality did not reach 60%, which is the desired goal for testing the potency of fish vaccines [36], the current model seems reproducible, since mortality was elevated in both full scale experiments.

In conclusion, the investigated model seems to be a good alternative to injections for studies requiring natural transmission of the pathogen without bypassing the outer barriers. Although the present study does not deal with the cause for the increased mortality induced by H₂O₂, it emphasizes and allows further investigation of the potential connection between routine non-medical treatments and pathogen outbreaks in aquaculture.

Acknowledgments

The technical assistance provided by Lene Gertman has been greatly appreciated as well as advice given regarding statistics by both Henrik Spliid and Claes Eno. Thanks to Kurt Buchmann and Per Walther Kania from the University of Copenhagen, Department of Veterinary Disease Biology, for allowing the use of their experimental facilities during the first three experiments. Finally, comments and suggestions given by Rune Lindgreen were also greatly appreciated.

Author Contributions

Proofreading: LM ID. Conceived and designed the experiments: MH LM ID. Performed the experiments: MH. Analyzed the data: MH. Contributed reagents/materials/analysis tools: LM ID. Wrote the paper: MH.

9. Jokumsen A, Svendsen LM (2010) Farming of freshwater rainbow trout in denmark. Charlottenlund. National Institute of Aquatic Resources, Technical University of Denmark. DTU Aqua report no.219–2010.
10. Pedersen LF (2010) Investigations of environmental benign aquaculture therapeutics replacing formalin (in Danish). DTU Aqua, Charlottenlund. DTU Aqua report, no.218–2010.
11. Sortkær O, Bovbjerg P, Steinfeldt SJ, Bruun MS, Dalsgaard I, et al. (2000) Undersøgelse af eventuelle miljøpåvirkninger ved anvendelse af hjælpestoffer og medicin i ferskvandsdambrug samt metoder til at reducere/eliminere sådanne påvirkninger [in Danish]. DFU report 79–00: 206.
12. Avendaño-Herrera R, Magariños B, Irgang R, Toranzo AE (2006) Use of hydrogen peroxide against the fish pathogen *Tenacibaculum maritimum* and its effect on infected turbot (*Scophthalmus maximus*). *Aquaculture* 257: 104–110.
13. Thomas-Jinu S, Goodwin AE (2004) Acute columnaris infection in channel catfish, *Ictalurus punctatus* (Rafinesque): Efficacy of practical treatments for warmwater aquaculture ponds. *Journal of Fish Diseases* 27: 23–28.
14. Toyama T, Kita-Tsukamoto K, Wakabayashi H (1994) Identification of *Cytophaga psychrophila* by PCR targeted 16S ribosomal RNA. *Fish Pathology* 29: 271–275.
15. Wiklund T, Madsen L, Bruun MS, Dalsgaard I (2000) Detection of *Flavobacterium psychrophilum* from fish tissue and water samples by PCR amplification. *Journal of Applied Microbiology* 88: 299–307.
16. R Development Core Team (2011) R: A language and environment for statistical computing, reference index version 2.13.0. R Foundation for Statistical Computing, Vienna.
17. Tort MJ, Jennings-Bashore C, Wilson D, Wooster GA, Bowser PR (2002) Assessing the effects of increasing hydrogen peroxide dosage on rainbow trout gills utilizing a digitized scoring methodology. *Journal of Aquatic Animal Health* 14: 95–103.
18. Gaikowski MP, Rach JJ, Ramsay RT (1999) Acute toxicity of hydrogen peroxide treatments to selected lifestages of cold-, cool-, and warmwater fish. *Aquaculture* 178: 191–207.
19. Rach JJ, Schreier TM, Howe GE, Redman SD (1997) Effect of species, life stage, and water temperature on the toxicity of hydrogen peroxide to fish. *Progressive Fish-Culturist* 59: 41–46.
20. Arndt RE, Wagner EJ (1997) The toxicity of hydrogen peroxide to rainbow trout *Oncorhynchus mykiss* and cutthroat trout *Oncorhynchus clarki* fry and fingerlings. *Journal of the World Aquaculture Society* 28: 150–157.
21. Harper C, Wolf JC (2009) Morphologic effects of the stress response in fish. *Ilar Journal* 50: 387–396.
22. Kiemer MCB, Black KD (1997) The effects of hydrogen peroxide on the gill tissues of atlantic salmon, *Salmo salar* L. *Aquaculture* 153: 181–189.
23. Bruno DW, Raynard RS (1994) Studies on the use of hydrogen peroxide as a method for the control of sea lice on atlantic salmon. *Aquaculture International* 2: 10–18.
24. Bowers JM, Speare DJ, Burka JF (2002) The effects of hydrogen peroxide on the stress response of Atlantic salmon (*salmo salar*). *J Journal of Veterinary Pharmacology and Therapeutics* 25: 311–313.
25. Roque A, Yildiz HY, Carazo I, Duncan N (2010) Physiological stress responses of sea bass (*Dicentrarchus labrax*) to hydrogen peroxide (H₂O₂) exposure. *Aquaculture* 304: 104–107.
26. Speare DJ, Arsenaault GJ (1997) Effects of intermittent hydrogen peroxide exposure on growth and columnaris disease prevention of juvenile rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences* 54: 2653–2658.
27. Nematollahi A, Decostere A, Pasmans F, Haesebrouck F (2003) *Flavobacterium psychrophilum* infections in salmonid fish. *Journal of Fish Disease* 26: 563–574.
28. Covert JB, Reynolds WW (1977) Survival value of fever in fish. *Nature* 267: 43–45.
29. Raida MK, Buchmann K (2007) Temperature-dependent expression of immune-relevant genes in rainbow trout following *Yersinia ruckeri* vaccination. *Diseases of Aquatic Organisms* 77: 41–52.
30. Gräns A, Rosengren M, Niklasson L, Axelsson M (2012) Behavioural fever boosts the inflammatory response in rainbow trout *Oncorhynchus mykiss*. *Journal of Fish Biology* 81: 1111–1117.
31. Lorenzen E, Dalsgaard I, From J, Hansen EM, Hørlyck V, et al. (1991) Preliminary investigations of fry mortality syndrome in rainbow trout. *Bulletin of the European Association of Fish Pathologists* 11: 77–79.
32. Bruno DW (1992) *Cytophaga psychrophila* (= '*Flexibacter psychrophilus*') (borg), histopathology associated with mortalities among farmed rainbow trout, *Oncorhynchus mykiss* (Walbaum) in the UK. *Eur Assoc Fish Pathol* 12: 215–216.
33. Decostere A, Lammens M, Haesebrouck F (2000) Difficulties in experimental infection studies with *Flavobacterium psychrophilum* in rainbow trout (*Oncorhynchus mykiss*) using immersion, oral and anal challenges. *Research in Veterinary Science* 69: 165–169.
34. Chua FHC (1991) A study on the rainbow trout fry syndrome. MSc thesis, Institute of Aquaculture, University of Stirling, Stirling.
35. Garcia C, Pozet F, Michel C (2000) Standardization of experimental infection with *Flavobacterium psychrophilum*, the agent of rainbow trout *Oncorhynchus mykiss* fry syndrome. *Diseases of Aquatic Organisms* 42: 191–197.
36. Amend DF (1981) Potency testing of fish vaccines. *Developments in Biological Standardization* 49: 447–454.