



RESEARCH ARTICLE

REVISED Continuous Storage Root Formation and Bulking in Sweetpotato [version 4; peer review: 2 approved, 1 not approved]

Astere Bararyenya ^{id}1, Phinehas Tukamuhabwa¹, Paul Gibson ^{id}1, Wolfgang Grüneberg², Reuben Ssali ^{id}3, Jan Low⁴, Thomas Odong¹, Mildred Ochwo-Ssemakula ^{id}1, Herbert Talwana¹, Natasha Mwila¹, Robert Mwangi ^{id}3

¹Department of Agricultural Production, College of Agricultural and Environmental Sciences, Makerere University, Kampala, Central Uganda, Box 7062, Uganda

²Crop Improvement, International Potato Center (CIP), Avenida La Molina 1895, Apartado 1558, Lima 12, Peru

³Crop Improvement, International Potato Center (CIP), Kampala, Central Uganda, Box 22274, Uganda

⁴Economics, International Potato Center (CIP), Nairobi, Nairobi, ILRI Campus Naivasha Rd, 25171-00603 Lavington, Kenya

v4 **First published:** 11 Feb 2019, 3:83
<https://doi.org/10.12688/gatesopenres.12895.1>
Second version: 14 Oct 2019, 3:83
<https://doi.org/10.12688/gatesopenres.12895.2>
Third version: 08 Apr 2020, 3:83
<https://doi.org/10.12688/gatesopenres.12895.3>
Latest published: 09 Jul 2020, 3:83
<https://doi.org/10.12688/gatesopenres.12895.4>

Abstract

This study investigated the phenotypic variation of continuous storage root formation and bulking (CSRFAB) growth patterns underlying the development of sweetpotato genotypes for identification of potential varieties adapted to piecemeal harvesting for small scale farmers. The research was conducted between September 2016 and August 2017 in Uganda. Genotypes from two distinct sweetpotato genepool populations (Population Uganda A and Population Uganda B) comprising 130 genotypes, previously separated using 31 simple sequence repeat (SSR) markers were used. Measurements (4 harvest times with 4 plants each) were repeated on genotypes in a randomized complete block design with 2 replications in 2 locations for 2 seasons. We developed a scoring scale of 1 to 9 and used it to compare growth changes between consecutive harvests. Data analysis was done using residual or restricted maximum likelihood (REML). Data showed a non-linear growth pattern within and between locations, seasons, and genotypes for most traits. Some genotypes displayed early initiation and increase of bulking, while others showed late initiation. Broad sense heritability of CSRFAB was low due to large GxE interactions but higher in other traits probably due to high genetic influence and the effectiveness of the methodology. A high level of reproducibility (89%) was observed comparing 2016B and 2017A seasons (A and B are first and second season, respectively) at the National Crops Resources

Open Peer Review

Reviewer Status ✗ ✓ ✓

	Invited Reviewers		
	1	2	3
version 4 (revision) 09 Jul 2020			
version 3 (revision) 08 Apr 2020			✓ report
version 2 (revision) 14 Oct 2019			↑ ? report
version 1 11 Feb 2019	✗ report	✓ report	

1 **Arthur Villordon** ^{id}, LSU Agcenter Sweet Potato Research Station, Chase, USA

Research Institute (NaCRRI), Namulonge, Uganda. Choosing CSRFB genotypes can more than double the sweetpotato production (average maximum yield of 13.1 t/ha for discontinuous storage root formation and bulking (DSRFAB) versus 28.6 t/ha for CSRFB, demonstrating the importance of this underresearched component of storage root yield.

Keywords

Sweetpotato, yield, continuous storage root formation and bulking, growth pattern, phenotypic variation

2 **Hussein Shimelis**, University of KwaZulu-Natal (UKZN), Pietermaritzburg, South Africa

3 **Alfonso del Rio** , University of Wisconsin-Madison, Sturgeon Bay, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Astere Bararyenya (barastere@gmail.com)

Author roles: **Bararyenya A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Tukamuhabwa P:** Conceptualization, Investigation, Methodology, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Gibson P:** Data Curation, Formal Analysis, Methodology, Validation, Writing – Review & Editing; **Grüneberg W:** Data Curation, Methodology, Writing – Review & Editing; **Ssali R:** Formal Analysis, Methodology, Writing – Review & Editing; **Low J:** Funding Acquisition, Project Administration, Supervision, Writing – Review & Editing; **Odong T:** Formal Analysis, Methodology, Writing – Review & Editing; **Ochwo-Ssemakula M:** Methodology, Supervision, Writing – Review & Editing; **Talwana H:** Supervision, Writing – Review & Editing; **Mwila N:** Formal Analysis, Writing – Review & Editing; **Mwanga R:** Conceptualization, Investigation, Methodology, Project Administration, Supervision, Validation, Visualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Bill and Melinda Gates Foundation [OPP1019987] as part of the SASHA II project, and the CGIAR Research Program on Roots, Tubers and Bananas (RTB) under the CGIAR Fund Donors (<http://www.cgiar.org/about-us/our-funders/>). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2020 Bararyenya A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bararyenya A, Tukamuhabwa P, Gibson P *et al.* **Continuous Storage Root Formation and Bulking in Sweetpotato [version 4; peer review: 2 approved, 1 not approved]** Gates Open Research 2020, 3:83 <https://doi.org/10.12688/gatesopenres.12895.4>

First published: 11 Feb 2019, 3:83 <https://doi.org/10.12688/gatesopenres.12895.1>

REVISED Amendments from Version 3

We thank Dr. Alfonso del Rio for critically reviewing our manuscript and for providing very useful comments and suggestions for improving the quality of the manuscript to reach a broad audience. We are in agreement with his suggestions for improvement. We have to the extent possible revised the manuscript as recommended and a detailed report of the responses is provided. The main differences between this version and the previous version are:

We have reduced the content by one table and one figure and reorganized some portions of the manuscript in a simpler way and improved the general flow of the information to enable the broad audience to follow the content of the article.

The abstract has been revised and the information synthesized in a simpler way.

In the experimental design, the form of planting material has been clarified.

In the Results and discussion section, information was synthesized as recommended by Dr. Alfonso del Rio.

We hope that the modifications and shortening of the text to improve the flow and improve clarity have improved the quality of the manuscript.

Any further responses from the reviewers can be found at the end of the article

Introduction

Sweetpotato (*Ipomoea batatas* (L.) Lam, family Convolvulaceae) is one of the most important food crops worldwide, with approximately 106 million tons produced in almost 120 countries from an area of about 8 million ha and an average global yield of 11.1 tons/ha (FAO, 2016). Asia is the world's largest sweetpotato producing continent, with 79 million tons, followed by Africa (FAOstat, 2016). About 75% of this global production is from China alone. A total of 21.3 million tons is produced in Africa, with 48% from the Great Lakes region. In East Africa, the crop is the second most important root crop after cassava and has played an important role as a famine-relief crop during its long history and has recently been reevaluated as a health-promoting

food (Low *et al.*, 2017). Uganda ranks as the fourth largest sweetpotato producer in the world after China, Nigeria and Tanzania, with a production of 2.1 million t. In Africa, Uganda is ranked third after Nigeria and Tanzania. Sweetpotato is one of the main staple crops in the food systems of Uganda, Rwanda, and Burundi with a per capita consumption of 50.9, 80.1 and 57.0 kg, respectively (Table 1).

On average and across the 30-year period, the population across East Africa increased by two-and-half-fold while sweetpotato production increased by two-fold. The increase in production was due to a combination of factors varying in different countries but mainly due to increase in area and breeding efforts. This trend resulted in a decrease in per capita production from 86 to 56 kg per person. Statistics in general underestimate production in most annual plants since not all crop production is recorded. Usually, only the main planting season is recorded even though crops are grown over two to three growing seasons per year (Bararyanya *et al.*, 2018a). The most noticeable production increase took place in Tanzania with more than ten-fold increase of tonnage following an almost two-fold increase in growing area and almost four-fold increase in yields. The highest productivity is recorded in Kenya, with more than 17 t/ha/yr, followed by Tanzania (12.5 t/ha/yr). In other East African countries, productivity has more or less stagnated. The yield increase in the region over the 30-year period can be attributed to breeding and release of improved varieties by national breeding programs, and to the introduction of new varieties mainly, by the International Potato Center (CIP). The increase in yield can also be explained by the importance and recent interest in sweetpotato as a food and nutritious crop compared to the early 1990s when it was hardly known.

Despite the giant strides made through breeding and release of 27 high yielding and disease resistant varieties (Mwanga *et al.*, 2016), Uganda has over the last 30 years consistently reported extremely low yields of 4.5 t/ha (Table 1) compared to the achievable yields of over 40 t/ha under improved conditions. The low yield could be attributed partly to the growing of low

Table 1. Increase in sweetpotato production in East and Central African countries, production data for 1985 and 30 years later for 2015.

Country	Area		Yield		Production per year		Population		Per annum	
	x 1000 ha		tons per ha		x 1000 tons		millions		kg per capita	
	1985	2015	1985	2015	1985	2015	1985	2015	1985	2015
Burundi	87.0	58.6	6.4	9.9	555.0	580.9	4.7	10.2	118.0	57.0
Kenya	50.0	72.2	9.9	17.1	488.0	1232.3	19.7	47.2	24.8	26.1
Rwanda	135.9	139.7	7.1	6.7	979.5	931.0	6.1	11.6	160.1	80.1
Uganda	358.7	454.5	4.6	4.5	1664.0	2045.2	14.7	40.1	113.6	50.9
Tanzania	290.0	746.6	4.6	12.5	303.0	3454.5	21.8	53.9	13.9	64.1
Average	184.3	294.3	6.5	10.1	797.9	1648.8	13.4	32.6	86.1	55.6
Increase		1.60		1.06		2.07		2.44		30.45

Source: FAOstat, 2016

yielding and highly disease susceptible landraces by small-scale farmers. [Abidin \(2004\)](#) reported high farmer preference for their landraces, while [Sseruwu \(2012\)](#) associated this preference with lack of farmer desired attributes in the newly released varieties. In Uganda and other East African countries, piecemeal harvesting, characterized by repeated harvesting from the same sweetpotato plants on a mound, ridge or other seedbed over time, is the predominant mode of harvesting among subsistence and commercial sweetpotato farmers. Importantly, this practice is also known in other root and tuber crops including potato, cassava and yam. Recently, [Tadesse *et al.* \(2017\)](#) reported that piecemeal harvesting was the dominant practice among poor potato farmers (60% of respondents) in Ethiopia, whereas the majority of the wealthy and medium-wealthy farmers combined piecemeal harvesting with harvesting all at once. Farmers harvest enough for one or few meal(s), or enough for one market. This is because these crops are very difficult to store, and storage for fresh sweetpotato produce is virtually non-existent. Therefore, the practice allows for in-ground storability and market partitioning during the cropping season. It also creates room for new storage roots to initiate and enlarge for the next harvest. However, breeding and selection have been based on a single harvest and no breeding research has tried to understand the genetics underlying traits associated with these common practices in sweetpotato farming systems. Continuous storage root formation and bulking CSRFAB in sweetpotato is associated with the perennial nature of the crop and allows for longer photosynthetic activity resulting from persistent canopy development leading to increased plant productivity. This is supported in the case of practices in Uganda by the fact that more than 90% ($n = 350$) of the farmers have no knowledge of the maturity periods of the local sweetpotato varieties they grow ([Bashaasha *et al.*, 1995](#)).

Maturity periods in sweetpotato vary with genotype and the environmental conditions under which they are grown. Common external signs of maturity such as senescing and yellowing of leaves do not apply to all varieties. In addition, storage roots do not all mature at the same time, while the storage root formation period is highly variable among genotypes ([Belehu, 2003](#); [Belehu *et al.*, 2004](#); [Lowe & Wilson, 1974](#); [Wilson & Lowe, 1973](#); [Yanfu *et al.*, 1989](#)). It is reported that the onset of storage root initiation can occur as early as 7 to 13 days after planting (DAP) ([Du Plooy, 1989](#)), and the total storage root number to form varies from 30 to 112 DAP depending on the genotype and the environmental conditions under which they are grown ([Yanfu *et al.*, 1989](#)). Furthermore, sweetpotato storage roots can undergo periods of arrested growth during unfavorable conditions and then continue active growth upon favorable conditions ([Ravi *et al.*, 2009](#)). There are many reported studies on the effects of storage root formation and bulking under controlled conditions on sweetpotato yields ([Gajanayake & Reddy, 2016](#); [Meyers *et al.*, 2013](#); [Solis *et al.*, 2014](#); [Villordon, 2015](#)) and under field experiments ([Agata, 1982](#); [Belehu *et al.*, 2004](#); [Du Plooy, 1989](#); [Enyi, 1977](#); [Gajanayake *et al.*, 2014](#); [Lowe & Wilson, 1974](#);

[Wilson & Lowe, 1973](#); [Wilson, 1982](#); [Yanfu *et al.*, 1989](#)). However, most of the studies were undertaken on single season harvest basis, and no study attempted to understand the CSRFAB patterns overtime under field conditions to focus on improving the trait through breeding.

Knowledge of the growth patterns of storage root formation and bulking traits is critical for crop yield improvement, crop management, and specifically for timing fertilizer application and irrigation. Sweetpotato has been for centuries selected for its starchy roots on seasonal basis and may have been progressively losing its perennial feature. It is well known in other root and tuber crops that yields are a result of the rate and duration of tuberization, which in turn depends on longevity of the leaves, the beginning of storage root formation, and duration of the growth cycle. In almost all experiments, the end time of storage root formation was not defined or properly assessed; number of storage roots was infrequently recorded and maximum number of storage roots was never established. The effect of longer vegetative maintenance periods of green leaves observed in some genotypes has never been investigated in sweetpotato, but it is reported to influence greater productivity in potato ([Çalışkan *et al.*, 2007](#)). Despite these deficiencies, the storage root formation and bulking patterns are widely regarded as a key developmental stage in the crop's life, having profound implications for subsequent growth and development. It was therefore hypothesized that longevity of green foliage due to genetic properties in CSRFAB sweetpotato genotypes is greater than in discontinuous storage root formation and bulking (DSRFAB) genotypes. Consequently, the extended period of green leaves for CSRFAB genotypes will impact storage root formation and bulking leading to a significant increase in yield. This expected variation in yield and yield components is mainly due to longer duration of photosynthetic activity and the great availability of photo-synthesizing material, mostly its green leaves ([Paltridge *et al.*, 1984](#)). Thus, the amount of change in the mean value of expected responses associated with a unit increase in growth time, holding all other variables constant, varies with increased growth time periods of some sweetpotato cultivars and produces higher amounts of storage root number and weight. To really understand the evolution of a trait, we need to know whether any variability in that trait can be assigned to genetic effects. If so, and if there is fitness variation associated with the trait; it will be subject to natural selection ([Croston *et al.*, 2015](#)). This study investigated genetic variability of CSRFAB and, characterized growth patterns at different development stages to identify possible CSRFAB sweetpotato genotypes in the germplasm collection in Uganda for use as parents in improvement of the trait.

Methods

Plant materials and experimental sites

Plant materials. This study utilized 130 genotypes currently used for population improvement in Uganda for various breeding purposes and were screened for CSRFAB traits ([Table 2](#)).

Table 2. List of genotypes screened for continuous storage root formation and bulking in Uganda, 2016 to 2017.

Code	Name	Code	Name	Code	Name	Code	Name	Code	Name
A1	Carrot C	A27	Apa352	B3	Hma496	B29	Mkn1210	B55	K-566632
A2	Ejumula	A28	Luw1274	B4	Msd380	B30	NASPOT 5	B56	New Kawogo
A3	Mayai	A29	Sponge	B5	Luw1230	B31	Kre723	B57	Bitambi
A4	Naspot5/58	A30	NASPOT 7	B6	Srt43	B32	Ara236	B58	Kbl611
A5	Kbl619	A31	NASPOT 10 O	B7	Srt28	B33	SPK004(CIP)	B59	Mkn171
A6	Kmi61	A32	Kml872	B8	Mle199	B34	Mpg1158	B60	Mle191
A7	Kbl631	A33	Msk1040	B9	Mary	B35	Jonathan	B61	Iga998
A8	Pal133	A34	Dimbuka-Bukulula	B10	Kml942	B36	Kml960	B62	Pal134
A9	Pal94 silk	A35	Oguoilwe	B11	Lir302	B37	Bsh741	B63	Wt-237
A10	Rak786	A36	Kmi159	B12	Kbl650	B38	Bnd145l	B64	Iga994
A11	Mpg1128	A37	Mkn1168	B13	Mpg1151	B39	Srt41	B65	Naveto
A12	Tanzania	A38	Apa335	B14	Mkn1180	B40	Srt01	B66	Xushu18
A13	Kala	A39	Apa323	B15	Luw1257	B41	Wagabolige	B67	Yanshu1
A14	Kbl648	A40	Otada	B16	Kre691	B42	Lir258	B68	Zambezi
A15	Kml956	A41	Ukerewe	B17	Mbr536	B43	Pal148	B69	Jewel
A16	Bsh740	A42	Kmi88	B18	Ksr662	B44	Rak865	B70	Caromex
A17	Silk(1254)	A43	Mpg1148	B19	Mle179	B45	Mpg1122	B71	Resisto cip
A18	Rak819	A44	Iga983	B20	Msd431	B46	Msk1079	B72	Baeuregard
A19	SPK004	A45	Pal108	B21	Mle163	B47	Rak848	B73	Kyabafuruki
A20	Nk259l	A46	Rak835	B22	Magabali	B48	Mugande	B74	Tainung 64
A21	Kml881	A47	Msd384	B23	Kbl618	B49	Ara224	B75	Kre696
A22	NASPOT 9 O	A48	Srt27	B24	Mbr552	B50	Dlp3163	B76	Raihna
A23	Tororo 3	A49	Nk318l	B25	Iga978	B51	Msk1094	B77	Tis9265
A24	NASPOT 1	A50	Mle194	B26	Sowola	B52	Tis9101	B78	199062.1
A25	Rak808	B1	Resisto	B27	Hma490	B53	NASPOT 3	B79	Santa amaro
A26	Mpg1146	B2	Ara209	B28	Apa356	B54	NK1081L	B80	Huarmeyano

The 130 genotypes included two distinct sweetpotato gene pool populations (Uganda A and Uganda B) that were formed to reflect similarity within and divergence between the populations based on 31 simple sequence repeat (SSR) markers (David *et al.*, 2018). The genotypes are maintained by the International Potato Center (CIP) at the National Crops Resources Research Institute (NaCRRI) at Namulonge in Uganda.

Experimental sites and duration of the study. Grüneberg *et al.* (2005) identified Uganda as environmentally diverse and suitable for sweetpotato selection for East African countries, and a breeding platform where crossings and selection in early and later breeding stages were conducted. The results from such environments are extended to all East African countries with one or two more season evaluations for confirmation. Two locations, NaCRRI-Namulonge and National Semi-Arid Resources Research Institute (NaSARRI-Serere) were therefore selected to host the trials for screening potential CSRFAB genotypes adapted to the East African agroecologies. The trials were planted on September 22, 2016 and September 29, 2016, respectively at Namulonge and

Serere and harvested from January through April 2017 for the first season (2016B). In the second season (2017A), trials were planted on March 10, 2017 and March 29, 2017, respectively, at Namulonge and Serere and harvested from June to September 2017. The altitude of the sites was around 1,150 meters above sea level with an average day temperature of 22.2°C. Crops were not irrigated and often suffered from low rainfall 4 months after planting. Namulonge is characterized by a tropical rain forest zone with a bimodal rainfall average of 1,270 mm annually and high sweetpotato virus disease (SPVD) pressure, while Serere is in the tall grassland savanna zone with low rainfall and high weevil population (Table 3).

Experimental design

Vine cuttings of approximately 30 cm of 3 to 4 nodes each were planted; 20 cuttings of each of the 130 sweetpotato genotypes were planted in a randomized complete block design in 2 locations (Namulonge and Serere) for 2 seasons, 2016B and 2017A (A and B are first and second season, respectively). Each genotype in a plot of five plants had four replications,

Table 3. Description of study sites used for screening sweetpotato genotypes for continuous storage root formation and bulking in Uganda.

Location	Elevation (masl)	Temperature (°C)	Rainfall (mm)	Soil			
				pH	OM	K	P
					(%)	Cmol/kg	Cmol/kg
NaCRRRI (2016B)	1150	24–30	1400–1600	4.8	3.8	0.11	Trace
NaSARRI (2016B)	1140	25–32	1100–1300	4.9	3.7	0.19	Trace
NaCRRRI (2017A)	1150	22–29	1550	5.2	3.4	0.76	1.66
NaSARRI (2017A)	1140	23–31	1150–1350	4.9	3.6	0.56	0.92

Note. NaCRRRI: National Crops Resources Research Institute; 2016B: 2016 second season; NaSARRI: National Semi-Arid Resources Research Institute; 2017A: 2017 first season; masl: meters above sea level; °C: degrees celcius; mm: millimeters; OM: organic matter; K: potassium; P: phosphorus

but for ease of data collection and analysis, the samples were collected as two replications. Plant density was 1 m between rows and 0.3 m between plants within the row. The genotypes were sequentially and destructively harvested at 3, 4, 5 and 6 months after planting (MAP) to allow identification of storage root formation and bulking patterns following the repeated measures design (Causton, 1994; Hesser, 2015). During each harvest period, four plants were uprooted for above and below ground part data collection. Each row was bordered by two sweetpotato plants at the extreme ends. The plots were kept weed free and no fertilizer or other agro-chemicals were applied.

Data collection methods

Growth and development measurements. Traits that are known to characterize growth in sweetpotato storage roots were selected. These included: (i) number of harvested plants (NPH), allowing calculation of average values, (ii) total storage root number (SRN), (iii) Total root weight (TRW) measured with a round spring balance scale (Hanson, 8 in x 8 in x 3 in), was used to calculate yield as tons per hectare, (iv) commercial and non-commercial storage root number (CRN & NCRN, respectively) and commercial and noncommercial storage root weights (CRW & NCRW, respectively), allowing the estimation of storage root formation, bulking rate, (v) vine weight (VW), (vi) root system weight (RSW) used to calculate biomass yield (BMY), and (vii) harvest index (HI). Senescence (SEN) was estimated using a scale of 1 to 9, where 1 = no senescence, 9 = severe senescence marked by death/drying (CIP/AVRDC/IPGR, 1991).

Development of a scale for continuous storage root formation and bulking (CSRFB). To capture the possible overtime initiation and bulking in sweetpotato, we developed a rating scale of 1 to 9. The scale rates the expression of CSRFB trait for an individual genotype. CSRFB scores were therefore developed throughout the 2016B growing season. Detailed observation of storage root formation and bulking over 6 months provided comprehensive knowledge of the development of a scale for measuring changes in the CSRFB trait. The scale was set to measure the changes in root formation and bulking of a given individual genotype. The change in scores would reflect the potential of roots to bulk into storage roots (fleshy or lignified),

newly initiated storage roots, their status of bulking and their maturity levels (see Figure 1, scoring scale).

Data analysis

Growth model analysis. Growth models were estimated using a multilevel, mixed model framework described by Payne (2009) in GenStat 18th Edition. Each of the 130 genotypes was maximally compared (4160 times) with the other genotypes in a model with two replications, four harvest times, two seasons and two locations.

To better understand the details in the storage root development patterns across the four harvest times, we analyzed piecewise, storage root growth patterns in different phases of the trial and selected individual plants to represent different growth pattern characteristics. Means were plotted to visualize the growth patterns using Genstat 18th edition.

Restricted maximum likelihood (REML) variance components analysis of the phenotypic data was performed using the following general linear mixed model;

$$Y_{ijklcqm} = \mu + S_i + L_j + SL_{ij} + SLR_{ijk} + H_l^1 + H_q^2 + H_c^3 + SH_{il} + LH_{ij} + SLH_{ijl} + E_m + EH_{lm}^1 + EH_{qm}^2 + EH_{cm}^3 + SE_{im} + LE_{jm} + SLE_{ijm} + \epsilon_{iml} + LEH_{jml} + SLEH_{ijml} + \epsilon_{ijklcqm}$$

Where

Y_{ijklm} is the observed overtime response of genotypes across location and season.

μ , is the overall mean, S_i is the effect of the i th season, L_j is the effect of the j th location, SLR_{ijk} is the effect of the interaction between the i th season in the j th location and the k th replication, HT_l^1 is the effect of linear term l in polynomial model, HT_q^2 is the effect of the quadratic term q in polynomial model, HT_c^3 is the random effect of the cubic term in the polynomial model, SH_{il} is the interaction between season and harvest time, LH_{ij} is the random effect of interaction between j th location and i th harvest time, SLH_{ijl} is the effect of the interaction between the i th season and the j th location at the l th harvest time, E_m is the effect of the m th genotype, EH_{lm}^1 is the effect of interaction between m th genotype and l th linear term, EH_{qm}^2 is the effect of




Figure 1. A 1 to 9 scoring scale for continuous storage root formation and bulking in which score: 1 = no visual detectable storage root initiation (SRI) and no visually detectable bulking; 2 = Storage root formation and bulking detectable; 3 = no visually SRI and has unclear levels of bulking; 4 = distinct SRI and 2 clear levels of bulking; 5 = distinct SRI and 3 clear levels of bulking; 6 = distinct SRI and 4 clear levels of bulking; 7 = distinct SRI and 5 clear levels of bulking; 8 = distinct SRI and 6 clear levels of bulking; and 9 = distinct SRI and 7 clear levels of bulking.

the interaction between *i*th genotype and *q*th quadratic term in the polynomial model, EH_{cm}^3 is the effect of interaction between *i*th genotype and *c*th cubic term in the polynomial model, SE_{im} is the effect of interaction between season and genotype, LE_{jm} is the effect of interaction between location and genotype, SLE_{ijm} is the random effects of the interaction between season, location and genotype. ϵSEH_{iml} is the effect of the interaction between season, harvest time and genotypes, LEH_{jml} is the random effect of the interaction between location, harvest time and harvest time, $SLEH_{ijml}$ is the interaction season, location and harvest time, $\epsilon_{ijklcqm}$ is the error associated with all factors involved in the polynomial model.

The F-test denominator for genotypes tested in different locations and seasons (Bararyenya *et al.*, 2018b) was calculated following Satterthwaite’s formula for the “composite” F-test denominator for genotypes:

$$Den MS = MS(GL) + MS(GS) + MS(GSL)$$

$$Dendf = \frac{[MS(GL) + MS(GS) + MS(GSL)]^2}{\frac{MS(GL)^2}{df(GL)} + \frac{MS(GS)^2}{df(GS)} + \frac{MS(GSL)^2}{df(GSL)}}$$

giving Satterthwaite-type approximation of mean squares (MS) and degrees of freedom (df), where MS = mean square,

G = genotype, S = season, L = location, df = degree of freedom, Den df = denominator degrees of freedom (Satterthwaite, 1946; Snedecor & Cochran, 1967).

Results and discussion

For growth pattern analysis and phenotypic variability across selected sweetpotato traits, we used a linear mixed model to decompose phenotypic variance (P) into different components: genetic (G) and environmental (E) sources, and their interaction effects (GxE). The interactions between genotype, location and season were significant as were most of the three-way interactions, implying that the growth curve must be optimized for the three factors together, because they modify each other’s effect. Thus, the interaction between season, location and replication (SLR) was highly significant across all the parameters. This is because the replication differences were large in some environments as previously observed by Tumwegamire *et al.* (2016). The variability in replication for an experiment involving clonal crops can easily occur. The main effects for season and location were not significant, as were their interactions. The interaction between season, location and harvest time (SLH) was highly significant (P<0.001) for all the parameters in this study, resulting in non-significance of SL, SH and LH interactions. This implies that harvest time effect differs depending on the level of the location and season on storage root yield (SRY), CSRFAB, storage root diameter (SRD), storage root length (SRL), CRW, CRN, HI and SEN (Table 4).

Table 4. Mean squares and F-test of significance for continuous storage root formation and bulking (CSRFAB), storage root yield (SRY) (tons/ha), storage root number (SRNO), vine yield (VY) (tons/ha), storage root diameter (SRD) (mm), storage root length (SRL) (mm), commercial root number (CRN) and weight (CRW), harvest index (HI) and senescence (SEN) of 130 sweetpotato genotypes from Uganda.

SOV	d.f.	CSRFAB	SRY	SRNO	VY	SRD	SRL	CRW	CRN	HI	SEN
S	1	1223.56 ^{ns}	2263.89 ^{ns}	15284.37 ^{ns}	73832.70 ^{ns}	2821.05 ^{ns}	89273.20 ^{ns}	125.34 ^{ns}	4381.50 ^{ns}	383.10 ^{ns}	21.90 ^{ns}
L	1	243.40 ^{ns}	415.52 ^{ns}	1609.44 ^{ns}	77317.50 ^{ns}	1246.97 ^{ns}	16297.00 ^{ns}	42.99 ^{ns}	0.024 ^{ns}	22744.50 ^{ns}	7.57 ^{ns}
SL	1	0.037 ^{ns}	906.99 ^{ns}	249.05 ^{ns}	28398.30*	4147.13 ^{ns}	9217.60 ^{ns}	92.14*	281.46 ^{ns}	1855.40 ^{ns}	9.24 ^{ns}
SLR	4	13.14***	595.92***	147.08***	3278.10***	1021.48***	4815.60***	7.57***	42.20***	1235.5***	1.65***
HLin	1	134.56 ^{ns}	5039.56 ^{ns}	661.59 ^{ns}	3801.60 ^{ns}	276.73 ^{ns}	2692.20 ^{ns}	176.72*	107.95 ^{ns}	22840.60 ^{ns}	0.01 ^{ns}
HQ	1	0.037 ^{ns}	310.48 ^{ns}	43.89 ^{ns}	1670.50 ^{ns}	52.76 ^{ns}	34.80 ^{ns}	14.44 ^{ns}	45.44 ^{ns}	60.00 ^{ns}	6.35 ^{ns}
HC	1	33.212 ^{ns}	270.43 ^{ns}	35.27 ^{ns}	204.70 ^{ns}	0.22 ^{ns}	1170.50 ^{ns}	7.39 ^{ns}	4.39 ^{ns}	4655.80 ^{ns}	8.29 ^{ns}
SH	3	21.58 ^{ns}	172.93 ^{ns}	203.05 ^{ns}	13015.70 ^{ns}	100.56 ^{ns}	1998.90 ^{ns}	13.28 ^{ns}	11.99 ^{ns}	5152.80 ^{ns}	6.14 ^{ns}
LH	3	61.12 ^{ns}	1113.99 ^{ns}	1062.79 ^{ns}	1937.90 ^{ns}	4904.45 ^{ns}	34388.40 ^{ns}	3.83 ^{ns}	847.82 ^{ns}	7276.10 ^{ns}	3.55 ^{ns}
SLH	3	27.55***	242.29**	561.79***	1626.00**	728.56***	6705.80***	17.01***	134.57***	111.8***	2.90***
E	129	12.26***	262.85 ^{ns}	246.97***	1370.60*	503.38***	3801.90***	4.23***	103.32***	2569.60***	2.13***
HLinE	129	2.73***	71.90 ^{ns}	39.94***	613.30***	96.35*	522.20**	1.24*	19.72***	269.70 ^{ns}	1.13 ^{ns}
HQ.E	129	0.99 ^{ns}	38.08 ^{ns}	11.87 ^{ns}	215.40 ^{ns}	39.75 ^{ns}	141.40 ^{ns}	0.81 ^{ns}	7.32 ^{ns}	154.80 ^{ns}	0.46 ^{ns}
HC.E	129	1.40 ^{ns}	46.53 ^{ns}	17.32 ^{ns}	329.70 ^{ns}	38.37 ^{ns}	192.40 ^{ns}	0.69 ^{ns}	7.78 ^{ns}	183.10 ^{ns}	0.41 ^{ns}
SE	129	4.98***	175.22 ^{ns}	78.07***	1089.30 ^{ns}	194.50**	1451.00*	1.94**	36.73**	1115.90***	1.35*
L.E	128	4.02*	177.72 ^{ns}	81.34***	988.80 ^{ns}	189.38**	1263.80 ^{ns}	1.95**	33.26*	1009.30***	0.97***
SLE	126	2.90***	119.02 ^{ns}	45.69***	995.00***	112.38***	1029.10***	1.25***	24.18***	807.70***	0.89***
SHE	385	1.24 ^{ns}	50.31*	15.50 ^{ns}	399.80 ^{ns}	54.21 ^{ns}	289.10 ^{ns}	0.74 ^{ns}	7.79 ^{ns}	202.40 ^{ns}	0.77***
LHE	377	1.19 ^{ns}	57.48*	17.24 ^{ns}	367.00 ^{ns}	72.13 ^{ns}	364.60*	0.93 ^{ns}	8.67 ^{ns}	254.50***	0.50 ^{ns}
SLHE	331	1.44 ^{ns}	52.67 ^{ns}	19.74*	363.80 ^{ns}	59.75 ^{ns}	295.50 ^{ns}	0.82***	7.84 ^{ns}	177.40 ^{ns}	0.31 ^{ns}
Residual	1983	1.36	51.05	16.32	412.50	52.54	339.60	0.45	7.35	193.4 ^{ns}	0.46

Note: SOV = source of variation; df: degrees of freedom; CSRFAB = continuous storage root formation and bulking (scored on a scale of 1 to 9, where 1 = no storage initiation and no bulking, and 9 = high storage root initiation & bulking); SRY = storage root yield; SRNO = storage root number per plant; VY = vine yield; SRD = storage root diameter; SRL = storage root length; CRW = commercial storage root weight; CRN = commercial storage root number; HI = harvest index; SEN: senescence (scored on a scale of 1 to 9, where 1 = no senescence, 9 = severe senescence, and death/drying); S = Season; L = location; H = harvest time; E = genotype; HLin: Harvest time linear; HQ = harvest time squared; HC = harvest time cubic (ie. lack-of-fit); SL = season by location interaction; SLR = season by location by replication interaction; SH = season by harvest time interaction; LH = location by harvest time interaction; SLH = season by location by harvest time interaction; SE = season by genotype interaction; L.E = location by genotype interaction; SLE = season by location by genotype interaction; SHE = season by harvest time by genotype interaction; LHE = location by harvest time by genotype interaction; SLHE = season by location by harvest time by genotype; * = significant at 0.05; ** = significant at 0.01; *** = significant at 0.001; ^{ns} = non-significant.

These results require further investigation to identify the level of influence (Table 4). For all the traits in this study, the interaction between season, location and genotype (SLE) was highly significant ($P < 0.001$). This indicates that there was wide variability of genotype across locations and seasons and this wide variability can be used for sweetpotato yield improvement in a specific location. However, there is a need to study the stability across locations and seasons for easy selection of the CSRFAB trait.

The derived overall means (Table 5) suggest the existence of determinate and extended growth maturity stages among genotypes. High variation ($P < 0.001$) of storage root formation (root initiation) and bulking was observed using the 1 to 9 scoring scale for CSRFAB. This suggests that the scale can be used to differentiate and evaluate the CSRFAB trait among sweetpotato genotypes.

Variance component analysis and heritability estimates of selected sweetpotato growth traits

Variance component analysis and heritability estimates of selected sweetpotato growth traits in a nonlinear model structure. The genotypic variance among the nine traits associated with CSRFAB varied from 0.9 (CRW) to 7.7 (SRNO). Genotypic variance was high for SRNO (7.7), CRN (5.0) and CRW (5.0) (Table 5). The phenotypic variance was high for VY (449.8), SRL (378.5) and SRY (84.6) and varied from 2.4 (CSRFAB) to 449.8 (VY). The residual variance was extremely high for VY (412.5) and SRL (339.6). These two particular traits are influenced by continuous growth. While some genotypes continue to increase in biomass weight, others die by senescence, likewise for vine length. The error and the GxE variances for CSRFAB are not big compared to its genetic variance. This implies that the scale used to measure the trait is precise and the trait is not influenced much by the environment. Heritability was relatively

high for SRNO (67.2%) and CSRFAB (50.5%). The low residual variance and high heritability observed for CSRFAB confirmed the precision of the method used to study the CSRFAB trait in sweetpotato.

Genetic effects on yield of CSRFAB genotypes in sweetpotato

We performed a harvest basis analysis and calculated the corresponding heritability for each trait to understand the

dynamics of breeding values across harvest times. We targeted the traits that can affect the final yield. Thus, the genotypic variances among seven main traits associated with CSRFAB varied from 0 to 72.46 (Table 6). Genotypic variance was high for HI at 120 DAP (72.46) and VY at 180 DAP (59.91) (Table 6). Zero variance was recorded for VY (120 DAP), HI (150 DAP), SEN (180 DAP) and weevils (180 DAP). The zero (or negative) variance components could be due to an artifact of the optimization algorithm that includes a non-negative constraint; a negative variance component could also represent competition effects between adjacent plots. These results imply that there is no significant change in VY, HI, SEN and weevil damage within the population for the respective growth periods. Since heritability is influenced by genetic variance, it was directly affected at the same traits and harvest periods, (ranging from 0 to 67%). The 0% heritability implies that the effect of the traits is moving to fixation, that is, its frequency in the population is very high. This is because polyploidy crops are dominated by large mutations with small effects and the genetic fixation pattern is affected by interactions between cosegregating alleles (Kopp & Hermisson, 2009). If we score the effect of the allele(s) regulating the trait having no heritability in the population, the result would mean in genetic terms, that the allele frequency in the population is 100%, therefore the genetic variance at the loci of these genes is zero, so any variance in the corresponding phenotypic traits cannot be attributed to the non-existent genetic variance. These results need further investigation especially in the case of weevil resistance mechanisms as no significant increase in weevil infestation was observed in late harvest (Bararyenya et al., 2018b). The rate in storage root bulking resulting in the available high green biomass may compete with weevil infestation. Broad sense heritability was relatively high for CSRFAB (50.5%) compared to other yield component traits indicating a better yield prediction using the scale.

We found temporal dynamics of genotypic influence on overall trait development (Table 6). In the early growth phase, genotypic variance was mostly low. As plants grew, genotypic

Table 5. Variance components, within and across environment heritability estimation for nine characters associated with continuous storage root formation and bulking in sweetpotato.

Trait	σ^2_R	σ^2_{SLG}	σ^2_{LG}	σ^2_{SG}	σ^2_G	H1	H2	H(%)
CSRFAB	1.4	0.2	0.1	0.1	0.2	68.2	53.7	50.5
SRY	51.1	8.8	3.8	3.7	0.9	77.2	30.9	11.1
SRNO	16.3	3.9	2.4	2.2	7.7	74.9	53.2	67.2
VY	412.5	8.8	3.8	3.7	4.7	68.2	24.9	19.9
SRD	52.5	9.2	4	3.8	4.9	67.2	63.1	38.3
SRL	339.6	9.2	4	3.8	4.9	73.5	60.8	22.5
CRW	0.4	9.4	4.1	3.9	5	69.9	69.4	44
CRN	7.4	9.4	4.1	3.9	5	73.1	56.8	43.2
SEN	0.5	9	3.9	3.7	4.8	46.7	58.0	44

Note. H: broad sense heritability across location and season; H1: within environment broad sense heritability across 2 seasons for NaCRR1; H2: within environment broad sense heritability across 2 seasons for NaSARR1; σ^2_R : residual variance; σ^2_G : genotypic variance; σ^2_P : phenotypic variance; σ^2_{SLG} : variance due to season by location by genotype interaction; σ^2_{LG} : variance due to location by genotype interaction; σ^2_{SG} : variance due to season by genotype interaction; σ^2_{GE} : genetic by environmental variance; SRNO: storage root number per plant; SRY: storage root yield; VY: vine yield; SRDIA: storage root diameter; SRLG: storage root length; CRW: commercial root weight; CRN: commercial root number; VW: vine weight; CSRFAB: continuous storage root formation and bulking; SEN: senescence

Table 6. Estimates of genotypic (σ^2g) variance and broad sense heritability (H) within harvest time across environments for seven traits associated with continuous storage root formation and bulking.

Trait	Genetic variance (σ^2g)					Broad sense heritability H (%)				
	90DAP	120DAP	150DAP	180DAP	Comb.	90DAP	120DAP	150DAP	180DAP	Comb.
SRNO	3.29	10.08	6.46	8.06	8.84	47	67	62	58	70
SRY	2.84	2.32	4.04	0.81	2.43	26	20	23	4	24
VY	14.53	0.00	4.57	59.91	26.50	19	00	6	35	46
HI	29.80	72.46	0.00	36.25	75.26	43	61	0	29	61
SEN	0.01	0.02	0.07	0.00	0.07	12	19	22	0	56
WEEVIL	0.02	0.00	0.24	0.00	0.08	31	0	13	0	45
CSRFAB	0.23	0.51	0.10	0.43	0.45	35	62	49	55	69

Note. σ^2g : Genetic variance; DAP: days after planting; Comb.: combined estimate; SRNO: storage root number; SRY: storage root yield; VY: vine yield; HI: harvest index; CSRFAB: continuous storage root formation and bulking; SEN: senescence.

factors became in general more important. The increasing genetic effect was observed up to about 120 DAP and decreased thereafter. After 120 DAP, the genetic effect became relatively less important. This can be partly explained by the drought stress which became more important 4 months after planting (MAP). Although less obvious, the opposite pattern was seen in the growth recovery phase (after 5 MAP), likely due to growth resumption resulting in the decline in overall phenotypic differences between CSRFB and DCSRFB plants. The investigated highly correlated traits with CSRFB showed dynamic changes in heritability during the entire plant growth stage (Table 6). SRNO and CSRFB showed similar patterns of heritability over time. We found that heritability of SRNO and CSRFB increased in early growth stages and then decreased during drought stress, occurring between two consecutive rainy seasons and then increased thereafter during the growth recovery period occurring with the onset of the rainy season. These results are supported by Tuberosa (2012); quantitative traits reflecting the performance of crops under drought conditions tend to have low to modest heritability (Table 6).

Identification of storage formation and bulking growth patterns and their characterization

We have proposed in this study a method that can be used to estimate changes (Figure 2 to Figure 5) in the CSRFB trait and therefore estimate the potential maximum yield of a sweetpotato genotype. The scoring method developed in this study effectively measured changes in the CSRFB trait. This is supported by its low residual variance (1.4) compared to SRN (16.3) and SRY

(51.1) (Table 4) and its high broad sense heritability of 50.5% compared with heritability of SRN (67.2%) and yield (11.1%) (Table 5). Two types of growth patterns were identified according to the lifetime of sweetpotato genotypes. The first type which has determinate storage root formation and bulking is characterized by a rapid and short vegetative growth period, followed by a senescence period until the leaves die off. This period is also characterized by a quick maximum yield obtained about 90 to 120 DAP (Figure 2) for most DCSRFB genotypes. Respective CSRFB scores for the same genotypes declined overtime due to lack of new root formation and similar level of bulking for mature storage roots (Figure 3). Changes between successive harvest times were generally negative for DCSRFB genotypes. The cumulative yield changes, as were the cumulative score changes of DCSRFB genotypes were therefore lower than the respective maximum single harvest yields and the rate of CSRFB decreases in late stages. The second type of growth is characterized by a prolonged vegetative growth and a late switch-over time to reproductive phase (Figure 3 and Figure 5). The cumulative yield changes over time, as were the cumulative score changes, of CSRFB genotypes displayed similar results with the last harvest (180 DAP) confirming the accuracy of the scale to predict yield changes over time. The change responses were positive for CSRFB genotypes and the cumulative responses increased overtime (Figure 4) for yield and the increase was not significant (Figure 5) for the CSRFB scores suggesting a possibility of predicting CSRFB genotypes responses from early growth stages. During this period the yield increase starts later and increases drastically after 150 DAP (Figure 4) for




Figure 2. Growth pattern for discontinuous storage root formation and bulking (DCSRFB) of four sweetpotato genotypes representing different growth patterns over four harvest times (averaged across four environments). Growth trend for genotypes A3, A25, A43 and B72 showing a fully DCSRFB trait which generally increases the yield up to 120 days after planting (DAP) and then decreases overtime. The cumulative yield is generally lower than the maximum yield due to the yield decreasing in late stages. Note: DAP = days after planting, Cum. = cumulative yield.




Figure 3. Growth trend of scores over four harvest times for discontinuous storage root formation and bulking genotypes A3, A25, A43 and B72 decreases due to progressive reduction of newly formed storage roots and similar bulking across all storage roots of the same genotype. Note: DAP = days after planting; CSRFB = Continuous storage root formation and bulking; Cum. = Cumulative scores. DCSRFB = discontinuous storage root formation and harvesting. Trend of scores over four harvest times for DCSRFB genotypes A3, A25, A43 and B72 decreases due to progressive reduction of newly formed storage roots and similar bulking across all storage roots of the same genotype. The cumulative scores are generally lower than the respective maximum scores due to the rate of CSRFB decreasing in late stages.




Figure 4. Growth pattern of continuous storage root formation and bulking of four sweetpotato genotypes representing different growth patterns over four harvest times (averaged across four environments). Note: DAP = days after planting; Cum. = Cumulative yield. Growth trend for genotypes A30, A14, B1 and B80 shows CSRFB with increased yield overtime. The yield increase starts slowly and drastically increases at a late stage (150 DAP). The cumulative yield is the same with the last harvest (180 DAP).

most CSRFB genotypes. Respective CSRFB scores of the same genotypes did not change overtime due to continuous root formation and bulking (Figure 5) suggesting the ability of the scoring scale to predict CSRFB genotypes at early growth stages. This period increased drastically yields of CSRFB genotypes, however, the maximum yields were not observed in this trial due to continued vegetative growth. According to Paltridge & Denholm (1974) the variation in yield observed may be due

to the variation in their actual rates of dry matter production and the individual switchover time of the genotype to raise its maximum yield, referred to as the optimum growth pattern. The optimum growth pattern for determinate sweetpotato genotypes is of a two-phase plant growth, with the switchover occurring at the instant the plant becomes limited by restriction of its access to light (senescence). Subsequent values of the growing cycle which give maximum yield per unit time of harvest occur




Figure 5. Growth pattern for continuous storage root formation and bulking (CSRFB) of four sweetpotato genotypes representing different growth patterns over four harvest times (averaged across four environments). Note: DAP = days after planting; CSRFB = Continuous storage root formation and bulking; Cum. = Cumulative scores. Scores show a trend which is similar across the four harvest times suggesting a possibility of predicting CSRFB genotypes from early stages upwards. Genotypes A30, B1, B14 and B80 exhibit continuous bulking over the four harvest times. Cumulative scores were the same with 180DAP confirming the precision of the scale.

after these periods for indeterminate growth which are functions of above ground vegetative (leaves) lifetime. Each successive maximum is larger than the last, so one might expect the plants to evolve longer lifetimes and correspondingly longer periods of indeterminate (CSRFB) growth. The intensity of senescence strongly varied ($P < 0.001$) among genotypes and was negatively correlated with CSRFB (Table 7). The yield rate of change between the 120 DAP and the 150 DAP was always negative. For the CSRFB (Figure 3 & Figure 5), all the mean predictions and changes were positive. Yield increased up to more than 20-fold and the increase was genotype dependent.

These results are in agreement with the basic growth curves in many crop plants (Schurr *et al.*, 2006; Yanfu *et al.*, 1989). The rapid growth in the first stage for determinate or DSRFB genotypes could be attributed to early maturing genotype properties in which there is more energy invested in biomass production for early remobilization for storage root bulking and this plant growth expression is also reported in other annual crops (Wenk & Falster, 2015). However, the maximum yield in this period was low and was limited by the available amount of green biomass which affected the bulking rate. Paltridge *et al.* (1984) reported similar common characteristics in which many plants have a rather sharp transition between the vegetative and the reproductive stages of growth. If the time of switch over is small, the amount of green leaf would be small and the subsequent rate of production and final yield would be also low. In this phase, the time of switchover was before 90 DAP. Sweetpotato continues to grow and branch if environmental conditions are

favorable, due to its perennial habit, but the leaves formed earlier in the growing season start to fall and the total number of leaves and leaf area decrease toward the end of the growing season (Somda & Kays, 1990). Determinate or DSRFB genotypes lost their ability to initiate new shoots at around 120 DAP and the VW decreased drastically. These results agree with findings of Paltridge & Denholm (1974) and Bhagsari (1990) in which most annuals exhibit a single reproductive phase, often with a sudden onset.

The senescence phenotype was less observed in indeterminate types, although greenness intensity was reduced at 120 DAP which resumed and drastically increased (high positive quadratic slope) from 150 DAP when rains came back. Bhagsari (1990) reported similar results that sweetpotato cultivars maintained leaf area index to intercept a major portion of sunlight until harvest, and leaf area growth significantly differed depending on cultivar.

The slow increase in most of the yield component parameters in CSRFB genotypes resulted in low storage root yield in the first growth phases (Figure 3 & Figure 5). This can be explained by the fact that indeterminate genotypes invest resources for maintenance at earlier stages including root elongation for nutrition purposes and biomass production to maintain vegetative growth. This resulted in strong vegetative biomass increase for CSRFB genotypes. For these genotypes the time of switchover to reproduction was very late (around 150 DAP). This was the last growth phase where yield increased drastically in

Table 7. Yield means of 10 most distinct discontinuous (DSRFAB) and continuous (CSRFAB) genotypes at different harvest times showing overall best yielding genotypes in combined environments.

DSRFAB genotypes					CSRFAB genotypes				
Genotype	90DAP	120DAP	150DAP	180DAP	Genotype	90DAP	120DAP	150DAP	180DAP
MPG1146	8.23	10.49	17.46	13.17	SPK004	6.47	4.43	10.34	28.64
Ndimbuka	7.18	3.47	6.29	5.83	Kala	4.18	6.72	7.41	24.26
NASPOT 1	6.56	3.19	7.63	3.99	BSH740	5.31	5.67	8.15	18.62
Otada	6.49	3.56	9.03	4.08	KML872	4.31	6.59	12.16	21.44
MPG1128	6.11	6.41	17.33	5.29	NASPOT 9 O	3.52	8.83	4.17	10.37
MSK1040	5.97	9.12	11.97	11.83	Mayai	9.69	11.6	3.63	9.72
RAK786	5.13	2.8	6.36	2.96	Ukerewe	5.47	5.1	5.5	11.59
KMI88	5.03	6.4	7.91	8.52	NASPOT 7	2.68	6.98	7.89	13.88
KBL648	4.6	9.54	8.95	9.33	APA352	1.12	2.95	2.95	8.87
Jonathan	4.6	7.62	17.36	9.54	RAK819	2.92	8.3	5.19	11
Mean	6.29	4.48	7.35	4.87	Mean	3.54	5.95	7.09	9.83
Max	16.71	12.98	17.6	13.17	Max	9.82	15.45	19.57	28.64
Min	1.24	0.1	0.97	0.3	Min	0.21	0.65	0.22	2.41
SED	7.67	6.4	7.43	8.55	SED	7.67	6.4	7.43	8.55

DAP: Days after planting; DSRFAB: discontinuous storage root formation and bulking; CSRFAB: continuous storage formation and bulking; Comb.: combined; Max: maximum; Min: minimum; SED: standard error difference.

CSRFAB corresponding to the switch to reproduction (onset of flower and seed) and storage root remobilization in sweetpotato in which the plant invests its vegetative resources in increasing the sink capacity. Because of high availability of resource/energy in upper biomass, sink strength is increased leading to increased productivity.

For sweetpotato breeders and practitioners, harvest index (HI) is difficult to estimate due to the problem of measuring its components. For instance, the time of harvest influences greatly the value of HI because storage root bulking is likely to vary progressively from the onset of storage bulking and increases with the biomass translocation into storage roots. However, the senescence reduces the above-ground biomass progressively leading to high and unrealistic values of harvesting index. Therefore, HI of sweetpotato varies greatly with the time of harvest (Bararyenya *et al.*, 2018b). This variation is also influenced by other environmental factors including wet or dry conditions during harvest which increase or decrease weight due high or less moisture. Similar results in other root and tuber crops are reviewed by Hay (1995).

Overall, the starting yield was low at 90 DAP (overall mean) and increased progressively as growth time increased (Table 7). This relationship is well explained by the variation in the data (R^2 varied from 38% to 65%). High yielding genotypes showed consistently maximum yields in the population (Bararyenya *et al.*, 2018b) while low yielding genotypes showed inconsistent relationship. The maximum yield recorded across locations and seasons was 16.7 t/ha and 28.6 t/ha at the first

harvest and the last harvest, respectively (Table 7). This implies that the frequency of genotypes that increase yield overtime is high. In this trial, some genotypes reached maximum growth earlier, others late, leading to high variability in growth related parameters.

On average, there was a two-fold increase in yield from the 90 DAP to 180 DAP. The 10 most distinct DSRFAB genotypes were, MPG1146 (A26), Dimbuka-Bukulula (A34), NASPOT 1 (A24), Otada (A40), MPG1128 (A11), MSK1040 (A33), RAK786 (A10), KMI88 (A42), KBL648 (A14) and the 10 most distinct CSRFAB were, SPK004 (A19), Kala (A13), BSH740 (A16), KML872 (A32), NASPOT 10 O (A22), Mayai (A3), Ukerewe (A41), NASPOT 7 (A30), APA352 (A27) and RAK819 (A18). These contrasting genotypes for CSRFAB can be used in breeding programs to study and improve sweetpotato yield in CSRFAB genotypes. The average yield at two locations in two seasons shows that you can more than double sweetpotato production by choosing high yielding genotypes (Table 7) with maximum yield of 13.17 t/ha for DSRFAB versus maximum of 28.64 t/ha for CSRFAB). Similar results have been reported (Çalişkan *et al.*, 2007) in which there were significant differences among the locations, 60 to 70 t/ha of storage root yield being obtained by choosing high-yielding varieties.

Variability and distribution of 130 genotypes over 4 harvest times

For SRY and VY the first two box plots are comparatively equally shorter than the third and fourth box plots (Figure 6). This suggests that, overall, genotypes have little variation over the




Figure 6. Two seasons (2016B & 2017A) boxplot comparison showing overall variability and dispersion of storage root yield (SRY), vine yield (VY) and continuous storage root formation and bulking (CSRFB) over 4 harvesting times (HT) among 130 sweetpotato genotype at the National Crops Resources Research Institute (NaCRRI), Namulonge. Note: SRY = storage root yield; VY = vine yield; HT = harvesting time; 1 = 90DAP; 2 = 120 DAP; 3 = 150 DAP; 4 = 180 DAP.

first two harvest times. It also suggests low variation between time of harvest during this phase. The third and fourth box plots are higher than the first two box plots for SRY and VY. This suggests that genotypes have quite different growth patterns from the two first harvest times and the two last ones. It also suggests high variation among the genotypes. The last two box plots in each sub-figure show obvious differences within each box plot and the two first box plots. This suggests an area of difference that could be explored further. The four sections of the box plots are uneven in size. This shows that many genotypes have a similar growth pattern during the early growth phase, but in a later phase genotypes are more variable in their growth pattern. The long upper whisker shows that genotype growth varied amongst the most positive quartile group, and very similar for the least positive quartile group. This suggests a need for further exploration. The medians of the two first harvest time plots (which generally will be close to the average) are at the same level. However, the box plot of the last two harvest times shows very different distribution means. Variation and distribution of CSRFB scores across location and

seasons consistently remained almost the same (Figure 6 and Figure 7), suggesting an effective response prediction of the scores from the beginning.

Relationship between associated growth traits and CSRFB in sweetpotato

CSRFB scores were highly and positively correlated with most of the yield component traits (Table 8). This implies that CSRFB is also a component of yield. In other words, the higher the CSRFB scores of a sweetpotato genotype, the higher the yield of the genotype. CSRFB scores were negatively correlated with SEN which confirms our hypothesis in which a CSRFB genotype should maintain vegetative growth to continuously provide source/inputs for sink storage roots. The negative correlation with VW needs further investigation, however, the competition of source and sink activities in the plant, effects of pests and diseases such as nematodes and *Alternaria stem blight* (*Alternaria bataticola*) and SPVD could be among the causal factors. It is possible to cease vegetative growth and continue survival for an extended period, but most varieties reduce




Figure 7. Two seasons (2016B & 2017A) boxplot comparison showing overall variability and dispersion of storage root yield (SRY), vine yield (VY) and continuous storage root formation and bulking (CSRFAB) over 4 harvest times (HT) among 130 sweetpotato genotypes at the National Semi-Arid Resources Research Institute (NaSARRI), Serere. Note: SRY = storage root yield; VY = vine yield; HT = harvesting time; 1 = 90DAP; 2 = 120 DAP; 3 = 150 DAP; 4 = 180 DAP.

Table 8. Pearson’s correlation coefficients of CRN, CRW, CSRFAB, HI, SEN, SRD, SRL, SRY, VW and VY for 130 sweetpotato genotypes across two locations (Namulonge and Serere) and two seasons (2016B and 2017A) (N = 4160).

	CRN	CRW	CSRFAB	SEN	SRD	SRL	SRNO	SRY	VY
CRN	-								
CRW	0.84***	-							
CSRFAB	0.67***	0.46***	-						
SEN	0.08	-0.05	0.01	-					
SRD	0.74***	0.70***	0.63***	-0.12	-				
SRL	0.75***	0.68***	0.71***	-0.12	0.86***	-			
SRNO	0.90***	0.65***	0.82***	0.09	0.68***	0.72***	-		
SRY	0.74***	0.74***	0.53***	-0.08	0.80***	0.72***	0.70***	-	
VY	-0.29***	-0.25***	-0.16*	-0.21*	-0.17*	-0.18**	-0.25**	-0.14	-

Note. CRN: commercial storage root number; CRW = commercial storage root weight; CSRFAB = continuous storage root formation and bulking (scored 1–9); HI = harvesting index; SEN = senescence (scored 1 to 9); SRD: storage root diameter; SRL: storage root length; SRN storage root number; SRY = storage root yield; VW = vine weight; VY = vine yield.

their vegetative weight following the storage root bulking peak (Venus & Causton, 1979). The comparison of a classic method of evaluating growth change (analyzing yield mean change between two consecutive harvests) and the developed scale for measuring CSRFB traits produced consistent results with CSRFB scores. This is supported by high heritability and low residual variance observed for CSRFB versus yield and other component parameters. Using the developed scale, 48 genotypes were clustered among CSRFB genotypes, whereas 62 genotypes clustered among DSRFB (Figure 5). These results were reproduced in the following season, and 41% were common to the two methods in 2016B versus 46% in 2017A (Figure 8).

Accuracy of CSRFB scoring method

We characterized in this study growth patterns associated with CSRFB in sweetpotato and compared the accuracy of CSRFB scores and the classic method used to measure growth change overtime (compared yield change overtime) (Figure 8). It was hypothesized that CSRFB genotypes potentially increase yield over time, therefore, the trait can be screened for by measuring yield change overtime. This method, holding other factors constant, shall produce the same results of scoring for the trait using the developed 1 to 9 scale. We investigated this relationship in our data. Overall, the yield change analysis over four harvest times grouped the 110 genotypes (without missing data) into 39 CSRFB and 71 DSRFB genotypes. Using the developed scale, 48 genotypes clustered among the CSRFB genotypes, whereas 62 genotypes clustered together as DSRFB (Figure 8).

Conclusion

This study highlights three important results: 1) the CSRFB trait can be exploited to provide additional yield in sweetpotato 2) the 1 to 9 scale developed provided consistent scores across

replications, and reflected well the growth patterns observable phenotypically, 3) the method of analyzing growth variables over harvest times revealed distinct growth patterns among genotypes. These patterns identify which sweetpotato genotypes are likely to be suited to piecemeal harvesting. The methodology introduced here is expected to be useful in other root crops as well.

This study showed that CSRFB genotypes differentially increased yields up to 779% and discontinuous genotypes reduced yield after crop maturity up to 85% for determinate genotypes. Five months after planting (150 DAP) is proposed as the ideal scoring time for this scale, however, there is need for further work in different agroecologies to validate the reliability of the results. The sweetpotato genotypes used in this study are highly variable for CSRFB and breeding to improve the trait should be feasible due to its high heritability. Genotypes most distinct for CSRFB were, SPK004, Kala, BSH740, KML872, NASPOT 9 O, Mayai, Ukerewe, NASPOT 7, APA352 and RAK819, and those distinct for DCSRFB were, MPG1146, Dimbuka-Bukulula, NASPOT 1, Otada, MPG1128, MSK1040, RAK786, KMI88, and KBL648. The highest CSRFB yielder, SPK004 (28.6 t/ha) outperformed by 15.4 t/ha (117%) the highest DCSRFB yielder, MPG1146 (13.2 t/ha) across locations and seasons. Delaying in harvest leads to final yield losses for DCSRFB while it enhances final yield of CSRFB genotypes. These genotypes can be used in conventional sweetpotato breeding for yield improvement. For sweetpotato breeding, the CSRFB genotypes are recommended for the improvement of sweetpotato varieties suitable for piecemeal harvesting in small scale farming systems, while the discontinuous genotypes are recommended for the development of early maturing sweetpotato varieties. However, the trait needs much more understanding of the physiology of sweetpotato. Therefore, linking these phenotypic patterns with causal genomic variations can provide a clear understanding of selection




Figure 8. Comparative accuracy of screening continuous growth using yield increase overtime evaluation and using a 1 to 9 scale (discontinuous storage root formation and bulking (DSRFAB) and continuous storage root formation and bulking (CSRFB) genotypes). Note. SRY: storage root yield.

scenarios and speed up the breeding for this important trait in sweetpotato.

Data availability

The data underlying this study is available from International Potato Center (CIP) Dataverse.

CIP Dataverse: Dataset 1. Dataset for: Continuous Storage Root Formation and Bulking in Sweetpotato <http://dx.doi.org/10.21223/P3/IC6ZEY> (Bararyenya *et al.*, 2018b)

License: [Attribution 4.0 International \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Acknowledgements

The authors thank the National Agricultural Research Organization (NARO) of Uganda for providing sites for the trials, and CIP and NARO for kindly providing technical support and the sweetpotato genotypes. This study was supported by the CGIAR Research Program on Roots, Tubers and Bananas.

References

- Abidin PE: **Sweetpotato breeding for northeastern Uganda: Farmer varieties, farmer-participatory selection, and stability of performance.** PhD Thesis, Wageningen University, The Netherlands, 2004; ISBN: 90-8504-033-7.
[Reference Source](#)
- Agata W: **The characteristics of dry matter and yield production in sweet potato under field conditions.** In: R L. Villareal & T D. Griggs (eds.). *Sweet potato. Proceeding of the 1st. Int. Symp. AVRDC*, Taiwan, China. 1982.
[Reference Source](#)
- Bararyenya A, Nahayo PC, Nduwimana A, *et al.*: **Assessment of Opportunities for Burundian Small-Scale Potato Farmers to Increase Productivity and Income.** *Potato Research.* 2018a; **61**(1): 73–88.
[Publisher Full Text](#)
- Bararyenya A, Tukamuhabwa P, Gibson P, *et al.*: **Dataset for: Continuous Storage Root Formation and Bulking in Sweetpotato.** International Potato Center, V1, 2018b.
<http://www.doi.org/10.21223/P3/IC6ZEY>
- Bashaasha B, Ewell PT, Mwanga ROM, *et al.*: **Sweetpotato in the farming and food systems of Uganda: A farm survey report.** Lima, Peru: International Potato Center (CIP), National Agricultural Research Organization (NARO). 1995; 63.
[Reference Source](#)
- Belehu T: **Agronomical and physiological factors affecting growth, development and yield of sweet potato in Ethiopia.** (Doctoral dissertation, University of Pretoria). 2003.
[Reference Source](#)
- Belehu T, Hammes PS, Robbertse PJ: **The origin and structure of adventitious roots in sweet potato (*Ipomoea batatas*).** *Aust J Bot.* 2004; **52**(4): 551–558.
[Publisher Full Text](#)
- Bhagsari AS: **Photosynthetic evaluation of sweet potato germplasm.** *J Amer Soc Hort Sci.* 1990; **115**(4): 634–639.
[Publisher Full Text](#)
- Çalışkan ME, Söğüt T, Boydak E, *et al.*: **Growth, yield, and quality of sweet potato (*Ipomoea batatas* (L.) Lam.) cultivars in the southeastern Anatolian and east Mediterranean regions of Turkey.** *Turk J Agric For.* 2007; **31**(4): 213–227.
[Reference Source](#)
- Causton DR: **Plant growth analysis: a note on the variability of unit leaf rate (net assimilation rate) within a sample.** *Ann Bot.* 1994; **74**(5): 513–518.
[Publisher Full Text](#)
- CIP/AVRDC/IPGR, International Potato Center, Asian Vegetable Research and Development Center, International Board for Plant Genetic Resources: **Descriptors for sweetpotato.** In: Z. Huaman (ed.). *Intl. Board for Plant Genetic Resources*, Rome. 1991.
[Reference Source](#)
- Croston R, Branch CL, Kozlovsky DY, *et al.*: **Heritability and the evolution of cognitive traits.** *Behav Ecol.* 2015; **26**(6): 1447–1459.
[Publisher Full Text](#)
- David MC, Diaz FC, Mwanga ROM, *et al.*: **Gene pool subdivision of East African sweetpotato parental material.** *Crop Sci.* 2018; **58**(6): 2302–2314.
[Publisher Full Text](#)
- Du Plooy CP: **Storage root morphogenesis of the sweet potato (*Ipomoea batatas* (L.) Lam.).** Abstr, PhD thesis, University of Pretoria, South Africa. 1989.
[Reference Source](#)
- Enyi BAC: **Analysis of growth and tuber yield in sweet potato (*Ipomoea batatas*) cultivars.** *J Agric Sci.* 1977; **88**(2): 421–430.
[Publisher Full Text](#)
- Food and Agriculture Organization of the United Nations: **FAOSTAT statistics database.** [Rome], 2016; accessed May 2018.
[Reference Source](#)
- Gajanayake B, Reddy KR: **Sweetpotato responses to mid-and late-season soil moisture deficits.** *Crop Sci.* 2016; **56**(4): 1865–1877.
[Publisher Full Text](#)
- Gajanayake B, Raja Reddy K, Shankle MW, *et al.*: **Quantifying storage root initiation, growth, and developmental responses of sweetpotato to early season temperature.** *Agron J.* 2014; **106**(5): 1795–1804.
[Publisher Full Text](#)
- Grüneberg WJ, Manrique K, Zhang D, *et al.*: **Genotypic environment interactions for a diverse set of sweetpotato clones evaluated across varying ecogeographic conditions in Peru.** *Crop Sci.* 2005; **45**(6): 2160–2171.
[Publisher Full Text](#)
- Hay RKM: **Harvest index: a review of its use in plant breeding and crop physiology.** *Ann Appl Biol.* 1995; **126**(1): 197–216.
[Publisher Full Text](#)
- Hesser H: **Modeling individual differences in randomized experiments using growth models: recommendations for design, statistical analysis and reporting of results of internet interventions.** *Internet Interv.* 2015; **2**(2): 110–120.
[Publisher Full Text](#)
- Kopp M, Hermisson J: **The Genetic Basis of Phenotypic Adaptation I: Fixation of Beneficial Mutations in the Moving Optimum Model.** *Genetics.* 2009; **182**(1): 233–249.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lowe SB, Wilson LA: **Comparative Analysis of Tuber Development in Six Sweet Potato (*Ipomoea batatas* (L.) Lam) Cultivars: 1. Tuber initiation, tuber growth and partition of assimilate.** *Ann Bot.* 1974; **38**(2): 307–317.
[Publisher Full Text](#)
- Low JW, Mwanga ROM, Andrade M, *et al.*: **Tackling vitamin A deficiency with biofortified sweetpotato in sub-Saharan Africa.** *Glob Food Sec.* 2017; **14**: 23–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Meyers SL, Arancibia RA, Shankle MW, *et al.*: **Sweet Potato Storage Root Initiation.** Publication: 2809, Publication. 2013.
[Reference Source](#)
- Mwanga RO, Kyalo G, Ssemakula GN, *et al.*: **'NASPOT 12 O' and 'NASPOT 13 O' Sweetpotato.** *HortScience.* 2016; **51**(3): 291–295.
[Publisher Full Text](#)
- Paltridge GW, Denholm JV: **Plant yield and the switch from vegetative to reproductive growth.** *J Theor Biol.* 1974; **44**(1): 23–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Paltridge GW, Denholm JV, Connor DJ: **Determinism, senescence and the yield of plants.** *J Theor Biol.* 1984; **110**(3): 383–398.
[Publisher Full Text](#)
- Payne RW: **GenStat.** *Wiley Interdiscip Rev Comput Stat.* 2009; **1**(2): 255–258.
[Publisher Full Text](#)
- Ravi V, Naskar SK, Makesh Kumar T, *et al.*: **Molecular physiology of storage root formation and development in sweetpotato (*Ipomoea batatas* (L.) Lam.).** *J Root Crops.* 2009; **35**(1): 1–27. Indian Society for Root Crops, ISSN 0378-2409.
[Reference Source](#)
- Satterthwaite FE: **An approximate distribution of estimates of variance components.** *Biometrics.* 1946; **2**(6): 110–114.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schurr U, Walter A, Rascher U: **Functional dynamics of plant growth and photosynthesis—from steady-state to dynamics—from homogeneity to**

heterogeneity. *Plant Cell Environ.* 2006; **29**(3): 340–352.

[PubMed Abstract](#) | [Publisher Full Text](#)

Snedecor GW, Cochran WG: **Statistical Methods.** Iowa State University Press, 1967.

[Reference Source](#)

Solis J, Villordon A, Baisakh N, *et al.*: **Effect of drought on storage root development and gene expression profile of sweetpotato under greenhouse and field conditions.** *J Am Soc Hort Sci.* 2014; **139**(3): 317–324.

[Publisher Full Text](#)

Somda ZC, Kays SJ: **Sweet potato canopy architecture: branching pattern.** *J Amer Soc Hort Sci.* 1990; **115**(1): 33–38.

[Reference Source](#)

Sseruwu G: **Breeding of sweetpotato (*Ipomoea batatas* (L.) Lam.) for storage root yield and resistance to *Alternaria* leaf petiole and stem blight (*Alternaria* spp.) in Uganda.** *Breeding Major Food Staples.* 2012; 333–363.

[Reference Source](#)

Tadesse Y, Almekinders CJ, Schulte RP, *et al.*: **Understanding farmers' potato production practices and use of improved varieties in Chench, Ethiopia.** *J Crop Improv.* 2017; **31**(5): 673–688.

[Publisher Full Text](#)

Tuberosa R: **Phenotyping for drought tolerance of crops in the genomics era.** *Front Physiol.* 2012; **3**: 347.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tumwegamire S, Rubaihayo PR, Grüneberg WJ, *et al.*: **Genotype × environment**

interactions for East African orange-fleshed sweetpotato clones evaluated across varying ecogeographic conditions in Uganda. *Crop Sci.* 2016; **56**(4): 1628–1644.

[Publisher Full Text](#)

Venus JC, Causton DR: **Plant growth analysis: a re-examination of the methods of calculation of relative growth and net assimilation rates without using fitted functions.** *Ann Bot.* 1979; **43**(5): 633–638.

[Publisher Full Text](#)

Villordon A: **Progression of sweetpotato root architecture development in an experimental aeroponics culture system Sweetpotato Aeroponics Project LSU AgCenter Sweet Potato Research Sta?** 2015.

<http://www.doi.org/10.6084/m9.figshare.1472959>

Wenk EH, Falster DS: **Quantifying and understanding reproductive allocation schedules in plants.** *Ecol Evol.* 2015; **5**(23): 5521–5538.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wilson LA: **Tuberisation in sweet potato.** In: R L. Villareal & T D. Griggs (eds.). *Sweet potato. Proceeding of the 1st. Int. Symp. AVRDC, Taiwan, China.* 1982.

Wilson LA, Lowe SB: **The Anatomy of the Root System in West Indian Sweet Potato (*Ipomoea Batatas* (L.) Lam.) Cultivars.** *Ann Bot.* 1973; **37**(3): 633–643.

[Publisher Full Text](#)

Yanfu Y, Jialan T, Yunchu Z, *et al.*: **Breeding for early-maturing sweet potato varieties.** In: Mackay, K.T.; Palomer, M.K.; Sanitco, R.T.ed. *Sweetpotato research and development for small farmers.* Philippines, College Laguna, SEAMEO-SEARCA, 1989; 67–98.

[Reference Source](#)

Open Peer Review

Current Peer Review Status:



Version 3

Reviewer Report 02 June 2020

<https://doi.org/10.21956/gatesopenres.14302.r28757>

© 2020 del Rio A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Alfonso del Rio 

Department of Horticulture, US Potato Genebank, University of Wisconsin-Madison, Sturgeon Bay, WI, USA

I still find lots of details that obscure the data and the important information of the paper. It looks a little better but not quite completed in a way that can allow readers to capture the science and potential applications expected from this study. Some parts are still difficult to read since they do not connect the data with the narrative and the findings. In some cases are repetitive concepts scattered in diverse places.

The number of figures and tables are too many, 18 in total. In particular I do not think you need that much information.

I have checked some parts of the paper and have placed some examples of how the authors can synthesize information. My suggestion is that the authors go again all over the manuscript and work on organizing information in a simpler way. Because of the practical value in sweet potato agriculture and sustainability, this study should be easy to read for large audiences in diverse areas.

To start please note an example in one part of the abstract below. Everything highlighted in bold provides the information. Words between parenthesis are not needed since you can explain those details in results and discussion. I also added some wording suggestions in italics ---

...**Data analysis was done using residual or restricted maximum likelihood (REML)**. (in GenStat 18th Edition). (There were strong linear and quadratic trends over time ($P < 0.001$) indicating a) **Data showed non-linear growth pattern within and between locations, seasons, and genotypes for most traits. Some genotypes displayed early initiation and** (a determinate linear) **increase of bulking, while others showed late initiation** (following a quadratic growth pattern). **Broad sense heritability of CSRFAB was** (would be) **low probably due to large GxE interactions but higher in** (, however, it was relatively high (50.5%) compared to other yield related) **other traits probably due to** (indicating) **high genetic influence and the effectiveness of the methodology** (accuracy of the developed method to quantify yield overtime)....

In Experimental Design:

The plants of each of the 130 sweet potato genotype were propagated as clonal forms or segregating seeds?

In Results and Discussion:

In Growth pattern analysis and phenotypic variability across selected CSRFAB sweetpotato traits ---

What do you mean when you saidFor most of the traits, their three-way interactions were significant which means in reality that at least one of the 2-way interactions changes across the third factor. How is this important in the discussion?

Also it seems to be a contradiction saying that there was high genetic variability across locations and the need to study stability. Please clarify.

In Variance component analysis and heritability estimates of selected sweetpotato growth traits in a nonlinear model structure

You said the same in two parts of the same paragraph "This implies that the scale used to measure the trait is precise and the trait is not influenced much by the environment. Heritability was relatively high for SRNO (67.2%) and CSRFAB (50.5%). The low residual variance and high heritability observed for CSRFAB imply that the CSRFAB trait in this study was not greatly influenced by environment and the scale used to measure the trait is precise"

Genetic effects on yield of CSRFAB genotypes in sweetpotato:

I placed below another example on how you can be more straightforward in providing a more clear description of your results : "...We performed a harvest basis analysis and calculated the corresponding heritability for each trait to understand the dynamics of breeding values across harvest times. We targeted the traits that can affect final yield. Thus, the genotypic variances among seven main traits associated with CSRFAB varied from 0 to 72.46 (Table 6). Genotypic variance was high for HI at 120 DAP (72.46) and VY at 180 DAP (59.91) (Table 6). Zero variance was recorded for VY (120 DAP), HI (150 DAP), SEN (180 DAP) and weevils (180 DAP). The zero (or negative) variance could be due to an artifact of the algorithm or competition effects between adjacent plots. These results imply no significant change in VY, HI, SEN and weevil damage within population for the respective growth periods. Since heritability is influenced by genetic variance, it was directly affected at the same traits and harvest periods. (ranging from 0 to 67%)..."

The paragraphs below are highly speculative and possibly wrong if you assume your comments on artifacts in the algorithm. You need to support this better --- *The 0% heritability implies that the effect of the traits is moving to fixation, that is, its frequency in the population is close to 100%. If we score the effect of the allele (s) regulating the trait having no heritability in the population, the result would mean in genetic terms, that the allele frequency in the population is 100%, therefore the genetic variance at the loci of these genes is zero, so any variance in the corresponding phenotypic traits cannot be attributed to the non-existent genetic variance...and the next paragraph.*

In: Identification of storage formation and bulking growth patterns and their characterization:

I don't find the comparative data from the classic method ---

"We characterized in this study growth patterns associated with CSRFAB in sweetpotato and compared the accuracy of CSRFAB scores and the classic method used to measure growth change overtime (compared yield change overtime)....

Was this measuring rather **consistently** or **effectively** than **accurately**? I don't see data that supports this assertion.

...The scoring method developed in this study accurately measured changes in the CSRFAB trait..."

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plant breeding and Genetics and, Conservation Genetics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 28 Jun 2020

Robert Mwanga, International Potato Center (CIP), Kampala, Uganda

RESPONSE BY AUTHORS (Signed in By Robert Mwanga) TO REVIEWER (Dr. Alfonso del Rio)

I still find lots of details that obscure the data and the important information of the paper. It looks a little better but not quite completed in a way that can allow readers to capture the science and potential applications expected from this study. Some parts are still difficult to read since they do not connect the data with the narrative and the findings. In some cases are repetitive concepts scattered in diverse places.

RESPONSE: *We have modified the text and removed repetitive concepts in the results section.*

The number of figures and tables are too many, 18 in total. In particular I do not think you need that much information.

RESPONSE: *We have removed one table and one figure.*

I have checked some parts of the paper and have placed some examples of how the authors can synthesize information. My suggestion is that the authors go again all over the manuscript and work on organizing information in a simpler way. Because of the practical value in sweetpotato agriculture and sustainability, this study should be easy to read for large audiences in diverse areas.

RESPONSE: *We have revised the text to make it simpler to read and understand wherever it was not clear. All the suggested areas to be modified have been revised as recommended.*

To start please note an example in one part of the abstract below. Everything highlighted in bold provides the information. Words between parenthesis are not needed since you can explain those

details in results and discussion. I also added some wording suggestions in italics ---

....**Data analysis was done using residual or restricted maximum likelihood (REML)**. (in GenStat 18th Edition). (There were strong linear and quadratic trends over time ($P < 0.001$) indicating a) **Data showed non-linear growth pattern within and between locations, seasons, and genotypes for most traits. Some genotypes displayed early initiation and** (a determinate linear) **increase of bulking, while others showed late initiation** (following a quadratic growth pattern). **Broad sense heritability of CSRFB was** (would be) **low probably due to large GxE interactions but higher in** (, however, it was relatively high (50.5%) compared to other yield related) **other traits probably due to** (indicating) **high genetic influence and the effectiveness of the methodology** (accuracy of the developed method to quantify yield overtime)....

RESPONSE: *Text has been modified as suggested.*

In Experimental Design:

The plants of each of the 130 sweet potato genotype were propagated as clonal forms or segregating seeds?

RESPONSE: *Each genotype was propagated as a clone (not as a botanical seed).*

In Results and Discussion:

In Growth pattern analysis and phenotypic variability across selected CSRFB sweetpotato traits ---

What do you mean when you saidFor most of the traits, their three-way interactions were significant which means in reality that at least one of the 2-way interactions changes across the third factor. How is this important in the discussion?

Also it seems to be a contradiction saying that there was high genetic variability across locations and the need to study stability. Please clarify.

RESPONSE: *The interaction between genotype, location and season were significant as were most of the three-way interactions implying that you have to optimize your growth curve for the 3 factors together, because they modify each other's effect - see modified text.*

In Variance component analysis and heritability estimates of selected sweetpotato growth traits in a nonlinear model structure

You said the same in two parts of the same paragraph "This implies that the scale used to measure the *trait is precise and the trait is not influenced much by the environment*. Heritability was relatively high for SRNO (67.2%) and CSRFB (50.5%). The low residual variance and high heritability observed for CSRFB *imply that the CSRFB trait in this study was not greatly influenced by environment and the scale used to measure the trait is precise*"

RESPONSE: *This has been corrected in the text.*

Genetic effects on yield of CSRFB genotypes in sweetpotato:

I placed below another example on how you can be more straightforward in providing a more clear description of your results : "...We performed a harvest basis analysis and calculated the corresponding heritability for each trait to understand the dynamics of breeding values across harvest times. We targeted the traits that can affect final yield. Thus, the genotypic variances among seven main traits associated with CSRFB varied from 0 to 72.46 (Table 6). Genotypic variance was high for HI at 120 DAP (72.46) and VY at 180 DAP (59.91) (Table 6). Zero variance was recorded for VY (120 DAP), HI (150 DAP), SEN (180 DAP) and weevils (180 DAP). The zero (or negative) variance could be due to an artifact of the algorithm or competition effects between adjacent plots. These results imply no significant change in VY, HI, SEN and weevil damage within population for the respective growth periods. Since heritability is influenced by genetic variance, it was directly affected at the same traits and harvest periods. (ranging from 0 to 67%)..."

RESPONSE: *The recommended modification has been followed.*

The paragraphs below are highly speculative and possibly wrong if you assume your comments on artifacts in the algorithm. You need to support this better --- *The 0% heritability implies that the effect of the traits is moving to fixation, that is, its frequency in the population is close to 100%. If we score the effect of the allele (s) regulating the trait having no heritability in the population, the result would mean in genetic terms, that the allele frequency in the population is 100%, therefore the genetic variance at the loci of these genes is zero, so any variance in the corresponding phenotypic traits cannot be attributed to the non-existent genetic variance...and the next paragraph.*

RESPONSE: *The recommended modification has been incorporated.*

In: Identification of storage formation and bulking growth patterns and their characterization:**I don't find the comparative data from the classic method ---**

"We characterized in this study growth patterns associated with CSRFB in sweetpotato and compared the accuracy of CSRFB scores and the classic method used to measure growth change overtime (compared yield change overtime)...."

RESPONSE:

1. *Please, look at Figure 2 growth pattern for classic method using yield against Figure 3 using DCSRFB scores for the DCSRFB genotypes.*
2. *Please, also look at Figure 4 growth pattern for classic method using yield against Figure 5 using CSRFB scores for CSRFB genotypes.*

Was this measuring rather **consistently** or **effectively** than **accurately**? I don't see data that supports this assertion.

...The scoring method developed in this study accurately measured changes in the CSRFB trait..."

RESPONSE: *The conclusion has been modified as suggested.*

Competing Interests: The authors declare that they have no competing interests.

Version 2

Reviewer Report 20 January 2020

<https://doi.org/10.21956/gatesopenres.14205.r28324>

© 2020 del Rio A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Alfonso del Rio 

Department of Horticulture, US Potato Genebank, University of Wisconsin-Madison, Sturgeon Bay, WI, USA

This study was aimed to determine how continuous storage root formation and bulking vary over diverse genotypes of sweet potatoes from Uganda. This remarks that some genotypes with good CSRFAB can be identified and have the potential to be used to enhance sweet potato productivity.

It is evident that identifying sources of good traits is significant to promote advances in breeding and agriculture. From that standpoint this study has merit. However, the reviewer finds this manuscript doesn't have good flow and it is difficult to follow.

One criticism is that the authors build the discussion including excessive statistics and excessive technical details of the statistical analysis. In many cases it was not clear what the findings were as information and concepts become unclear. I would advise the authors to make it simpler, you have to take into consideration that this paper will reach a broad audience with different backgrounds. This study has practical implications, giving a clear view of the results, discussions and analyses could enhance its outreach and practical application from breeding groups in the region.

In summary, this manuscript needs major revision. It needs to simplify the statistical analysis to a point of showing what is truly relevant with respect to the variation in traits assessed to explain CSRFAB and the variation among genotypes. This manuscript is too long.

Some additional comments:

- In Introduction: The second paragraph, it seems to me that the authors should clarify better if yield increase was because of the effect of breeding or just an increase of planting areas. If the latter this doesn't help much justifying the addition of new breeding forms.
- In Methods: What is a mega-environment for selection? I assume it was as reference to environmental diversity but the term *mega-environment* is a bit misleading in my view.

- Results: It is not new to report that clonal forms can respond differently in different environments. Variation levels across multiple environments is often high in field experiments in any crop. How is this result original? Then, you indicate that there is high variability among genotypes across locations + seasons. Wasn't this expected from a variable set of genotypes?
- When the authors indicate that "*this can be explained by the large population effect combined with sample size*", I would suggest clarification in that sentence since I understand the same number of plants and populations were used in the experiments. How were the effects compared on that basis?
- The parameters that the authors have used are related to storage root development, it seems that indicating that genotype performance increases overtime and is linearly explained is a logical outcome and fully expected. Storage root formation is related to maturity and developmental stages (=time). Maybe you need to clarify that result and explain better.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plant breeding and Genetics and, Conservation Genetics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Version 1

Reviewer Report 20 February 2019

<https://doi.org/10.21956/gatesopenres.13991.r26950>

© 2019 Shimelis H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Hussein Shimelis

African Centre for Crop Improvement, School of Agricultural, Earth and Environmental Sciences , College of Agriculture, Engineering and Science, University of KwaZulu-Natal (UKZN), Pietermaritzburg, South Africa

I suggest the following points for possible consideration:

Title:

- It can be adjusted to read: "Variation for Continuous Storage Root Formation and Bulking in Sweetpotato".

Abstract:

- Some of the acronyms must be written in full during the first mention e.g. CSRFAB, NaCRRRI etc.
- The conclusion is not clear, e.g., which genotypes were selected and recommended for continuous storage root formation and bulking and what yield advantage?

Introduction:

- The authors have provided long and detailed information on the overall productivity issues of sweetpotato. Instead, the authors should refocus on the challenges and opportunities of piecemeal harvesting under field conditions, and data and literature on continuous storage root formation and bulking. Problems such as termites, terminal drought stress, multiple and intercropping systems, and limited agricultural lands etc. can be pressing issues for piecemeal harvesting.

Materials and methods:

- Provide information on the maturity period (in months) of the test genotypes in Table 2. This will influence genotype comparison and recommendation for piecemeal harvesting.
- The 1 to 9 senescence scale should be described in detail, e.g. what is 1, 2, 3, 4, 6, 6, 7, 8 and 9.

Results:

- In all the Figures it will be interesting if the final yield/score (cumulative yield) of test genotypes are included, not only at 90, 120, 150 and 180 DAP.
- Is there a control harvest (e.g. normal harvest/once off harvest of a given genotype) to compare with cumulative yields over 90, 120, 150 and 180 DAP?

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plant Breeding, Plant Genetics, Crop Improvement, Quantitative Genetics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 15 February 2019

<https://doi.org/10.21956/gatesopenres.13991.r26921>

© 2019 Villordon A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Arthur Villordon 

LSU Agcenter Sweet Potato Research Station, Chase, LA, USA

Introduction:

This work addresses the subject of what the authors have termed “continuous storage root formation and bulking” in sweetpotato. The stated objective of this study was to investigate the genetic variability of “continuous storage root formation and bulking” and characterize growth patterns at different development stages to identify possible “continuous storage root formation and bulking” sweetpotato genotypes in a Ugandan germplasm collection for possible use as parents in trait improvement.

Merits:

The work addresses an important issue of relatively low sweetpotato storage root yields in Sub Saharan Africa where the crop is considered an important component of food security. Therefore, any work that seeks to understand the biological and environmental constraints of storage root yield has merit for advancing the fundamental understanding of the problem and development of solutions to improve productivity.

Critique:

Although the premise of the study was reasonable, the manuscript has a number of general and specific concerns, chief among this is the apparent oversight of previous work on the subject of storage root formation. There is general scientific consensus of the definition of “storage root formation” within the context of storage root developmental stages. It is unfortunate that the manuscript failed to acknowledge this scientific consensus and how it could have informed the current work and led to the advancement of the understanding of storage root formation in general, and within the context of the study location in particular. A short list of such work is provided in the references section below. I do encourage the authors to review some or all of the listed references, particularly the work of Lowe and Wilson (1974¹). There are many similarities and parallels between the current work and Lowe and Wilson’s methodology and approach. I do believe there is valuable data to be published from this research, but as it stands, I am recommending revision for the manuscript, as I believe that conceptual and structural changes need to be made on the manuscript. I hope that the following critiques will aid the authors in the refining of this manuscript to allow for revision in the future.

There has been prior work looking at abiotic and biotic constraints on early storage root development (between 30 and 50 days) wherein storage roots were anatomically examined to confirm storage root formation and then storage root number was used to measure the response (please see Solis *et al.* (2014²) and Gajanayake *et al.* (2014³) below). A common thread in these prior reports is the confirmation of the anatomical markers of storage root formation or lignification (generally lignified adventitious roots typically do not undergo storage root formation). These anatomical observations are relatively quick and require very basic tools and equipment. In contrast, the present work used visual assessments to indicate “storage root formation” or lignification (third paragraph and figure on page 6). For example, a score of 1 is based on visual observations of “no storage root initiation and no bulking.” Unfortunately, as prior work will show, non-swollen or non-thickened adventitious roots cannot be visually classified as undergoing storage root formation or lignification without anatomical confirmation. As it stands, any analyses or models based on this scale is based on the assumption that a score of 1 or 2=“no storage root initiation”, which may not be accurate. Lowe and Wilson (1974¹) used the same approach as the current work and used an index to quantify thickened storage roots. They made very clear assumptions about their rating scale, in part as a result of their detailed anatomical work on storage root formation, as detailed in another study by Wilson and Lowe (1973⁴). The authors are encouraged to review Wilson and Lowe (1974¹) and consider revising along these lines. I encourage the authors to reconsider using the “CSRFAB index” and instead use the quantitative measurements that they are already collected to describe the variation in storage root development and yield among the genotypes.

If there is one single problem that stands out, one that I find very unfortunate, is this statement found on the 2nd paragraph on page 4: “no study attempted to understand the storage root formation and bulking patterns overtime under field conditions.” Lowe and Wilson (1974¹), referenced above and listed below, conducted such a study, not very different in its conceptual approach from the current work. Lowe and Wilson grew six sweetpotato cultivars in field plots, and subsequently sampled and assessed storage formation and storage root development at weekly intervals during the first 8 weeks of growth, and then at monthly intervals after 8 weeks, for up to 24 weeks. Lowe and Wilson were very keen to recognize that storage root formation can only be determined anatomically, so they developed an index to quantify storage root development based on thickness of roots.

Finally, I looked up Yanfu *et al.* (1989) and was unable to find the data about storage root initiation occurring as early as 7 days. At this stage, adventitious roots are typically still undergoing primary growth, even in high yielding cultivars. Please cite cultivar or genotype or primary source, if available. In addition, I am unable to locate the data on storage root number ranging from 30 to 112 depending on genotype. Please verify this citation. Perhaps this refers to 30 to 112 g/plant (average root yield)?

Discussion:

The research on which this manuscript is based has generated valuable data on the variability of storage root yields of the sweetpotato genotypes in the collection and once the recommended structural and conceptual changes are made, the authors are encouraged to revise. Lowe and Wilson's (1974¹) work was titled "Comparative Analysis of Tuber Development in Six Sweet Potato (*Ipomoea batatas* [L.] Lam) Cultivars." Other than the unfortunate use of "tuber" in this case, perhaps the authors will take a cue from this very relevant prior work. Given that the subject of phenotypic variability of storage root development is under-researched in this geographic area, these findings will have undoubtedly have fundamental and applied merit.

However, if the authors insist on applying the term "storage root formation" in the context of their work, then they must implicitly define the contextual usage and exceptions relative to what the scientific community has adopted as the generally accepted conceptual and functional definition (please see a sampling of references below). In particular, they need to make the needed assumption that all non-swollen adventitious roots (Score 1) and not all slightly swelling adventitious roots (Score 2) are not undergoing storage root initiation and that anatomical examinations were not conducted. However, I strongly urge the authors to avoid this route as it does not advance our understanding of sweetpotato storage root formation and unnecessarily deviates from the scientific consensus.

References

1. LOWE S, WILSON L: Comparative Analysis of Tuber Development in Six Sweet Potato (*Ipomoea batatas* (L.) Lam) Cultivars. *Annals of Botany*. 1974; **38** (2): 307-317 [Publisher Full Text](#)
2. Solis J, Villordon A, Baisakh N, LaBonte D, et al.: Effect of Drought on Storage Root Development and Gene Expression Profile of Sweetpotato under Greenhouse and Field Conditions. *Journal of the American Society for Horticultural Science*. 2014; **139** (3): 317-324 [Publisher Full Text](#)
3. Gajanayake B, Reddy K, Shankle M, Arancibia R, et al.: Quantifying Storage Root Initiation, Growth, and Developmental Responses of Sweetpotato to Early Season Temperature. *Agronomy Journal*. 2014; **106** (5). [Publisher Full Text](#)
4. WILSON L, LOWE S: The Anatomy of the Root System in West Indian Sweet Potato (*Ipomoea Batatas* (L.) Lam.) Cultivars. *Annals of Botany*. 1973; **37** (3): 633-643 [Publisher Full Text](#)
5. Belehu T, Hammes P, Robbertse P: The origin and structure of adventitious roots in sweet potato (*Ipomoea batatas*). *Australian Journal of Botany*. 2004; **52** (4). [Publisher Full Text](#)
6. Firon N, LaBonte D, Villordon A, McGregor C, et al.: Botany and Physiology: Storage Root Formation and Development. 2009. 13-26 [Publisher Full Text](#)
7. Togari Y: A Study of Tuberous Root Initiation in Sweetpotato. *Bulletin of National Agricultural Experimental Station*. 1950; **68**: 1-96

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Relationship between sweetpotato root architecture and storage root development.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Comments on this article

Version 2

Author Response 25 Mar 2020

Robert Mwangi, International Potato Center (CIP), Kampala, Uganda

Response to Reviewer's (Dr. Alfonso del Rio) Report (responses are in italics).

In general, and in specific instances the reviewer's comments were very helpful and have been responded to in order to help readers to follow the work as suggested by the reviewer.

This study was aimed to determine how continuous storage root formation and bulking vary over diverse genotypes of sweet potatoes from Uganda. This remarks that some genotypes with good CSRFAB can be identified and have the potential to be used to enhance sweet potato productivity.

It is evident that identifying sources of good traits is significant to promote advances in breeding and agriculture. From that standpoint this study has merit. However, the reviewer finds this manuscript doesn't have good flow and it is difficult to follow.

Significant revisions have been made to improve on the flow; all changes have been highlighted.

One criticism is that the authors build the discussion including excessive statistics and excessive technical details of the statistical analysis. In many cases it was not clear what the findings were as information and concepts become unclear. I would advise the authors to make it simpler, you have to take into consideration that this paper will reach a broad audience with different backgrounds. This study has practical implications, giving a clear view of the results, discussions and analyses could enhance its outreach and practical application from breeding groups in the region.

Details on statistics and technical details have been drastically reduced while maintaining content wherever it makes the flow and the text easier to understand.

In summary, this manuscript needs major revision. It needs to simplify the statistical analysis to a point of showing what is truly relevant with respect to the variation in traits assessed to explain CSRFAB and the variation among genotypes. This manuscript is too long.

Major revisions have been made on the manuscript; methods section and experimental details have been modified and statistical analysis details reduced drastically.

Some additional comments:

- In Introduction: The second paragraph, it seems to me that the authors should clarify better if yield increase was because of the effect of breeding or just an increase of planting areas. If the latter this doesn't help much justifying the addition of new breeding forms.

A modification has been made to the text: "The increase in production was due to a combination of factors varying in different countries but mainly due to increase in area and breeding efforts."

- In Methods: What is a mega-environment for selection? I assume it was as reference to environmental diversity but the term mega-environment is a bit misleading in my view.

A modification has been made to the text to reflect environmental diversity; "mega-environment" has been dropped.

- Results:

It is not new to report that clonal forms can respond differently in different environments. Variation levels across multiple environments is often high in field experiments in any crop. How is this result original? Then, you indicate that there is high variability among genotypes across locations + seasons. Wasn't this expected from a variable set of genotypes?

The confusing, rather long text has been deleted and the remaining text modified to refer to the CSRFAB trait.

- When the authors indicate that "this can be explained by the large population effect combined with sample size", I would suggest clarification in that sentence since I understand the same number of plants and populations were used in the experiments. How were the effects compared on that basis?

The details have been deleted and the remaining text modified for clarity.

- The parameters that the authors have used are related to storage root development, it seems that indicating that genotype performance increases overtime and is linearly explained is a logical outcome and fully expected. Storage root formation is related to maturity and developmental stages (=time). Maybe you need to clarify that result and explain better.

The terms used in the manuscript are "storage root formation" or "storage root initiation" and "bulking". Bulking is related to development and maturity. However, storage root formation or initiation, when

continuous in the life of the genotype is a different component which has been identified in some genotypes in this study and can be exploited to benefit farming systems where there is piecemeal harvesting. A small modification has been made in the text by using both storage root formation and storage root initiation.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Modifications have been made under the Methods section for the reader to follow the design.

Are sufficient details of methods and analysis provided to allow replication by others?

No

The Methods and Analysis sections have been modified while avoiding expanding technical details.

If applicable, is the statistical analysis and its interpretation appropriate?

No

Details of the statistical analysis and technical details have been reduced for clarity.

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results? Partly

The text in the Methods and analysis and results have been modified for smooth flow and clarity.

Competing Interests

No competing interests were disclosed.

A statement on competing interests has been added to the text.

Reviewer Expertise

Plant breeding and Genetics and, Conservation Genetics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Competing Interests: The authors declare that they have no competing interests.