



Vascular endothelial cell development and diversity

Emily Trimm^{1,2} and Kristy Red-Horse^{3,4,5}✉

Abstract | Vascular endothelial cells form the inner layer of blood vessels where they have a key role in the development and maintenance of the functional circulatory system and provide paracrine support to surrounding non-vascular cells. Technical advances in the past 5 years in single-cell genomics and in in vivo genetic labelling have facilitated greater insights into endothelial cell development, plasticity and heterogeneity. These advances have also contributed to a new understanding of the timing of endothelial cell subtype differentiation and its relationship to the cell cycle. Identification of novel tissue-specific gene expression patterns in endothelial cells has led to the discovery of crucial signalling pathways and new interactions with other cell types that have key roles in both tissue maintenance and disease pathology. In this Review, we describe the latest findings in vascular endothelial cell development and diversity, which are often supported by large-scale, single-cell studies, and discuss the implications of these findings for vascular medicine. In addition, we highlight how techniques such as single-cell multimodal omics, which have become increasingly sophisticated over the past 2 years, are being utilized to study normal vascular physiology as well as functional perturbations in disease.

The development of functional vasculature is crucial for the health of every cell in the body. Endothelial cells, which line the inside of all blood and lymphatic vessels, have key roles in delivering oxygen and nutrients, regulating blood flow, modulating immune cell trafficking and maintaining tissue homeostasis. Vascular endothelial cell dysfunction is central to the progression of most chronic conditions, as well as ischaemic heart disease and stroke, the top two global causes of mortality¹. However, many questions remain about the fundamental biology of endothelial cells; for example, how do endothelial cell progenitors differentiate into distinct subtypes; how do they achieve organotypic heterogeneity; and how is the function of specialized endothelial cells perturbed in disease? In the past 5 years, the advent of single-cell RNA sequencing (scRNA-seq) technologies, combined with increasingly sophisticated strategies for genetic lineage tracing, have allowed the field to address these questions^{2,3}. In this Review, we highlight new discoveries that can be generalized to blood vessel development across all tissues, such as the role of cell cycling in the differentiation of endothelial cell subtypes as well as the phenomenon of ‘pre-artery’ specification. In addition, we discuss new findings for organ-specific endothelial cells in several key organs such as the heart, brain, lungs, kidneys and liver. Lymphatic endothelial cell development and heterogeneity have been reviewed elsewhere⁴ and are

beyond the scope of this Review. While this Review is not intended to comprehensively cover all aspects of organ-specific vasculature, we aim to demonstrate the wealth of new insights into endothelial cell biology that can be gained from scRNA-seq and integrated multimodal omic atlases. Particularly groundbreaking discoveries include the identification of functionally distinct capillary subpopulations in the lung, kidney and liver, as well as capillary zonation in the brain and heart. The identification of novel endothelial cell subtypes and molecular markers, as well as cell–cell interactions, has implications for tissue engineering and the development of novel therapeutic strategies for cardiovascular disease.

Arterial and venous differentiation

During embryogenesis, blood vessels must first arise de novo in a process known as vasculogenesis. The differentiation of angioblasts (endothelial cell progenitors) and their transition to endothelial cells are tightly regulated by transcription factors and signalling pathways that have been extensively studied^{5–9}. The first blood vessels arise from the rapid proliferation of endothelial cells, which subsequently form vascular tubes to create separate arteries and veins. Much research over the past two decades has described the crucial role of vascular endothelial growth factor (VEGF) and Notch signalling during arteriovenous differentiation. In the broader context, increased VEGF and Notch signalling

¹Stanford Medical Scientist Training Program, Stanford University School of Medicine, Stanford, CA, USA.

²Stanford Biophysics Program, Stanford University School of Medicine, Stanford, CA, USA.

³Department of Biology, Stanford University, Stanford, CA, USA.

⁴Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.

⁵Howard Hughes Medical Institute, Stanford University, Stanford, CA, USA.

✉e-mail: kredhors@stanford.edu

<https://doi.org/10.1038/s41569-022-00770-1>

Key points

- Artery differentiation is coupled to cell cycle arrest, and new arteries expand during development by recruiting endothelial cells from capillaries and veins.
- Single-cell RNA sequencing (scRNA-seq) data and multimodal omics atlases provide insights into the heterogeneity of endothelial cells across tissues, as well as conserved transcription codes during development.
- Novel methods of endothelial cell lineage tracing combined with scRNA-seq has facilitated the identification of functionally distinct capillary populations in the lung, liver and kidney, as well as their transcriptional response to disease.
- Capillary zonation, or the gradual phenotypic continuum of endothelial cells along an axis, has been identified in multiple organs including the brain, heart and liver, and communication between endothelial cells and mural cells has a key role in maintaining capillary zonation and important structures such as the blood–brain barrier.
- Comparison of scRNA-seq atlases from humans and mice reveals that endothelial cell populations are largely conserved between species across multiple organs and that although unique species-specific gene expression patterns exist for endothelial subtypes, the high degree of similarity between species demonstrates the utility of mice as a model organism for vascular biology.
- Advances in bioengineering have led to the creation of organoids with increasingly functional vasculature; key insights from single-cell studies are expected to facilitate improved differentiation protocols for human pluripotent stem cells as well as the *in vitro* development of organ-specific vasculature.

supports arterial specification, whereas reduced activation of VEGF and COUP transcription factor 2 (COUP-TFII) (also known as NR2F2) is required for venous specification^{10–15}. Although angiogenesis (the formation of new vessels from existing vessels) is mechanistically distinct from vasculogenesis, angiogenesis is also regulated by VEGF and Notch signalling. VEGF activates angiogenesis, whereas Notch restricts it^{13,16–27}. Despite the decades-old knowledge that the VEGF and Notch signalling pathways are required for arterial differentiation, evidence indicating that these pathways directly regulate the expression of genes associated with arterial specification is scarce. However, studies from the past 5 years have begun to elucidate the underlying mechanisms involved in how these pathways alter the fate of endothelial cells.

Cell cycling and endothelial cell fate

Insights from scRNA-seq and mosaic analyses have spurred the advances in our understanding of how Notch and COUP-TFII signalling support and restrict arterial differentiation, respectively^{28–31}. Surprisingly, the mechanism seems to involve the effects of Notch and COUP-TFII signalling on cell cycling and metabolism, rather than the direct regulation of arterial and venous cell fate determinants. The first study describing the relationship between cell cycling and arterial differentiation used the retina as a model system²⁸. During blood vessel formation, initiation of blood flow resulted in shear stress-induced Notch signalling and subsequent upregulation of gap junction $\alpha 4$ protein (also known as connexin 37). Gap junction $\alpha 4$ protein signalling led to flow-induced endothelial cell cycle arrest via p27 expression, enabling the expression of arterial specification genes²⁸. A similar relationship between arterial differentiation and cell cycling was subsequently described in the developing heart, although in the heart, arterial differentiation was initiated in single endothelial cells within the immature vascular plexus before the initiation

of blood flow²⁹. These individual pre-artery endothelial cells were lineage-traced to mature coronary arteries, establishing these endothelial cells as pre-specified progenitors. Pre-arterial differentiation was augmented by Notch activation, but blocked by COUP-TFII overexpression, owing to the transcriptional activation of cell cycle genes^{29,30}.

Numerous epigenetic studies support the link between cell cycle genes and the inhibitory effects of COUP-TFII on arterial differentiation. In a genome-wide analysis comparing COUP-TFII DNA binding with artery-specific and vein-specific transcriptional enhancers, COUP-TFII was over-represented at vein-specific enhancers, the majority of which were near actively expressed cell cycle genes³². The association between proliferating endothelial cells and capillary clusters with enriched expression of genes characteristic of venous endothelial cells had previously been identified in the brain as well as in regenerating aortas, adding further evidence supporting the idea that actively cycling cells are skewed towards a venous fate^{33,34}. Investigators are also beginning to identify the molecular mechanisms underlying the role of the cell cycle in arterial endothelial cell differentiation³¹. Genomic analyses of mice expressing the fluorescent ubiquitination cell cycle indicator reporter specifically in endothelial cells showed that endothelial cells in the late G1 state are more sensitive to transforming growth factor- $\beta 1$ (TGF $\beta 1$)-stimulated arterial specification-associated gene transcription than those in the early G1 state³¹. These newer studies correlating arterial endothelial differentiation with cell cycle exit contribute to our mechanistic understanding of arterial differentiation during development and form an important embryological parallel to older observations that mature arterial endothelial cells rarely proliferate in adults^{20,26}.

Pre-specification of arterial endothelial cells has also been described in other vascular beds such as the zebrafish trunk and the mouse developing retina in cell populations with low proliferative rates and high Notch activity^{35–39}. Live imaging in zebrafish revealed activation of Notch in tip cells, which upregulated *cxc4a* (encoding the chemokine C-X-C motif receptor 4a) expression in endothelial cells, enabling them to connect with and join arterial vessels³⁶. In the mouse retina, a subset of cells within the tip cell subpopulation similarly receives higher Notch signalling, which facilitates arterial differentiation (as indicated by the expression of *Cxcr4*) and migration of cells to form arteries^{35,40}. Pre-specification at the tip cell location is supported by data from scRNA-seq analyses showing that tip cells and arterial endothelial cells are closely related at the transcriptional level³³. In addition to Notch, increased VEGF–ERK signalling in both the retinal tip cells and coronary vessel plexus is another upstream trigger of cell cycle exit and arterial differentiation^{35,41,42}. VEGF is normally expressed where arterial pre-specification occurs, notably the retinal tip cell region and the intramyocardium^{41,42}. A mosaic analysis that involved genetic alteration of individual cells to express high levels of either the VEGF receptor 2 or Notch showed that both stimuli caused endothelial cells to exit the cell cycle and increased their propensity to

colonize arteries^{30,43}. To further investigate the downstream effects of Notch signalling on endothelial cell differentiation, an RNA-seq analysis was performed in mice with normal, diminished or increased Notch signalling³⁰. Notch-induced arterial endothelial cell differentiation was not dependent on upregulation of genes that determine arterial fate, but instead was dependent on the suppression of MYC-induced transcription of cell cycle and/or metabolism genes.

Arterial expansion via cell recruitment

The findings discussed above support a model in which arterial differentiation is coupled to cell cycle arrest (FIG. 1), which presents an intriguing question: how do arteries expand in size during development if their endothelial cell population does not proliferate? Numerous lineage-tracing studies from the past 5 years have shown that arteries grow by recruiting cells from veins and capillaries, rather than by expansion via endothelial cell division^{26,34,39,44,45}. In the mouse heart, timed labelling of *Apj*-expressing capillary plexus cells resulted in progressive pre-arterial specification and

accumulation of endothelial cells in arteries during permissive developmental and postnatal time windows^{29,46}. Three studies have assessed lineage-labelled arteries (BMXCreER), capillaries (Mfsd2aCreER) and veins (Gm5127(BAC)CreER) during postnatal retinal vasculature development in mice and showed that endothelial cells migrate in a vein-to-capillary-to-artery direction as the retinal vascular network expands and matures, but not in the opposite direction^{39,44,47}. Interestingly, these migration patterns are restricted to development and not seen in the mature eye. Furthermore, another study combined scRNA-seq analysis with endothelial cell lineage tracing and showed that endothelial cell migration from the vein to the artery is a common theme during development⁴⁵. The trajectory analysis performed on scRNA-seq data from mice at embryonic days 8–11 and matched human data predicted differentiation from vein to artery at multiple sites in the embryo in this early period⁴⁵. Computational predictions were validated by overlaying scRNA-seq data from lineage-labelled mouse vein endothelial cells (Nr2f2CreER)⁴⁵. This propensity of endothelial cells to migrate into arteries might also explain why endocardium-derived vessels preferentially populate arterial segments in the outer wall of postnatal hearts⁴⁸.

Endothelial cells migrate against blood flow

The physical forces of blood flow drive endothelial cell migration towards arteries because endothelial cells are programmed to move against the direction of flow^{26,49}. Many questions remain regarding how endothelial cells are guided to perform this task, but several mechanisms have been proposed, including involvement of the bone morphogenetic protein (BMP) signalling pathway. Inhibiting the BMP receptor signalling components activin receptor type 1 (ALK1), endoglin or SMAD4 blocks blood flow-guided migration of endothelial cells in many settings^{50–53}. Loss of these components and subsequent loss of proper endothelial cell migration leads to arteriovenous malformations that form direct shunts between arteries and veins, bypassing capillaries^{39,44,54,55}. Zebrafish intersomitic vessel development is balanced by vein endothelial cell proliferation and polarization or migration against flow into arterial segments^{38,56–58}, a process that requires the actin cytoskeletal regulator Wasb (the homologue of human Wiskott–Aldrich syndrome protein)⁵⁸ and flow-activated Notch^{38,57}.

Overactivation of endothelial cell migration against blood flow and into arteries has been achieved in mice by overexpressing *Eng* (encoding endoglin) or *Dach1* (encoding dachshund homologue 1)^{51,59,60}, the latter of which results in increased numbers of artery branches. Interestingly, loss of endothelial cell-derived non-canonical WNT5A and WNT11 caused endothelial cells to become over-responsive to flow, increasing their polarization and migration against flow⁶¹. In the plexus, this augmented polarization and migration against flow increased pruning (the regression of a subset of microvessels within a growing vasculature), although the effect on arterial branching was not investigated⁶². Further investigation into the mechanisms that drive




Fig. 1 | Artery and vein endothelial cell differentiation in a developing vascular bed. Artery differentiation is coupled to cell cycle arrest. Shear stress activation of Notch signalling upregulates the levels of gap junction α4 protein (Gjα4), which subsequently inhibits the expression of cell cycle genes via p27 expression. Notch signalling also inhibits MYC-induced transcription of genes related to cell cycle and metabolism. Together, Notch signalling indirectly supports arterial cell fate via cell cycle inhibition. Pre-artery endothelial cells then migrate against the direction of blood flow in the vein-to-artery direction. Adjacent endothelial cells exchange positions in a process known as ‘neighbour exchange’ to expand arteries. By contrast, COUP transcription factor 2 (COUP-TFII) (also known as NR2F2) activates the expression of cell cycle genes, inhibits the expression of certain genes associated with arterial specification and induces vein-specific gene expression. VEGF, vascular endothelial growth factor.

endothelial cell migration towards arteries for arterial expansion might provide the foundation for therapeutic applications in ischaemic diseases⁶⁰.

Compared with arterial endothelial cells, less is known about the development and maintenance of endothelial cells in veins and capillaries. Veins have transcriptional profiles that are distinct from those of arteries and the signals involved in inducing the venous cell state have mostly been studied in the early embryo^{6–8,11}. The establishment of veins requires cell autonomous endothelial cell phosphoinositide 3-kinase signalling and expression of COUP-TFII⁶. Venous gene expression has been shown to be induced directly by BMP–BMP receptor type 1A (BMPRI1A)–SMAD signalling in both mice and fish, ultimately leading to the formation of the first veins of the embryo⁶³. This discovery was made through the identification of a vein-specific enhancer in *Ephb4* (encoding ephrin type B receptor 4) and SMAD1 and SMAD5 binding sites as regulators of venous gene expression. Further experiments showed that *Bmp4* is expressed preferentially in cells around early veins that express *Bmpr1a*⁶⁴. *Smad4* or *Bmpr1a* deletion in mouse endothelial cells abolished the expression of *Ephb4* and *Nr2f2* (which encodes COUP-TFII), repressing the development of the veins⁶⁴. Less is known about the induction of capillary fates; however, emerging evidence indicates that capillary endothelial cells are less of an established cell state than artery and vein endothelial cells, and might instead exist along a continuum, known as zonation^{29,45,65–69}. This zonation has been observed in capillaries in the brain and heart of both mice and humans.

scRNA-seq and multimodal omics

Endothelial cells have heterogeneous phenotypes that match the diverse structures and metabolic needs of each tissue and organ. For example, capillary endothelial cells of the brain form the blood–brain barrier (BBB) with tight intercellular junctions to restrict transport, whereas glomerular capillary endothelial cells in the kidney are highly fenestrated to support filtration of blood to Bowman's space⁷⁰. The anatomical heterogeneity of blood vessels has been well characterized and extensively reviewed⁷⁰. Technical innovations in genomics in the past decade have provided insights into the mechanisms underlying this molecular heterogeneity.

Despite substantial functional differences across organ systems, all endothelial cells share the same genome. Heterogeneity must, therefore, stem from how and when transcriptional networks are activated, as well as from epigenetic modifications. Transcriptional profiling of endothelial cells, especially at the single-cell level through scRNA-seq, has greatly informed our understanding of endothelial cell heterogeneity. The pace of technical innovations for single-cell level genomic measurements is staggering, such that it is now possible to profile both gene expression and chromatin accessibility in each cell with the use of 'multiome' platforms^{71,72}. In this section, we discuss studies that have incorporated these techniques to create atlases for individual tissues or entire organisms^{29,33,65–69,73–101} (TABLE 1).

Global insights from multi-organ studies

Bulk RNA-seq provided the first transcriptomic insights into the molecular heterogeneity of endothelial cells among different tissues, facilitating the identification of several tissue-specific endothelial markers for the heart, muscle, brain, liver and lung^{33,102}. scRNA-seq corroborates the finding that endothelial cells from different tissues have specialized transcription programmes, which can be observed when endothelial cells in large multi-tissue datasets typically computationally separate (or cluster) into organ-specific subtypes^{2,103,104}. A detailed summary of scRNA-seq techniques that are commonly used in cardiovascular research is available elsewhere¹⁰⁵. Interestingly, organotypic endothelial cell specialization in mammals might be more exaggerated in blood endothelial cells because lymphatic endothelial cells cluster together regardless of the tissue of origin². Single-cell analyses also facilitate the characterization of heterogeneity according to endothelial cell subtype (that is, artery, arteriole, capillary, venule and vein) (TABLES 2, 3). Endothelial cell heterogeneity across tissues is largely attributable to differential transcription in capillary endothelial cells; the endothelial cells in arteries and veins from different organs are more similar at the transcriptional level^{104,106}. Accordingly, ageing in the brain also affects capillary endothelial cells to a greater extent than arterial or venous endothelial cells^{107,108}. The level of resolution afforded by single-cell analyses also establishes that organs with highly specialized vasculature, such as the brain, liver, lungs and kidneys, have more distinct transcriptomic profiles, whereas the gene expression of heart, skeletal muscle, diaphragm, mammary gland and adipose tissue tends to overlap^{93,95}. In addition, whereas endothelial cells from different organs share several major developmental pathways, including those relating to WNT signalling, cytokine regulation and metabolism, the expression patterns of the majority of genes associated with these pathways are unique to each organ^{2,104}.

Numerous studies have incorporated chromatin architecture information to gain further insights into the tissue-specific transcriptional regulation in endothelial cells, first by identifying DNA binding motifs in accessible chromatin regions and then comparing these motifs with the expression of genes encoding transcription factors to computationally predict endothelial transcription codes^{109–111}. Some consistent observations have emerged from these studies. Transcription factor motifs from the ETS transcription factor family, including transcriptional regulator ERG and Friend leukaemia integration 1 transcription factor, are universally enriched in endothelial cells^{33,109}. This finding is consistent with data showing that heterogeneous endothelial cell gene expression requires the combined function of both the endothelial cell subtype-specific transcription factors and ETS factors at enhancers¹¹⁰. The source of heterogeneity in gene expression between tissues might stem from differences in distal, rather than proximal, *cis*-regulatory elements. In a study comparing assay for transposase-accessible chromatin using sequencing (ATAC-seq) peaks in endothelial cells from the liver, lung, brain and kidney, the majority of ATAC-seq

Table 1 | Multi-organ and organ-specific endothelial cell scRNA-seq and multiome atlases

Organ system	Species	Developmental stage	Dataset contents	Single-cell capture and scRNA-seq method	Refs.
Multi-organ	Mouse	Adult	scRNA-seq	Smart-seq2 and 10× Genomics 3'	95
			scRNA-seq	Microwell-seq	73
			scRNA-seq	Smart-seq2	74
			scRNA-seq, RNA-seq, ATAC-seq, MethylC-seq	10× Genomics GemCode	33
			sci-ATAC-seq	sci-ATAC-seq	75
Human	Embryonic (E72–E129)	sci-RNA-seq, sci-ATAC-seq	sci-RNA-seq3	76,92	
		Adult (22–74 years)	Smart-seq2 and 10× Genomics 3'	93	
Heart	Mouse	Embryonic (E12.5–E14.5)	scRNA-seq	10× Genomics 3'	29
	Mouse, human	Embryonic mouse (E12, E17.5), embryonic human (11 weeks, 14 weeks, 22 weeks), adult mouse	scRNA-seq	10× Genomics 3'	66
	Human	Embryonic (5–25 weeks)	scRNA-seq	STRT-seq	77
		Adult (40–75 years)	scRNA-seq	10× Genomics 3'	78
Lung	Mouse	Embryonic (E12.5–E17.5), postnatal (P3–P42)	scRNA-seq, scATAC-seq	10× Genomics 3'	79
		Postnatal (P1)	scRNA-seq	Drop-seq	80
		Postnatal (P3, P7, P14)	scRNA-seq	MULTI-seq	81
	Human	Adult	scRNA-seq	Smart-seq2 and 10× Genomics 3'	82
		Adult	scRNA-seq	10× Genomics 3'	83
Brain	Mouse	Embryonic (E14.5)	RNA-seq (TRAP-seq), scRNA-seq	TruSeq v2	84
		Adult	scRNA-seq	Smart-seq2	65
	Mouse, human	Adult mouse, paediatric and adult human	snRNA-seq	10× Genomics 3'	67
	Human	Fetal, adult	scRNA-seq	10× Genomics 3'	69
		Adult	scRNA-seq, RNA-seq	10× Genomics 3'	68
Kidney	Mouse	Embryonic	scRNA-seq	10× Genomics 3'	85
		Embryonic E14.5	scRNA-seq	10× Genomics 3'	87
		Embryonic E14.5	scRNA-seq	Drop-seq, Chromium 10× Genomics and Fluidigm C1	86
		Embryonic, adult	snRNA-seq, sn-ATAC-seq	Smart-seq2 and 10× Genomics 3'	88
	Human	Embryonic, paediatric, adult	scRNA-seq	10× Genomics 3'	89
		Embryonic (12–18 weeks)	scRNA-seq	Drop-seq	90
		Embryonic (9–19 weeks)	scRNA-seq	10× Genomics 3'	91
Adult	snRNA-seq, sn-ATAC-seq	10× Genomics 3' v2	96		
Liver	Mouse	Embryonic (E12–E18), postnatal (P2–P30)	scRNA-seq, bulk RNA-seq	10× Genomics 3'	97
		Adult	scRNA-seq	MARS-seq	98
			scRNA-seq	10× Genomics 3'	99
	Human	Adult	scRNA-seq	CEL-seq2	100
			scRNA-seq	10× Genomics 3'	101

ATAC-seq, assay for transposase-accessible chromatin using sequencing; CEL-seq2, cell expression by linear amplification and sequencing; Drop-seq, droplet sequencing; E, embryonic day; MARS-seq, massively parallel RNA single-cell sequencing; MULTI-seq, multiplexing using lipid-tagged indices for single-cell and single-nucleus RNA sequencing; P, postnatal day; scRNA-seq, single-cell RNA sequencing; snRNA-seq, single-nucleus RNA sequencing; sci-ATAC-seq, single-cell combinatorial indexing assay for transposase-accessible chromatin using sequencing; sci-RNA-seq, single-cell combinatorial indexing RNA sequencing; Smart-seq2, switching mechanism at the 5' end of the RNA template sequencing; sn-ATAC-seq, single-nucleus combinatorial indexing assay for transposase-accessible chromatin using sequencing; STRT-seq, single-cell tagged reverse transcription sequencing; TRAP-seq, targeted purification of polysomal mRNA.

peaks at <2 kb from transcription start sites (proximal peaks) were shared between tissues, whereas the majority of ATAC-seq peaks at >2 kb from transcription start sites (distal peaks) were unique to individual tissues³³. Predicted tissue-specific endothelial cell transcriptional

networks include lymphoid enhancer-binding factor or T cell factor family members in the brain, which are well-known to be downstream effectors of canonical WNT signalling. GATA transcription factors seem to regulate heart-specific and liver-specific endothelial

Table 2 | Representative selection of marker genes for mouse organ-specific vascular subtypes

Organ system	Developmental stage	Vascular EC subtype	Marker genes	Refs.
Multi-organ	Embryonic or adult	Pan-endothelial	<i>Pecam1, Kdr, Cldn5, Emcn, Cdh5, Tie1, Egfl7</i>	2,80,85,131
		Vein	<i>Apj, Nr2f2, Vwf</i>	29,65,66
		Artery	<i>Gja4, Mecom</i>	29,66
Heart	Embryonic	Artery	<i>Gja4, Gja5</i>	66
		Pre-artery or arteriole	<i>Gja4, Gja5, Unc5b, Hey1, Slc45a4</i>	29,66
		Vein	<i>Nr2f2, Apj, Nr2f2</i>	29,66
		Capillary	<i>Aplnr, Car4</i>	66
	Adult	Tissue-specific EC	<i>Wt1, Slc28a2, Eepd1, Kcna5, Car8, Fbln1, Meox2, Rftn1, Lamb1, Myadm</i>	2
		Artery	<i>Apln, Dll4, Notch1, Mecom, Igfbp3, Cxcr4, Cx40</i>	29
Lung	Adult	Artery	<i>Mgp, Cdh13, Htra1, Bmx, Fbln2, Sulf1</i>	131
		Vein	<i>Slc6a2, Bst1, Car8, Amigo2, Mutn1, Csrp2</i>	131
		Tissue-specific EC	<i>Grtp1, Adrb1, Scn7a, Tmem100, Hpgd, Foxf1a, Nckap5, Rasgef1a, Fendrr, Prx</i>	2
		Capillary (aCap)	<i>Car4, Ednrb, Fibin, Tbx2, Rprml, Chst1, Apln</i>	81,131
		Capillary (gCap)	<i>Aplnr, Gpihbp1, Plvap, Cd93, Ptpnb, Cemip2, Tek, Cxcl12</i>	81,131
Brain	Embryonic	Tissue-specific EC	<i>Foxf2, Foxl2, Foxq1, Lef1, Ppard, Zfp551, Zic3</i>	84
		Adult	Tissue-specific EC	<i>Slco1c1, Slco1a4, Slc22a8, Mfsd2a, Slc38a3, Spock2, Foxf2, Edn3, Stra6, Slc38ak</i>
	Adult	Artery	<i>Gkn3, Sema3g, Efnb2</i>	65
		Venule	<i>Slc38a5</i>	65
		Capillary or vein	<i>Slc16a1</i>	65
Kidney	Adult	Tissue-specific EC	<i>Egfl7, Dram1, Dkk2, Esm1, Igfbp5, Pbx1, Boc, Igfbp3, Irx3, Tnfrsf25, Ptpn22</i>	87,88
		Glomerular EC	<i>Pi16, Plat, Ehd3, Cyb4b1, Tspan7, Lpl</i>	143
		Cortical EC	<i>Igfbp3, Plvap, Npr3</i>	143
		Medullary EC	<i>Igf1, Cryab, Igfbp7, Cd36, Aqp1, Ifi27l2a</i>	143
Liver	Embryonic or adult	Portal vein	<i>Ly6a, Cd34, Cd9, Ephb2, Gja5, Sox17</i>	97
		LSEC	<i>Mrc1, Fcgr2b, Stab1, Stab2</i>	97
	Adult	Tissue-specific EC	<i>c-Maf, Clec4g, Fcgr2b, Stab2, Oit3, Bmp2, Aass, Mrc1, Plxnc1, Wnt2</i>	2,136
		LSEC zone 1	<i>Dll4, Efnb2, Ltbp4, Msr1, Ntn4</i>	98,99
		LSEC zone 2 or 3	<i>Rspo3, Wnt2, Wnt9b</i>	98,99

aCap, aerocyte capillary; EC, endothelial cell; gCap, general capillary; LSEC, liver sinusoidal endothelial cell.

cell transcription, whereas homeobox transcription factor family members are enriched in the lungs and kidneys^{33,109}.

Multi-organ datasets have provided support for earlier observations that surrounding cells induce tissue-specific behaviour in endothelial cells, whereas endothelial cells in turn affect the transcriptomic profiles of their neighbours. For example, subpopulations of cardiac endothelial cells express cardiac muscle-specific genes and sinusoidal endothelial cells have similar gene expression profiles to those of adjacent hepatocytes^{2,98,111}. Numerous studies have begun to investigate the functional implications of these interactions across a number of organ systems. A study combining scRNA-seq analysis with bone marrow transplantation experiments characterized the crucial role of arterial endothelial cells in

establishing haematopoietic stem cell colonization of the bone marrow^{112,113}. Another study similarly used scRNA-seq to probe the complex response of endothelial cells to pericyte loss in the brain, a phenomenon seen in a wide range of neurological diseases¹¹⁴.

Finally, sexual dimorphisms are known to contribute to differences in organ function, susceptibility to disease and ageing, processes that involve blood vessels that permeate almost every tissue¹¹⁵. Transcriptomic and proteomic analyses have begun to establish that these differences, at least in the heart, are attributable to both hormonal control and sex chromosome composition¹¹⁵. Many organism-wide scRNA-seq analyses control for the possibility of sex-dependent differences in gene expression by utilizing tissues from a single sex, but only a few studies have assessed the effects of sex on organ-specific

transcriptomic profiles^{2,106}. Using the *Tabula Muris* dataset, sex was found to be an important factor underlying endothelial cell transcriptome heterogeneity, but only in some tissues, including the aorta, brain and lungs^{2,95}. Future cell atlas studies should incorporate sex as a parameter to better understand its influence on vascular physiology.

Organ-specific findings at high resolution

Cardiac vasculature. The majority of coronary blood vessels develop via convergent differentiation of endothelial cells from two distinct progenitor sources: the inlet vein known as the sinus venosus and the

endocardium that lines the heart chamber^{42,116–120}. Cells from these two populations ultimately localize to complementary yet overlapping regions of the heart. The sinus venosus mostly forms vessels in the outer ventricular walls and the endocardium mostly forms vessels in the septum and inner ventricular wall. Despite this spatial divergence, endothelial cells from both progenitor sources begin to sprout between embryonic days 11 and 12.5 in mice¹¹⁶. The sinus venosus and endocardium initially sprout in response to distinct molecular signals, specifically epicardial VEGFC–apelin receptor early endogenous ligand signalling and SMAD signalling for sinus venosus-derived cells, and intramyocardial

Table 3 | Representative selection of marker genes for human organ-specific vascular subtypes

Organ system	Developmental stage	Vascular EC subtype	Marker genes	Refs.
Multi-organ	Embryonic or adult	Pan-endothelial	PECAM1, CDH5, VWF, KDR, FLT1, TEK, CLDN5	68,78,82,91,93
		Artery	GJA4, GJA5, HEY1, GATA2, CXCR4, SOX17, MECOM	66,67,93
		Vein	ACKR1, NR2F2, PLVAP	68,69,78
Heart	Embryonic	Artery	GABBR2, GRIA2, SSUH2, JAG1	66
		Vein	ACKR1, LHX6, SELE	66
		Capillary	PRDM1, INMT, APLNR, CA4	66
	Adult	Tissue-specific EC	SLC14A1	93
		Artery	SEMA3G, EFNB2, DLL4,	78
		Vein	NR2F231, ACKR1	78
		Capillary	RGCC, CA4	78
Lung	Adult	Tissue-specific EC	VIPR1	93
		Artery	DKK2, SERPINE2	131
		Vein	CPE, PTGDS, C7, PLA1A	131
		Bronchial EC	VWA1, HSPG2, PLVAP, MYC, HBEGF	82
		Capillary (aCap)	ADIRF, S100A4, EMCN, HPGD, EDNRB, SOSTDC1, IL1RL1, APLN	131
		Capillary (gCap)	FCN3, EDN1, SLC6A4, GPIHBP1, CD36, IL7R, VWF, PTPRB, PLVAP	131
Brain	Adult	Large artery	LTBP4	69
		Artery	INTS6, HSPA1A, JUNB, MECOM, ARL15, TXNIP, MGP, ADAMTS1	68,69
		Arteriole	VEGFC, ARL15, BMX, EFNB2, WISR, AIF1L, CD320	67
		Venule	TSHZ2, ADGRG6, SLC38A5, LRRC1, BNC2, ETV6, TMEM132C, ATP10A, JAM2, PRCP, PRSS23, RAMP3	67–69
		Vein	ACKR1, IL1R1, TSHZ2, PTGDS, POSTN, DNASE1	68,69
		Large vein	CCL2	69
		Capillary	MFS2A, SLC7A5, TFRC, SLC38A5, SRARP, RGCC, SLC3A2, BSG, SLC16A1, SLC01A2	67–69
Kidney	Embryonic	Tissue-specific EC	NOTCH4	90
	Paediatric or adult	Glomerular EC	SEMA3G, CLDN5	89
Liver	Adult	Tissue-specific EC	OIT3	93
		Portal vein	AQP1, CD9, IFITM1, RAMP3, INMT, DNASE1L3, LIFR	97,101
		LSEC	CLEC4G, STAB1, STAB2, CD14	97
		LSEC zone 1	MGP, SPARCL1, TM4SF1, CLEC14A, BTNL9, ANPEP	100,101
		LSEC zone 2 or 3	CCL14, CLEC1B, FCN2, S100A13, LYVE1, FCN3	100,101
Central vein	SELP, SELE, VWF	97		

aCap, aerocyte capillary; EC, endothelial cell; gCap, general capillary; LSEC, liver sinusoidal endothelial cell.

VEGFA for endocardium-derived cells^{64,117,118,120}. After sprouting onto the developing heart, previously fully differentiated sinus venosus vein endothelial cells or endocardium will undergo extensive cell state changes within the coronary plexus and remodel into coronary arteries, capillaries and veins¹¹⁸.

After birth, neonatal cardiac ventricles undergo rapid growth and remodelling. The remaining trabecular myocardium, which lacks intramyocardial vessels during embryogenesis, undergoes compaction and acquires rich coronary vascularization¹²¹. Endocardium-derived vessels have been shown to expand into this region in the weeks after birth, and new vessels have been suggested to arise from postnatal lineage conversion of endocardial cells to coronary vessel endothelial cells⁴². Stage-specific lineage-tracing experiments have further refined this model by showing that late fetal and neonatal endocardial cells contribute minimally to the formation of postnatal coronary vessels, even with injury^{48,122,123}. Differentiation of endocardial cells into coronary endothelial cells occurs mostly during early heart development (before embryonic day 13.5 in mice) and endocardium-derived vascular tunnels do not adopt a mature artery fate until postnatal day 7 in mice^{9,48}. Instead of endocardial conversion, postnatal endothelial cells from coronary vessels in the inner myocardium express high levels of VEGFR3 and undergo angiogenic expansion in a delta-like protein 4–NOTCH 1-dependent manner to vascularize the newly compacted myocardium and extend the endocardium-derived vascular tunnels into mature arteries^{48,122}. Several studies have found that this endocardium-directed coronary growth can be enhanced under certain conditions such as with VEGFB transduction or overexpression of *Kdr* (encoding VEGFR2) in adult mice^{124,125}. Comparing scRNA-seq data of all cardiac endothelial cells from early (embryonic day 12.5) and late (embryonic day 17.5) stages revealed that *Bmp2* expression is specific to the embryonic time period when angiogenesis occurs¹²³. BMP2 expression could also reactivate endocardial angiogenesis in neonatal mouse models of myocardial infarction¹²³.

Direct and deep comparisons of capillary endothelial cell transcriptional states from lineage-traced hearts indicated that adult coronary endothelial cells originating from distinct sources do not retain the molecular signatures of their progenitors⁶⁶. This phenomenon of ‘convergent differentiation’ has also been seen in olfactory projection neurons¹²⁶, but not in certain populations of haematopoietic cells or tissue-resident macrophages, which retain markers of primed fate potential^{127,128}. Furthermore, no functional differences have been observed in endothelial cells from either lineage, such that they have largely identical proliferative capacity in response to injury⁶⁶.

Techniques such as reference mapping, as well as comparisons of mouse and human scRNA-seq datasets through robust integration, provide information about the transcriptional similarities and differences between species. These analyses of developing mouse and human hearts revealed that the developmental environment and endothelial cell subtype transcriptional states are similar

in both species^{66,129}, validating the use of mice as a model for many aspects of human coronary development (FIG. 2a). Follow-up functional experiments used these data in an induced pluripotent stem cell culture model to determine that the histone–lysine *N*-methyltransferase MECOM is crucial for human artery endothelial cell differentiation¹²⁹. Furthermore, trajectory analyses of human data predicted a transitory cell population from the endocardium¹²⁹, which had a gene expression pattern validated using lineage tracing that overlapped with that of the transitory cell population in mice¹²³. Therefore, despite the dearth of lineage-tracing techniques for human studies, overlapping human and mouse datasets provide some evidence that coronary vessels from humans also arise from the endocardium.

Lung vasculature. Alveoli, the terminal airspaces in the mammalian lung and a major site of gas exchange, are surrounded by a dense network of capillaries. Single-cell studies have identified heterogeneities in alveolar vasculature that are crucial for structural integrity. Alveolar development beginning during late embryogenesis is accompanied by the emergence of a specialized capillary subtype called *CAR4*⁺ aeryocyte capillaries (aCaps), which intermingle with the ‘general capillaries’ (gCaps)^{130–132}. *CAR4* encodes carbonic anhydrase 4, which catalyses the reversible hydration of carbon dioxide into bicarbonate and protons. Although gCaps have similar gene markers to capillary endothelial cells from other organs, aCaps express markers such as *ICAM1* (encoding intercellular adhesion molecule 1) and *EDNRB* (encoding endothelin receptor type B), and are unique to the lung. aCaps are also atypically large with a ‘Swiss cheese’-like morphology, spreading over the thin alveolar type 1 epithelium where gas exchange occurs^{130,131}. This location and their gene expression patterns indicate that aCaps have specialized roles in gas exchange and leukocyte trafficking.

Trajectory analysis of scRNA-seq and lineage-tracing data revealed that gCaps are progenitors for aCaps during alveolar development^{79,130–132} and after injury in adults^{131,133}. Development and maintenance of aCaps specifically require VEGF from airway epithelium, given that *Vegf* deletion results in depleted aCap numbers but does not affect gCap numbers. aCaps also seem to be required for lung maturation because lack of aCaps in mice results in enlarged alveoli despite the presence of myofibroblasts¹³⁰.

Together, this deeper understanding of capillary heterogeneity has informed human pathology. As in the heart, intraspecies comparisons have revealed that all endothelial cell subtypes found in mice are present in humans, and that *CAR4*⁺ aCaps also emerge during late embryogenesis^{79,83,131}. Analysis of a human lung adenocarcinoma sample revealed a loss of aCaps and gain of a transcriptionally intermediate population that co-expressed markers of aCaps and gCaps¹³¹. This same phenomenon was observed in single-cell atlases of lungs from patients who died of coronavirus disease 2019 (COVID-19)¹³⁴ (FIG. 2b). Interestingly, endothelial cells were the second-most abundant site of viral RNA load in these patients, despite low-to-absent expression of the entry receptor for SARS-CoV-2 (REF.¹³⁵), consistent with the




Fig. 2 | scRNA-seq reveals novel endothelial cell types. a | Comparison of human and mouse single-cell RNA sequencing (scRNA-seq) datasets for heart and brain vasculature reveals similar endothelial cell states between species. Although expression of unique genes exists for endothelial cell subtypes within each species, these analyses support the use of mouse models for studying human development and pathology. Key findings from these datasets reveal that the heart has conserved endothelial cell populations and similar capillary endothelial cell states in mice and humans and unique human artery specification genes. Furthermore, the brain also shows conserved endothelial cell populations and zonation between mice and humans and a greater diversity of human perivascular cells. **b** | Two

transcriptionally distinct populations of capillary endothelial cells exist in the lung: general capillaries (gCap) and aerocyte capillaries (aCap). These two capillary populations have different roles in lung maturation and mature lung function. In disease states such as adenocarcinoma, a transcriptional intermediate between gCap and aCap arises in both humans and mice. This transcriptional intermediate has also been seen in the lungs of patients with coronavirus disease 2019 (COVID-19). Overlapping features of inflammation and cellular stress can also be seen in both capillary subtypes in advanced chronic obstructive pulmonary disease. AV1, alveolar type 1 cell; RBC, red blood cell; t-SNE, t-distributed stochastic neighbour embedding.

knowledge that endothelial cells are a major site of pathology in patients with COVID-19 (REF. 136). A comparison of hyperoxia-induced bronchopulmonary dysplasia in mice with matched human samples showed that aCaps mediate the increased expression of *INHBA*, which encodes inhibin- β A chain⁸¹, a member of the TGF β superfamily that has been suggested to contribute to bronchopulmonary dysplasia. Furthermore, the study showed that capillary endothelial cell subtypes, rather than artery and vein endothelial cells, contain the most transcriptional changes that are potentially pathological in nature. Capillary endothelial cell subtypes also show the greatest change in inflammatory signalling and stress response in patients with advanced chronic obstructive pulmonary disease¹³⁷. These and other similar findings demonstrate the power of single-cell level atlases to provide a more granular understanding of pulmonary disease.

Brain vasculature. Given the prominent role of the central nervous system blood vessels in human disease and ageing¹³⁸, these vessels have been the subject of many

single-cell transcriptomic studies in the past 5 years. Indeed, single-cell analyses of brain-specific capillaries led to the identification of a previously unappreciated feature of many capillary beds. Capillaries comprise the vast majority of blood vessels and are the main site of oxygen exchange, yet multi-organ studies have thus far failed to identify a unifying capillary molecular marker². This lack of a unifying capillary marker might be attributable to the observation that instead of being a homogeneous entity, cells within a single capillary bed exist along an arterial-venous transcriptional continuum, referred to as zonation, which depicts gradual cellular phenotypic changes along an anatomical axis (FIG. 1). This feature of capillaries was first confirmed in the brain (via an approach that assessed and visualized data using distance matrices) and later in the heart (using artery or vein gene scores)^{29,65}. Interestingly, zonation is not a phenotype shared by endothelial cell-adjacent mural cells in the central nervous system vasculature. Arteriole smooth muscle cells abruptly transition to become pericytes at the arteriole-capillary boundary in both mice and humans^{65,139}.

Other studies have also identified capillary zonation in human brains, early mouse and human embryos and developing human hearts^{45,66–69} (FIG. 2a). Loss of communication between pericytes — a cell type reported to regulate the formation of the BBB^{140,141} — and endothelial cells was also found to skew capillary zonation towards a venous identity rather than causing widespread loss of BBB function and ectopic tip cells¹¹⁴.

Finally, single-cell analyses can also provide insights into the source of disease that results from genetic variants. Findings from the cross-referencing of cell subtype gene expression in mice with gene variants associated with Alzheimer disease or genes with altered expression in brain tissue from patients with Alzheimer disease suggest that capillary endothelial cells are a potential source of pathology, possibly through changes in BBB-related genes¹⁰⁸. The same analysis using human data pinpointed endothelial cells and mural cells as potential sites of disease¹³⁹.

Kidney vasculature. Given that the kidney has diverse and distinctive endothelial cell populations, the renal vasculature is a good candidate for single-cell analysis^{96,142}. The renal microvasculature regulates blood flow, facilitates filtration, modulates inflammation and maintains physiological blood pressure. These functions are compartmentalized to specific regions: glomerular endothelial cells make up the filtration barrier, whereas tubular and peritubular capillary endothelial cells absorb and secrete fluids and other substances^{96,142}. Single-cell analyses of mouse models of kidney disease have provided insights into the functional divisions between cortical, medullary and glomerular endothelial cells and identified cell type-specific responses to pathology. Cortical endothelial cells express high levels of *Igfbp3* (encoding insulin-like growth factor binding protein 3) and *Npr3* (encoding natriuretic peptide receptor 3) among other genes, whereas medullary endothelial cells express *Igfl1* (encoding insulin-like growth factor 1) and *Cd36* (encoding platelet glycoprotein 4). Glomerular endothelial cell-enriched genes include *Ehd3* (encoding EH domain-containing protein 3), *Cyp4b1* (encoding cytochrome P450 4B1) and *Tspan7* (encoding tetraspanin 7)^{143–146} (FIG. 3a). In response to acute water deprivation, medullary endothelial cells show the greatest transcriptional changes, upregulating the expression of genes related to oxidative phosphorylation in an effort to promote survival and concentrate urine¹⁴³. A mouse model of nephritis (via nephrotoxic serum administration) showed that the disease involves an endothelial cell trajectory towards an activated cell state, whereas endothelial cell states in a mouse model of diabetes mellitus (*ob/ob*) overlapped completely with that of wild-type mice^{147,148}.

Liver vasculature. Liver endothelial cells have long been known to have crucial roles in liver function, including supporting its dual blood supply from the hepatic artery and portal vein, scavenging macromolecules and regulating the host immune response to pathogens¹⁴⁹. scRNA-seq has provided insights into the transcriptome that support these diverse functions in both mice and

humans, and new studies have demonstrated how liver sinusoidal endothelial cells become dysregulated in the context of cirrhosis and hepatocarcinoma^{99,100}.

In a landmark study, investigators developed a strategy for spatially resolved scRNA-seq in the liver by mapping the positions of attached hepatocyte–endothelial cell pairs back to their expression-inferred tissue coordinates¹⁵⁰. Molecular signatures for subpopulations of pericentral liver endothelial cells were identified and the zonation of endothelial cells across lobules was characterized⁹⁸ (FIG. 3b). An analysis of the spatially resolved proteome revealed a high level of concordance with the transcriptomic data. Interestingly, although zonation in the brain corresponds to a continuum of arterial–venous transcriptional states, several studies have found that zonation in the liver relates to functional differences across the lobule axis^{98,101}. Endothelial cell zonation was found to drive much of this functional zonation in hepatocytes via a TIE–WNT signalling axis¹⁵⁰. The initial establishment of zonation in embryonic endothelial cell progenitors depends on MAF, a transcription factor crucial for sinusoid (or liver capillary) identity⁹⁷.

As with capillaries in the kidney, liver sinusoid endothelial cells have zone-specific susceptibility to injury inflicted during disease, such as sinusoid endothelial cells in lobule zone 3 that are particularly affected in cirrhotic liver of mice⁹⁹. Just as aCaps and gCaps in the lung lose their specialized transcriptional states in disease states, cirrhosis causes zone 3 sinusoid endothelial cells to lose their specialized fenestrae through a process known as capillarization⁹⁹.

Advances in organoid vasculature

Recent developments in single-cell genomics have revolutionized our understanding of the heterogeneity of vascular cells and their exquisitely regulated development within tissue microenvironments. As well as facilitating the identification of rare and novel endothelial cell types, such as specialized capillary subtypes in the lungs, liver, kidneys and retina, scRNA-seq has offered a higher resolution window into the transcriptional changes that occur during disease^{47,98,131,143}. Furthermore, scRNA-seq data have provided new insights into the interactions between endothelial cells and adjacent cell types, such as smooth muscle cells, pericytes and immune cells, and the roles that these interactions have in both the development and the maintenance of organ-specific vasculature. These findings, combined with results from sophisticated lineage-tracing studies, raise important considerations for tissue engineering. A lack of physiologically accurate perfusable vasculature is one of the most important challenges limiting the expansion of multicellular ‘mini-organs’ known as organoids¹⁵¹. Advances in differentiation protocols and culture techniques for human pluripotent stem cells (hPSCs)¹⁵² have resulted in the development of 3D self-organizing human blood vessel organoids, which are capable of forming complex vascular networks that recapitulate many of the mural cell–endothelial cell interactions known to be crucial for supporting structures such as the BBB¹⁵³. These organoids, as well as




Fig. 3 | scRNA-seq reveals inter-organ heterogeneity and heterogeneous endothelial cell responses to disease.
a | Single-cell RNA sequencing (scRNA-seq) has revealed transcriptomic heterogeneity between endothelial cells in each compartment of the kidney. Cortical endothelial cells, glomerular endothelial cells and medullary endothelial cells have differential responses to injury. Marker genes and enriched transcriptional factor networks (TFN) for each subtype are indicated. Mouse models have been developed to assess kidney injury during disease states including immune-mediated injury (nephritis and systemic lupus erythematosus), diabetic kidney injury and dehydration. Key pathways that are altered in endothelial cells from the different renal compartments of mice are highlighted. **b** | Endothelial cell zonation in the liver has been characterized in both humans and mice. Broadly, liver sinusoid endothelial cells (LSECs) can be grouped into three zones based on their localization with respect to the portal vein and central vein. The top five differentially expressed genes for human LSECs are indicated as marker genes for each zone. Studies in mice have revealed that LSECs in zone 3 are the most susceptible to damage associated with cirrhosis. EC, endothelial cell; HCC, hepatocellular carcinoma.

several hPSC-derived liver^{154,155}, kidney¹⁵⁶ and brain¹⁵⁷ organoids, have shown progress promising, though often partial, levels of host vascularization in vivo after transplantation into immunodeficient mice.

Although in vivo transplants are an exciting demonstration of the therapeutic potential of organoids, the

generation of endogenous organ-specific vasculature remains crucial. By using scRNA-seq data, one group demonstrated that hPSC-derived intestinal organoids are capable of inducing the differentiation of endothelial cells with an intestinal-specific transcriptional signature¹⁵⁸. Other efforts have focused on increasing

the complexity and functionality of organ-specific vasculature by modulating key signalling pathways and exposing in vitro systems to physiologically relevant physical forces¹⁵⁹. Through a combination of these methods, several groups have developed increasingly sophisticated cardiac organoids complete with internal lumens and vascular networks^{160–162}. Other groups have shown that modulating morphogen gradients and fluid shear stress conditions using microfluidic systems improves the expansion of endothelial progenitors in the kidney¹⁶³ and brain¹⁶⁴, although further analysis is required to determine whether faithful endothelial cell subtype differentiation has been achieved with these methods.

Finally, 3D bioprinting offers an alternative approach to embedding perfusable vascular channels for larger regions of tissue. In a newly developed biomanufacturing method, multiple hPSC-derived organoids, embryoid bodies or spheroids can be combined into a granular tissue matrix. Sacrificial ink is injected and then removed to create channels within the tissue that can be seeded with endothelial cells¹⁶⁵. This approach has resulted in the successful vascularization of a contractile cardiac organoid matrix¹⁶⁵ and might be capable of producing more structurally accurate organ-specific vasculature than other methods.

Conclusions

In this Review, we discuss the current molecular understanding of the differentiation of endothelial progenitors into arteries and veins. New insights are used to form a model in which artery differentiation is coupled to cell cycle arrest, whereas vein differentiation is associated with cell proliferation and direct induction of the BMP–BMPRIA–SMAD pathway. Arteries then expand by recruiting cells from capillaries and veins, which migrate against the direction of blood flow to expand the arterial network. Numerous multi-organ scRNA-seq and single-cell multi-omic atlases developed in the past 5 years have led to the discovery of novel endothelial cell-specific findings for several critical cardiovascular organs. Capillary endothelial cells are the primary source of vascular heterogeneity across organs, and findings from the past 5 years have shown that they are also the primary source of transcriptional changes in multiple disease states. A general trend observed in the liver, lungs and brain is that disease promotes a loss of specialized capillary function and a reversal to a more general transcriptional state. These findings have implications for the development of targeted therapies and for the differentiation of improved organ-specific vasculature for tissue-engineering applications.

Published online: 05 October 2022

1. World Health Organization. *The Top 10 Causes of Death* (WHO, 2020).
2. Paik, D. T. et al. Single-cell RNA sequencing unveils unique transcriptomic signatures of organ-specific endothelial cells. *Circulation* **142**, 1848–1862 (2020).
3. Elmentaite, R., Domínguez Conde, C., Yang, L. & Teichmann, S. A. Single-cell atlases: shared and tissue-specific cell types across human organs. *Nat. Rev. Genet.* <https://doi.org/10.1038/s41576-022-00449-w> (2022).
4. den Braanker, H., van Stigt, A. C., Kok, M. R., Lubberts, E. & Bisoendial, R. J. Single-cell RNA sequencing reveals heterogeneity and functional diversity of lymphatic endothelial cells. *Int. J. Mol. Sci.* <https://doi.org/10.3390/ijms222111976> (2021).
5. Corada, M., Morini, M. F. & Dejana, E. Signaling pathways in the specification of arteries and veins. *Arterioscler. Thromb. Vasc. Biol.* **34**, 2372–2377 (2014).
6. Fish, J. E. & Wythe, J. D. The molecular regulation of arteriovenous specification and maintenance. *Dev. Dyn.* **244**, 391–409 (2015).
7. Dejana, E., Hirschi, K. K. & Simons, M. The molecular basis of endothelial cell plasticity. *Nat. Commun.* **8**, 14361 (2017).
8. Qiu, J. & Hirschi, K. K. Endothelial cell development and its application to regenerative medicine. *Circ. Res.* **125**, 489–501 (2019).
9. Apte, R. S., Chen, D. S. & Ferrara, N. VEGF in signaling and disease: beyond discovery and development. *Cell* **176**, 1248–1264 (2019).
10. Phng, L.-K. & Gerhardt, H. Angiogenesis: a team effort coordinated by notch. *Dev. Cell* **16**, 196–208 (2009).
11. You, L.-R. et al. Suppression of Notch signalling by the COUP-TFII transcription factor regulates vein identity. *Nature* **435**, 98–104 (2005).
12. Lawson, N. D., Vogel, A. M. & Weinstein, B. M. sonic hedgehog and vascular endothelial growth factor act upstream of the Notch pathway during arterial endothelial differentiation. *Dev. Cell* **3**, 127–136 (2002).
13. Benedito, R. & Hellström, M. Notch as a hub for signaling in angiogenesis. *Exp. Cell Res.* **319**, 1281–1288 (2013).
14. Kohli, V. et al. Arterial and venous progenitors of the major axial vessels originate at distinct locations. *Dev. Cell* **25**, 196–206 (2013).
15. Casie Chetty, S. et al. Vegf signaling promotes vascular endothelial differentiation by modulating etv2 expression. *Dev. Biol.* **424**, 147–161 (2017).
16. Zygmunt, T. et al. In parallel interconnectivity of the dorsal longitudinal anastomotic vessels requires both VEGF signaling and circulatory flow. *J. Cell Sci.* **125**, 5159–5167 (2012).
17. Bahary, N. et al. Duplicate VegfA genes and orthologues of the KDR receptor tyrosine kinase family mediate vascular development in the zebrafish. *Blood* **110**, 3627–3636 (2007).
18. Covassin, L. D., Villefranc, J. A., Kacergis, M. C., Weinstein, B. M. & Lawson, N. D. Distinct genetic interactions between multiple Vegf receptors are required for development of different blood vessel types in zebrafish. *Proc. Natl Acad. Sci. USA* **103**, 6554–6559 (2006).
19. Bussmann, J., Wolfe, S. A. & Siekmann, A. F. Arterial-venous network formation during brain vascularization involves hemodynamic regulation of chemokine signaling. *Development* **138**, 1717–1726 (2011).
20. Ehling, M., Adams, S., Benedito, R. & Adams, R. H. Notch controls retinal blood vessel maturation and quiescence. *Development* **140**, 3051–3061 (2013).
21. Hellström, M. et al. Dll4 signalling through Notch 1 regulates formation of tip cells during angiogenesis. *Nature* **445**, 776–780 (2007).
22. Leslie, J. D. et al. Endothelial signalling by the Notch ligand delta-like 4 restricts angiogenesis. *Development* **134**, 839–844 (2007).
23. Lobov, I. B. et al. Delta-like ligand 4 (Dll4) is induced by VEGF as a negative regulator of angiogenic sprouting. *Proc. Natl Acad. Sci. USA* **104**, 3219–3224 (2007).
24. Noguera-Troise, I. et al. Blockade of Dll4 inhibits tumour growth by promoting non-productive angiogenesis. *Nature* **444**, 1032–1037 (2006).
25. Suchting, S. et al. The Notch ligand delta-like 4 negatively regulates endothelial tip cell formation and vessel branching. *Proc. Natl Acad. Sci. USA* **104**, 3225–3230 (2007).
26. Red-Horse, K. & Siekmann, A. F. Veins and arteries build hierarchical branching patterns differently: bottom-up versus top-down. *BioEssays* **41**, 1800198 (2019).
27. Siekmann, A. F., Covassin, L. & Lawson, N. D. Modulation of VEGF signalling output by the Notch pathway. *BioEssays* **30**, 303–313 (2008).
28. Fang, J. S. et al. Shear-induced Notch-Cx37-p27 axis arrests endothelial cell cycle to enable arterial specification. *Nat. Commun.* <https://doi.org/10.1038/s41467-017-01742-7> (2017).
29. Su, T. et al. Single-cell analysis of early progenitor cells that build coronary arteries. *Nature* **559**, 356–362 (2018).
30. Luo, W. et al. Arterialization requires the timely suppression of cell growth. *Nature* **589**, 437–441 (2021).
31. Chavkin, N. W. et al. *Endothelial Cell Cycle State Determines Propensity for Arterial-Venous Fate* (Cold Spring Harbor Laboratory, 2020).
32. Sissaoui, S. et al. Genomic characterization of endothelial enhancers reveals a multifunctional role for NR2F2 in regulation of arteriovenous gene expression. *Circ. Res.* **126**, 875–888 (2020).
33. Sabbagh, M. F. et al. Transcriptional and epigenomic landscapes of CNS and non-CNS vascular endothelial cells. *eLife* <https://doi.org/10.7554/elife.36187> (2018).
34. McDonald, A. I. et al. Endothelial regeneration of large vessels is a biphasic process driven by local cells with distinct proliferative capacities. *Cell Stem Cell* **23**, 210–225.e6 (2018).
35. Pitulescu, M. E. et al. Dll4 and Notch signalling couples sprouting angiogenesis and artery formation. *Nat. Cell Biol.* **19**, 915–927 (2017).
36. Hasan, S. S. et al. Endothelial Notch signalling limits angiogenesis via control of artery formation. *Nat. Cell Biol.* **19**, 928–940 (2017).
37. Quillien, A. et al. Distinct Notch signaling outputs pattern the developing arterial system. *Development* **141**, 1544–1552 (2014).
38. Geudens, I. et al. Artery-vein specification in the zebrafish trunk is pre-patterned by heterogeneous Notch activity and balanced by flow-mediated fine tuning. *Development* **146**, dev181024 (2019).
39. Lee, H.-W. et al. Role of venous endothelial cells in developmental and pathologic angiogenesis. *Circulation* **144**, 1308–1322 (2021).
40. Xu, C. et al. Arteries are formed by vein-derived endothelial tip cells. *Nat. Commun.* **5**, 5758 (2014).
41. Blanco, R. & Gerhardt, H. VEGF and Notch in tip and stalk cell selection. *Cold Spring Harb. Perspect. Med.* **3**, a006569 (2013).
42. Tian, X. et al. De novo formation of a distinct coronary vascular population in neonatal heart. *Science* **345**, 90–94 (2014).
43. Pontes-Quero, S. et al. High mitogenic stimulation arrests angiogenesis. *Nat. Commun.* <https://doi.org/10.1038/s41467-019-09875-7> (2019).
44. Park, H. et al. Defective flow-migration coupling causes arteriovenous malformations in hereditary hemorrhagic telangiectasia. *Circulation* **144**, 805–822 (2021).
45. Hou, S. et al. Heterogeneity in endothelial cells and widespread venous arterialization during early vascular development in mammals. *Cell Res.* <https://doi.org/10.1038/s41422-022-00615-z> (2022).

46. Das, S. et al. A unique collateral artery development program promotes neonatal heart regeneration. *Cell* **176**, 1128–1142.e18 (2019).
47. Zarkada, G. et al. Specialized endothelial tip cells guide neuroretina vascularization and blood-retina-barrier formation. *Dev. Cell* **56**, 2237–2251.e6 (2021).
48. Tang, J. et al. Extension of endocardium-derived vessels generate coronary arteries in neonates. *Circ. Res.* **130**, 352–365 (2022).
49. Roman, B. L. & Hinck, A. P. ALK1 signaling in development and disease: new paradigms. *Cell. Mol. Life Sci.* **74**, 4539–4560 (2017).
50. Rochon, E. R., Menon, P. G. & Roman, B. L. Alk1 controls arterial endothelial cell migration in lumenized vessels. *Development* **143**, 2593–2602 (2016).
51. Jin, Y. et al. Endoglin prevents vascular malformation by regulating flow-induced cell migration and specification through VEGFR2 signalling. *Nat. Cell Biol.* **19**, 639–652 (2017).
52. Poduri, A. et al. Endothelial cells respond to the direction of mechanical stimuli through SMAD signaling to regulate coronary artery size. *Development* **144**, 3241–3252 (2017).
53. Capasso, T. L. et al. BMP10-mediated ALK1 signaling is continuously required for vascular development and maintenance. *Angiogenesis* **23**, 203–220 (2020).
54. Sugden, W. W. & Siekmann, A. F. Endothelial cell biology of Endoglin in hereditary hemorrhagic telangiectasia. *Curr. Opin. Hematol.* **25**, 237–244 (2018).
55. Deng, H. et al. Activation of Smad2/3 signaling by low fluid shear stress mediates artery inward remodeling. *Proc. Natl Acad. Sci. USA* <https://doi.org/10.1073/pnas.2105339118> (2021).
56. Kwon, H.-B. et al. In vivo modulation of endothelial polarization by Apelin receptor signalling. *Nat. Commun.* **7**, 11805 (2016).
57. Weijts, B. et al. Blood flow-induced Notch activation and endothelial migration enable vascular remodeling in zebrafish embryos. *Nat. Commun.* **9**, 5314 (2018).
58. Rosa, A. et al. WASp controls oriented migration of endothelial cells to achieve functional vascular patterning. *Development* **149**, dev200195 (2022).
59. Chang, A. H. et al. DACH1 stimulates shear stress-guided endothelial cell migration and coronary artery growth through the CXCL12-CXCR4 signaling axis. *Genes Dev.* **31**, 1308–1324 (2017).
60. Raffrey, B. et al. Dach1 extends artery networks and protects against cardiac injury. *Circ. Res.* <https://doi.org/10.1161/circresaha.120.318271> (2021).
61. Franco, C. A. et al. Non-canonical Wnt signalling modulates the endothelial shear stress flow sensor in vascular remodelling. *eLife* **5**, e07727 (2016).
62. Franco, C. A. et al. Dynamic endothelial cell rearrangements drive developmental vessel regression. *PLoS Biol.* **13**, e1002125 (2015).
63. Neal, A. et al. Venous identity requires BMP signalling through ALK3. *Nat. Commun.* **10**, 453 (2019).
64. Payne, S. et al. Regulatory pathways governing murine coronary vessel formation are dysregulated in the injured adult heart. *Nat. Commun.* <https://doi.org/10.1038/s41467-019-10710-2> (2019).
65. Vanlandewijck, M. et al. A molecular atlas of cell types and zonation in the brain vasculature. *Nature* **554**, 475–480 (2018).
66. Phansalkar, R. et al. Coronary blood vessels from distinct origins converge to equivalent states during mouse and human development. *eLife* <https://doi.org/10.7554/eLife.70246> (2021).
67. Garcia, F. J. et al. Single-cell dissection of the human brain vasculature. *Nature* <https://doi.org/10.1038/s41586-022-04521-7> (2022).
68. Winkler, E. A. et al. A single-cell atlas of the normal and malformed human brain vasculature. *Science* **375**, eabi7377 (2022).
69. Wälchli, T. et al. *Molecular Atlas of the Human Brain Vasculature at the Single-Cell Level* (Cold Spring Harbor Laboratory, 2021).
70. Augustin, H. G. & Koh, G. Y. Organotypic vasculature: from descriptive heterogeneity to functional pathophysiology. *Science* **357**, eaal2379 (2017).
71. Stoeckius, M. et al. Simultaneous epitope and transcriptome measurement in single cells. *Nat. Methods* **14**, 865–868 (2017).
72. Zhu, C., Preissl, S. & Ren, B. Single-cell multimodal omics: the power of many. *Nat. Methods* **17**, 11–14 (2020).
73. Han, X. et al. Mapping the mouse cell atlas by Microwell-Seq. *Cell* **172**, 1091–1107.e17 (2018).
74. He, L. et al. Single-cell RNA sequencing of mouse brain and lung vascular and vessel-associated cell types. *Sci. Data* **5**, 180160 (2018).
75. Cusanovich, D. A. et al. A single-cell atlas of in vivo mammalian chromatin accessibility. *Cell* **174**, 1309–1324.e18 (2018).
76. Domcke, S. et al. A human cell atlas of fetal chromatin accessibility. *Science* <https://doi.org/10.1126/science.aba7612> (2020).
77. Cui, Y. et al. Single-cell transcriptome analysis maps the developmental track of the human heart. *Cell Rep.* **26**, 1934–1950.e5 (2019).
78. Litviňuková, M. et al. Cells of the adult human heart. *Nature* **588**, 466–472 (2020).
79. Zepp, J. A. et al. Genomic, epigenomic, and biophysical cues controlling the emergence of the lung alveolus. *Science* **371**, eabc3172 (2021).
80. Guo, M. et al. Single cell RNA analysis identifies cellular heterogeneity and adaptive responses of the lung at birth. *Nat. Commun.* <https://doi.org/10.1038/s41467-018-07770-1> (2019).
81. Hurskainen, M. et al. Single cell transcriptomic analysis of murine lung development on hyperoxia-induced damage. *Nat. Commun.* <https://doi.org/10.1038/s41467-021-21865-2> (2021).
82. Travaglini, K. J. et al. A molecular cell atlas of the human lung from single-cell RNA sequencing. *Nature* **587**, 619–625 (2020).
83. Schupp, J. C. et al. Integrated single-cell atlas of endothelial cells of the human lung. *Circulation* **144**, 286–302 (2021).
84. Hupe, M. et al. Gene expression profiles of brain endothelial cells during embryonic development at bulk and single-cell levels. *Sci. Signal.* **10**, eaag2476 (2017).
85. Park, J. et al. Single-cell transcriptomics of the mouse kidney reveals potential cellular targets of kidney disease. *Science* **360**, 758–763 (2018).
86. Magella, B. et al. Cross-platform single cell analysis of kidney development shows stromal cells express Gdnf. *Dev. Biol.* **434**, 36–47 (2018).
87. Bais, A. S. et al. Single-cell RNA sequencing reveals differential cell cycle activity in key cell populations during nephrogenesis. *Sci. Rep.* **11**, 22434 (2021).
88. Miao, Z. et al. Single cell regulatory landscape of the mouse kidney highlights cellular differentiation programs and disease targets. *Nat. Commun.* **12**, 2277 (2021).
89. Young, M. D. et al. Single-cell transcriptomes from human kidneys reveal the cellular identity of renal tumors. *Science* **361**, 594–599 (2018).
90. Menon, R. et al. Single-cell analysis of progenitor cell dynamics and lineage specification in the human fetal kidney. *Development* <https://doi.org/10.1242/dev.164038> (2018).
91. Hochane, M. et al. Single-cell transcriptomics reveals gene expression dynamics of human fetal kidney development. *PLoS Biol.* **17**, e3000152 (2019).
92. Cao, J. et al. A human cell atlas of fetal gene expression. *Science* <https://doi.org/10.1126/science.aba7721> (2020).
93. The Tabula Sapiens Consortium The Tabula Sapiens: a multiple-organ, single-cell transcriptomic atlas of humans. *Science* **376**, eabi4896 (2022).
94. Calvanese, V. et al. Mapping human haematopoietic stem cells from haemogenic endothelium to birth. *Nature* **604**, 534–540 (2022).
95. Tabula Muris Consortium et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367–372 (2018).
96. Muto, Y. et al. Single cell transcriptional and chromatin accessibility profiling redefine cellular heterogeneity in the adult human kidney. *Nat. Commun.* **12**, 2190 (2021).
97. Gómez-Salineró, J. M. et al. Specification of fetal liver endothelial progenitors to functional zoned adult sinusoids requires c-Maf induction. *Cell Stem Cell* **29**, 593–609.e7 (2022).
98. Halpern, K. B. et al. Paired-cell sequencing enables spatial gene expression mapping of liver endothelial cells. *Nat. Biotechnol.* **36**, 962–970 (2018).
99. Su, T. et al. Single-cell transcriptomics reveals zone-specific alterations of liver sinusoidal endothelial cells in cirrhosis. *Cell. Mol. Gastroenterol. Hepatol.* **11**, 1139–1161 (2021).
100. Aizarani, N. et al. A human liver cell atlas reveals heterogeneity and epithelial progenitors. *Nature* **572**, 199–204 (2019).
101. MacParland, S. A. et al. Single cell RNA sequencing of human liver reveals distinct intrahepatic macrophage populations. *Nat. Commun.* <https://doi.org/10.1038/s41467-018-06318-7> (2018).
102. Nolan, D. J. et al. Molecular signatures of tissue-specific microvascular endothelial cell heterogeneity in organ maintenance and regeneration. *Dev. Cell* **26**, 204–219 (2013).
103. Feng, W., Chen, L., Nguyen, P. K., Wu, S. M. & Li, G. Single cell analysis of endothelial cells identified organ-specific molecular signatures and heart-specific cell populations and molecular features. *Front. Cardiovasc. Med.* <https://doi.org/10.3389/fcvm.2019.00165> (2019).
104. Kalucka, J. et al. Single-cell transcriptome atlas of murine endothelial cells. *Cell* **180**, 764–779.e20 (2020).
105. Paik, D. T., Cho, S., Tian, L., Chang, H. Y. & Wu, J. C. Single-cell RNA sequencing in cardiovascular development, disease and medicine. *Nat. Rev. Cardiol.* **17**, 457–473 (2020).
106. Brulois, K. et al. A molecular map of murine lymph node blood vascular endothelium at single cell resolution. *Nat. Commun.* **11**, 3798 (2020).
107. Chen, M. B. et al. Brain endothelial cells are exquisite sensors of age-related circulatory cues. *Cell Rep.* **30**, 4418–4432.e4 (2020).
108. Zhao, L. et al. Pharmacologically reversible zonation-dependent endothelial cell transcriptomic changes with neurodegenerative disease associations in the aged brain. *Nat. Commun.* <https://doi.org/10.1038/s41467-020-18249-3> (2020).
109. Gutierrez, M. E. C., Hill, M. C., Largoza, G., Martin, J. F. & Wythe, J. D. Defining the transcriptional and epigenetic basis of organotypic endothelial diversity in the developing and adult mouse. *bioRxiv* <https://doi.org/10.1101/2021.11.15.468651> (2021).
110. Neal, A. et al. ETS factors are required but not sufficient for specific patterns of enhancer activity in different endothelial subtypes. *Dev. Biol.* **473**, 1–14 (2021).
111. Yucel, N. et al. Cardiac endothelial cells maintain open chromatin and expression of cardiomyocyte myofibrillar genes. *eLife* <https://doi.org/10.7554/eLife.55730> (2020).
112. Liu, Y. et al. A specialized bone marrow microenvironment for fetal haematopoiesis. *Nat. Commun.* **13**, 1327 (2022).
113. Tikhonova, A. N. et al. The bone marrow microenvironment at single-cell resolution. *Nature* **569**, 222–228 (2019).
114. Mäe, M. A. et al. Single-cell analysis of blood-brain barrier response to pericyte loss. *Circ. Res.* <https://doi.org/10.1161/circresaha.120.317473> (2021).
115. Shi, W. et al. Cardiac proteomics reveals sex chromosome-dependent differences between males and females that arise prior to gonad formation. *Dev. Cell* **56**, 3019–3034.e3017 (2021).
116. Red-Horse, K., Ueno, H., Weissman, I. L. & Krasnow, M. A. Coronary arteries form by developmental reprogramming of venous cells. *Nature* **464**, 549–553 (2010).
117. Wu, B. et al. Endocardial cells form the coronary arteries by angiogenesis through myocardial-endocardial VEGF signaling. *Cell* **151**, 1085–1096 (2012).
118. Chen, H. I. et al. The sinus venosus contributes to coronary vasculature through VEGFC-stimulated angiogenesis. *Development* **141**, 4500–4512 (2014).
119. Zhang, H. et al. Endocardium minimally contributes to coronary endothelium in the embryonic ventricular free walls. *Circ. Res.* **118**, 1880–1893 (2016).
120. Sharma, B. et al. Alternative progenitor cells compensate to rebuild the coronary vasculature in Elabela- and Apj-deficient hearts. *Dev. Cell* **42**, 655–666.e3 (2017).
121. Sedmera, D., Pexieder, T., Vuillemin, M., Thompson, R. P. & Anderson, R. H. Developmental patterning of the myocardium. *Anat. Rec.* **258**, 319–337 (2000).
122. Lu, P. et al. Perinatal angiogenesis from pre-existing coronary vessels via DLL4–NOTCH1 signalling. *Nat. Cell Biol.* **23**, 967–977 (2021).
123. D'Amato, G. et al. *Endocardium-to-coronary Artery Differentiation During Heart Development and Regeneration Involves Sequential Roles of Bmp2 and Cxcl12/Cxcr4* (Cold Spring Harbor Laboratory, 2021).
124. Jiang, Z. et al. Overexpression of Kdr in adult endocardium induces endocardial neovascularization and improves heart function after myocardial infarction. *Cell Res.* **31**, 485–487 (2021).
125. Räsänen, M. et al. VEGF-B promotes endocardium-derived coronary vessel development and cardiac regeneration. *Circulation* **143**, 65–77 (2021).
126. Li, H. et al. Classifying Drosophila olfactory projection neuron subtypes by single-cell RNA sequencing. *Cell* **171**, 1206–1220.e22 (2017).
127. Dick, S. A. et al. Self-renewing resident cardiac macrophages limit adverse remodeling following myocardial infarction. *Nat. Immunol.* **20**, 29–39 (2019).
128. Weinreb, C., Rodriguez-Fraticelli, A., Camargo, F. D. & Klein, A. M. Lineage tracing on transcriptional

- landscapes links state to fate during differentiation. *Science* **367**, eaaw3381 (2020).
129. McCracken, I. R. et al. Mapping the developing human cardiac endothelium at single-cell resolution identifies MECOM as a regulator of arteriovenous gene expression. *Cardiovasc. Res.* <https://doi.org/10.1093/cvr/cvac023> (2022).
 130. Vila Ellis, L. et al. Epithelial Vegfa specifies a distinct endothelial population in the mouse lung. *Dev. Cell* **52**, 617–630.e6 (2020).
 131. Gillich, A. et al. Capillary cell-type specialization in the alveolus. *Nature* **586**, 785–789 (2020).
 132. Negretti, N. M. et al. A single-cell atlas of mouse lung development. *Development* <https://doi.org/10.1242/dev.199512> (2021).
 133. Niethamer, T. K. et al. Defining the role of pulmonary endothelial cell heterogeneity in the response to acute lung injury. *eLife* <https://doi.org/10.7554/elife.53072> (2020).
 134. Wang, S. et al. A single-cell transcriptomic landscape of the lungs of patients with COVID-19. *Nat. Cell Biol.* **23**, 1314–1328 (2021).
 135. McCracken, I. R. et al. Lack of evidence of angiotensin-converting enzyme 2 expression and replicative infection by SARS-CoV-2 in human endothelial cells. *Circulation* **143**, 865–868 (2021).
 136. Delorey, T. M. et al. COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature* **595**, 107–113 (2021).
 137. Sauler, M. et al. Characterization of the COPD alveolar niche using single-cell RNA sequencing. *Nat. Commun.* <https://doi.org/10.1038/s41467-022-28062-9> (2022).
 138. Watanabe, C. et al. Aging of the vascular system and neural diseases. *Front. Aging Neurosci.* <https://doi.org/10.3389/fnagi.2020.557384> (2020).
 139. Yang, A. C. et al. A human brain vascular atlas reveals diverse mediators of Alzheimer's risk. *Nature* <https://doi.org/10.1038/s41586-021-04369-3> (2022).
 140. Daneman, R., Zhou, L., Kebede, A. A. & Barres, B. A. Pericytes are required for blood–brain barrier integrity during embryogenesis. *Nature* **468**, 562–566 (2010).
 141. Armulik, A. et al. Pericytes regulate the blood–brain barrier. *Nature* **468**, 557–561 (2010).
 142. Jourde-Chiche, N. et al. Endothelium structure and function in kidney health and disease. *Nat. Rev. Nephrol.* **15**, 87–108 (2019).
 143. Dumas, S. J. et al. Single-cell RNA sequencing reveals renal endothelium heterogeneity and metabolic adaptation to water deprivation. *J. Am. Soc. Nephrol.* **31**, 118–138 (2020).
 144. Brunskill, E. W. & Potter, S. S. Gene expression programs of mouse endothelial cells in kidney development and disease. *PLoS ONE* **5**, e12034 (2010).
 145. Karaiskos, N. et al. A single-cell transcriptome atlas of the mouse glomerulus. *J. Am. Soc. Nephrol.* **29**, 2060–2068 (2018).
 146. Takemoto, M. et al. Large-scale identification of genes implicated in kidney glomerulus development and function. *EMBO J.* **25**, 1160–1174 (2006).
 147. Fu, J. et al. Single-cell RNA profiling of glomerular cells shows dynamic changes in experimental diabetic kidney disease. *J. Am. Soc. Nephrol.* **30**, 533–545 (2019).
 148. Chung, J.-J. et al. Single-cell transcriptome profiling of the kidney glomerulus identifies key cell types and reactions to injury. *J. Am. Soc. Nephrol.* **31**, 2341–2354 (2020).
 149. Aird, W. C. Phenotypic heterogeneity of the endothelium. *Circ. Res.* **100**, 174–190 (2007).
 150. Inverso, D. et al. A spatial vascular transcriptomic, proteomic, and phosphoproteomic atlas unveils an angiocrine Tie–Wnt signaling axis in the liver. *Dev. Cell* **56**, 1677–1693.e10 (2021).
 151. Shirure, V. S., Hughes, C. C. W. & George, S. C. Engineering vascularized organoid-on-a-chip models. *Annu. Rev. Biomed. Eng.* **23**, 141–167 (2021).
 152. Ang, L. T. et al. Generating human artery and vein cells from pluripotent stem cells highlights the arterial tropism of Nipah and Hendra viruses. *Cell* **185**, 2523–2541.e30 (2022).
 153. Wimmer, R. A. et al. Human blood vessel organoids as a model of diabetic vasculopathy. *Nature* **565**, 505–510 (2019).
 154. Takebe, T. et al. Vascularized and functional human liver from an iPSC-derived organ bud transplant. *Nature* **499**, 481–484 (2013).
 155. Takebe, T. et al. Massive and reproducible production of liver buds entirely from human pluripotent stem cells. *Cell Rep.* **21**, 2661–2670 (2017).
 156. Van Den Berg, C. W. et al. Renal subcapsular transplantation of PSC-derived kidney organoids induces neo-vasculogenesis and significant glomerular and tubular maturation in vivo. *Stem Cell Rep.* **10**, 751–765 (2018).
 157. Mansour, A. A. et al. An in vivo model of functional and vascularized human brain organoids. *Nat. Biotechnol.* **36**, 432–441 (2018).
 158. Holloway, E. M. et al. Differentiation of human intestinal organoids with endogenous vascular endothelial cells. *Dev. Cell* **54**, 516–528.e7 (2020).
 159. Garreta, E. et al. Rethinking organoid technology through bioengineering. *Nat. Mater.* **20**, 145–155 (2021).
 160. Lewis-Israeli, Y. R. et al. Self-assembling human heart organoids for the modeling of cardiac development and congenital heart disease. *Nat. Commun.* **12**, 5142 (2021).
 161. Hofbauer, P. et al. Cardioids reveal self-organizing principles of human cardiogenesis. *Cell* **184**, 3299–3317.e22 (2021).
 162. Drakhlis, L. et al. Human heart-forming organoids recapitulate early heart and foregut development. *Nat. Biotechnol.* **39**, 737–746 (2021).
 163. Homan, K. A. et al. Flow-enhanced vascularization and maturation of kidney organoids in vitro. *Nat. Methods* **16**, 255–262 (2019).
 164. Campisi, M. et al. 3D self-organized microvascular model of the human blood-brain barrier with endothelial cells, pericytes and astrocytes. *Biomaterials* **180**, 117–129 (2018).
 165. Skylar-Scott, M. A. et al. Biomanufacturing of organ-specific tissues with high cellular density and embedded vascular channels. *Sci. Adv.* **5**, eaaw2459 (2019).

Acknowledgements

E.T. is supported by the Fannie and John Hertz Foundation.

Author contributions

The authors contributed substantially to all aspects of the article.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Cardiology thanks Bin Zhou and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

RELATED LINKS

Tabula Muris: <https://tabula-muris.ds.czbiohub.org/>

© Springer Nature Limited 2022