




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Narrative review

COVID-19 pandemic—a focused review for clinicians

M. Cevik^{1,*}, C.G.G. Bamford², A. Ho³¹) Division of Infection and Global Health Research, School of Medicine, University of St Andrews, UK²) Wellcome–Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK³) MRC-University of Glasgow Centre for Virus Research, University of Glasgow, Glasgow, UK

ARTICLE INFO

Article history:

Received 8 April 2020

Received in revised form

19 April 2020

Accepted 21 April 2020

Available online 25 April 2020

Editor: L Leibovici

Keywords:

Coronavirus

COVID-19

Novel coronavirus

SARS-CoV-2

ABSTRACT

Background: The COVID-19 pandemic caused by SARS-CoV-2 remains a significant issue for global health, economics and society. A wealth of data has been generated since its emergence in December 2019, and it is vital for clinicians to keep up with this data from across the world at a time of uncertainty and constantly evolving guidelines and clinical practice.

Objectives: Here we provide an update for clinicians on the recent developments in the virology, diagnostics, clinical presentation, viral shedding, and treatment options for COVID-19 based on current literature.

Sources: We considered published peer-reviewed papers and non-peer-reviewed pre-print manuscripts on COVID-19 and related aspects with an emphasis on clinical management aspects.

Content: We describe the virological characteristics of SARS-CoV-2 and the clinical course of COVID-19 with an emphasis on diagnostic challenges, duration of viral shedding, severity markers and current treatment options.

Implications: The key challenge in managing COVID-19 remains patient density. However, accurate diagnosis as well as early identification and management of high-risk severe cases are important for many clinicians. For improved management of cases, there is a need to understand test probability of serology, qRT-PCR and radiological testing, and the efficacy of available treatment options that could be used in severe cases with a high risk of mortality. **M. Cevik, Clin Microbiol Infect 2020;26:842**

© 2020 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Introduction

The first cases of atypical pneumonia of unidentified aetiology were reported on December 30, 2019, from Wuhan, China. By January 7, 2020, a novel betacoronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), was identified, while the disease has been named COVID-19. COVID-19 has now been declared a pandemic, affecting nearly every country, with over 2.3 million confirmed cases and >160 000 deaths. The initial clinical case series from China largely comprised hospitalized patients with severe pneumonia. Further data suggested that approximately 80% of the patients have mild disease, 20% require hospital admission, and approximately 5% require intensive care admission [1]. Mortality rates are higher among people over 60 years of age and with

coexisting conditions (most commonly hypertension, diabetes and cardiovascular disease). Here we provide an update for clinicians on the recent developments in virology, diagnostics, clinical presentation, and treatment options for COVID-19 based on current literature.

Virology

Metagenomic sequencing and targeted real-time polymerase chain reaction (qRT-PCR) assays identified a novel human CoV (SARS-CoV-2) in bronchoalveolar lavage fluid taken from the initial patient cluster in Wuhan [2]. Infectious SARS-CoV-2 has been cultured on monkey Vero, human Huh7 and primary human airway epithelial cells [3], where it is cytopathic. Furthermore, serum antibodies (IgM and IgG) from cases neutralized SARS-CoV-2 in cell culture and detected virus-infected cells by indirect immunofluorescence [3].

Phylogenetic analysis reveals that SARS-CoV-2 is closely related to SARS-CoV (~80% similar) in the *Sarbecovirus* subfamily (genus

* Corresponding author. M. Cevik, Division of Infection and Global Health Research, School of Medicine, University of St Andrews, Fife, KY16 9TF, UK.

E-mail address: mc349@st-andrews.ac.uk (M. Cevik).

Betacoronavirus) [2]. While an intermediate host has yet to be determined, it shares strong genetic similarity (>95%) to known bat coronaviruses from China, suggesting a likely bat origin. Relatively similar coronaviruses have been found in pangolins, but in these the receptor-binding domain (RBD) of the spike (S) glycoprotein is more like that of SARS-2-CoV-2 than that of known bat viruses [4].

SARS-CoV-2 shares most of its gene content with SARS-CoV, including the S glycoprotein, the RNA-dependent RNA polymerase (Nsp12), and two proteases: papain-like protease (PLpro) and 3C-like protease (3CLpro) [3]. There is also substantial antigenic cross-reactivity between SARS-CoV-2 and SARS-CoV [3,5]. A recent study confirmed that the angiotensin-converting enzyme 2 (ACE2), expressed in the human respiratory tract epithelium, is the entry receptor for SARS-CoV-2, similar to SARS-CoV, and has been shown to cause pneumonia in lab mice only expressing human ACE2 [6,7]. This is likely mediated by the RBD of the S glycoprotein [8]. Although there is obvious homology between SARS-CoV and SARS-CoV-2, and cross neutralization has been observed [9], significant biological differences—specifically in the S glycoprotein—have been noted [5,10,11].

Clinical presentation

A key difference between COVID-19 and seasonal influenza-associated pneumonia is the potential severity of the disease even in young adults without comorbidities [12]. In a study that compared three well-conducted Chinese case series to a reference group of patients with influenza-associated pneumonia from 73 German sentinel hospitals, the severity of the pneumonia—even in adults aged <60 years without chronic preconditions—was significantly greater in COVID-19. For instance, 28% of COVID-19 patients treated on the ICU had no reported comorbidity. The rate of acute respiratory distress syndrome (ARDS) and mechanical ventilation was markedly higher among COVID-19 patients. The median duration of ventilation was 9 days for non-invasive and 17 days for invasive ventilation [12].

Across all studies, the most common symptoms at onset of illness were fever, cough, fatigue, and myalgia. However, available data suggest that only half the patients are febrile at the time of admission [13,14]. Gastrointestinal symptoms—including anorexia, nausea, vomiting and diarrhoea—are also common, reported in nearly 40% of patients in some cohorts [15,16]. Furthermore, up to 10% of patients present with gastrointestinal symptoms without respiratory symptoms or fever [17]. COVID-19 has been associated with a hypercoagulable state with increased risk of venous thromboembolism [18]. Neurological manifestations (including headache, dizziness, and altered consciousness), ischaemic and haemorrhagic strokes, as well as muscle injury have also been reported [19]. A third of patients in a small Italian cohort reported taste or olfactory disorders, including anosmia [20]. Other extrapulmonary manifestations include skin and ocular manifestations. An Italian study reported cutaneous manifestations in 20% of patients [21]. Ocular manifestations consistent with conjunctivitis were reported in 32% of COVID patients in a Chinese case series [22].

The estimated mean incubation period is reported to be 3–6 days (range 1.3–11.3 days) [12]. The duration from symptom onset to dyspnoea was 5–6 days [13,17]. On average, disease progresses further, requiring hospitalization, at 7–8 days from symptom onset. Patients may initially appear relatively stable, but they often deteriorate rapidly with severe hypoxia [13,17]. The key feature seen in these cases is ARDS [13,17]. The interval from symptom onset to the development of ARDS is approximately 8–12 days [13]. In addition, the incidence of cardiovascular manifestations such as myocardial injury seems to be high, likely due to

the systemic inflammatory response and immune system disorders during disease progression [23].

Illness severity and development of ARDS are associated with older age and underlying medical conditions [17]. Additionally, neutrophilia and raised lactate dehydrogenase and D-dimer, lymphocyte counts, CD3 and CD4 T cell counts, aspartate aminotransferase (AST), prealbumin, creatinine, glucose, low-density lipoprotein, serum ferritin, and prothrombin time were also associated with higher risk of severe disease and ARDS [17]. In a cohort of 191 patients with a definitive clinical outcome (137 discharged and 54 died), mortality was independently associated with older age, higher qSOFA score, and d-dimer >1 µg/mL on admission, and the majority had severe disease and experienced complications such as ARDS, acute kidney injury and sepsis [13]. Factors most associated with critical illness were admission oxygen saturation <88%, first d-dimer >2500, first ferritin >2500, and first CRP >200 [24]. Furthermore, patients with cardiovascular disease were shown to be more likely to develop severe symptoms [23] in keeping with the picture seen in MERS-CoV and SARS.

In comparison, most children appear to have mild disease. Among 1391 asymptomatic and symptomatic children (median age: 6.7 years) with known COVID-19 contact in Wuhan Children's Hospital [25], 171 (12.3%) were SARS-CoV-2-positive; 27 (15.8%) had no symptoms or radiological features of pneumonia, 33 (19.3%) had upper respiratory symptoms, and 64.9% had pneumonia. Three patients (with coexisting conditions) required intensive care and one died.

In terms of co-infections, a pre-print examining >8000 samples of COVID-19 contacts tested for SARS-CoV2 in China reported viral co-infections in 5.8% of COVID-19-positive individuals (including seasonal coronaviruses, influenza A virus and rhinoviruses) [26]. Another study of 1206 patients identified viral co-infection in 24 of 116 (21%) SARS-CoV2-positive patients; rhino/enterovirus, respiratory syncytial virus, and seasonal CoVs were the most common [27]. Bacterial and fungal co-infections with SARS-CoV-2 have been documented, especially in the ICU setting, including *Acinetobacter baumannii* and *Klebsiella pneumoniae* [28]. Among 191 patients, non-survivors were more likely to have sepsis based on qSOFA score and secondary infection, although detailed bacteriology results were not reported [13]. Secondary infection and positive association between steroid administration and secondary infection should be explored further.

Molecular and serological diagnosis

The first genome sequence for SARS-CoV-2 was released on virological.org on 10 January 2020 (GenBank accession number MN908947). This allowed the rapid development of several sensitive and specific qRT-PCR assays [29]. Many laboratories worldwide are now able to test for SARS-CoV-2. Assays have been described that can detect fewer than ten copies of SARS-CoV-2 per reaction and will not cross-react with SARS-CoV or other human coronaviruses [29]. However, the sensitivity and specificity of these tests remain unknown and there is no clear consensus on which is preferred.

Viral RNA loads by qRT-PCR were substantially higher in sputum than in throat swabs [3,30,31], suggesting that the type of sample may also influence the outcome of the test. Therefore, submission of both lower and upper respiratory tract samples is currently advised.

Precise molecular detection is hampered by the variability in viral loads in the upper respiratory tract, especially at later stages of infection. In a study from China, among 241 COVID-19 patients with at least one positive SARS-CoV-2 qRT-PCR test result, in the first test 384 (63.0%) were negative [32]. In addition, several tests at

different points from the same patient varied during the course of diagnosis and treatment [32]. Therefore, a single positive test should be confirmed by a second qRT-PCR assay targeting a different SARS-CoV-2 gene. However, similar studies in Taiwan and Hong Kong reported fewer false negatives [33]. Second, a single negative SARS-CoV-2 test (especially if from an upper respiratory tract specimen) or a positive test result for another respiratory pathogen should not be used to exclude COVID-19 infection. These findings indicate that qRT-PCR has a low probability of ruling out an infection, and in clinically highly suspicious cases repeat sampling and also CT imaging may need to be used to guide the diagnosis.

Antibody-based methods to detect seroconversion in serum or plasma based upon enzyme-linked immunosorbent assays (ELISA), indirect immunofluorescence or virus neutralization have been reported [34–36]. Around 40–50% of patients develop an antibody response to SARS-CoV-2 infection after 7 days, and the majority by 14 days [35,37]. S1 has been shown to be more specific than S as an antigen for SARS-CoV-2 in serological diagnosis [36]. The commercial S1 IgG and IgA assays have lower specificity, but with IgA showing higher sensitivity [36]. Recently, an ELISA assay based on detection of recombinant S protein by serum antibodies demonstrated robust and scalable determination of seroconversion that will facilitate screening of potentially exposed individuals for evidence of past infection [38]. Since seroconversion occurs relatively late in infection, rapid antibody tests have a limited role in the diagnosis of acute infection; qRT-PCR remains the reference standard.

Work is ongoing to understand protective antibody levels and immunological markers. Among 175 recovered laboratory-confirmed COVID-19 patients, neutralizing antibodies (NAb) peaked at 10–15 days after disease onset. However, approximately 30% failed to develop a good level of NAb titres (ID50 < 500) [39]. In addition, patients who did not generate NAb at the time of discharge did not develop NAb thereafter. These results highlight the fact that some patients with SARS-CoV-2 will recover without developing high titres of virus-specific NAb. These findings have some implications for vaccine development and also for convalescent plasma treatment as the donor plasma should be titrated before use in passive therapy. There is less information available on the T-cell response during SARS-CoV-2 infection and how it correlates with the NAb titres.

Duration of viral shedding and isolation period

SARS-CoV-2 RNA has been identified by qRT-PCR in respiratory tract samples 1–2 days prior to symptom onset and can persist for 7–12 days in moderate cases and for up to 2 weeks in severe cases [35,40]. SARS-CoV-2 has also been detected in whole blood [41], saliva [42], faeces [43], and urine [44] by qRT-PCR (Table 1). In several case series with serial sampling, viral loads were highest soon after symptom onset [35,45]. Patients with severe COVID-19

had a significantly higher viral load and a longer period of viral shedding than mild cases [46]. Prolonged viral RNA shedding has been reported from throat swabs for up to 37 days among adult patients [13], and in faeces for over 1 month after illness onset in children [40,47]. However, detection of viral RNA by qRT-PCR does not necessarily equate with infectious virus. No live virus was cultured from throat swabs or sputum from nine mild COVID-19 cases beyond day 8 after symptom onset despite an ongoing high viral load [35]. Persistently high levels of RNA were also identified in the stool of the mild cases, but no live virus was cultured [35]. These findings suggest that patients may continue to shed RNA in various samples for a long period, but this does not equate to infectiousness potential (Table 1). This supports the current guidance of 7–14 days self-isolation from symptom onset. Certain hospitals following a protocol to confirm viral clearance prior to transfer out of dedicated COVID-19 wards; however, this may not be required given the prolonged RNA shedding without evidence of viable virus. However, further exploration is required on whether faecal–oral or faecal–respiratory transmission occurs, and the role of shedding in transmission in severe cases.

Transmission patterns

A review of modelling studies based on Chinese case numbers reported a median basic reproduction number (R_0) of 2.79 [48], although an R_0 as high as 5.7 has been reported [49]. These estimates are substantially higher than the reproduction number for seasonal influenza (~1.3) [50], and indicate that control measures would need to prevent >60% transmission to stop the epidemic. Of note, R_0 will vary by setting, and can be substantially reduced by countermeasures, as observed in China [51].

It is now clear that a significant proportion of individuals with COVID-19 have very mild or no symptoms. Asymptomatic infection at the time of laboratory testing has been reported [52,53], although a large proportion go on to develop symptoms. For instance, among 55 asymptomatic carriers with positive qRT-PCR for SARS-CoV-2 in pharyngeal swab samples, 14 went on to develop mild, 39 ordinary, and two severe COVID-19 [54]. There have been several reports of SARS-CoV-2 transmission from asymptomatic or pre-symptomatic persons [55,56], which poses significant challenges to contact tracing. Nevertheless, the relative contribution of asymptomatic or pre-symptomatic transmission on the overall transmission dynamics of the pandemic remains uncertain. Thus, household studies of secondary human transmission of SARS-CoV-2 and serosurveys to determine the incidence of asymptomatic and subclinical infections are needed.

A further consideration is the superspreading event, whereby a small number of cases are responsible for a disproportionate number of secondary cases. This was a feature of both SARS- and MERS-CoV, and was responsible for multiple nosocomial outbreaks [57,58]. Several superspreading events have been reported for

Table 1
Transmission routes

Source	Mode of transmission	RNA by PCR (Days since onset of symptoms)	Viable virus (Days since onset of symptoms)
Nasopharynx	Droplet	Up to 37 days	Up to 7 days (in mild cases)
Sputum	Droplet/airborne during aerosol-producing procedures	Up to 37 days	Up to 7 days (in mild cases)
Stool	No evidence of faecal–oral transmission	>30 days	Only 1 report; uncertain
Blood	No viable virus to date	Up to 14 days	No
Urine	No viable virus to date	No	No
Conjunctiva	No viable virus to date		
Macaques with corneal inoculation develop infection	Yes	No	
Vertical	No strong evidence of vertical transmission to date	No	N/A

COVID-19 [17]. Rapid identification and mitigation of these events will be crucial to controlling this pandemic.

Treatment options in clinical trials

At present, there are no approved antivirals for SARS-CoV-2. Several antivirals that have shown promise against SARS- or MERS-CoV *in vitro* and *in vivo* are currently being evaluated in clinical trials for COVID-19. Lopinavir/ritonavir (LPV/r), a protease inhibitor used as an antiretroviral, showed inconclusive results for the treatment of SARS, but demonstrated strong *in vitro* and *in vivo* antiviral activity against MERS-CoV when combined with interferon- β (IFN β) [59]. The first of a number of clinical trials involving LPV/r was recently published [60]. Among 199 seriously ill laboratory-confirmed COVID-19 patients, no significant difference in clinical improvement, mortality or viral clearance was observed between the LPV/r ($n = 99$) and standard care ($n = 100$) arms. However, treatment was instituted late in infection; median time from symptom onset to treatment was 13 days, and >40% of patients had undetectable viral load before or during treatment. The results were complicated by the variable use of other treatments, including interferon, glucocorticoids and antibiotics. Of note, day 28 mortality was lower (not significantly) in those with early treatment (19% versus 27%), and those who received LPV/r also had lower vasopressor and non-invasive ventilation use.

Another promising drug is remdesivir, a novel nucleotide analogue that interferes with nsp12 polymerase [61]. It has shown *in vitro* activity against a wide range of RNA viruses, including SARS and MERS-CoV [62,63], and has also demonstrated superior antiviral activity compared to LPV/r-IFN β against MERS-CoV in a mouse model [59]. Against SARS-CoV-2, it has shown promising antiviral activity in Vero E6 cells and Huh7 cells [64]. Remdesivir has been given to a small number of patients with severe COVID-19 as compassionate use; however, given the lack of randomization and a control group, the findings are difficult to interpret [65]. There are ongoing RCTs worldwide assessing its efficacy and safety in patients with COVID-19, and a study in France is evaluating its impact on viral shedding in high- and moderate-risk contacts in confirmed COVID-19 cases (NCT04259892).

Other candidate antivirals being studied in RCTs include favipiravir and hydroxychloroquine, which has been shown to inhibit virus cell entry *in vitro* [66]. Hydroxychloroquine (HCQ), an analogue of chloroquine, has demonstrated anti-SARS-CoV-2 activity *in vitro* [67]. Among a small open-label non-randomized study, patients treated with HCQ and HCQ + azithromycin showed viral load reduction compared to controls. However, there have been significant concerns and ethical issues about the trial content, its ethical approval, and the peer review process prior to publication raised by several physicians and also by the International Society of Antimicrobial Chemotherapy. In a small RCT of HCQ ($n = 30$) there was no change in viral load or clinical outcome after 7 days [68]. Currently, there are 45 trials evaluating chloroquine or HCQ for the treatment and prophylaxis of COVID-19, including multi-centre RCTs in the UK (RECOVERY, ISRCTN50189673), Europe (Dis-CoVeRY, NCT04315948) and globally (involving >70 countries) (SOLIDARITY, ISRCTN83971151).

Host-targeted therapeutic options are also being explored, such as inhibition of human cytokine interleukin-6 (IL-6), the abundance of which has been associated with a worse prognosis [69]. A pre-print including 21 patients who received tocilizumab (an IL-6 receptor inhibitor) reported improvement in symptoms, hypoxaemia and CT changes in the majority of patients [70]. There are ongoing RCTs evaluating tocilizumab and sarilumab, also an IL-6 receptor inhibitor. With insufficient evidence of efficacy for any existing treatments, the Infectious Diseases Society of America (IDSA)

recommends that experimental therapies should only be offered to patients in the context of a clinical trial [71].

There is no licenced vaccine to protect against COVID-19. However, a number of experimental candidates are in development, with some already in early clinical trials. Most vaccine candidates focus on immunization with only the S glycoprotein, which is the major target for neutralization antibodies. Candidate vaccines differ in the mode of S delivery, and platforms dependent on recombinant protein, mRNA or viral vectored approaches are being tested. Passive immunization through transfusion of convalescent sera or plasma containing neutralizing antibodies from recovered donors have been reported in several case series, with clinical improvement reported in recipients [72,73]. Clinical trials evaluating convalescent plasma as treatment for severe COVID-19 are ongoing.

Conclusion

A wealth of data has already been generated on COVID-19 since early January 2020. Nevertheless, key questions remain regarding understanding the population at risk and age groups, the proportion of individuals who have had asymptomatic infections and their transmission potential, the endemicity and seasonality of COVID-19, and whether stringent physical distancing measures will be effective in countries outside China. The main challenge in managing COVID-19 remains the patient density; however, accurate diagnoses as well as early identification and management of high-risk severe cases remains a daily battle for many clinicians. For improved management of cases, there is a need to understand test probability of serology, qRT-PCR and radiological testing, and the efficacy of available treatment options that could be used in severe cases with high risk of mortality.

Author contributions

MC, CM, AH drafted the first and subsequent versions of the manuscript, and all authors provided critical feedback and contributed to the manuscript.

Transparency declaration

The authors have declared no conflicts of interest. No funding was received for this review.

References

- [1] Team TNCPERE. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) — China. 2020. Available at: <http://weekly.chinacdc.cn/fileCCDCW/journal/article/ccdcw/newcreate/COVID-19.pdf>. [Accessed 18 February 2020].
- [2] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020;382:727–33.
- [3] Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–3.
- [4] Lam TT-Y, Shum MH-H, Zhu H-C, Tong Y-G, Ni X-B, Liao Y-S, et al. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature* 2020. epub ahead of print.
- [5] Tian X, Li C, Huang A, Xia S, Lu S, Shi Z, et al. Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *bioRxiv* 2020. 2020.01.28.923011. epub ahead of print.
- [6] Bao L, Deng W, Huang B, Gao H, Ren L, Wei Q, et al. The pathogenicity of 2019 novel coronavirus in hACE2 transgenic mice. *bioRxiv* 2020. 2020.02.07.939389. epub ahead of print.
- [7] Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, et al. Severe acute respiratory syndrome-related coronavirus: the species and its viruses — a statement of the Coronavirus Study Group. *bioRxiv* 2020. 2020.02.07.937862. epub ahead of print.

- [8] Letko M, Munster V. Functional assessment of cell entry and receptor usage for lineage B β -coronaviruses, including 2019-nCoV. *bioRxiv* 2020. 2020.01.22.915660. epub ahead of print.
- [9] Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181:271–80.
- [10] Lei C, Fu W, Qian K, Li T, Zhang S, Ding M, et al. Potent neutralization of 2019 novel coronavirus by recombinant ACE2-Ig. *bioRxiv* 2020. 2020.02.01.929976. epub ahead of print.
- [11] Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *bioRxiv* 2020. 2020.02.11.944462. epub ahead of print.
- [12] Tolksdorf K, Buda S, Schuler E, Wieler LH, Haas W. Influenza-associated pneumonia as reference to assess seriousness of coronavirus disease (COVID-19). *Euro Surveill* 2020;25.
- [13] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
- [14] Kluytmans M, Buiting A, Pas S, Bentvelsen R, van den Bijllaardt W, van Oudheusden A, et al. SARS-CoV-2 infection in 86 healthcare workers in two Dutch hospitals in March 2020. *medRxiv* 2020. 2020.03.23.20041913. epub ahead of print.
- [15] Liang W, Feng Z, Rao S, Xiao C, Xue X, Lin Z, et al. Diarrhoea may be underestimated: a missing link in 2019 novel coronavirus. *Gut* 2020. [gutjnl-2020-320832](https://doi.org/10.1136/gutjnl-2020-320832). epub ahead of print.
- [16] Zhang J-j, Dong X, Cao Y-y, Yuan Y-d, Yang Y-b, Yan Y-q, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 2020. <https://doi.org/10.1111/all.14238>.
- [17] Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 Hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061–9.
- [18] Danzi GB, Loffi M, Galeazzi G, Gherbesi E. Acute pulmonary embolism and COVID-19 pneumonia: a random association? *Eur Heart J* 2020. <https://doi.org/10.1093/eurheartj/ehaa25>.
- [19] Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, et al. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol* 2020. <https://doi.org/10.1001/jamaneurol.2020.1127>.
- [20] Giacomelli A, Pezzati L, Conti F, Bernacchia D, Siano M, Oreni L, et al. Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa330>.
- [21] Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16387>.
- [22] Wu P, Duan F, Luo C, Liu Q, Qu X, Liang L, et al. Characteristics of ocular findings of patients with coronavirus disease 2019 (COVID-19) in Hubei Province, China. *JAMA Ophthalmol* 2020. <https://doi.org/10.1001/jamaophthalmol.2020.1291>.
- [23] Zheng Y-Y, Ma Y-T, Zhang J-Y, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol* 2020;17:259–60.
- [24] Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell LF, Chernyak Y, et al. Factors associated with hospitalization and critical illness among 4,103 patients with COVID-19 disease in New York City. *medRxiv* 2020. 2020.04.08.20057794. epub ahead of print.
- [25] Lu X, Zhang L, Du H, Zhang J, Li YY, Qu J, et al. SARS-CoV-2 Infection in children. *N Engl J Med* 2020;382:1663–5.
- [26] Wang M, Wu Q, Xu W, Qiao B, Wang J, Zheng H, et al. Clinical diagnosis of 8274 samples with 2019-novel coronavirus in Wuhan. *medRxiv* 2020. 2020.02.12.20022327. epub ahead of print.
- [27] Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of co-infection between SARS-CoV-2 and other respiratory pathogens. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.6266>.
- [28] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507–13.
- [29] Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020;25.
- [30] Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;395:514–23.
- [31] Han H, Luo Q, Mo F, Long L, Zheng W. SARS-CoV-2 RNA more readily detected in induced sputum than in throat swabs of convalescent COVID-19 patients. *Lancet Infect Dis* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30174-2](https://doi.org/10.1016/S1473-3099(20)30174-2).
- [32] Li Y, Yao L, Li J, Chen L, Song Y, Cai Z, et al. Stability issues of RT-PCR testing of SARS-CoV-2 for hospitalized patients clinically diagnosed with COVID-19. *J Med Virol* 2020. <https://doi.org/10.1002/jmv.25786>.
- [33] Chan JF-W, Yip CC-Y, To KK-W, Tang TH-C, Wong SC-Y, Leung K-H, et al. Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/HEP real-time reverse transcription-polymerase chain reaction assay validated in vitro and with clinical specimens. *J Clin Microbiol* 2020. [JCM.00310-20](https://doi.org/10.1128/JCM.00310-20). epub ahead of print.
- [34] Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *bioRxiv* 2020. 2020.01.22.914952. epub ahead of print.
- [35] Wolfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Muller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020.
- [36] Okba NMA, Muller MA, Li W, Wang C, GeurtsvanKessel CH, Corman VM, et al. SARS-CoV-2 specific antibody responses in COVID-19 patients. *medRxiv* 2020. 2020.03.18.20038059. epub ahead of print.
- [37] Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *medRxiv* 2020. 2020.03.02.20030189. epub ahead of print.
- [38] Amanat F, Nguyen T, Chromikova V, Strohmeier S, Stadlbauer D, Javier A, et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *medRxiv* 2020. 2020.03.17.20037713. epub ahead of print.
- [39] Wu F, Wang A, Liu M, Wang Q, Chen J, Xia S, et al. Neutralizing antibody responses to SARS-CoV-2 in a COVID-19 recovered patient cohort and their implications. *medRxiv* 2020. 2020.03.30.20047365. epub ahead of print.
- [40] Cai J, Xu J, Lin D, Yang Z, Xu L, Qu Z, et al. A case series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa198>.
- [41] Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA* 2020;323:1488–94. <https://doi.org/10.1001/jama.2020.3204>.
- [42] To KK-W, Tsang OT-Y, Yip CC-Y, Chan K-H, Wu T-C, Chan JM-C, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa149>.
- [43] Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.3786>.
- [44] Peng L, Liu J, Xu W, Luo Q, Deng K, Lin B, et al. 2019 Novel coronavirus can be detected in urine, blood, anal swabs and oropharyngeal swabs samples. *medRxiv* 2020. 2020.02.21.20026179. epub ahead of print.
- [45] Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382:1177–9.
- [46] Liu Y, Yan L-M, Wan L, Xiang T-X, Le A, Liu J-M, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30232-2](https://doi.org/10.1016/S1473-3099(20)30232-2).
- [47] Xu Y, Li X, Zhu B, Liang H, Fang C, Gong Y, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 2020;26:502–5.
- [48] Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* 2020;27.
- [49] Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High contagiousness and rapid spread of severe acute respiratory syndrome coronavirus 2. *Emerg Infect Dis* 2020;26.
- [50] Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic review of the literature. *BMC Inf Dis* 2014;14:480.
- [51] Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *Lancet Infect Dis* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30144-4](https://doi.org/10.1016/S1473-3099(20)30144-4).
- [52] Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020. *Euro Surveill* 2020;25.
- [53] Ki M. Epidemiologic characteristics of early cases with 2019 novel coronavirus (2019-nCoV) disease in Korea. *Epidemiol Health* 2020;42. e2020007-0.
- [54] Wang Y, Liu Y, Liu L, Wang X, Luo N, Li L. Clinical outcomes in 55 patients with severe acute respiratory syndrome coronavirus 2 who were asymptomatic at hospital admission in Shenzhen, China. *J Infect Dis* 2020. <https://doi.org/10.1093/infdis/jiaa119>.
- [55] Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, et al. Asymptomatic and presymptomatic SARS-CoV-2 infections in residents of a long-term care skilled nursing facility—king County, Washington, March 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:377–81.
- [56] Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic transmission of SARS-CoV-2—Singapore, January 23—march 16, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:411–5.
- [57] Shen Z, Ning F, Zhou W, He X, Lin C, Chin DP, et al. Superspreading SARS events, Beijing, 2003. *Emerg Infect Dis* 2004;10:256–60.
- [58] Kim KH, Tandil TE, Choi JW, Moon JM, Kim MS. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak in South Korea, 2015: epidemiology, characteristics and public health implications. *J Hosp Infect* 2017;95:207–13.
- [59] Sheahan TP, Sims AC, Leist SR, Schafer A, Won J, Brown AJ, et al. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nat Commun* 2020;11:222.
- [60] Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020. <https://doi.org/10.1056/NEJMoa2001282>.
- [61] Agostini ML, Andres EL, Sims AC, Graham RL, Sheahan TP, Lu X, et al. Coronavirus susceptibility to the antiviral remdesivir (GS-5734) is mediated by the viral polymerase and the proofreading exoribonuclease. *mBio* 2018;9. e00221-18.

- [62] Warren TK, Jordan R, Lo MK, Ray AS, Mackman RL, Soloveva V, et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature* 2016;531:381–5.
- [63] Sheahan TP, Sims AC, Graham RL, Menachery VD, Gralinski LE, Case JB, et al. Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. *Sci Transl Med* 2017;9.
- [64] Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* 2020;30:269–71.
- [65] Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, et al. Compassionate use of remdesivir for patients with severe Covid-19. *New Eng J Med* 2020. <https://doi.org/10.1056/NEJMoa2007016>.
- [66] Keyaerts E, Vijgen L, Maes P, Neyts J, Van Ranst M. In vitro inhibition of severe acute respiratory syndrome coronavirus by chloroquine. *Biochem Biophys Res Commun* 2004;323:264–8.
- [67] Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020. ciaa237. epub ahead of print.
- [68] Chen JLD, Liu L, Liu P, Xu Q, Xia L, Ling Y, et al. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J Zhejiang Univ (Med Sci)* 2020;49.
- [69] Liu T, Zhang J, Yang Y, Ma H, Li Z, Zhang J, et al. The potential role of IL-6 in monitoring severe case of coronavirus disease 2019. medRxiv 2020. 2020.03.01.20029769. epub ahead of print.
- [70] Gritti G, Raimondi F, Ripamonti D, Riva I, Landi F, Alborghetti L, et al. Use of siltuximab in patients with COVID-19 pneumonia requiring ventilatory support. medRxiv 2020. 2020.04.01.20048561. epub ahead of print.
- [71] Adarsh B, Shumaker AH, Lavergne V, Baden L, Chi-Chung Cheng V, Edwards KM, et al. Infectious diseases society of America guidelines on the treatment and management of patients with COVID-19. IDSA; 2020.
- [72] Duan K, Liu B, Li C, Zhang H, Yu T, Qu J, et al. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proc Nat Acad Sci* 2020: 202004168.
- [73] Shen C, Wang Z, Zhao F, Yang Y, Li J, Yuan J, et al. Treatment of 5 critically ill patients with COVID-19 with convalescent plasma. *JAMA* 2020;323:1582–9.