

Air pollution, traffic noise, greenness, and temperature and the risk of incident type 2 diabetes

Results from the KORA cohort study

Mahnaz Badpa^{a,b,*}, Alexandra Schneider^a, Lars Schwettmann^{c,d}, Barbara Thorand^{a,b,e}, Kathrin Wolf^a, Annette Peters^{a,b,e}

Introduction: Type 2 diabetes (T2D) is a major public health concern, and various environmental factors have been associated with the development of this disease. This study aimed to investigate the longitudinal effects of multiple environmental exposures on the risk of incident T2D in a German population-based cohort.

Methods: We used data from the KORA cohort study (Augsburg, Germany) and assessed exposure to air pollutants, traffic noise, greenness, and temperature at the participants' residences. Cox proportional hazard models were used to analyze the associations with incident T2D, adjusting for potential confounders.

Results: Of 7736 participants included in the analyses, 10.5% developed T2D during follow-up (mean: 15.0 years). We found weak or no association between environmental factors and the risk of T2D, with sex and education level significantly modifying the effects of air pollutants.

Conclusion: Our study contributes to the growing body of literature investigating the impact of environmental factors on T2D risks and suggests that the impact of environmental factors may be small.

Keywords: Air pollution; Greenness; Traffic noise; Temperature; Type 2 diabetes; Environmental epidemiology; Population-based cohort

^aInstitute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Munich, Germany; ^bInstitute for Medical Information Processing, Biometry and Epidemiology (IBE), Faculty of Medicine, Pettenkofer School of Public Health, LMU Munich, Munich, Germany; ^cInstitute of Health Economics and Health Care Management, Helmholtz Zentrum München, Munich, Germany; ^dDepartment of Health Services Research, School of Medicine and Health Sciences, Carl von Ossietzky University of Oldenburg, Oldenburg, Germany; and ^eGerman Center for Diabetes Research (DZD), Partner München-Neuherberg, Neuherberg, Germany

Kathrin Wolf and Annette Peters shared last authorship.

The KORA study was initiated and financed by the Helmholtz Zentrum München – German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research (BMBF) and by the State of Bavaria. Data collection in the KORA study is done in cooperation with the University Hospital of Augsburg. The funders had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

The data are subject to national data protection laws. Therefore, data cannot be made freely available in a public repository. However, data can be requested through an individual project agreement with KORA. To obtain permission to use KORA data under the terms of a project agreement, please use the digital tool KORA.PASST (<https://www.helmholtz-munich.de/en/epi/cohort/kora>).

M.B. did the data analyses and wrote the manuscript. M.B., A.S., L.S., B.T., K.W., and A.P. have verified the underlying data and reviewed the manuscript. A.S., K.W., and A.P. were involved in the study design, results interpretation, and review of the manuscript. A.P. provided oversight on the KORA study design. K.W. was involved in the acquisition of the KORA data, the modeling of air pollution concentrations as well as the assignment of all exposure data. All authors read and approved the final manuscript.

SDC Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.environepidem.com).

*Corresponding Author, Address: Institute of Epidemiology, Helmholtz Munich, Ingolstädter Landstrasse 1, 85764 Neuherberg, Germany.
E-mail: Mahnaz.badpa@campus.lmu.de (M. Badpa).

Introduction

Type 2 diabetes (T2D) is projected to become the 7th highest cause of mortality by 2030.^{1,2} Alongside well-known risk factors such as obesity, physical inactivity, and genetic predisposition, several environmental factors have been linked to T2D risk in previous studies.^{3,4}

Four reviews^{5–8} compiled data from epidemiological studies and suggest a positive association between long-term exposure to particulate matter (PM) and nitrogen dioxide (NO₂), and prevalence and incidence of T2D. Similarly, meta-analyses suggested a positive correlation between exposure to traffic noise, particularly during nighttime, and a higher risk of T2D incidence.^{9,10} Moreover, access to green spaces, as measured by either the proportion of greenness or the proximity to green areas, has shown beneficial effects on reducing the likelihood of developing T2D.¹¹ The association of temperature with T2D risk and fasting plasma glucose levels is, however, complex, with some studies suggesting increased risk with higher temperatures,¹² while others indicated U-shaped relationships involving both high and low temperatures.¹³

Despite these findings, studies assessing multiple environmental exposures are still scarce, and there is a lack of longitudinal investigations examining the link with T2D incidence. In this article, we

What the study adds

This study provides critical insights into the link between environmental exposures and T2D. It emphasizes the significance of exposure to PM_{2.5}, highlights the need for cautious interpretation due to uncertainties, and underscores the sex-specific variations in pollutant effects on T2D risk. These findings contribute to our understanding of T2D risk factors.

aimed to investigate the longitudinal effects of various exposures on T2D incidence in a German population-based cohort.

Methods

Study population

We used data from the third (KORA S3, 1994–1995, $n = 4856$) and fourth (KORA S4, 1999–2001, $n = 4260$) survey of the KORA (Cooperative Health Research in the Region of Augsburg) study¹⁴ and follow-up information until 2016. For our analysis, we excluded participants who requested data withdrawal ($N = 80$), with a diabetes diagnosis at baseline ($N = 381$), lacked follow-up data ($N = 521$), with unknown residential address ($N = 270$), or had missing values in the main model's covariates ($N = 128$) (Figure S1, <http://links.lww.com/EE/A267>).

Incident T2D assessment

T2D incidence was assessed through self-reported, clinically diagnosed diabetes assessed through follow-up questionnaires that were validated by physicians, medical chart review, or self-reported use of glucose-lowering medication.^{15,16} Self-reported dates of diagnosis, were verified through medical records or physician contact. (Text S1, <http://links.lww.com/EE/A267>).

Exposure assessment

Annual mean concentrations of air pollutants, including nitrogen oxides (NO_2 , NO_x), ozone (O_3), PM in different size classes (PM_{10} [$\leq 10 \mu\text{m}$], $\text{PM}_{2.5}$ [$\leq 2.5 \mu\text{m}$], $\text{PM}_{2.5\text{absorbance}}$ [$\text{PM}_{2.5\text{sabs}}$] as an indicator for soot, $\text{PM}_{\text{coarse}}$ [$2.5\text{--}10 \mu\text{m}$]), and particle number concentration (PNC) as an indicator for ultrafine particles were estimated using land-use regression models for 2014–2015.¹⁷ Traffic noise exposure was estimated for 2011 using the noise and air pollution information system for Augsburg city and georeferenced pictures for rural areas.¹⁸ Greenness within 300 m and 1000 m buffers was assessed using normalized difference vegetation index (NDVI) from Landsat 5 Thematic Mapper satellite images for 1994–1995 (S3) and 1999–2001 (S4).¹⁹ Annual mean and standard deviation (SD) air temperature were extracted from temperature maps developed within a multi-stage modeling approach for 2000.²⁰ All exposures were assigned to the participants' residential baseline addresses.

Statistical methods

We used the Cox proportional hazards model to analyze the association between environmental factors and incident T2D, with follow-up time as the underlying timescale and an indicator variable for subcohort (S3/S4). Participants were censored at the time of the event, withdrawal request, death, emigration, loss to follow-up, or the end of the study period, whichever came first.

Three models with varying degrees of confounder adjustment were specified a priori based on previous research²¹ and data availability. The minimum model included age, sex, and subcohort indicator, the main model was further adjusted for other baseline characteristics (body mass index [BMI], smoking behavior, alcohol consumption, education level, physical activity, and dietary score²²), and the extended model additionally included clinical information (cardiovascular diseases (CVD), waist-hip ratio, and cholesterol level) (Text S2, <http://links.lww.com/EE/A267>).

In premodels, we tested nonlinearity but observed no considerable deviance from linearity. Therefore, exposure variables were included as linear terms (Figure S2, <http://links.lww.com/EE/A267>).

Effect modification was assessed by including an interaction term between each exposure variable and potential effect modifiers (sex, age, obesity, educational level, and physical activity) (Figure S3, <http://links.lww.com/EE/A267>). Results are presented as hazard ratios (HRs) per interquartile range (IQR) increase in exposure variables with 95% confidence intervals (CIs).

We used R 3.6.1 with “survival,” “mgcv,” and “raster” packages for all statistical analyses.

Sensitivity analysis

To evaluate the robustness of our findings, we used an alternative statistical model (Poisson regression) and applied the main model separately for the two subcohorts.

Results

Altogether, 7864 (86.3%) participants out of 9116 participants were included in our study (Figure S1, <http://links.lww.com/EE/A267>). Of these, 10.5% developed T2D until the end of follow-up (Table 1). At baseline, the mean age was 49.2 years, almost half of the participants were male (49.2%), and the mean BMI was 27.0 kg/m². About a quarter of the participants reported being active smokers, and 47.0% reported being physically active (Table 1).

Annual average concentrations of NO_2 , PM_{10} , and $\text{PM}_{2.5}$ at participants' residences were below the European air quality standard values but exceeded the current WHO guideline values. The mean levels of traffic noise were above the European-recommended maximum values (Table 2).

Although some of the environmental exposures showed positive effect estimates (e.g. $\text{PM}_{2.5}$ showing the strongest effect estimate) confidence limits were large as estimated with the main model (Figure 1). The incorporation of additional covariates in the models had little to moderate impact on the HRs and 95% CIs compared to the minimum adjusted model. Sex modified the effect of NO_x , O_3 , PM_{10} , and PNC on T2D incidence, with O_3 being more pronounced in females and the others in males (Figure S3, <http://links.lww.com/EE/A267>). The effect of $\text{PM}_{2.5\text{sabs}}$ was modified by educational level.

Poisson models provided similar associations (Figure S4, <http://links.lww.com/EE/A267>) as the main Cox model. Also, stratified analyses by subcohort showed comparable results (Figure S5, <http://links.lww.com/EE/A267>) to the main pooled model.

Discussion

Our longitudinal analysis showed weak or no associations between the various environmental factors and T2D incidence, after adjusting for confounding factors. These results were consistent across different analytical approaches and stratified analyses of the two subcohorts. Considering the lack of significant associations in the single-exposure models, we refrained from conducting multi-exposure models.

Copyright © 2024 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The Environmental Epidemiology. All rights reserved. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Environmental Epidemiology (2024) 8:e302

Received 17 October, 2023; Accepted 12 February, 2024

Published online 11 March 2024

DOI: 10.1097/EE9.0000000000000302

Table 1.
Baseline characteristics of study participants

| Variable | KORA S3 + S4 (n = 7864) ^a | | KORA S3 (n = 4042) ^a | | KORA S4 (n = 3822) ^a | |
|-------------------------------|--------------------------------------|---------------------|---------------------------------|---------------------|---------------------------------|--|
| | Missing n (%) | Mean ± SD/ n (%) | Mean ± SD/ n (%) | Mean ± SD/ n (%) | Mean ± SD/ n (%) | |
| Incident T2D during follow-up | 0 (0.0) | 829 (10.5) | 486 (12.0) | 343 (9.0) | | |
| Age (years) | 0 (0.0) | 49.2 ± 13.8 | 49.6 ± 13.9 | 48.8 ± 13.7 | | |
| Sex (male) | 0 (0.0) | 3,873 (49.2) | 2020 (50.0) | 1852 (48.5) | | |
| BMI (kg/m ²) | 71 (0.9) | 27.0 ± 4.5 | 27.0 ± 4.4 | 27.0 ± 4.6 | | |
| Waist-hip ratio | 55 (0.7) | 0.9 ± 0.1 | 0.9 ± 0.1 | 0.9 ± 0.1 | | |
| Education level | 8 (0.1) | | | | | |
| Low | | 1148 (14.6) | 676 (16.7) | 472 (12.4) | | |
| Middle | | 5671 (72.2) | 2895 (71.6) | 2776 (72.8) | | |
| High | | 1037 (13.2) | 470 (11.6) | 567 (14.9) | | |
| Smoking status | 5 (0.1) | | | | | |
| Current smoker | | 2012 (25.6) | 1013 (25.1) | 999 (26.2) | | |
| Ex-smoker | | 2486 (31.6) | 1243 (30.8) | 1243 (32.6) | | |
| Never smoker | | 3361 (42.8) | 1786 (44.2) | 1575 (41.3) | | |
| Alcohol intake (g/day) | 17 (0.2) | 16.3 ± 22.1 | 16.8 ± 22.8 | 15.9 ± 21.3 | | |
| Physical activity (active) | 18 (0.2) | 3687 (47.0) | 1784 (44.1) | 1903 (50.0) | | |
| Dietary score | 21 (0.3) | 15.3 ± 3.6 | 15.3 ± 3.6 | 15.2 ± 3.6 | | |
| Cardiovascular disease (yes) | 17 (0.2) | 3057 (39.0) | 1674 (41.4) | 1383 (36.3) | | |
| Total cholesterol (mg/dL) | 139 (1.8) | 229.2 ± 43.7 | 230.8 ± 44.0 | 227.4 ± 43.3 | | |

^an refers to the sample size before excluding participants with missing confounders used in the main model.

BMI indicates body mass index; KORA, Cooperative Health Research in the Region of Augsburg; S3, Third cross-sectional health survey of the KORA cohort; S4, Fourth cross-sectional health survey of the KORA cohort; SD, standard deviation; T2D, type 2 diabetes.

Table 2.
Descriptive statistics of annual average air pollutant concentrations, traffic noise, NDVI, and air temperature at residences

| Variable | | KORA S3 + S4 (n = 7864) ^a | | KORA S3 (n = 4042) ^a | | KORA S4 (n = 3822) ^a | |
|---------------|---|--------------------------------------|---------------------|---------------------------------|---------------------|---------------------------------|--|
| | | Mean ± SD | Mean ± SD/ n (%) | Mean ± SD/ n (%) | Mean ± SD/ n (%) | | |
| Air pollutant | NO ₂ (µg/m ³) | 14.2 ± 4.4 | 14.1 ± 4.3 | 14.5 ± 4.5 | | | |
| | NO _x (µg/m ³) | 22.1 ± 7.1 | 22.0 ± 6.9 | 22.2 ± 7.4 | | | |
| | O ₃ (µg/m ³) | 38.8 ± 2.4 | 38.8 ± 2.4 | 38.9 ± 2.5 | | | |
| | PM ₁₀ (µg/m ³) | 16.6 ± 1.5 | 16.5 ± 1.5 | 16.7 ± 1.5 | | | |
| | PM _{2.5} (µg/m ³) | 11.8 ± 1.0 | 11.8 ± 1.0 | 11.8 ± 1.0 | | | |
| | PM _{2.5,abs} (10 ⁻⁵ /m) | 1.2 ± 0.2 | 1.2 ± 0.2 | 1.2 ± 0.2 | | | |
| | PM _{coarse} (µg/m ³) | 4.9 ± 1.0 | 4.8 ± 1.0 | 5.0 ± 1.0 | | | |
| | PNC (10 ⁹ /cm ³) | 7.3 ± 1.8 | 7.2 ± 1.8 | 7.3 ± 1.8 | | | |
| Noise | Daily average traffic noise (dB) | 54.7 ± 6.6 | 54.6 ± 6.6 | 54.9 ± 6.6 | | | |
| | Nighttime average traffic noise (dB) | 45.7 ± 6.4 | 45.6 ± 6.3 | 45.8 ± 6.4 | | | |
| Greenness | NDVI in 300 m buffer | 0.5 ± 0.1 | 0.4 ± 0.1 | 0.5 ± 0.1 | | | |
| | NDVI in 1000 m buffer | 0.5 ± 0.1 | 0.5 ± 0.1 | 0.5 ± 0.1 | | | |
| Temperature | Annual mean temperature (°C) | 10.6 ± 0.4 | 10.6 ± 0.4 | 10.6 ± 0.4 | | | |
| | Annual SD temperature (°C) | 6.1 ± 0.2 | 6.0 ± 0.2 | 6.1 ± 0.2 | | | |

The EU air quality standard values are 40 µg/m³ for PM₁₀ and NO₂, and 25 µg/m³ for PM_{2.5}. The WHO air quality guideline values are 10 µg/m³ for NO₂, 15 µg/m³ for PM₁₀, and 5 µg/m³ for PM_{2.5}. The WHO air quality guideline value for O₃ is 60 µg/m³, which is calculated based on the peak season. The EU-recommended maximum value for traffic noise during night is 40 dB in residential areas.

^an refers to the sample size before excluding participants with missing confounders used in the main model.

NDVI values below 0 were excluded since they represent water or bare rocks.

KORA indicates Cooperative Health Research in the Region of Augsburg; NDVI, normalized difference vegetation index; NO₂, nitrogen dioxide; NO_x, nitrogen oxide; O₃, ozone; PM₁₀, particulate matter with an aerodynamic diameter ≤10 µm; PM_{2.5}, particulate matter with an aerodynamic diameter ≤2.5 µm; PM_{2.5,abs}, PM_{2.5} absorbance; PM_{coarse}, particulate matter with an aerodynamic diameter of 2.5–10 µm; PNC, particle number concentration; S3, Third cross-sectional health survey of the KORA cohort; S4, Fourth cross-sectional health survey of the KORA cohort; SD, standard deviation.

Systematic reviews and meta-analyses have reported both positive and null associations between environmental exposures and T2D risk, indicating the complexity of these relationships.^{3,11,23} They indicated robust results for T2D prevalence and heterogeneity between studies for incident T2D. A meta-analysis of cohort studies found higher risks of T2D associated with exposure to PM_{2.5} (risk ratio [RR]: 1.39 [1.14–1.68]), PM₁₀ (RR: 1.34 [1.22–1.47]), and NO₂ (RR: 1.11 [1.07–1.16]),³ while another meta-analysis only reported an association for PM_{2.5}.³ Our study found positive but nonsignificant associations for PM_{2.5} (HR: 1.07 [0.97–1.17]) and weak associations

for other air pollutants. Air pollution may contribute to insulin resistance and chronic inflammation, which are key mechanisms in the development of T2D.²⁴

Similarly, a higher diabetes risk has been associated with higher levels of traffic noise.^{9,23} However, this was not confirmed in a meta-analysis of four studies (pooled odds ratio [OR]: 1.49 [0.78–2.82]).¹¹ In our study, we observed weak and nonsignificant associations for traffic noise during the day (HR: 1.02 [0.93–1.11]) or at night (HR: 1.02 [0.94–1.11]). Noise-induced sleep disturbance and chronic stress may contribute to insulin resistance and an increased risk of diabetes.²⁵




Figure 1. Hazard ratios and 95% CIs for the associations between environmental factors, and the risk of incident T2D. Minimum model included age, sex, and subcohort indicator. Main model was further adjusted for BMI, smoking status, alcohol consumption, education level, physical activity, and dietary score. Extended model additionally included cardiovascular diseases, waist-hip ratio, and total cholesterol level. HRs are expressed per IQR increase for each exposure variable. The IQRs were as follows: NO₂: 7.0 µg/m³, NO_x: 8.1 µg/m³, O₃: 3.6 µg/m³, PM₁₀: 2.2 µg/m³, PM_{2.5}: 1.3 µg/m³, PM_{2.5,abs}: 0.3 10⁻⁵/m, PM_{coarse}: 1.4 µg/m³, PNC: 1.9 10³/cm³, Noise_{Day}: 8.2 dB, Noise_{Night}: 7.9 dB, NDVI₃₀₀:0.12, NDVI₁₀₀₀: 0.14, Temp_{annual mean}: 0.6 °C, Temp_{annual SD}: 0.2 °C. NDVI values below 0 were excluded since they represent water or bare rocks. Note: Minimum and main models were based on data for 7696 participants. In extended models we excluded 141 participants with missing clinical information, resulting in 7555 participants. BMI indicates body mass index; CI, confidence interval; HR, hazard ratio; IQR, interquartile range; NDVI, normalized difference vegetation index; NO₂, nitrogen dioxide; NO_x, nitrogen oxide; O₃, ozone; PM₁₀, particulate matter with an aerodynamic diameter ≤10 µm; PM_{2.5}, particulate matter with an aerodynamic diameter ≤2.5 µm; PM_{2.5,abs}, PM_{2.5} absorbance; PM_{coarse}, particulate matter with an aerodynamic diameter of 2.5–10 µm; PNC, particle number concentration; SD, standard deviation; T2D, type 2 diabetes; Temp, temperature.

Systematic reviews reported that green spaces may have a protective effect on T2D.^{11,26,27} A meta-analysis provided further evidence that exposure to greenspace is associated with wide-ranging health benefits, including a significant reduction in T2D incidence (pooled OR: 0.72 [0.61–0.85]).²⁸ However, we observed only weak associations. Green spaces play a crucial role in promoting physical activity and protecting against air and noise pollution, all of which may contribute to mitigating chronic inflammation processes.²⁹

Studies on the effect of air temperature on T2D risk are limited. In a study, diabetes incidence increased by 0.31 [0.19–0.43] per 1 °C increase in annual mean temperature.³⁰ In this study, we observed a weak effect of annual mean air temperature increase (HR: 1.02 [0.92–1.12]). In previous research, both higher temperatures³¹ and lower exposure to cold³² were associated with increased insulin resistance. However, our study found no evidence of this effect.

As previously reported,^{33–36} our study found that sex and education level significantly modified air pollutant effects. However, our results showed opposite directions, possibly due to population variations, exposure misclassification, or chance findings. Lifestyle and physiological differences between sexes or education levels may contribute to these variations.^{34,37}

Strengths and limitations

Our study’s strengths include the use of two datasets from a large and well-characterized population-based prospective cohort, comprehensive exposure assessment, adjustment for important confounders, and application of various analytical approaches. However, limitations such as single-time point exposure assessment, potential exposure misclassification, not accounting for residential mobility from baseline to follow-up, and lack of information on other lifestyle and clinical factors should be acknowledged. Potentially, measurement error may have resulted in underestimation of the underlying associations and insufficient statistical power may be responsible for the wide CI. Our analyses were also restricted to a specific population and geographic region.

Conclusions

Although we observed weaker associations than previous studies indicated, our study contributes to the literature on environmental factors on incident T2D. Future research should continue exploring the role of the environment in T2D development.

Conflicts of interest statement

The authors declare that they have no conflicts of interest with regard to the content of this report

References

- Organization WH. *Global Status Report on Noncommunicable Diseases 2010*. World Health Organization; 2011.
- Organization WH. *Global Diffusion of eHealth: Making Universal Health Coverage Achievable: Report of the Third Global Survey on eHealth*. World Health Organization; 2017.
- Liu F, Chen G, Huo W, et al. Associations between long-term exposure to ambient air pollution and risk of type 2 diabetes mellitus: a systematic review and meta-analysis. *Environ Pollut*. 2019;252:1235–1245.
- Shin S, Bai L, Oiamo TH, et al. Association between road traffic noise and incidence of diabetes mellitus and hypertension in Toronto, Canada: a population-based cohort study. *J Am Heart Assoc*. 2020;9:e013021.
- Wang B, Xu D, Jing Z, Liu D, Yan S, Wang Y. Effect of long-term exposure to air pollution on type 2 diabetes mellitus risk: a systematic review and meta-analysis of cohort studies. *Eur J Endocrinol*. 2014;171:R173–R182.
- Eze IC, Hemkens LG, Bucher HC, et al. Association between ambient air pollution and diabetes mellitus in Europe and North America: systematic review and meta-analysis. *Environ Health Perspect*. 2015;123:381–389.
- He D, Wu S, Zhao H, et al. Association between particulate matter 2.5 and diabetes mellitus: a meta-analysis of cohort studies. *J Diabetes Investig*. 2017;8:687–696.
- Yang BY, Fan S, Thiering E, et al. Ambient air pollution and diabetes: a systematic review and meta-analysis. *Environ Res*. 2020;180:108817.
- Sakhvidi MJZ, Sakhvidi FZ, Mehrparvar AH, Foraster M, Dadvand P. Association between noise exposure and diabetes: a systematic review and meta-analysis. *Environ Res*. 2018;166:647–657.
- Wang H, Sun D, Wang B, et al. Association between noise exposure and diabetes: meta-analysis. *Environ Sci Pollut Res Int*. 2020;27:36085–36090.
- Den Braver NR, Lakerveld J, Rutters F, Schoonmade LJ, Brug J, Beulens JW. Built environmental characteristics and diabetes: a systematic review and meta-analysis. *BMC Med*. 2018;16:1–26.
- Speakman JR, Heidari-Bakavoli S. Type 2 diabetes, but not obesity, prevalence is positively associated with ambient temperature. *Sci Rep*. 2016;6:30409.
- Luo J, He G, Xu Y, et al. The relationship between ambient temperature and fasting plasma glucose, temperature-adjusted type 2 diabetes prevalence and control rate: a series of cross-sectional studies in Guangdong Province, China. *BMC Public Health*. 2021;21:1534.
- Holle R, Happich M, Löwel H, Wichmann HE; Group null for the MS. KORA—a research platform for population based health research. *Gesundheitswesen*. 2005;67:19–25.
- Huth C, Bauer A, Zierer A, et al. Biomarker-defined pathways for incident type 2 diabetes and coronary heart disease—a comparison in the MONICA/KORA study. *Cardiovasc Diabetol*. 2020;19:1–14.
- Thorand B, Zierer A, Büyükköçkan M, et al. A panel of 6 biomarkers significantly improves the prediction of type 2 diabetes in the MONICA/KORA study population. *J Clin Endocrinol Metab*. 2021;106:1647–1659.
- Wolf K, Cyrus J, Harciníková T, et al. Land use regression modeling of ultrafine particles, ozone, nitrogen oxides and markers of particulate matter pollution in Augsburg, Germany. *Sci Total Environ*. 2017;579:1531–1540.
- Pitchika A, Hampel R, Wolf K, et al. Long-term associations of modeled and self-reported measures of exposure to air pollution and noise at residence on prevalent hypertension and blood pressure. *Sci Total Environ*. 2017;593–594:337–346.
- U.S. Geological Survey. Available at: <https://earthexplorer.usgs.gov/>. Accessed May 10, 2023.
- Nikolaou N, Dallavalle M, Stafoggia M, et al. High-resolution spatio-temporal modeling of daily near-surface air temperature in Germany over the period 2000–2020. *Environ Res*. 2023;219:115062.
- Zhang S, Mwiberi S, Pickford R, et al. Longitudinal associations between ambient air pollution and insulin sensitivity: results from the KORA cohort study. *Lancet Planet Health*. 2021;5:e39–e49.
- Pestoni G, Riedl A, Breuninger TA, et al. Association between dietary patterns and prediabetes, undetected diabetes or clinically diagnosed diabetes: results from the KORA FF4 study. *Eur J Nutr*. 2021;60:2331–2341.
- Dzhambov AM. Long-term noise exposure and the risk for type 2 diabetes: a meta-analysis. *Noise Health*. 2015;17:23–33.
- Kelishadi R, Mirghaffari N, Poursafa P, Gidding SS. Lifestyle and environmental factors associated with inflammation, oxidative stress and insulin resistance in children. *Atherosclerosis*. 2009;203:311–319.
- Roswall N, Raaschou-Nielsen O, Jensen SS, Tjønneland A, Sørensen M. Long-term exposure to residential railway and road traffic noise and risk for diabetes in a Danish cohort. *Environ Res*. 2018;160:292–297.
- De la Fuente F, Saldías MA, Cubillos C, et al. Green space exposure association with type 2 diabetes mellitus, physical activity, and obesity: a systematic review. *Int J Environ Res Public Health*. 2021;18:97.
- Dendup T, Feng X, Clingan S, Astell-Burt T. Environmental risk factors for developing type 2 diabetes mellitus: a systematic review. *Int J Environ Res Public Health*. 2018;15:78.
- Twohig-Bennett C, Jones A. The health benefits of the great outdoors: a systematic review and meta-analysis of greenspace exposure and health outcomes. *Environ Res*. 2018;166:628–637.
- Groenewegen PP, Zock JP, Spreeuwenberg P, et al. Neighbourhood social and physical environment and general practitioner assessed morbidity. *Health Place*. 2018;49:68–84.
- Blauw LL, Aziz NA, Tannemaat MR, et al. Diabetes incidence and glucose intolerance prevalence increase with higher outdoor temperature. *BMJ Open Diabetes Res Care*. 2017;5:e000317.
- Wallwork RS, Colicino E, Zhong J, et al. Ambient fine particulate matter, outdoor temperature, and risk of metabolic syndrome. *Am J Epidemiol*. 2017;185:30–39.
- Valdés S, Doulatram-Gangaram V, Lago A, et al. Ambient temperature and prevalence of diabetes and insulin resistance in the Spanish population: Di@bet.es study. *Eur J Endocrinol*. 2019;180:273–280.
- Andersen ZJ, Raaschou-Nielsen O, Ketznel M, et al. Diabetes incidence and long-term exposure to air pollution: a cohort study. *Diabetes Care*. 2012;35:92–98.
- Brook RD, Jerrett M, Brook JR, Bard RL, Finkelstein MM. The relationship between diabetes mellitus and traffic-related air pollution. *J Occup Environ Med*. 2008;50:32–38.
- Chen H, Burnett RT, Kwong JC, et al. Risk of incident diabetes in relation to long-term exposure to fine particulate matter in Ontario, Canada. *Environ Health Perspect*. 2013;121:804–810.
- Eze IC, Schaffner E, Fischer E, et al. Long-term air pollution exposure and diabetes in a population-based Swiss cohort. *Environ Int*. 2014;70:95–105.
- Yang M, Cheng H, Shen C, et al. Effects of long-term exposure to air pollution on the incidence of type 2 diabetes mellitus: a meta-analysis of cohort studies. *Environ Sci Pollut Res Int*. 2020;27:798–811.