

RESEARCH ARTICLE

Open Access



# Prevalence and factors associated with PTSD among female urban slum dwellers in Ibadan, Nigeria: a cross-sectional study

Olutoyin Sekoni<sup>1,2\*</sup> , Sumaya Mall<sup>2†</sup> and Nicola Christofides<sup>2†</sup>

## Abstract

**Background:** Little is known about the prevalence of and factors associated with PTSD among adult females in Nigeria, particularly those who live in slums. PTSD is a mental health condition that develops among some individuals who experience or witness a traumatic event. Several other factors could place individuals at heightened risk of PTSD including stress and comorbid mental disorders. Therefore, this study aimed to examine the prevalence and factors associated with PTSD among female urban slum dwellers in Ibadan, Nigeria.

**Methods:** We conducted a cross sectional survey using multistage sampling of 550 women aged 18 and above from selected slums. Interviewer administered questionnaires were used to elicit information on experience of childhood trauma, recent stressors, intimate partner violence, other mental disorders, sociodemographic characteristics and PTSD. PTSD was measured using the Harvard Trauma Questionnaire (HTQ) which is based on DSM IV. A multivariable linear regression model was built to test associations between PTSD and independent variables.

**Results:** The prevalence for PTSD was found to be 4.18% and the mean PTSD score was  $5.80 \pm 7.11$ . Sexual abuse in childhood, past year intimate partner violence and anxiety were significantly associated with higher PTSD scores. PTSD was not significantly associated with a history of recent stressors. Education, employment and marital status were not associated with PTSD however, age and wealth index showed marginal association with PTSD.

**Conclusion:** The prevalence of PTSD among women living in Ibadan slums was relatively low. Both child sexual abuse and intimate partner violence can be prevented. We also recommend longitudinal studies to better understand risk and protective factors.

**Keywords:** Post-traumatic stress disorder, Urban slum, Traumatic events, Nigeria

## Background

Post-Traumatic Stress Disorder (PTSD) is a disabling mental disorder consisting of three core symptoms: avoidance and numbing, re-experiencing and hyperarousal [1, 2]. PTSD can occur after the experience or witnessing of one or more Traumatic Events (TE) during

childhood or adulthood [3, 4]. Traumatic events can include physical, sexual or emotional abuse during childhood [5, 6]. In adulthood these events include the experience or witnessing, of violent personal assault such as physical or sexual violence perpetrated by a partner, terrorism, serious illness or death of a loved one [7, 8]. The Diagnostic and Statistical Manual of mental disorders V (DSM V) highlights specific traumatic events at least one of which must be experienced prior to a diagnosis of PTSD [9].

\* Correspondence: [t1toyin@yahoo.com](mailto:t1toyin@yahoo.com); [oosekoni@com.ui.edu.ng](mailto:oosekoni@com.ui.edu.ng)

<sup>1</sup>Department of Community Medicine, College of Medicine, University of Ibadan, Queen Elizabeth Road, Ibadan, Nigeria

<sup>2</sup>School of Public Health, University of the Witwatersrand, 27 St Andrews Road, Parktown 2193, South Africa



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Traumatic and stressful life events, can lead to the development of PTSD [10–13]. Traumatic stressors can include serious injury, illness or assault happening to a close relative or the death of a parent, child or spouse. The experience of several negative life events occurring during an individual's lifespan is not unusual in the general population and this has been associated with negative mental health outcomes [12]. Even when the occurrence of negative life events are not strictly characterised as being of sufficient impact as to be traumatic, the degree to which an individual perceives such an event as traumatic can precipitate the development of mental health concerns such as PTSD [14].

PTSD has been found to be associated with mental disorders such as depression and anxiety [15]. This could be the result of co-morbidity or due to overlapping symptoms [16]. PTSD experienced together with other mental disorders, as well as PTSD on its own may contribute to a significant reduction in quality of life [17–19].

Research examining the prevalence and risk factors associated with PTSD suggests that it is associated with a range of socio-demographic characteristics [20–22]. Studies have found PTSD is more commonly reported among women [23–25]. Individuals with higher education levels are less likely to experience PTSD [26, 27]. However, there is inconsistent evidence with regards to marital status. Some studies suggest that marital status is protective against developing PTSD while others suggest that it increases the risk for PTSD [28–31]. Age has been found to play a role in the development of PTSD, with middle-aged women being at greatest risk [32, 33]. Unemployment after the experience of trauma has also been found to be significantly associated with PTSD [34].

Most studies examining PTSD have been conducted in the US and Europe (with earlier research focusing on service members and veterans) [35–37]. A few studies have been conducted on the African continent. In South Africa PTSD has been measured in a population-based mental health epidemiological survey, the South African Stress and Health Study (SASH). SASH suggested a lifetime and 12 month PTSD prevalence of 2.3 and 0.7% respectively while the risk of PTSD was found to be greatest among those who had lost a loved one [7]. Another population-based survey of adult women in one province in South Africa reported a prevalence of 11.6% [38]. High rates of PTSD (87%) were reported among female rape survivors surveyed from health facilities in South Africa [39]. The prevalence of PTSD among adolescents from low socioeconomic communities also in South Africa was about 6% [39, 40]. PTSD was reported among Eritrean refugees in Ethiopia and Rwanda among children of rape survivors [41, 42].

In Nigeria, where historically and currently there have been a wide range of TE and the prevalence of PTSD is thought to range widely between 2.7 and 66.7% [43–46]. Though reasons for such wide range variation in prevalence of PTSD are not known, it is possible that the use of varying study designs, tools and samples studied may have influenced this variation [47]. For example the cross-sectional study reported by Sheikh et al. reported a PTSD prevalence of 6.8%, while the case control study of Asuquo et al. reported a prevalence of 41.3% [43, 45]. Predictors of PTSD in Nigeria include a wide spectrum of events e.g. political and ethno-religious violence to road traffic accidents [43–46, 48]. However, slum environments tend to be associated with a greater number of TE, therefore the risk of developing PTSD in such settings may be higher [49–51]. To our knowledge, there are limited prevalence data for PTSD from community-based samples and poor urban slum settings in Nigeria.

Slum areas in Ibadan, which is the setting of this study, are located in the central and oldest parts of the city and represent areas of little or no identifiable sanitation facilities or social infrastructure, physical deterioration, and very high population density [52, 53]. The slums developed in the wake of the colonial era which was followed by an increase in economic activities making it possible for individuals to attain economic independence at earlier ages and as a result the desire to build personal property [54]. In a bid to remain within the extended family setting, such houses could only be built within the open space of the family compound. This ultimately has led to wide spread overcrowding and poor access to infrastructure such as water, electricity and sewerage due to the un-planned nature of the neighbourhoods and almost non-existent road networks [55, 56].

The purpose of this study was to estimate the prevalence of PTSD, and to describe associated factors including socio demographic characteristics, experience of childhood trauma, intimate partner violence, recent stressors, as well as comorbid mental disorders among women in a slum setting in South West Nigeria. Our study addresses gaps in the literature on PTSD with particular reference to trauma across the life course and mental health in community samples as well as among women in deprived circumstances such as a slum setting in Ibadan, Nigeria.

The results potentially contribute to the body of literature on PTSD among women in low income urban neighbourhoods and the role that neighbourhood context plays in the development of PTSD which are research gaps that have been highlighted by other researchers [57]. It will also contribute to our understanding of the prevalence of comorbid mental disorders in such contexts [58, 59].

## Methods

### Participants

A community based cross-sectional household survey was conducted in slum areas of Ibadan in Oyo State. Participants were adult women above the age of 18 years who were resident members of the selected households during the period October to November 2018. Regular residence was defined as spending at least four nights a week in the household for at least 1 year [60].

### Procedure

Six female research assistants with tertiary education were recruited and trained. Training of research assistants took place over 5 days and covered data collection and ethical considerations relevant for the study. These included method of administration of questionnaires, review of questions for completeness and maintenance of ethical standards with particular emphasis on the sensitive nature of the questions. The training also included the World Health Organization requirements for collecting data for trauma research [61]. Training on the use of the Open Data Kit (ODK) was also carried out to teach research assistants how to use the software and upload the data once obtained. ODK is a mobile technology that enhances the data collection and data management process by ensuring real time data entry through the use of the internet. It was developed at the University of Washington and has been found to be an effective data collection tool [62, 63]. Role plays were utilised in order to enhance skills and evaluate the training.

A sample of 550 eligible women were selected based on a multistage sampling technique. Simple random sampling was used to select five of thirteen slum settlements in Ibadan through the use of a table of random numbers. The selected slums were: Beere, Oje, Oritamerin/Agbeni, Ojaba and Mapo located within the central core of the city occupied by the indigenous people of Ibadan and are also known as “traditional slums” [52]. Enumeration Areas (EAs) as demarcated by the National Population Commission [64] were identified by using already existing maps prepared by the Commission. One third of the EAs were selected from each of the five slums using a table of random numbers. This yielded a total of 42 EAs. However this number was later increased to 68. This ensured that there were sufficient households available per EA to select one eligible respondent per household while adhering to the required sample size. This approach has been utilised in the conduct of other population based surveys on violence research in order to ensure safety of both the respondent and the interviewer [61]. The Nigerian Demographic and Health Survey (NDHS) indicates 48 households per EA [60]. However this figure was not specified for a slum area where higher number of households would be

expected. We assumed an average of 60 households per EA (four households per dwelling in 15 dwellings per EA). One household was selected randomly by balloting per house where there were at least two households. One person in the selected household was selected randomly by balloting. Data collection was carried out through the use of face to face interviews by the research assistants using standardized questionnaires uploaded on mobile devices with ODK software.

### Measures

#### Outcome variable

The outcome variable for this study was PTSD assessed using the Harvard Trauma Questionnaire (HTQ) which is based on DSM-IV PTSD criteria [65]. The HTQ is a reliable screening instrument that has been widely used and is acceptable across cultures [65–67]. The trauma symptom items of the HTQ were used in this study. Items included an individual’s perception of their ability for daily functioning. For example, “*feeling detached or withdrawn from people*” and”, “*feeling as though the event is happening again*”.

The HTQ utilizes a 4 point Likert scale of 16 items with responses of “not at all”, “a little bit”, “quite a bit” or “extremely” to determine the occurrence of post-traumatic symptoms within the previous week [65, 68]. A score of 1 was assigned to the responses of “not at all”, a score of 2 was assigned to the responses of “a little bit”, a score of 3 was assigned to the responses of “quite a bit” and 4 was assigned to responses of “extremely”. The minimum score was 1 and the maximum possible score was 64. The sum of the items were then divided by the total number of items. A cut point of 2.0 was used in interpreting the score which has been recommended for use when doing community based research as opposed to the cut point of 2.5 recommended for use among clinic based population samples [65, 67, 69]. We adopted this lower cut point as population characteristics have been suggested to influence the performance of the HTQ with differences seen among highly traumatised populations as opposed to those that are not as highly traumatised [67]. Based on this, the outcome variable was dichotomized as either ‘No PTSD symptoms’ or as having ‘PTSD symptoms’. We found good internal consistency with a Cronbach’s alpha of 0.88.

#### Explanatory variables

Experiences of childhood trauma were assessed using the Childhood Trauma Questionnaire (CTQ). The CTQ is an instrument consisting of 28 items covering a range of childhood experiences [70]. Domains include emotional abuse and neglect, physical abuse and neglect and sexual abuse [70, 71]. For the purposes of the analyses presented in this paper we used only the sexual, physical

and emotional abuse sub-scales as they are more likely to be associated with PTSD [72, 73]. The CTQ has been used previously on the African continent [74, 75]. We found good internal consistency with a Cronbach's alpha of 0.82. Examples of items on the CTQ include: "People in my family called me things like 'stupid', 'lazy' or 'ugly'", "I got hit so hard by someone in my family that I had to see a doctor or go to the hospital", "Someone tried to touch me in a sexual way or tried to make me touch them". Each subscale was assessed as a dichotomous variable with any affirmative answer to an item as an experience.

Intimate partner violence was assessed using the WHO Multi-country Study on Women's Health and Domestic Violence Core Questionnaire [61]. Experiences of intimate partner violence were asked for within the various domains of physical and sexual abuse. The internal consistency of the scale was good at a Cronbach's alpha of 0.84. In our analysis, we identified past year experience of intimate partner violence with variable responses being having had an experience of intimate partner violence in the year prior to the study or not having had an experience of intimate partner violence in the year prior to the study.

Recent stressors were assessed using the Life Events Questionnaire [76]. The scale considers recent life events that tend to be stressful and threatening and have occurred within the preceding 6 months. It is a 12 item self-report questionnaire which assesses the quality of life of an individual before, during and after the occurrence of a stressful life event. We utilised four items of the twelve item questionnaire as these four items were the only ones that fit the PTSD criteria for a traumatic experience [77, 78]. The items are as follows: "Did you suffer serious illness, injury or assault?", "Did your parent, child or spouse die?", "Did a serious illness, injury or assault happen to a close relative?" and "Did a close family friend or another relative (aunt, cousin, grandparent) die?" For each item a "yes" (with a score of 1) or "no" (with a score of 0) response was given which has been identified as a valid method of assessment [79]. The frequency of each of the items has been presented. In addition to this, a dichotomous score was computed from the total sum of all the items and was categorized into "no stressors" or "any stressors".

Depression and anxiety was assessed using the DASS-21 (Depression, Anxiety and Stress Scale-21) which has been used previously in Africa [80]. It contains three subscales: one subscale for depression, one for anxiety and one for stress. Each subscale is composed of 7 separate items using a 4 point Likert scale with the respondent identifying the extent to which each of the statements on the instrument applied over the past 1 week [81]. Examples of items on the depression scale

include; "I found it difficult to work up the initiative to do things" and "I felt that I had nothing to look forward to".

Examples of items on the anxiety scale include; "I experienced trembling (e.g in the the hands)" and "I was worried about situations in which I might panic and make a fool of myself".

Scores for each scale are calculated by summing the scores for the items on each scale. The cutoff values serve as a means of interpretation with respect to the dimensions of severity. Responses for each of the scales are indicated as 0 (did not apply to me at all), 1(applied to me to some degree, or some of the time), 2 (applied to me to a considerable degree or a good part of time), 3 (applied to me very much or most of the time). Scores on the DASS 21 are multiplied by 2 to give the final score which is categorized accordingly into "normal", "mild", "moderate", "severe", and "extremely severe".

Due to very small numbers of participants with DASS scores in the moderate and severe categories and none in the extremely severe category, we collapsed categories for the depression and anxiety scales. We generated a dichotomous variable for depression and anxiety with categories of "no anxiety" representing the initial "normal" and "mild" and "anxiety" represented by collapsing the initial categories of "moderate", "severe" and "extremely severe". No participants had stress scores that were in the "moderate", "severe" or "extremely severe" category so we did not include the assessment of stress in this study.

Validity, reliability and internal consistency of the DASS has been found to be satisfactory across various populations of both clinical and non - clinical samples [82–84]. We found satisfactory internal consistency with a Cronbach's alpha of 0.73.

Sociodemographic characteristics: included age, educational status, marital status, employment status, occupation and wealth index. Age was categorized into three age groups as follows: 18–29 years, 30–44 years and age 45 years or older. Educational status was categorized as none- for those who had not undergone any form of formal education, primary- which represents the first 6 years of schooling and secondary- which represents the subsequent 6 years of schooling. In our study, marital status was categorized as "married"- for those in a formal marriage union, "single"- for those not in any formal marriage union and "others"- for those that did not fall into the category of married or single such as those who were widowed, separated or divorced. Employment status was determined by whether or not the respondent had been employed within the last 3 months or not with responses being "yes" or "no". Respondents were categorized into three major occupations: trader, artisans and professionals. Our assessment of wealth was done



utilizing the 11 item Simplified Asset Index, a previously published tool which assesses the relative wealth profile of urban dwellers in comparison to other urban dwellers making it a more accurate assessment of wealth for the urban poor [85]. The Simplified Asset Index is based on Demographic and Health Survey (DHS) data. The DHS has been implemented in more than 90 countries since 1984 [86] and allows for reliable sub-group analyses in urban areas. The index is a shorter country specific alternative to the original DHS wealth index variables but has been found to equally effective and easier to administer [85]. Examples of items from the Simplified Asset Index include; “*What type of fuel does your household mainly use for cooking?*” and “*Does any member of this household have a bank account?*”

The original Simplified Asset Index is divided into 5 quintiles but due to small cell numbers we had to merge these into three. The three urban wealth index categories were assigned as follows: the lowest category 1 being “poorest”, the middle category 2 being “poor” and the highest category 3 being “slightly wealthier”.

The instrument was translated to the local language [Yoruba] and back translated to ensure that original meanings were preserved and was pre-tested in an enumeration area within a slum area that had not been selected for the study. Such a process of translation has been described by other researchers [87]. Based on the feedback received, we made corrections to the questionnaire.

Ethical approval for this study was sought from the University Of Witwatersrand Faculty Of Health Science Human Ethics Committee as well as the Ethics Review Board of the Oyo State Ministry of Health, Oyo State, Nigeria.

#### **Data analysis**

Data were analyzed using STATA Version 14.0 software (Stata Corp., College Station, TX, USA). There were no missing data on the outcome measure of interest and hence no case dropping was implemented. We adjusted for the cluster sampling during analysis using the *svyset* command in STATA. We also weighted the data to ensure that it was representative of the Ibadan slums using the command *pweight*. The outcome variable under consideration was the mean PTSD score while the explanatory variables included the experience of childhood trauma (sexual, physical and / or emotional abuse), sexual and /or physical intimate partner violence, recent stressors, depression, anxiety and sociodemographic characteristics. T-tests were used to compare mean PTSD symptomatology scores with explanatory variables such as child sexual abuse, intimate partner violence, depression and anxiety which were dichotomous variables. Oneway ANOVA was used for categorical variables such

as educational attainment, marital status and wealth index. Subsequently, a multivariable linear regression analysis was built to examine associations between PTSD score and explanatory variables which were associated in the univariate analyses or considered theoretically important based on the literature. Statistical significance was set at  $p < 0.05$ .

#### **Results**

All 550 women in selected households agreed to participate in the study giving a 100% response rate. This high response rate is not uncommon in the Ibadan context. The median age of respondents was 40 years with an interquartile range of 30–55 years. About a fifth had never attended school (22.4%). Almost two thirds were married or cohabiting with a partner (67.8%). The majority of the sample had been employed 3 months prior to the interview (77.3%) and came from a household that was in the poor and poorest urban wealth categories (71.6%). Twenty three participants from our sampled population reported PTSD symptoms and this yielded a PTSD prevalence of 4.18%.

With respect to the univariate association with PTSD, being a woman from a household in the poorest category was associated with the occurrence of PTSD even though the association was not strong. No other socio-demographic characteristics were associated with PTSD (Table 1).

The prevalence of the trauma experiences during childhood ranged from 8.9% (sexual abuse), 50.4% physical abuse and 70.4% emotional abuse. Women without a history of child sexual abuse were less likely to report PTSD symptoms and this was statistically significant. The experience of recent stressors was not associated with reports of PTSD symptoms among the women. Women who were not depressed and not anxious were also significantly less likely to report PTSD symptoms (Table 2).

We conducted a regression analysis to examine the relationship between PTSD symptoms, traumatic events and mental disorders while controlling for sociodemographic characteristics. Child sexual abuse, recent intimate partner violence and symptoms of anxiety were all associated with a higher PTSD score. For age and wealth status, there were marginal associations with a higher PTSD score. The age of women who were 30–44 compared to those who were in the 18–29 year age group were marginally more likely to have higher PTSD scores. Women in the slightly wealthier category were less likely to report PTSD symptoms compared to the poorest women. PTSD was not significantly associated with a history of other recent stressors among the sample (Table 3).

**Table 1** Associations between sociodemographic characteristics and PTSD

	PTSD symptoms n (%)	No PTSD symptoms n (%)	P value
<b>Age at last birthday</b>			
18–29	5 (21.74)	132 (25.05)	0.63
30–44	10 (43.48)	178 (33.78)	
≥ 45	8 (34.78)	217 (41.18)	
<b>Educational status</b>			
None	6 (26.09)	117 (22.20)	0.74
Any Primary	7 (30.43)	158 (29.98)	
Some Secondary	1 (4.35)	66 (12.52)	
Completed secondary/Tertiary	9 (39.13)	186 (35.29)	
<b>Marital status</b>			
Married/Cohabiting	15 (65.22)	358 (67.93)	0.59
Single/Never married	3 (13.04)	40 (7.59)	
Separated/widowed/divorced	5 (21.74)	129 (24.48)	
<b>Employed in the last 3 months</b>			
No	8 (34.78)	117 (22.20)	0.15
Yes	15 (65.22)	410 (77.80)	
<b>Wealth index</b>			
Poorest	13 (56.52)	171 (32.45)	0.05 <sup>a</sup>
Poor	5 (21.74)	205 (38.90)	
Slightly wealthier	5 (21.74)	151 (28.65)	

<sup>a</sup>Marginal finding

## Discussion

Our study set out to estimate the prevalence of PTSD and associated factors among women in a slum setting in South West Nigeria. We found a prevalence of 4.18% for PTSD with only 23 women among our sample of 550 women reporting symptoms of PTSD. Sexual abuse during childhood, recent experience of intimate partner violence as well as anxiety were significantly associated with PTSD. A history of recent stressful life events was not associated with PTSD. None of the sociodemographic characteristics considered (age, education, marital status or wealth index) were associated with PTSD. However, in the multivariable linear regression analysis there were marginal associations with age and wealth index only.

We observed low prevalence of PTSD among our respondents despite their precarious circumstances in the slum environment, which included high levels of adverse life events, very limited resources and services. In urban slums elsewhere, researchers have described greater prevalence of PTSD among residents of as a result of the adversity experienced within such settings. Mbwanyo et al. reported a PTSD prevalence of 15.4% among respondents in urban slums in Kenya [88] while Ndugu et al. reported 21% among women in informal settings in South Africa [89]. A high prevalence of 29 and 20% was also reported among women in low income

neighbourhoods in the United States [51, 59]. There is evidence in literature to suggest that resilience may play a role in moderating the effect of trauma and subsequent PTSD experienced. This has been seen among populations that are at particular risk for PTSD such as police officers and fire-fighters [90, 91], as well as among undergraduate women [92]. Resilience may explain the low prevalence we observed.

We found that a history of sexual abuse in childhood was associated with PTSD in adults, a finding which has been reported previously [93–95]. Child sexual abuse has also been reported widely on the African continent [96–100]. We did not find an association between physical and emotional abuse during childhood and PTSD although other studies have reported an association. McLaughlin and colleagues who analysed data from the World Mental Health Surveys suggest that 4 childhood adversities out of a potential of 12 were associated with the onset of PTSD. Childhood physical abuse and sexual abuse were amongst these [101]. In another analysis from the WMHS led by Koenen and colleagues, the odds of lifetime PTSD among trauma exposed adults was 2.5 [102].

The experience of sexual and / or physical intimate partner violence represents a trauma in adulthood. We found that recent intimate partner violence was

**Table 2** Association between traumatic events, mental disorders and PTSD

Childhood trauma	PTSD symptoms n (%)	No PTSD symptoms n (%)	P value
<b>Sexual abuse</b>			
None	15 (65.22)	486 (92.22)	0.000*
Any sexual abuse	8 (34.78)	41 (7.78)	
<b>Emotional abuse</b>			
None	4 (17.39)	159 (30.17)	0.246 <sup>a</sup>
Any emotional abuse	19 (82.61)	368 (69.83)	
<b>Physical abuse</b>			
None	8 (34.78)	265 (50.28)	0.146
Any physical abuse	15 (65.22)	262 (49.72)	
<b>Recent Stressors (Items)</b>			
<b>Suffered serious illness, injury or assault</b>			
No	18 (78.26)	450 (85.39)	0.34
Yes	5 (21.74)	77 (14.61)	
<b>Serious illness, injury or assault happened to a close relative</b>			
<b>No</b>	21 (91.30)	492 (93.36)	0.66 <sup>a</sup>
Yes	2 (8.70)	35 (6.64)	
<b>Parent, child or spouse died</b>			
<b>No</b>	20 (86.96)	469 (88.99)	0.73 <sup>a</sup>
<b>Yes</b>	3 (13.04)	58 (11.01)	
<b>A close family friend or another relative (aunt, cousin, grandparent) died</b>			
<b>No</b>	22 (95.65)	454 (86.15)	0.19 <sup>a</sup>
<b>Yes</b>	1 (4.35)	73 (13.85)	
<b>Recent stressors (Total)</b>			
No stressors	14 (60.87)	375 (71.16)	0.28
Any stressors	9 (39.13)	152 (28.84)	
<b>Past year Intimate Partner Violence</b>			
No	18 (78.26)	441 (83.68)	0.49
Yes	5 (21.74)	86 (16.32)	
<b>Depression</b>			
No depression	22 (95.65)	524 (99.43)	0.15 <sup>a</sup>
Depression	1 (4.35)	3 (0.57)	
<b>Anxiety</b>			
No anxiety	16 (69.57)	496 (94.12)	0.00*
Anxiety	7 (30.43)	31 (5.88)	

<sup>a</sup>Fishers exact \* $p < 0.05$

significantly associated with a higher PTSD symptoms among our sample. Research from other sub-Saharan African countries has reported an association [38]. Mahenge and colleagues in Tanzania also found higher levels of PTSD among pregnant women who had experienced intimate partner violence [103]. A multicountry study among university students from 25 countries (7 countries from sub-Saharan Africa) also suggested that

PTSD was associated with intimate partner violence in all of the sub-Saharan African countries [104].

We found an association between PTSD and anxiety. This association has been reported by other authors [105, 106]. There are two possible explanations for this finding: it is possible that women with anxiety may be more susceptible to developing PTSD (or vice versa) or it could be that they share common symptoms [16].

**Table 3** Multivariable linear regression model of PTSD symptoms and traumatic events and mental disorders adjusting for socio demographic characteristics (N = 550)

Variable	Beta	95% CI	P value
<b>Childhood trauma</b>			
<b>Sexual abuse</b>			
None	Ref		
Any sexual abuse	3.82	1.75–5.88	0.00
<b>Emotional abuse</b>			
None	Ref		
Any emotional abuse	0.81	−0.56–2.19	0.24
<b>Physical abuse</b>			
None	Ref		
Any physical abuse	0.55	−0.70–1.82	0.38
<b>Recent Stressors (Total)</b>			
No stressors	Ref		
Any stressors	1.08	−0.17–2.34	0.09
<b>Past year Intimate Partner Violence</b>			
No	Ref		
Yes	1.83	0.20–3.46	0.02
<b>Depression</b>			
No Depression	Ref		
Depression	2.08	−4.71–8.88	0.54
<b>Anxiety</b>			
No Anxiety	Ref		
Anxiety	6.38	4.09–8.67	0.00
<b>Age at last birthday</b>			
18–29	Ref		
30–44	1.68	−0.04–3.41	0.05 <sup>a</sup>
≥ 45	0.95	−0.99–2.90	0.33
<b>Educational status</b>			
None	Ref		
Any Primary	−0.05	−1.94–1.83	0.95
Some Secondary	−0.41	−2.77–1.94	0.73
Completed secondary/Tertiary	0.24	−1.78–2.27	0.81
<b>Marital status</b>			
Married/Cohabiting	Ref		
Single/Never married	1.65	−0.92–4.22	0.20
Separated/widowed/divorced	0.87	−0.88–2.63	0.32
<b>Employed in the last 3 months</b>			
No	Ref		
Yes	0.46	−1.96–1.03	0.54
<b>Wealth index</b>			
Poorest	Ref		
Poor	−0.73	−2.17–0.70	0.31
Slightly wealthier	−1.53	−3.15–0.07	0.06 <sup>a</sup>

<sup>a</sup>Marginal findings CI Confidence Interval

There may be an overlap of symptoms of PTSD and the spectrum of both anxiety and depressive disorders which may mask anxiety and mood symptoms and as such must be borne in mind when assessing for PTSD [16]. A bidirectional association between anxiety and PTSD has been suggested in the literature with some researchers reporting that anxiety predisposes to PTSD in the face of trauma while others indicate that PTSD is a risk factor for anxiety [107, 108]. It is clear that anxiety occurs very commonly with PTSD and as such treatment measures for PTSD should factor in the need to screen for as well as treat this co-morbidity.

We did not find any association between the experience of stressful life events such as serious illness, injury or assault or death of a relative within the preceding 6 months and symptoms of PTSD. However, other studies conducted in different settings have shown that stressful life events can contribute to the development of PTSD [13, 109, 110]. The findings with respect to the association between sociodemographic characteristics and PTSD have been mixed in the literature. Age has generally not been associated with PTSD as has been reported by authors in Africa [7, 48, 111, 112]. However, Kobayashi in the United States reported an association between the middle aged subgroup of women and PTSD, with middle aged women being more likely to have PTSD compared to younger and older women after controlling for potential confounders [33]. Our results are similar to those of Kobayashi, with women aged 30 to 44 having statistically significant higher PTSD scores than women aged 18- to 29. Middle aged women are thought to carry the greater burden and stress of care-giving for both the younger and older individuals within the community and a disruption of that role due to trauma can be distressing to them and make them more likely to suffer from anxiety and depression [113]. Some authors have reported that educational level is associated with PTSD with individuals with higher levels of education less likely to develop PTSD [20, 23, 114–116]. We did not find an association between education and PTSD in our sample. Other researchers in Africa have reported that education level, employment status, religious affiliation and marital status were not associated with PTSD [117].

The aetiology of PTSD is multifactorial and differs depending on context [118]. The context of our study may have protected participants from exposure to trauma and the development of PTSD. The particular slum environment in urban Ibadan where the study took place is stable and homogenous. Most participants belonged to one ethnic group and had lived in the area for most or all of their lives. Literature has suggested that living in neighborhoods that have greater ethnic homogeneity is protective against mental disorders [119, 120]. This has been described as a function of the perception of being



in similar socioeconomic circumstances to everyone else in the environment and with it the a higher acceptance of income inequality and less mental stress [120]. With a majority of our slum sample falling within the same lower wealth categories, this explanation may not be out of place.

It is evident in that not everyone exposed to trauma develops PTSD. The results of our study among this sample has highlighted this clearly with unexpectedly low levels of PTSD in spite of a substantial history of childhood trauma as well as less than optimal living conditions of a slum setting. This suggests the need for further research to explore resilience to PTSD and individual and community-based factors that may provide protection from developing symptoms of PTSD in the slum setting in Ibadan, Nigeria. Factors such as social connectedness, self – esteem and social support have been found to be protective against the development and severity of PTSD [121–124].

The results of the study need to be interpreted in the light of several limitations. From a study design standpoint, the cross-sectional nature of our study precludes conclusions about the temporal relationship between intimate partner violence, anxiety and PTSD. The HTQ was developed in the United States and it is possible that cultural differences between our study population and the original population for which it was designed could have resulted in underreporting of PTSD symptoms. Several studies have suggested that there might be cultural explanations for mental disorders such as depression and that language and culture shape views of illness [125, 126].

We also cannot be sure whether translation of instruments might have resulted in the loss of meaning of items measuring mental disorders as has been suggested by seminal work on translation and mental health care in South Africa, a culturally and linguistically diverse society [127, 128].

Reports of childhood trauma and stressors could be affected by recall bias with participants underreporting the occurrence. However, retrospective accounts of trauma have been found to be acceptable for PTSD research [129]. Certain kinds of traumatic events were not measured in this study. We did not ask about experiences of witnessing a murder or witnessing of rape, to name a few. Our study was not sufficiently powered because of our expectation of a higher prevalence of PTSD in this sample based on previous research that had indicated higher prevalence of PTSD in a slum setting.

## Conclusions

Our findings shed light on the low levels of PTSD symptoms among this community based sample of poor urban women. However, there was a significant

relationship between the experience of childhood trauma and PTSD. In addition, intimate partner violence and anxiety were also associated with PTSD. Our findings add to the literature which suggests that the consequences of early childhood experiences of trauma especially childhood sexual abuse and intimate partner violence can have quite serious mental health consequences including PTSD. These results speak to the urgency to prevent both childhood abuse as well as intimate partner violence. Policy and actions geared towards these two issues would be particularly important as both are preventable and ultimately could reduce the incidence of PTSD.

## Abbreviations

CI: Confidence Interval; EAs: Enumeration Areas; DASS: Depression Anxiety and Stress Scale; PTSD: Post-Traumatic Stress Disorder; TE: Traumatic Events; ODK: Open Data Kit

## Acknowledgements

Not applicable.

## Authors' contributions

OS was involved in conception of the study, data collection, data analysis and interpretation and drafting the article. SM was involved in conception of the study, data analysis and interpretation and critical revision of the article. NC was involved in conception of the study, data analysis and interpretation and critical revision of the article. OS, SM and NC all approved of the final version of the manuscript to be published.

## Funding

OS was supported by" by the Consortium for Advanced Research Training in Africa (CARTA). CARTA is jointly led by the African Population and Health Research Center and the University of the Witwatersrand and funded by the Carnegie Corporation of New York (Grant No–B 8606.R02), Sida (Grant No:54100113), the DELTAS Africa Initiative (Grant No: 107768/Z/15/Z) and Deutscher Akademischer Austauschdienst (DAAD). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (UK) and the UK government.

SM has been supported by a Claude Leon award for early career researchers and a self-initiated research fellowship from the South African Medical Research Council (2018–2020).

The funders had no role in the design of the study, collection, analysis, and interpretation of data nor in writing the manuscript.

## Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## Declarations

### Ethics approval and consent to participate

Ethical approval for this study was sought from the University Of Witwatersrand Faculty Of Health Science Human Ethics Committee (reference number: M180249) as well as the Ethics Review Board of the Oyo State Ministry of Health, Oyo State, Nigeria (reference number: AD/13/479/762). Both ethics committees approved the study. Permission to conduct the study was also obtained from appropriate local community authorities and written informed consent was obtained from all participants. The principles of beneficence, justice & non-maleficence were observed.

### Consent for publication

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

Received: 10 August 2020 Accepted: 19 July 2021

Published online: 12 August 2021

**References**

- Scher CD, McCreary DR, Asmundson GJG, Resick PA. The structure of post-traumatic stress disorder symptoms in three female trauma samples: a comparison of interview and self-report measures. *J Anxiety Disord.* 2008; 22(7):1137–45. <https://doi.org/10.1016/j.janxdis.2007.11.012>.
- Radell ML, Beck KD, Gilbertson MW, Myers CE. Post-traumatic stress disorder symptom burden and gender each affect generalization in a reward- and punishment-learning task. *PLoS One.* 2017;12(2):e0172144. <https://doi.org/10.1371/journal.pone.0172144>.
- Iribarren J, Prolo P, Neagos N, Chiappelli F. Post-traumatic stress disorder: evidence-based research for the third millennium. *Evid Based Complement Alternat Med.* 2005;2(4):503–12. <https://doi.org/10.1093/ecam/neh127>.
- Stein DJ, Karam EG, Shahly V, Hill ED, King A, Petukhova M, et al. Post-traumatic stress disorder associated with life-threatening motor vehicle collisions in the WHO world mental health surveys. *BMC Psychiatry.* 2016; 16(1):1–14. <https://doi.org/10.1186/s12888-016-0957-8>.
- Chemtob CM, Gudiño OG, Luthra R, Yehuda R, Schmeidler J, Auslander B, et al. Child trauma exposure and posttraumatic stress disorder: identification in community mental health clinics. *Evid Based Pract Child Adolescent Mental Health.* 2016;1(2–3):103–15. <https://doi.org/10.1080/23794925.2016.1227949>.
- Sheriff RS, van Hooff M, Malhi G, Grace B, McFarlane A. Associations of childhood trauma and childhood mental disorder with past year mental disorder in military and civilian employed men. *Psychiatry Res.* 2019;280:1–36. <https://doi.org/10.1016/j.psychres.2019.112482>.
- Atwoli L, Stein DJ, Williams DR, McLaughlin KA, Petukhova M, Kessler RC, et al. Trauma and posttraumatic stress disorder in South Africa: analysis from the south African stress and health study. *BMC Psychiatry.* 2013;13(1):182. <https://doi.org/10.1186/1471-244X-13-182>.
- Kleber RJ. Trauma and public mental health: a focused review. *Front Psychiatry.* 2019;10(451):1–6. <https://doi.org/10.3389/fpsy.2019.00451>.
- APA. Highlights of changes from DSM-IV to DSM-5, highlights of changes from DSM-IV-TR to DSM-5. Arlington; 2013. <https://doi.org/10.1176/appi.focus.11.4.525>.
- Leserman J, Whetten K, Lowe K, Stangl D, Swartz MS, Thielman NM. How trauma, recent stressful events, and PTSD affect functional health status and health utilization in HIV-infected patients in the south. *Psychosom Med.* 2005;67(3):500–7. <https://doi.org/10.1097/01.psy.0000160459.78182.d9>.
- Gold SD, Marx BP, Soler-Baillo JM, Sloan DM. Is life stress more traumatic than traumatic stress? *Journal of Anxiety Disorders.* 2005;19(6):687–98. <https://doi.org/10.1016/j.janxdis.2004.06.002>.
- Schock K, et al. Impact of new traumatic or stressful life events on pre-existing PTSD in traumatized refugees: results of a longitudinal study. *Eur J Psychotraumatol.* 2016;7:1–11.
- Van Den Berg LJM, et al. A new perspective on PTSD symptoms after traumatic vs stressful life events and the role of gender. *Eur J Psychotraumatol.* 2017;8(1380470):1–12. <https://doi.org/10.1080/20008198.2017.1380470>.
- Willard VW, Long A, Phipps S. Life stress vs traumatic stress: the impact of life events on psychological functioning in children with and without serious illness. *Psychol Trauma Theory Res Pract Policy.* 2016;8(1):63–71. <https://doi.org/10.1037/tra0000017>.
- Flory JD, Yehuda R. Comorbidity between post-traumatic stress disorder and major depressive disorder: alternative explanations and treatment considerations. *Dialogues Clin Neurosci.* 2015;17(2):141–50 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26246789> (Accessed 17 May 2017).
- Carmassi C, Bertelloni CA, Cordone A, Cappelli A, Massimetti E, Dell'Oste V, et al. Exploring mood symptoms overlap in PTSD diagnosis: ICD-11 and DSM-5 criteria compared in a sample of subjects with bipolar disorder. *J Affect Disord.* 2020;276(July):205–11. <https://doi.org/10.1016/j.jad.2020.06.056>.
- Lohr JB, Palmer BW, Eidt CA, Aailaboyina S, Mausbach BT, Wolkowitz OM, et al. Is post-traumatic stress disorder associated with premature senescence? A review of the literature HHS public access. *Am J Geriatr Psychiatry.* 2015;23(7):709–25. <https://doi.org/10.1016/j.jagp.2015.04.001>.
- McMillan KA, Asmundson GJG, Sareen J. Comorbid PTSD and social anxiety disorder: associations with quality of life and suicide attempts. *J Nerv Ment Dis.* 2017;205(9):732–7. <https://doi.org/10.1097/NMD.0000000000000704>.
- Danielsson FB, Schultz Larsen M, Nørgaard B, Lauritsen JM. Quality of life and level of post-traumatic stress disorder among trauma patients: a comparative study between a regional and a university hospital. *Scand J Trauma Resusc Emerg Med.* 2018;26(1):1–9. <https://doi.org/10.1186/s13049-018-0507-0>.
- Olf M. Sex and gender differences in post-traumatic stress disorder: an update. *Eur J Psychotraumatol.* 2017;8(sup4):1351204. <https://doi.org/10.1080/20008198.2017.1351204>.
- Frankenberg E, Sikoki B, Sumantri C, Suriastini W, Thomas D. Education, vulnerability, and resilience after a natural disaster. *Ecol Soc.* 2013;18(2). <https://doi.org/10.5751/ES-05377-180216>.
- Ryb GE, et al. PTSD after severe vehicular crashes. *Ann Adv Automot Med.* 2009;53:177–93 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20184843> (Accessed 13 July 2019).
- Birkeland, M. S., Blix I., Solberg Ø., Heir T. 'Gender Differences in Posttraumatic Stress Symptoms after a Terrorist Attack: A Network Approach'. 2017;8:1–11. doi: <https://doi.org/10.3389/fpsyg.2017.02091>.
- Shalev AY, Gevonden M, Ratanatharathorn A, Laska E, van der Mei WF, Qi W, et al. Estimating the risk of PTSD in recent trauma survivors: results of the international consortium to predict PTSD (ICPP). *World Psychiatry.* 2019; 18(1):77–87. Available at: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/wps.20608> (Accessed 7 Aug 2019).
- Haag K, Fraser A, Hiller R, Seedat S, Zimmerman A, Halligan SL. The emergence of sex differences in PTSD symptoms across development: evidence from the ALSPAC cohort. *Psychol Med.* 2019;50(10):1–6. <https://doi.org/10.1017/S0033291719001971>.
- Lin W, Gong L, Xia M, Dai W. Prevalence of posttraumatic stress disorder among road traffic accident survivors. *Medicine.* 2018;97(3):e9693. <https://doi.org/10.1097/MD.0000000000009693>.
- Tang B, Deng Q, Glik D, Dong J, Zhang L. A meta-analysis of risk factors for post-traumatic stress disorder (PTSD) in adults and children after earthquakes. *Int J Environ Res Public Health.* 2017;14(1537):1–20. <https://doi.org/10.3390/ijerph14121537>.
- Grieger TA, et al. Posttraumatic stress disorder and depression in battle-injured soldiers. *Am J Psychiatr.* 2006;163(10):1777–83. <https://doi.org/10.1176/ajp.2006.163.10.1777>.
- Smith TC, Ryan MA, Wingard DL, Slymen DJ, Sallis JF, Kritz-Silverstein D, et al. New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *Bmj.* 2008;336(7640):366–71. <https://doi.org/10.1136/bmj.39430.638241.AE>.
- Dekel R, Monson CM. Military-related post-traumatic stress disorder and family relations: current knowledge and future directions. *Aggress Violent Behav.* 2010;15(4):303–9. <https://doi.org/10.1016/j.avb.2010.03.001>.
- Freedman SA, Gilad M, Ankri Y, Roziner I, Shalev AY. Social relationship satisfaction and PTSD: which is the chicken and which is the egg. *Eur J Psychotraumatol.* 2015;6(28864):1–7. <https://doi.org/10.3402/ejpt.v6.28864>.
- Ditlevsen DN, Elklit A. The combined effect of gender and age on post traumatic stress disorder: do men and women show differences in the lifespan distribution of the disorder? *Ann Gen Psychiatry.* 2010;9(1). <https://doi.org/10.1186/1744-859X-9-32>.
- Kobayashi I, Sledjeski EM, Delahanty DL. Gender and age interact to predict the development of posttraumatic stress disorder symptoms following a motor vehicle accident. *Psychol Trauma Theory Res Pract Policy.* 2019;11(3): 328–36. <https://doi.org/10.1037/tra0000366>.
- Harris IA, Young JM, Rae H, Jalaludin BB, Solomon MJ. Predictors of post-traumatic stress disorder following major trauma. *ANZ J Surg.* 2008;78(7): 583–7. <https://doi.org/10.1111/j.1445-2197.2008.04578.x>.
- Richardson LK, Frueh BC, Acierno R. Prevalence estimates of combat-related post-traumatic stress disorder: critical review. *Aust N Z J Psychiatry.* 2010; 44(1):4–19. <https://doi.org/10.3109/00048670903393597>.
- Dursa EK, Reinhard MJ, Barth SK, Schneiderman AI. Prevalence of a positive screen for PTSD among OEF/OIF and OEF/OIF-era veterans in a large population-based cohort. *J Trauma Stress.* 2014;27(5):542–9. <https://doi.org/10.1002/jts.21956>.
- Magruder K, Serpi T, Kimerling R, Kilbourne AM, Collins JF, Cypel Y, et al. Prevalence of posttraumatic stress disorder in Vietnam-era women veterans: the health of Vietnam-era women's study (HealthVIEWS). *JAMA Psychiatry.* 2015;72(11):1127–34. <https://doi.org/10.1001/jamapsychiatry.2015.1786>.

38. Machisa MT, Christofides N, Jewkes R. Mental ill health in structural pathways to women's experiences of intimate partner violence. *PloS one*. 2017;12(4):e0175240. <https://doi.org/10.1371/journal.pone.0175240>.
39. Mbalo N, Zhang M, Ntuli S. Risk factors for PTSD and depression in female survivors of rape. *Psychol Trauma Theory Res Pract Policy*. 2017;9(3):301–8. <https://doi.org/10.1037/tra0000228>.
40. Swain KD, Pillay BJ, Kiewer W. Traumatic stress and psychological functioning in a south African adolescent community sample. *S Afr J Psychiatry*. 2017;23(1):1–6. <https://doi.org/10.4102/sajpsychiatry.v23i0.1008>.
41. Getnet B, Medhin G, Alem A. Symptoms of post-traumatic stress disorder and depression among Eritrean refugees in Ethiopia: identifying direct, mediating and moderating predictors from path analysis. *BMJ Open*. 2019; 9(1):1–12. <https://doi.org/10.1136/bmjopen-2017-021142>.
42. Kagoyire MG, Richters A. "We are the memory representation of our parents": intergenerational legacies of genocide among descendants of rape survivors in Rwanda. *Torture J*. 2018;28(3):30–45. <https://doi.org/10.7146/torture.v28i3.111183>.
43. Sheikh TL, Mohammed A, Eseigbe E, Adekeye T, Nuhu FT, Lasisi M, et al. Descriptive characterization of psycho-trauma, psychological distress, and post-traumatic stress disorder among children and adolescent Internally Displaced persons in Kaduna, Nigeria. *Front Psychiatry*. 2016;7:179. <https://doi.org/10.3389/fpsy.2016.00179>.
44. Iteke O, Bakare MO, Agomoh AO, Uwakwe R, Onwukwe JU. Road traffic accidents and posttraumatic stress disorder in an orthopedic setting in south-Eastern Nigeria: a controlled study. *Scand J Trauma Resusc Emerg Med*. 2011;19(1):39. <https://doi.org/10.1186/1757-7241-19-39>.
45. Asuquo, J. E. et al. 'Depression and posttraumatic stress disorder among road traffic accident victims managed in a tertiary hospital in southern Nigeria'. 2017.
46. Mishara WL, Gbaden EA. The prevalence of depression among the youths as an aftermath of the internal insurgency attacks in Maiduguri, Nigeria. *IOSR J Hum Soc Sci (IOSR-JHSS)*. 2014;19(10):32–5. Available at: <http://www.iosrjournals.org/iosr-jhss/papers/Vol19-issue10/Version-7/F0191073235.pdf>. <https://doi.org/10.9790/0837-191073235>.
47. Wisco BE, Marx BP, Keane TM. Screening, Diagnosis, and Treatment of Post-Traumatic Stress Disorder. *Mil Med*. 2012;177:7–13.
48. Nwoga C, Audu M, Obembe A. Prevalence and correlates of posttraumatic stress disorder among medical students in the University of Jos, Nigeria. *Nigerian J Clin Pract*. 2016;19(5):595–9. <https://doi.org/10.4103/1119-3077.188704>.
49. Gapen M, Cross D, Ortigo K, Graham A, Johnson E, Evces M, et al. Perceived neighborhood disorder, community cohesion, and PTSD symptoms among low-income African Americans in an urban health setting. *Am J Orthopsychiatry*. 2011;81(1):31–7. <https://doi.org/10.1111/j.1939-0025.2010.01069.x>.
50. Goldmann E, Aiello A, Uddin M, Delva J, Koenen K, Gant LM, et al. Pervasive exposure to violence and PTSD in a predominantly African American Urban Community. *J Trauma Stress*. 2011;24(6):747–51. <https://doi.org/10.1002/jts.20705.Pervasive>.
51. Jain KM, Davey-Rothwell M, Crossnohere NL, Latkin CA. Post-traumatic stress disorder, neighborhood residency and satisfaction, and social network characteristics among underserved women in Baltimore, Maryland. *Womens Health Issues*. 2018;28(3):273–80. <https://doi.org/10.1016/j.whi.2018.02.004>.
52. Fourchard L. The case of Ibadan, Nigeria. Ibadan; 2003.
53. Coker AO. Challenges of urban housing quality and its associations with neighbourhood environments: insights and experiences of Ibadan city, Nigeria. *J Environ Health Res*. 2008. Available at: <http://tech.ui.edu.ng/sites/default/files/Challengesofurbanhousingqualityanditsassociationswithneighbourhoodenvironments.pdf> (Accessed 24 July 2019).
54. Mabogunje A. The growth of residential districts in Ibadan. *Geogr Rev*. 1962; 52(1):56–77. <https://doi.org/10.2307/213005>.
55. Neiderud CJ. How urbanization affects the epidemiology of emerging infectious diseases. *Afr J Disability*. 2015;5(1):1–9. <https://doi.org/10.3402/iee.v5.27060>.
56. Adiga A, Chu S, Eubank S, Kuhlman CJ, Lewis B, Marathe A, et al. Disparities in spread and control of influenza in slums of Delhi: findings from an agent-based modelling study. *BMJ Open*. 2018;8(1):1–12. <https://doi.org/10.1136/bmjopen-2017-017353>.
57. Lilford RJ, Oyeboode O, Satterthwaite D, Melendez-Torres GJ, Chen YF, Mberu B, et al. Improving the health and welfare of people who live in slums. *Lancet*. 2017;389(10068):559–70. [https://doi.org/10.1016/S0140-6736\(16\)31848-7](https://doi.org/10.1016/S0140-6736(16)31848-7).
58. Parto JA, Evans MK, Zonderman AB. Symptoms of posttraumatic stress disorder among urban residents. *J Nerv Ment Dis*. 2011;199(7):1–8. <https://doi.org/10.1097/NMD.0b013e3182214154>.
59. Golin CE, Haley DF, Wang J, Hughes JP, Kuo I, Justman J, et al. Post-traumatic stress disorder symptoms and mental health over time among low-income women at increased risk of HIV in the U. S HHS public access. *J Health Care Poor Underserved*. 2016;27(2):891–910. <https://doi.org/10.1353/hpu.2016.0093>.
60. Machisa MT, et al. The war at home: the Gauteng GBV indicators research study; 2011. <https://doi.org/10.13140/RG.2.1.4295.0007>.
61. Garcia-Moreno C, Jansen HAFM, Ellsberg M, Heise L, Watts CH. WHO multi-country study on Women's health and domestic violence against women. *World Health Organ Techn Rep Series*. 2005;368(9543):1–253. [https://doi.org/10.1016/S0140-6736\(06\)69523-8](https://doi.org/10.1016/S0140-6736(06)69523-8).
62. Maduka O, Akpan G, Maleghemi S. Using android and open data kit Technology in Data Management for research in resource-limited settings in the Niger Delta region of Nigeria: cross-sectional household survey. *JMIR mHealth and uHealth*. 2017;5(11):e171. <https://doi.org/10.2196/mhealth.7827>.
63. Steiner A, Hella J, Grüniger S, Mhalu G, Mhimbira F, Cercamondi CI, et al. Managing research and surveillance projects in real-time with a novel open-source e management tool designed for under-resourced countries. *J Am Med Inform Assoc*. 2016:1–8. <https://doi.org/10.1093/jamia/ocv185>.
64. NDHS NPC (NPC) [Nigeria] and I. I. 2014. N. D. and H. S. Abuja and Rockville: NPC and ICF International; 2013.
65. Mollica RF, et al. The Harvard trauma questionnaire: validating a cross-cultural instrument for measuring torture, trauma and posttraumatic stress disorder in Indochinese refugees. *J Nerv Ment Dis*. 1992;180(2):111–6. <https://doi.org/10.1097/00005053-199202000-00008>.
66. Shoeb M, Weinstein H, Mollica R. The Harvard trauma questionnaire: adapting a cross-cultural instrument for measuring torture, trauma and posttraumatic stress disorder in Iraqi refugees. *Int J Soc Psychiatry*. 2007; 53(5):447–63. <https://doi.org/10.1177/0020764007078362>.
67. Silove D, Manicavasagar V, Mollica R, Thai M, Khiek D, Lavelle J, et al. Screening for depression and PTSD in a Cambodian population unaffected by war: comparing the Hopkins symptom checklist and Harvard trauma questionnaire with the structured clinical interview. *J Nerv Ment Dis*. 2007; 195(2):152–7. <https://doi.org/10.1097/01.nmd.0000254747.03333.70>.
68. Rasmussen A, Verkuilen J, Ho E, Fan Y. Posttraumatic stress disorder among refugees: measurement invariance of Harvard trauma questionnaire scores across global regions and response patterns. *Psychol Assess*. 2015;27(4): 1160–70. <https://doi.org/10.1037/pas0000115>.
69. Housen T, Lenglet A, Ariti C, Shah S, Shah H, Ara S, et al. Prevalence of anxiety, depression and post-traumatic stress disorder in the Kashmir Valley. *BMJ Glob Health*. 2017;2(4):1–12. <https://doi.org/10.1136/bmjgh-2017-000419>.
70. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the childhood trauma questionnaire. *Child Abuse Negl*. 2003;27(2):169–90. [https://doi.org/10.1016/S0145-2134\(02\)00541-0](https://doi.org/10.1016/S0145-2134(02)00541-0).
71. Bernstein DP, Fink LA. CTQ: childhood trauma questionnaire: a retrospective self-report. San Antonio: Psychological Corporation; 1998.
72. Nöthling J, Suliman S, Martin L, Simmons C, Seedat S. Differences in abuse, neglect, and exposure to community violence in adolescents with and without PTSD and depression. *J Interpers Violence*. 2019;34(21–22):4357–83. <https://doi.org/10.1177/0886260516674944>.
73. Kisely S, Strathearn L, Najman JM. Child maltreatment and mental health problems in 30-year-old adults: a birth cohort study. *J Psychiatr Res*. 2020; 129:111–7. <https://doi.org/10.1016/j.jpsychires.2020.06.009>.
74. Ritacco G, Suffla S. A critical review of child maltreatment indices: psychometric properties and application in the south African context. *Afr Safety Promotion J Injury Violence Prev*. 2012;10(2):3–17.
75. Mall S, et al. The relationship between childhood trauma and schizophrenia in the genomics of schizophrenia in the Xhosa people (SAX) study in South Africa. 2019.
76. Brugha TS, Cragg D. The List of Threatening Experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand*. 1990;82(1): 77–81 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2399824>. Accessed 26 Nov 2017.

77. Sareen J. Posttraumatic stress disorder in adults: impact, comorbidity, risk factors, and treatment. *Can J Psychiatry*. 2014;59(9):460–7. <https://doi.org/10.1177/070674371405900902> *Revue canadienne de psychiatrie*.
78. Van Veldhoven LM, et al. Predictive ability of preinjury stressful life events and post-traumatic stress symptoms for outcomes following mild traumatic brain injury: analysis in a prospective emergency room sample. *J Neurol Neurosurg Psychiatry*. 2011;82(7):782–7. <https://doi.org/10.1136/jnnp.2010.228254>.
79. Motricó E, Moreno-Küstner B, de Dios Luna J, Torres-González F, King M, Nazareth I, et al. Psychometric properties of the list of threatening experiences-LTE and its association with psychosocial factors and mental disorders according to different scoring methods. *J Affect Disord*. 2013; 150(3):931–40. <https://doi.org/10.1016/j.jad.2013.05.017>.
80. Lovibond SH, Lovibond PF. Manual for the depression anxiety & stress scales. 2nd ed. Sydney: Psychology Foundation; 1995. Available at: <https://maic.qld.gov.au/wp-content/uploads/2016/07/DASS-21.pdf>
81. Parkitny L, McAuley J. The depression anxiety stress scale (DASS). *J Phys*. 2010;56(3):204. [https://doi.org/10.1016/S1836-9553\(10\)70030-8](https://doi.org/10.1016/S1836-9553(10)70030-8).
82. Henry JD, Crawford JR. The short-form version of the depression anxiety stress scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol*. 2005;44(2):227–39. <https://doi.org/10.1348/014466505X29657>.
83. Ng F, Trauer T, Dodd S, Callaly T, Campbell S, Berk M. The validity of the 21-item version of the depression anxiety stress scales as a routine clinical outcome measure. *Acta Neuropsychiatrica*. 2007;19(5):304–10. <https://doi.org/10.1111/j.1601-5215.2007.00217.x>.
84. Chan RCK, Xu T, Huang J, Wang Y, Zhao Q, Shum DHK, et al. Extending the utility of the depression anxiety stress scale by examining its psychometric properties in Chinese settings. *Psychiatry Res*. 2012;200(2–3):879–83. <https://doi.org/10.1016/j.psychres.2012.06.041>.
85. Chakraborty NM, Fry K, Behl R, Longfield K. Simplified asset indices to measure wealth and equity in health programs: a reliability and validity analysis using survey data from 16 countries. *Global Health Sci Pract*. 2016; 4(1):141–54. <https://doi.org/10.9745/GHSP-D-15-00384>.
86. Rutstein SO, Johnson K. The DHS wealth index. *DHS Comparative Rep*. 2004; 6:1–71. <https://doi.org/10.1017/CBO9781107415324.004>.
87. Menon B, Cherkil S, S. A. A. G. U. G. R. The process and challenges in the translation of World Health Organization quality of life (WHOQOL- BREF) to a regional language; Malayalam. *Indian J Psychol Med*. 2012;34(2):149–52. <https://doi.org/10.4103/0253-7176.101783>.
88. Mwayo AW, Ndeti DM, Mutiso V, Khasakhalo LI. Traditional healers and provision of mental health services in cosmopolitan informal settlements in Nairobi, Kenya. *Afr J Psychiatry (South Africa)*. 2013;16(2):134–40. <https://doi.org/10.4314/ajpsy.v16i2.17>.
89. Ndungu J, Ramsommar L, Willan S, Washington L, Ngcobo-Sithole M, Gibbs A. Depression, posttraumatic stress disorder (PTSD) and their comorbidity: implications of adversity amongst young women living in informal settlements in Durban, South Africa. *J Affect Disorders Rep*. 2020;1:100022. <https://doi.org/10.1016/j.jadr.2020.100022>.
90. Lee JS, Ahn YS, Jeong KS, Chae JH, Choi KS. Resilience buffers the impact of traumatic events on the development of PTSD symptoms in firefighters. *J Affect Disord*. 2014;162:128–33. <https://doi.org/10.1016/j.jad.2014.02.031>.
91. Yuan C, Wang Z, Inslicht SS, McCaslin SE, Metzler TJ, Henn-Haase C, et al. Protective factors for posttraumatic stress disorder symptoms in a prospective study of police officers. *Psychiatry Res*. 2011;188(1):45–50. <https://doi.org/10.1016/j.psychres.2010.10.034>.
92. Fergerson, A. K. and Brausch, A. M. (2020) 'Resilience mediates the relationship between PTSD symptoms and disordered eating in college women who have experienced sexual victimization', *J Interpers Violence*, pp. 1–18. doi: <https://doi.org/10.1177/0886260520918581>, 0886260520918581.
93. Powers A, Etkin A, Gyurak A, Bradley B, Jovanovic T. Associations between childhood abuse, posttraumatic stress disorder, and implicit emotion regulation deficits: evidence from a low-income Inner City population. *Psychiatry*. 2015;78(3):251–64. <https://doi.org/10.1080/00332747.2015.1069656>.
94. Simonelli A, et al. Child abuse predicts adult PTSD symptoms among individuals diagnosed with intellectual disabilities. *Front Psychol*. 2015;6:1–11. <https://doi.org/10.3389/fpsyg.2015.01600>.
95. Sanchez SE, Pineda O, Chaves DZ, Zhong QY, Gelaye B, Simon GE, et al. Childhood physical and sexual abuse experiences associated with PTSD among pregnant women. *Ann Epidemiol*. 2017;27(11):716–723.e1. <https://doi.org/10.1016/j.annepidem.2017.09.012>.
96. Haileye A. Psychopathological correlates of child sexual abuse : the case of female students in jimma zone , South West Ethiopia. *Ethiopian J Health Sci*. 2013;23:32–8.
97. Ombok CA, Obondo A, Kangethe R, Atwoli L. The prevalence of post-traumatic stress disorder among sexually abused children at Kenyatta National Hospital in Nairobi, Kenya. *East Afr Med J*. 2013;90(10):332–7.
98. Choi KW, Sikkema KJ, Velloza J, Marais A, Jose C, Stein DJ, et al. Maladaptive coping mediates the influence of childhood trauma on depression and PTSD among pregnant women in South Africa. *Arch Womens Ment Health*. 2015;18(5):731–8. <https://doi.org/10.1007/s00737-015-0501-8>.Maladaptive.
99. Mutavi T, Mathai M, Obondo A. Post-traumatic stress disorder (PTSD) in sexually abused children and educational status in Kenya: a longitudinal study. *J Child Adolescent Behav*. 2017;05(05). <https://doi.org/10.4172/2375-4494.1000357>.
100. Charak R, de Jong JTVM, Berckmoes LH, Ndayisaba H, Reis R. Assessing the factor structure of the childhood trauma questionnaire, and cumulative effect of abuse and neglect on mental health among adolescents in conflict-affected Burundi. *Child Abuse Negl*. 2017;72(September):383–92. <https://doi.org/10.1016/j.chiabu.2017.09.009>.
101. McLaughlin KA, Koenen KC, Bromet EJ, Karam EG, Liu H, Petukhova M, et al. Childhood adversities and post-traumatic stress disorder: evidence for stress sensitisation in the world mental health surveys. *Br J Psychiatry*. 2017;211(5): 280–8. <https://doi.org/10.1192/bjp.bp.116.197640>.
102. Koenen KC, Ratanatharathorn A, Ng L, McLaughlin KA, Bromet EJ, Stein DJ, et al. Posttraumatic stress disorder in the world mental health surveys. *Psychol Med*. 2017;47(13):2260–74. <https://doi.org/10.1017/S0033291717000708>.
103. Mahenge B, Likindikoki S, Stöckl H, Mbwambo J. Intimate partner violence during pregnancy and associated mental health symptoms among pregnant women in Tanzania: a cross-sectional study. *BJOG*. 2013;120(8): 940–7. <https://doi.org/10.1111/1471-0528.12185>.
104. Pengpid S, Peltzer K. Associations of physical partner violence and sexual violence victimization on health risk behaviours and mental health among university students from 25 countries; 2020. p. 1–10.
105. Spinhoven P, Penninx B. Childhood trauma questionnaire: factor structure, measurement invariance, and validity across emotional disorders. *Psychol Assess*. 2014;26(3):717–29. <https://doi.org/10.1037/pas0000002>.
106. Ginzburg K, Ein-Dor T, Solomon Z. Comorbidity of posttraumatic stress disorder, anxiety and depression: a 20-year longitudinal study of war veterans. *J Affect Disord*. 2010;123(1–3):249–57. <https://doi.org/10.1016/j.jad.2009.08.006>.
107. Gros DF, et al. Symptom overlap in PTSD and major depression. *Psychiatry Res*. 2012;195(2–3):267–70. <https://doi.org/10.1038/jid.2014.371>.
108. Spinhoven P, Penninx BW, van Hemert AM, de Rooij M, Elzinga BM. Comorbidity of PTSD in anxiety and depressive disorders: prevalence and shared risk factors. *Child Abuse Negl*. 2014;38(8):1320–30. <https://doi.org/10.1016/j.chiabu.2014.01.017>.
109. McLaughlin KA, Choi K, et al. Childhood adversity, adult stressful life events and risk of past year psychiatric disorder: a test of the stress sensitization hypothesis in a population based sample of adults. *Psychol Med*. 2010;40(10):1–23. <https://doi.org/10.1038/jid.2014.371>.
110. Kempainen JK, MacKain S, Alexander M, Reid P, Jackson MP. Posttraumatic stress disorder and stressful life events among rural women with HIV disease. *J Assoc Nurses AIDS Care*. 2017;28(2):216–25. <https://doi.org/10.1016/j.jana.2016.06.001>.
111. Mahenge B, Stöckl H, Likindikoki S, Kaaya S, Mbwambo J. The prevalence of mental health morbidity and its associated factors among women attending a prenatal clinic in Tanzania. *Int J Gynecol Obstet*. 2015;130(3): 261–5. <https://doi.org/10.1016/j.ijgo.2015.04.032>.
112. Nothling J, et al. Traumatic dissociation as a predictor of posttraumatic stress disorder in south African female rape survivors. *Medicine*. 2015;94(16): 1–9. <https://doi.org/10.1097/MD.0000000000000744>.
113. Soltys A, Tyburski E. Predictors of mental health problems in formal and informal caregivers of patients with Alzheimer's disease. *BMC Psychiatry*. 2020;20:1–11.
114. Xue C, Ge Y, Tang B, Liu Y, Kang P, Wang M, et al. A meta-analysis of risk factors for combat-related PTSD among military personnel and veterans. *PLoS One*. 2015;10(3):1–21. <https://doi.org/10.1371/journal.pone.0120270>.



115. Asnakew S, Shumet S, Ginbare W, Legas G, Haile K. Prevalence of post-traumatic stress disorder and associated factors among Koshe landslide survivors, Addis Ababa, Ethiopia: a community-based, cross-sectional study. *BMJ Open*. 2019;9(6):1–8. <https://doi.org/10.1136/bmjopen-2018-028550>.
116. Taru MY, et al. Posttraumatic stress disorder among internally displaced victims of boko haram terrorism in North-Eastern Nigeria. *Jos J Med*. 2018; 12(1):8–15.
117. Mugisha J, et al. Prevalence and factors associated with PTSD seven years after the conflict in three districts in northern Uganda. *BMC Psychiatry*. 2015; 15(170):1–12. Available at: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4513792/pdf/12888\\_2015\\_Article\\_551.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4513792/pdf/12888_2015_Article_551.pdf). Accessed 11 Sept 2019.
118. McLaughlin KA, Lambert HK. Child trauma exposure and psychopathology: mechanisms of risk and resilience. *Curr Opin Psychol*. 2017;14:29–34. <https://doi.org/10.1016/j.copsyc.2016.10.004>.
119. Gruebner O, Rapp MA, Adli M, Kluge U, Galea S, Heinz A. Cities and mental health. *Deutsches Ärzteblatt*. 2017;114(8):121–7. <https://doi.org/10.3238/aerztebl.2017.0121>.
120. De Vries E, et al. Housing index, urbanisation level and lifetime prevalence of depressive and anxiety disorders: a cross-sectional analysis of the Colombian national mental health survey. *BMJ Open*. 2018;8(6):1–11. <https://doi.org/10.1136/bmjopen-2017-019065>.
121. Adams RE, Boscarino JA. Predictors of PTSD and delayed PTSD after disaster: the impact of exposure and psychosocial resources. *J Nerv Ment Dis*. 2006; 194(7):485–93. <https://doi.org/10.1097/01.nmd.0000228503.95503.e9>.
122. Skogstad M, Skorstad M, Lie A, Conradi HS, Heir T, Weisaeth L. Work-related post-traumatic stress disorder. *Occup Med*. 2013;63(3):175–82. <https://doi.org/10.1093/occmed/kqt003>.
123. Schwartz E, Shrira A. Social connectedness moderates the relationship between warfare exposure, PTSD symptoms, and health among older adults. *Psychiatry (New York)*. 2018;00(00):1–15. <https://doi.org/10.1080/00332747.2018.1534521>.
124. Kintzle S, Barr N, Corletto G, Castro C. PTSD in U.S. veterans: the role of social connectedness, combat experience and discharge. *Healthcare*. 2018; 6(3):102. <https://doi.org/10.3390/healthcare6030102>.
125. Kirmayer LJ, Gomez-Carrillo A, Veissière S. Culture and depression in global mental health: an ecosocial approach to the phenomenology of psychiatric disorders. *Soc Sci Med*. 2017;183:163–8. <https://doi.org/10.1016/j.socscimed.2017.04.034>.
126. Mayston R, Frissa S, Tekola B, Hanlon C, Prince M, Fekadu A. Explanatory models of depression in sub-Saharan Africa: synthesis of qualitative evidence'. *Social Sci Med*. 2020;246:112760. <https://doi.org/10.1016/j.socscimed.2019.112760>.
127. Kilian S, Swartz L, Dowling T, Dlaki M, Chiliza B. The potential consequences of informal interpreting practices for assessment of patients in a south african psychiatric hospital. *Soc Sci Med*. 2014;106:159–67. <https://doi.org/10.1016/j.socscimed.2014.01.019>.
128. Kilian S, Swartz L, Hunt X, Benjamin E, Chiliza B. When roles within interpreter-mediated psychiatric consultations speak louder than words. *Transcult Psychiatry*. 2020;58(1):27–37. <https://doi.org/10.1177/1363461520933768>.
129. Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall R. The psychological risks of Vietnam for U.S. veterans: a revisit with new data and methods. *Science*. 2006;313(5789):979–82. <https://doi.org/10.1126/science.1128944>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

