

POSTER PRESENTATION

Open Access

The prevalence of metabolic syndrome up to 5 years post-partum in patients with a history of gestational diabetes mellitus

C Crowe^{1*}, E Noctor¹, LA Carmody¹, B Wickham¹, G Avalos¹, G Gaffney², P O'Shea³, F Dunne¹,
the ATLANTIC DIP Collaborators

From International Conference for Healthcare and Medical Students 2011
Dublin, Ireland. 4-5 November 2011

Introduction

Metabolic syndrome (MetS) is associated with cardiovascular mortality and increased risk of type 2 diabetes.

Methods

We examine the prevalence of MetS in a cohort of Caucasian women with previous gestational diabetes (GDM) (n=116), and those with normal glucose tolerance (NGT) during pregnancy (n=51). Fasting glucose alone (known DM/pre-diabetes post-partum patients) or 75g OGTT (other patients), lipid profile, insulin and c-peptide were performed. We calculated insulin resistance using the HOMA2-IR computer model.

Results

Conclusions

Metabolic syndrome and insulin resistance are significantly more prevalent in Caucasian patients with GDM progressing to post-partum DM/pre-diabetes than those who do not ($p < 0.01$), suggesting a need to target lifestyle changes in the early post-partum period to help prevent progression to T2DM.

Author details

¹Department of Medicine, University Hospital Galway and National University of Ireland, Galway, Ireland. ²Department of Obstetrics and Gynaecology, University Hospital Galway, Ireland. ³Department of Clinical Biochemistry, University Hospital Galway, Ireland.

Published: 9 July 2012

doi:10.1186/1753-6561-6-S4-P44

Cite this article as: Crowe et al.: The prevalence of metabolic syndrome up to 5 years post-partum in patients with a history of gestational diabetes mellitus. *BMC Proceedings* 2012 **6**(Suppl 4):P44.

Table 1

	Number with MetS (ATPIII criteria) (%)	Mean HOMA-2IR (+/- SD)	Number with HOMA-2IR>1.7 (%)
DM/pre-diabetes post GDM (n=33)	12 (36%)	1.21(+/-0.77)	18 (55%)
NGT post GDM (n= 83)	11 (13%)	1.93(+/-1.4)	18 (22%)
No history GDM (n=52)	3 (6%)	1.02(+/-0.85)	6 (13%)

¹Department of Medicine, University Hospital Galway and National University of Ireland, Galway, Ireland

Full list of author information is available at the end of the article