



Seronegative myasthenic crisis: a multicenter analysis

Philipp Mergenthaler^{1,2,3} · Henning R. Stetefeld⁴ · Christian Dohmen^{4,5} · Siegfried Kohler¹ · Silvia Schönenberger⁶ · Julian Bösel^{6,7} · Stefan T. Gerner^{8,9} · Hagen B. Huttner^{8,9} · Hauke Schneider^{10,11} · Heinz Reichmann¹⁰ · Hannah Fuhrer¹² · Benjamin Berger¹² · Jan Zinke¹³ · Anke Alberty¹⁴ · Ingo Kleiter^{15,16} · Christiane Schneider-Gold¹⁵ · Christian Roth^{17,18} · Juliane Dunkel¹⁷ · Andreas Steinbrecher¹⁹ · Andrea Thieme¹⁹ · De-Hyung Lee²⁰ · Ralf A. Linker²⁰ · Klemens Angstwurm²⁰ · Andreas Meisel^{1,2,21} · Bernhard Neumann^{20,22} · for The German Myasthenic Crisis Study Group

Received: 19 November 2021 / Revised: 7 February 2022 / Accepted: 10 February 2022 / Published online: 7 April 2022
© The Author(s) 2022

Abstract

Myasthenic crisis (MC) is a life-threatening condition for patients with myasthenia gravis (MG). Seronegative patients represent around 10–15% of MG, but data on outcome of seronegative MCs are lacking. We performed a subgroup analysis of patients who presented with MC with either acetylcholine-receptor-antibody-positive MG (AChR-MG) or seronegative MG between 2006 and 2015 in a retrospective German multicenter study. We identified 15 seronegative MG patients with 17 MCs and 142 AChR-MG with 159 MCs. Seronegative MCs were younger (54.3 ± 14.5 vs 66.5 ± 16.3 years; $p = 0.0037$), had a higher rate of thymus hyperplasia (29.4% vs 3.1%; $p = 0.0009$), and were more likely to be female (58.8% vs 37.7%; $p = 0.12$) compared to AChR-MCs. Time between diagnosis of MG and MC was significantly longer in seronegative patients (8.2 ± 7.6 vs 3.1 ± 4.4 years; $p < 0.0001$). We found no differences in duration of mechanical ventilation (16.2 ± 15.8 vs 16.5 ± 15.9 days; $p = 0.94$) and length of stay at intensive care unit (17.6 ± 15.2 vs 17.8 ± 15.4 days; $p = 0.96$), or in-hospital mortality (11.8% vs. 10.1%; $p = 0.69$). We conclude that MC in seronegative MG affects younger patients after a longer period of disease, but that crisis treatment efficacy and outcome do not differ compared to AChR-MCs.

Keywords Myasthenia gravis · Myasthenic crisis · Antibody status · Seronegative · Outcome

Introduction

Myasthenia gravis (MG) is an autoimmune disease with antibodies (Abs) targeting the postsynaptic neuromuscular junction. Ultimately, muscle fatigability and weakness are caused by disrupted neuromuscular signaling. Nearly 90% of all MG patients have positive test results for AChR, Muscle-specific kinase (MuSK), or low-density lipoprotein receptor-related protein (LRP4) autoantibodies, with the majority tested positive for AChR-Abs [1]. However, in around 10–15% of MG patients no specific autoantibodies can be found. This group of seronegative patients is also thought to include patients

with very low antibody titers, low-affinity antibodies and yet to be defined autoantigens [1].

Myasthenic crisis (MC) is the most severe form of MG and is potentially life threatening. MC is mostly provoked by infections, but also fever, aspiration, inadequate treatment, various medications, or following surgery [2]. In the first two years after diagnosis, around 15–20% of MG patients suffer from a MC [2, 3]. Characteristic symptoms are extensive weakness, dysphagia, and dyspnea which can result in respiratory insufficiency. The clinical management of MC is well defined and has led to a significant decline in mortality from around 40% in the early 1960s to 5% to 12% in recent studies [2–8].

However, to date little is known about the management of MC in seronegative MG. Here, we therefore investigated seronegative patients with MC and compared their crises to AChR-MCs regarding clinical features, therapeutic management, and outcome.

Bernhard Neumann: on behalf of the Initiative of German NeuroIntensive Trial Engagement (IGNITE) and with support from the German Neurocritical Care Society (DGNI)

✉ Bernhard Neumann
bernhard.neumann@donau-issar-klinikum.de

Extended author information available on the last page of the article

Methods

Study design and patient selection

We performed a subgroup analysis of seronegative MC needing mechanical ventilation (MV) compared to AChR-MC treated at eight German Departments of Neurology with specialized Neuro-Intensive Care Units (NICU) or neurologically associated interdisciplinary ICU [2]. For identification, records of all patients discharged with the diagnosis of MG according to the International Classification of Diseases (ICD10: G70.0–70.3) who were treated and ventilated on an ICU between 2006 and 2015 were reviewed. MC was defined as an exacerbation of myasthenic symptoms with bulbar and/or general weakness requiring MV. Seronegative MG was defined as absence of AChR and MuSK autoantibodies. Per protocol, antibody status was confirmed by routine laboratory testing using certified assays. Most AChR-Abs and MuSK-Abs were tested by radio-receptor assay, but the method is not known in all cases due to the retrospective character. Diagnosis of MG had to be established clinically according to national guidelines and confirmed by specific tests (antibody testing or repetitive stimulation or improvement after cholinergic medication) [9]. New episodes of MC were counted separately if patients were discharged in their prehospital status and if new triggers for the next crisis could be determined. For this analysis, we only included AChR-MCs treated at the same centers as the seronegative MCs to reduce treatment and data acquisition bias.

Data acquisition

Data on baseline demographics, clinical information, medication and comorbidities were obtained through review of medical records and institutional databases. Characteristics reviewed included antibody-status, evidence of thymoma and Myasthenia Gravis Foundation of America (MGFA) Score prior to MC. Assessed treatment regimens were intravenous immunoglobulins (IVIG), plasma exchanging therapy (PE), immunoadsorption (IA), use of intravenous pyridostigmine, and continuous potassium infusion. Analyzed data regarding the MC included time at intensive care unit (ICU-LOS), days in hospital, duration of MV, in-hospital mortality and referral/discharge. In addition, we performed a survey about LRP4- and Agrin-antibody-positive MGs in our study group in June 2021.

Statistics

GraphPad Prism 5® (GraphPad Software, La Jolla, USA) was used for statistical analysis. Data were presented as

mean with standard deviation or range (as indicated) or total number with percentage. Group comparison was tested with either Student's *t* test or Fisher's exact test (with odds ratios (OR)). The significance level was set to $\alpha=0.05$ both sided.

Results

Characteristics of study group

The cohort consisted of 15 patients with 17 seronegative MCs and 142 AChR antibody-positive patients with 159 MCs requiring MV. Patients from both groups were treated at the same centers (Table 1). Seronegative patients were responsible for 6.8% of the crises in our whole cohort ($n=250$ crises). Patients with seronegative MC were significantly younger (54.3 ± 14.5 vs 66.5 ± 16.3 ; $p=0.0037$) and more likely to be female (58.8 vs 37.7% ; $p=0.12$) than AChR-MCs. AChR-MCs were significantly more often late-onset MGs (85.5% vs 41.2% ; $p=0.0001$; OR = 0.12), whereas seronegative MCs belonged mainly to the early-onset group (Table 1) and had significantly more frequently a thymus hyperplasia (29.4% vs 3.1% ; $p=0.0009$; OR 12.83). Thymus hyperplasias were resected in all patients prior to crisis, except in one patient in the AChR-group. Importantly, the time between diagnosis of MG and onset of MC was significantly longer in seronegative patients (8.2 ± 7.6 vs 3.1 ± 4.4 years; $p<0.0001$) (Fig. 1A). Due to the higher age, patients with AChR-MCs had more comorbidities, yet without reaching statistical significance (Table 1). We also did not find statistically significant differences in the status before crisis, MGFA classification before crisis, dosage of pyridostigmine treatment before crisis (252.4 ± 243.3 vs 251.1 ± 206.6 mg/d; $p=0.99$) or number of myasthenic worsening/crises before present MC (Table 1). The number of days between first symptoms of MC and hospitalization were similar (9.9 ± 13.1 vs 9.6 ± 14.9 ; $p=0.95$).

To further characterize the patients with seronegative MCs, we surveyed all participating centers on retesting for LRP4 and Agrin antibodies in these seronegative patients. 6 of 15 seronegative MGs were tested for LRP4-antibodies and 5 of 15 for Agrin-antibodies. However, all tests remained negative. At the centers of our study group 28 patients with LRP4 and 0 patients with Agrin-antibodies are treated, but none developed an MC needing ICU-treatment within the period of observation.

Treatment and outcome

Seronegative MCs were significantly less frequently treated with IVIGs (23.5% vs 58.5% ; $p=0.009$; OR = 0.22) (Table 1), and although they received PE or IA more frequently than AChR-MCs, no statistically significant

Table 1 Comparison of episodes of myasthenic crisis with AChR-Abs and seronegative patients


Myasthenic crises	AChR-positive (<i>n</i> = 159)	Seronegative (<i>n</i> = 17)	<i>P</i> value	Odds ratio
Age	66.5 ± 16.3 (14–89)	54.3 ± 14.5 (25–81)	0.0037	
Male/female	99 (62.3%)/60 (37.7%)	7 (41.2%)/10 (58.8%)	0.12	0.42
Pulmonary disease	38 (23.9%)	2 (11.8%)	0.37	0.42
Heart disease	61 (38.4%)	5 (29.4%)	0.60	0.67
Diabetes mellitus	48 (30.2%)	2 (11.8%)	0.16	0.31
Tumour (other than thymoma)	24 (15.1%)	0 (0%)	0.13	0.16
Dialysis	1 (0.6%)	0 (0%)	1.00	3.02
Smoker	16 (10.1%)	2 (11.8%)	0.69	1.19
Alcohol addicted	5 (3.1%)	2 (11.8%)	0.14	4.11
≥ 3 diseases (kidney, heart, lung, diabetes, tumour)	21 (13.2%)	0 (0%)	0.23	0.18
Myasthenia gravis				
Early onset	22 (13.8%; 1 unknown)	10 (58.8%)	< 0.0001	8.90
Late onset	136 (85.5%)	7 (41.2%)	0.0001	0.12
Paraneoplastic MG (Thymoma)	53 (33.3%)	3 (17.6%)	0.27	0.43
Thymus hyperplasia	5 (3.1%)	5 (29.4%)	0.0009	12.83
Number of myasthenic worsenings/crises before present myasthenic crisis	0.7 ± 1.3	0.9 ± 1.3	0.49	
Time between first diagnosis and crisis (years)	3.1 ± 4.4 (0–18.2)	8.2 ± 7.6 (0–22)	< 0.0001	
MGFA-classification before crisis				
First manifestation of MG	38 (23.9%)	3 (17.6%)	0.77	0.68
Class I	8 (5.0%)	1 (5.9%)	1.00	1.18
Class II	44 (27.7%)	4 (23.5%)	1.00	0.80
Class III	40 (25.2%)	4 (23.5%)	1.00	0.92
Class IV	16 (10.1%)	2 (11.8%)	0.69	1.19
Unknown	13 (8.2%)	2 (11.8%)		
Status before crisis				
Independent at home	73 (45.9%)	7 (41.2%)	0.80	0.82
At home dependent on help	23 (14.5%)	3 (17.6%)	0.72	1.27
In a care facility or hospital	49 (30.8%)	7 (41.2%)	0.42	1.57
Unknown	14 (8.8%)	0 (0%)		
Cause of crisis				
Infection	80 (50.3%)	8 (47.1%)	n.s	
First episode	36 (22.6%)	3 (17.6%)		
Poor treatment compliance	14 (8.8%)	0 (0%)		
Intake of contraindicated medication	2 (1.2%)	0 (0%)		
Idiopathic/unknown	32 (20.1%)	6 (35.3%)		
Therapy				
IVIG	93 (58.5%)	4 (23.5%)	0.009	0.22
Plasma exchange/immunoabsorption in total	75 (47.2%)	10 (58.8%)	0.45	1.60
PE or IA as first line therapy	60 (37.7%)	10 (58.8%)	0.12	2.34
IVIG + plasma exchange or immunoabsorption	30 (18.9%)	1 (5.9%)	0.31	0.27
Continuous pyridostigmine infusion	61 (38.4%)	5 (29.4%)	0.60	0.67
Complications				
CPR	19 (11.9%)	2 (11.8%)	1.00	0.98
Pneumonia	86 (54.1%)	9 (52.9%)	1.00	0.95
Sepsis	32 (20.1%)	3 (17.6%)	1.00	0.85
Outcome				
Days of mechanical ventilation at ICU	19.2 ± 19.5 (1–119)	16.2 ± 15.8 (1–55)	0.54	
Days at ICU	22.0 ± 20.5 (1–135)	17.6 ± 15.2 (3–56)	0.42	
Days in hospital	30.8 ± 21.4 (3–144)	29.9 ± 16.5 (3–71)	0.87	
In-hospital mortality	16 (10.1%)	2 (11.8%)	0.69	1.19

Age, “Days of mechanical ventilation at ICU”, “Days at ICU”, “Days in hospital” and “Time between first diagnosis and crisis (years)” are depicted as mean ± Standard Deviation and range, other parameters are total number with percentage in brackets. *MGFA* Myasthenia Gravis Foundation of America, *MG* Myasthenia Gravis, *IVIG* Intravenous Immunoglobulin, *PE* Plasma exchange, *IA* Immunoabsorption, *CPR* Cardio

Table 1 (continued)

Pulmonary Resuscitation, *n.s.* not significant. *t* test was used for statistical analysis of age-differences and for comparison of “Days of mechanical ventilation at ICU”, “Days at ICU” and “Days in hospital”. For other parameters Fisher’s exact test with odds ratio was used. Significant result ($p \leq 0.05$) are shown in bold letters

Fig. 1 **A** Time between first diagnosis of MG to first MC in years. Every dot or square symbolizes one patient. Long line shows mean, short lines show SD (*t* test). *** $p < 0.001$ **B** Days of mechanical ventilation, **C** Days at ICU, **D** Days in hospital in 159 MCs with AChR-Abs and 17 MCs with MuSK-Abs. **B–D** Bars show mean \pm SD (*t* test). No significant results were found



difference could be found (58.8% vs 47.2%; $p = 0.45$; OR = 1.60). Likewise, although AChR-MCs were more frequently treated with the combination of PE or IA and IVIG, this was not statistically significant (18.9% vs 5.9%; $p = 0.31$; OR = 0.27). Furthermore, days of MV at ICU (16.2 ± 15.8 vs 19.2 ± 19.5 ; $p = 0.54$), ICU-LOS (17.6 ± 15.2 vs 22.0 ± 20.5 ; $p = 0.42$) and hospital-LOS (29.9 ± 16.5 vs 30.8 ± 21.4 ; $p = 0.87$) were not statistically significantly different (Table 1 and Fig. 1B–D). The in-hospital mortality was similar between both groups (11.8% vs 10.1%; $p = 0.69$; OR = 1.19). Importantly, seronegative patients were less frequently discharged while still needing MV (5.9% vs 20.8%; $p = 0.20$; OR = 0.24) compared to AChR-MG patients, but without a statistically significant difference. Consequently, after matching 17 seronegative to 51 AChR-positive MCs regarding age (54.3 ± 14.5 vs 53.9 ± 16.3 ; $p = 0.93$) and sex (58.8% vs 53.0% female; $p = 0.78$) we found no differences in days of MV (16.2 ± 15.8 vs 16.5 ± 15.9 ; $p = 0.94$) and

ICU-LOS (17.6 ± 15.2 vs 17.8 ± 15.4 ; $p = 0.96$). Treatment and outcome details for all 17 seronegative MCs are shown in Table 2. We found no difference between the treatment options IVIG and PE/IA or the additional use of intravenous pyridostigmine regarding the endpoint duration of MV in seronegative MCs.

Discussion

Here, we investigated clinical features of seronegative MC compared to AChR-MC requiring mechanical ventilation based on our multicenter cohort of MC [2]. In contrast to MuSK-MCs, which are as old as AChR-MCs [5], we found that in seronegative MC, patients were younger but that the time between diagnosis of MG and onset of MC was longer compared to AChR-MC. Interestingly, 8 of 17 seronegative MCs had thymic abnormalities. Although seronegative MCs

Table 2 Treatment and outcome details of all 17 seronegative MCs

Patient no.	Age	IVIG	PE/IA	Pyridostigmine intravenous	Days of mechanical ventilation	Death
1	80	5 × 30 g	No	No	38	No
2	66	No	5 cycles of PE	10,8 mg/24 h	39	No
3	25	No	5 cycles of PE	9,6 mg/24 h	20	No
4	45	3 × 30 g	No	No	3	No
5	48	No	No	7,2 mg/24 h	12	No
6	46	5 × 20 g	No	2,4 mg/24 h	13	No
7	58	No	No	No	30	Septic shock
8	55	No	5 cycles of PE	12 mg/24 h	5	No
9	81	5 × 30 g	5 cycles of IA	No	26	No
10	68	No	No	No	1	Septic shock
11	60	No	5 cycles of IA	No	55	No
12	45	No	7 cycles of PE	No	10	No
13	47	No	5 cycles of PE	No	2	No
14	66	No	unknown cycles of PE	No	3	No
15	52	No	6 cycles of PE	No	8	No
16	42	No	3 cycles of PE	No	4	No
17	39	No	5 cycles of PE	No	16	No

IVIG intravenous immunoglobulin, PE plasma exchange, IA immunoadsorption

were less frequently treated with IVIg, there was no difference in other MC treatments. Furthermore, we did not find any difference in baseline characteristics, in the rate of complications or outcome between the patient groups, which is in contrast to more severely affected MuSK MC patients [5].

Seronegative patients represent 10–15% of all MGs. However, there are only very limited data on the clinical management and outcome of MC. In our cohort of MCs, 17 of 250 (6.8%) events were from seronegative patients [2], which may indicate that MC is less prevalent in seronegative patients. Yet, our study was not designed to unambiguously address this question. While MC occurs in most MG patients within the first 2 years after diagnosis [3], seronegative patients in our cohort developed MC significantly later compared to AChR-MG thus suggesting a less severe disease onset in these patients.

Interestingly, LRP4-positive MGs treated at our centers until now never experienced a MC and all retested seronegative patients in our cohort (40%) were negative for LRP4. Moreover, we did not find any publication about a LRP4-MC suggesting that this subgroup is even less severely affected than seronegative MG.

Seronegative MG patients are a heterogeneous group of patients and although we had stringent inclusion criteria, we cannot rule out that some seronegative patients have low-affinity antibodies or would be positive for complement deposition at the neuromuscular junction [10, 11]. Especially the high portion of thymus hyperplasia in our cohort might argue for low-affinity antibodies since thymic pathologies in MG are known to produce AChR-antibodies [12], but 75% of

double negative patients (AChR and MuSK) showed lymph node-type infiltrates in thymus similar to AChR-MG [13].

Limitations of this study arise from its retrospective nature and the relatively small sample size. Nevertheless, our study is by far the largest analysis of seronegative MCs and therefore provides important evidence on the treatment of this understudied patient population. Nevertheless, large prospective multicenter studies are needed to further elucidate the character of seronegative myasthenic crisis and whether specific treatment is warranted compared to AChR-MC or MuSK-MC. Another limitation is that antibody tests were done in different labs and therefore false negative or false positive results due to unspecificity of the test technique or positive/negative results near the threshold range cannot be ruled out in every case, like in previous studies in myasthenia gravis.

We conclude that patients with seronegative MC are younger, with a longer course of disease until first crisis needing MV than AChR-MC but that there is no difference in outcome between these patient groups.

Acknowledgements Co-investigators: Kornelius Fuchs, (Department of Neurology, University Medical Center Regensburg, Regensburg, Germany): Data acquisition, supervision of personnel. Berthold Schalke (Department of Neurology, University Medical Center Regensburg, Regensburg, Germany): Data acquisition, supervision of personnel. Amelie Vidal (Department of Neurology, University Medical Center Regensburg, Regensburg, Germany): Data acquisition. Izabela Brachaczek, Jana Maidhof (Charité-Universitätsmedizin Berlin, Berlin, Department of Neurology, NeuroCure Clinical Research Center): Data acquisition. Arno Wenke (Charité-Universitätsmedizin Berlin, Berlin, Department of Neurology, NeuroCure Clinical Research Center): Data

acquisition, supervision of personnel. Manuel Hagen (Department of Neurology, University Hospital Erlangen, Erlangen, Germany): Data acquisition. Jan Rahmig (Department of Neurology, University Hospital, Technische Universität Dresden, Dresden, Germany): Data acquisition. Eik Schimmel (Department of Neurology, University Hospital, Technische Universität Dresden, Dresden, Germany. Now: Department of Neurology, Staedtisches Klinikum Dresden, Dresden, Germany): Data acquisition. Wolf Niesen (Department of Neurology, University of Freiburg, Freiburg, Germany): Data acquisition, supervision of personnel. Christine Fahrenndorf (Department of Neurology, St. Josef Hospital, Ruhr University Bochum, Bochum, Germany): Data acquisition, supervision of personnel.

Author contributions All authors have made a substantial contribution to the design, data collection and analysis of the research and the drafting of the manuscript and reviewed and accepted the contents of the manuscript prior to its submission. PM, HRS, AM, and BN: conception and design of the study, acquisition and analysis of data and drafting or revising a significant portion of the manuscript or figures. CD and KA: conception and design of the study and drafting or revising a significant portion of the manuscript or figures. SK, STG, HBH, HF, JZ, AA, IK, CS-G, CR, JD, AS, and AT: acquisition and analysis of data. SS, JB, and HS: acquisition and analysis of data and drafting or revising a significant portion of the manuscript or figures. HR, BB, D-HL, and RL: drafting or revising a significant portion of the manuscript or figures.

Funding Open Access funding enabled and organized by Projekt DEAL. PM is Einstein Junior Fellow of the Einstein Foundation Berlin and has been supported by the Charité-BIH Clinician Scientist Program and the Bundesministerium für Bildung und Forschung (Grant nos. 16GW0191 and NUM-COVID 19—Organo-Strat 01KX2021). This research did not receive further specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data availability Anonymized data will be made available upon reasonable request.

Declarations

Conflicts of interest Philipp Mergenthaler is on the Advisory Board of HealthNextGen Inc. and has equity interest in the company, within the scope of the present work, but not directly related; Henning R. Stetefeld reports no disclosures; Christian Dohmen reports no disclosures; Siegfried Kohler reports no disclosures; Silvia Schönenberger reports no disclosures; Julian Bösel reports personal fees from Medtronic, personal fees from Zoll, personal fees from Boehringer Ingelheim, personal fees from Sedana Medical, personal fees from PCORI, outside the submitted work. Stefan T. Gerner reports no disclosures; Hagen B. Huttner reports personal fees from Daiichi Sanyko, grants and personal fees from Novartis, grants and personal fees from UCB Pharma, grants and personal fees from Portola Pharmaceuticals, personal fees from Bayer AG, personal fees from Boehringer Ingelheim, personal fees from Prediction BioSciences, outside the submitted work; Hauke Schneider reports no disclosures; Heinz Reichmann reports no disclosures; Hannah Fuhrer reports no disclosures; Benjamin Berger received travel grants and/or training expenses from Bayer Vital GmbH, Ipsen Pharma GmbH, Novartis, Biogen GmbH, Merck Serono GmbH, and Genzyme, as well as lecture fees from Ipsen Pharma GmbH, Alexion Pharma Germany GmbH, Merck Serono GmbH, Sanofi Genzyme, and Roche Pharma AG outside the submitted work; Jan Zinke reports no disclosures; Anke Alberty reports no disclosures; Ingo Kleiter reports personal fees from Biogen, personal fees from Novartis, personal fees from Merck, personal fees from Sanofi Genzyme, personal fees from Roche, personal fees from Mylan, personal fees from Alexion,

personal fees from Celgene, grants and personal fees from Chugai, personal fees from IQVIA, outside the submitted work; Christiane Schneider-Gold has received consulting and/or speaker's honoraria from Alexion Pharmaceuticals, Amicus Therapeutics, Bayer Schering, CSL Behring, Grünenthal, Hormosan, Lupin Pharmaceuticals, and TEVA outside the submitted work; Christian Roth reports no disclosures; Juliane Dunkel reports no disclosures; Andreas Steinbrecher reports no disclosures; Andrea Thieme reports no disclosures; De-Hyung Lee reports no disclosures; Ralf Linker reports no disclosures; Klemens Angstwurm reports no disclosures; Andreas Meisel reports personal fees from Hormosan, personal fees from Grifols, grants and personal fees from Alexion, personal fees from Argnx, personal fees from Novartis, personal fees from UCB, grants from Octapharma, outside the submitted work. Bernhard Neumann reports no disclosures. The authors declare no financial or other conflicts of interest related to the manuscript. Statistical analysis was conducted by Dr. Bernhard Neumann, University Medical Center Regensburg/Donau-Isar-Klinikum Deggendorf.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. Local ethic committees and institutional review boards of the participating centers approved the study based on the central vote of the ethics committee at University of Regensburg (No. 15–101-0259). Due to the retrospective character of the study, patient's consent was not necessary according to the decisions of the ethic committees and institutional review boards.




Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Gilhus NE, Skeie GO, Romi F, Lazaridis K, Zisimopoulou P, Tzartos S (2016) Myasthenia gravis—autoantibody characteristics and their implications for therapy. *Nat Rev Neurol* 12(5):259–268. <https://doi.org/10.1038/nrneurol.2016.44>
2. Neumann B, Angstwurm K, Mergenthaler P, Kohler S, Schönenberger S, Bösel J, Neumann U, Vidal A, Huttner HB, Gerner ST, Thieme A, Steinbrecher A, Dunkel J, Roth C, Schneider H, Schimmel E, Fuhrer H, Fahrenndorf C, Alberty A, Zinke J, Meisel A, Dohmen C, Stetefeld HR, German Myasthenic Crisis Study Group (2020) Myasthenic crisis demanding mechanical ventilation: a multicenter analysis of 250 cases. *Neurology* 94(3):e299–e313. <https://doi.org/10.1212/WNL.00000000000008688>
3. Thomas CE, Mayer SA, Gungor Y, Swarup R, Webster EA, Chang I, Brannagan TH, Fink ME, Rowland LP (1997) Myasthenic crisis: clinical features, mortality, complications, and risk factors for prolonged intubation. *Neurology* 48(5):1253–1260. <https://doi.org/10.1212/wnl.48.5.1253>

4. Angstwurm K, Vidal A, Stetefeld H, Dohmen C, Mergenthaler P, Kohler S, Schönerberger S, Bösel J, Neumann U, Lee DH, Gerner ST, Huttner HB, Thieme A, Dunkel J, Roth C, Schneider H, Schimmel E, Reichmann H, Fuhrer H, Berger B, Kleiter I, Schneider-Gold C, Alberty A, Zinke J, Schalke B, Steinbrecher A, Meisel A, Neumann B (2020) Early tracheostomy is associated with shorter ventilation time and duration of ICU stay in patients with myasthenic crisis—a multicenter analysis. *J Intensive Care Med* 25:885066620967646. <https://doi.org/10.1177/0885066620967646>
5. König N, Stetefeld HR, Dohmen C, Mergenthaler P, Kohler S, Schönerberger S, Bösel J, Lee DH, Gerner ST, Huttner HB, Schneider H, Reichmann H, Fuhrer H, Berger B, Zinke J, Alberty A, Kleiter I, Schneider-Gold C, Roth C, Dunkel J, Steinbrecher A, Thieme A, Schlachetzki F, Linker RA, Angstwurm K, Meisel A, Neumann B (2021) MuSK-antibodies are associated with worse outcome in myasthenic crisis requiring mechanical ventilation. *J Neurol* 268(12):4824–4833. <https://doi.org/10.1007/s00415-021-10603-9>
6. Alsheklee A, Miles JD, Katirji B, Preston DC, Kaminski HJ (2009) Incidence and mortality rates of myasthenia gravis and myasthenic crisis in US hospitals. *Neurology* 72(18):1548–1554. <https://doi.org/10.1212/WNL.0b013e3181a41211>
7. Spillane J, Hirsch NP, Kullmann DM, Taylor C, Howard RS (2014) Myasthenia gravis—treatment of acute severe exacerbations in the intensive care unit results in a favourable long-term prognosis. *Eur J Neurol* 21(1):171–173. <https://doi.org/10.1111/ene.12115>
8. Ramos-Fransi A, Rojas-García R, Segovia S, Márquez-Infante C, Pardo J, Coll-Cantí J, Jericó I, Illa I (2015) Myasthenia gravis: descriptive analysis of life-threatening events in a recent nationwide registry. *Eur J Neurol* 22(7):1056–1061. <https://doi.org/10.1111/ene.12703>
9. Wiendl H (2015). Diagnostik und Therapie der Myasthenia gravis und des Lambert-Eaton-Syndroms. Leitlinien für Diagnostik und Therapie in der Neurologie
10. Leite MI, Jacob S, Viegas S, Cossins J, Clover L, Morgan BP, Beeson D, Willcox N, Vincent A (2008) IgG1 antibodies to acetylcholine receptors in ‘seronegative’ myasthenia gravis. *Brain* 131(Pt 7):1940–1952. <https://doi.org/10.1093/brain/awn092>
11. Hoffmann S, Harms L, Schuelke M, Rückert JC, Goebel HH, Stenzel W, Meisel A (2020) Complement deposition at the neuromuscular junction in seronegative myasthenia gravis. *Acta Neuropathol* 139(6):1119–1122. <https://doi.org/10.1007/s00401-020-02147-5>
12. Vincent A, Scadding GK, Thomas HC, Newsom-Davis J (1978) In-vitro synthesis of anti-acetylcholine-receptor antibody by thymic lymphocytes in myasthenia gravis. *Lancet* 1(8059):305–307. [https://doi.org/10.1016/s0140-6736\(78\)90073-9](https://doi.org/10.1016/s0140-6736(78)90073-9)
13. Leite MI, Ströbel P, Jones M, Micklem K, Moritz R, Gold R, Niks EH, Berrih-Aknin S, Scaravilli F, Canelhas A, Marx A, Newsom-Davis J, Willcox N, Vincent A (2005) Fewer thymic changes in MuSK antibody-positive than in MuSK antibody-negative MG. *Ann Neurol* 57(3):444–448. <https://doi.org/10.1002/ana.20386>

Authors and Affiliations

Philipp Mergenthaler^{1,2,3}  · Henning R. Stetefeld⁴ · Christian Dohmen^{4,5} · Siegfried Kohler¹ · Silvia Schönerberger⁶ · Julian Bösel^{6,7} · Stefan T. Gerner^{8,9} · Hagen B. Huttner^{8,9} · Hauke Schneider^{10,11} · Heinz Reichmann¹⁰ · Hannah Fuhrer¹² · Benjamin Berger¹² · Jan Zinke¹³ · Anke Alberty¹⁴ · Ingo Kleiter^{15,16} · Christiane Schneider-Gold¹⁵ · Christian Roth^{17,18} · Juliane Dunkel¹⁷ · Andreas Steinbrecher¹⁹ · Andrea Thieme¹⁹ · De-Hyung Lee²⁰ · Ralf A. Linker²⁰ · Klemens Angstwurm²⁰ · Andreas Meisel^{1,2,21}  · Bernhard Neumann^{20,22}  · for The German Myasthenic Crisis Study Group

Philipp Mergenthaler
philipp.mergenthaler@charite.de

Henning R. Stetefeld
henning.stetefeld@uk-koeln.de

Christian Dohmen
christian.dohmen@lvr.de

Siegfried Kohler
siegfried.kohler@charite.de

Silvia Schönerberger
Silvia.Schoenenberger@med.uni-heidelberg.de

Julian Bösel
julian.boesel@klinikum-kassel.de

Stefan T. Gerner
stefan.gerner@neuro.med.uni-giessen.de

Hagen B. Huttner
direktion@neuro.med.uni-giessen.de

Hauke Schneider
hauke.schneider@uk-augsburg.de

Heinz Reichmann
heinz.reichmann@uniklinikum-dresden.de

Hannah Fuhrer
hannah.fuhrer@uniklinik-freiburg.de

Benjamin Berger
benjamin.berger@uniklinik-freiburg.de

Jan Zinke
jan.zinke@med.uni-jena.de

Anke Alberty
anke.alberty@mariahilf.de

Ingo Kleiter
Ingo.Kleiter@ruhr-uni-bochum.de

Christiane Schneider-Gold
Christiane.Schneider-Gold@ruhr-uni-bochum.de

Christian Roth
roth@drk-nh.de

Juliane Dunkel
dunkel@drk-nh.de

Andreas Steinbrecher
andreas.steinbrecher@helios-gesundheit.de

Andrea Thieme
andrea.thieme@helios-gesundheit.de

De-Hyung Lee
dehyung.lee@medbo.de

Ralf A. Linker
ralf.linker@medbo.de

Klemens Angstwurm
klemens.angstwurm@medbo.de

Andreas Meisel
andreas.meisel@charite.de

- ¹ Charité - Universitätsmedizin Berlin, Department of Neurology with Experimental Neurology, Neuroscience Clinical Research Center, Berlin, Germany
- ² Charité - Universitätsmedizin Berlin, Center for Stroke Research Berlin, Berlin, Germany
- ³ Berlin Institute of Health at Charité-Universitätsmedizin Berlin, BIH Academy, 10117 Berlin, Germany
- ⁴ Faculty of Medicine, Department of Neurology, University of Cologne, University Hospital Cologne, Cologne, Germany
- ⁵ Department of Neurology, LVR-Klinik Bonn, Bonn, Germany
- ⁶ Department of Neurology, Heidelberg University Hospital, Heidelberg, Germany
- ⁷ Department of Neurology, Klinikum Kassel, Kassel, Germany
- ⁸ Department of Neurology, University Hospital Erlangen, Erlangen, Germany
- ⁹ Department of Neurology, Universitätsklinikum Gießen Und Marburg, Giesen, Germany
- ¹⁰ Department of Neurology, University Hospital, Technische Universität Dresden, Dresden, Germany

¹¹ Department of Neurology, University Hospital Augsburg, Augsburg, Germany

¹² Clinic of Neurology and Neurophysiology, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

¹³ Hans Berger Department of Neurology, Jena University Hospital, Jena, Germany

¹⁴ Department of Neurology, Kliniken Maria Hilf GmbH Moenchengladbach, Mönchengladbach, Germany

¹⁵ Department of Neurology, St. Josef-Hospital, Ruhr-University Bochum, Bochum, Germany

¹⁶ Marianne-Strauß-Klinik, Behandlungszentrum Kempfenhausen Für Multiple Sklerose Kranke gGmbH, Berg, Germany

¹⁷ Department of Neurology, DRK-Kliniken Nordhessen, Kassel, Germany

¹⁸ Department of Neurology, Philipps University of Marburg, Marburg, Germany

¹⁹ Department of Neurology, Helios Klinikum Erfurt, Erfurt, Germany

²⁰ Department of Neurology, University of Regensburg, Bezirksklinikum, Regensburg, Germany

²¹ Berlin Institute of Health at Charite-Universitätsmedizin Berlin, Berlin, Germany

²² Present Address: Department of Neurology, Donau-Isar-Klinikum Deggendorf, Deggendorf, Germany