

Severe enterovirus A71 associated hand, foot and mouth disease, Vietnam, 2018: preliminary report of an impending outbreak

Le Nguyen Thanh Nhan¹, Nguyen Thi Thu Hong², Le Nguyen Truc Nhu², Lam Anh Nguyet², Nguyen Thi Han Ny², Tran Tan Thanh², Do Duong Kim Han², Hoang Minh Tu Van², C Louise Thwaites^{2,3}, Tran Tinh Hien^{2,3}, Phan Tu Qui⁴, Pham Van Quang¹, Ngo Ngoc Quang Minh¹, H. Rogier van Doorn^{2,3}, Truong Huu Khanh¹, Nguyen Van Vinh Chau⁴, Guy Thwaites^{2,3}, Nguyen Thanh Hung¹, Le Van Tan²

1. Children's Hospital 1, Ho Chi Minh City, Vietnam
2. Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam
3. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom
4. Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam

Correspondence: Le Van Tan (tanlv@oucru.org)

Citation style for this article:

Nhan Le Nguyen Thanh, Hong Nguyen Thi Thu, Nhu Le Nguyen Truc, Nguyet Lam Anh, Ny Nguyen Thi Han, Thanh Tran Tan, Han Do Duong Kim, Van Hoang Minh Tu, Thwaites C Louise, Hien Tran Tinh, Qui Phan Tu, Quang Pham Van, Minh Ngo Ngoc Quang, van Doorn H. Rogier, Khanh Truong Huu, Chau Nguyen Van Vinh, Thwaites Guy, Hung Nguyen Thanh, Tan Le Van. Severe enterovirus A71 associated hand, foot and mouth disease, Vietnam, 2018: preliminary report of an impending outbreak. *Euro Surveill.* 2018;23(46):pii=1800590. <https://doi.org/10.2807/1560-7917.ES.2018.23.46.1800590>

Article submitted on 25 Oct 2018 / accepted on 15 Nov 2018 / published on 15 Nov 2018

Since January 2018, over 53,000 hospitalisations and six deaths due to hand, foot and mouth disease (HFMD) have occurred across Vietnam with most cases from September onward. In a large tertiary referral hospital, Ho Chi Minh City, enterovirus A71 subgenogroup C4 was predominant, while B5 was only sporadically detected. The re-emergence of C4 after causing a severe HFMD outbreak with >200 deaths in 2011–12 among susceptible young children raises concern of another impending severe outbreak

Since the beginning of January 2018, there has been a significant increase in the number of hospitalisations (including cases with neurological symptoms) due to hand, foot and mouth disease (HFMD) across Vietnam. By the end of September more than 53,000 clinical cases were reported, of whom six died [1]. Our aim was to characterise the epidemiology, virology and clinical characteristics of patients, especially those with severe HFMD, through an ongoing clinical study conducted at a large tertiary referral hospital for children in Ho Chi Minh City, Vietnam from January to September 2018.

Setting and outbreak investigations

Children's Hospital 1 (CH1) in Ho Chi Minh City, Vietnam is a 1,600-bed hospital, and is one of the three tertiary referral centres for children with HFMD in Ho Chi Minh City. The hospital admits children from southern Vietnam with a catchment population of over 40 million.

According to the Vietnamese Ministry of Health, HFMD is clinically divided into four major grades: Grade 1 is

assigned to patients presenting with mouth ulcers or vesicles/papules on hands, feet or buttocks, with or without mild fever ($<39^{\circ}\text{C}$); Grade 2 is further divided into Grade 2a (central nervous system (CNS) involvement, (myoclonus reported by parents or caregivers only, fever $>39^{\circ}\text{C}$ or ataxia)), Grade 2b1 (myoclonus observed by medical staff or history of myoclonus and lethargy or pulse higher than 130 bpm), and Grade 2b2 (ataxia, cranial nerve palsies, limb weakness, nystagmus, persistent high fever or pulse higher than 150 bpm); Grade 3 involves autonomic dysfunction with sweating, hypertension, tachycardia and tachypnoea and Grade 4 is for disease with additional cardio-pulmonary compromise with pulmonary oedema or shock syndrome [2].


The data presented here were either derived from the hospital database record or obtained from cases enrolled into an ongoing clinical study conducted at CH1 [3]. Since January 2018, our recruitment focus has been inpatients, especially those with severe HFMD, with a targeted ratio of 1:1 for Grade 2a and 2b1 vs 2b2 or above.

Recruited patients had throat and rectal swabs collected on admission alongside clinical data during hospitalisation. The study was approved by the Institutional Review Board of CH1 and the Oxford Tropical Research Ethics Committee (OxTREC).

Multiplex PCR for simultaneous detection of enteroviruses (EVs) and enterovirus A71 (EV-A71) was performed on rectal and throat swabs [4]. PCR positive swabs

FIGURE 1


Monthly distribution of HFMD cases and HFMD cases with severe disease admitted to Children's Hospital 1, Vietnam, January–September 2018 (n = 2,283)



HFMD: hand, foot and mouth disease.

FIGURE 2

Monthly distribution of clinical HFMD cases, including those who were PCR positive for EV-A71 or non-EV-A71 EVs, and clinical cases with severe HFMD enrolled in the study, Vietnam, January–September 2018 (n = 86 patients)



EV: enterovirus; HFMD: hand, foot and mouth disease.

For cases testing positive for EV-71, stratification of the numbers by subgenogroup B5 or C4 is shown.

TABLE

Demographics, clinical characteristics and aetiology of HFMD patients enrolled in the clinical study, Vietnam, January–September 2018 (n = 86 patients)

Characteristics	Total (n = 86)		EV-A71 (n = 59)		Non-EV-A71 EVs (n = 15)		PCR negative (n = 12)	
	n	%	n	%	n	%	n	%
Median age in months (range)	21.7 (2.4–60.3)		24.3 (5.4–60.3)		17.6 (2.4–34.6)		15.6 (6.2–33.7)	
Sex ratio, male/female	54/32		36/23		12/3		6/6	
Median illness day on admission (range)	3 (1–8)		3 (2–8)		3 (2–6)		3 (1–7)	
Origin	n	%	n	%	n	%	n	%
Ho Chi Minh City	46	54	33	56	7	47	6	50
Other provinces	40	47	26	44	8	53	6	50
Clinical grade^a	n	%	n	%	n	%	n	%
2a	31	36	14	24	10	67	7	58
2b1	11	13	8	14	1	7	2	17
2b2	14	16	12	20	2	13	0	0
3	26	30	21	36	2	13	3	25
4	4	5	4	7	0	0	0	0

EV: enterovirus; HFMD: hand, foot and mouth disease.

^a According to the Vietnamese Ministry of Health, HFMD is clinically divided into four major Grades, i.e. 1, 2, 3 and 4 according to increasing disease severity. Grade 2 is further divided in Grade 2a, 2b1, 2b2 [2]. The four Grades are described in more detail in the main text of this report.

were then subjected to viral protein 1 (VP1) nucleotide sequencing using previously described PCR primers [2,5,6]. The obtained PCR amplicons were sequenced in both directions using corresponding PCR primers and BigDye Terminator V3.1 Cycle Sequencing kit in an ABI 3130XL DNA sequencer (Applied Biosystems, Carlsbad, CA, United States (US)). EV-A71 subgenogroup determination and EV serotyping were then carried out using the obtained VP1 sequences and an online EV typing tool [7].

Description of the epidemic and characteristics of the patients

CH1 recorded 2,283 hospitalisations due to HFMD since January 2018, of whom, 113 (5%) had severe HFMD (Grade 2b1, 2b2, 3 or 4) (Figure 1). A total of 1,376 (60%) hospitalisations concerned boys, and 2,250 (99%) were in children ≤5 years old. The number of admissions (n=1,411) and cases with severe disease (n=91) in August and September combined was 1.6 and 4.1 times, respectively, that of the first 7 months of the year combined (Figure 1).

Of 86 clinically diagnosed HFMD patients enrolled in the clinical study, 55 (64%) had severe HFMD, 54 (63%) were boys, and 85 (99%) were ≤5 years old. As for CH1, the number of cases with neurological symptoms and/or autonomic dysfunction enrolled in our study sharply increased compared with previous months, in August (n=11) and September (n=28), accounting for 39/55 (71%) of all cases enrolled with severe HFMD in 2018 (Figure 2). One patient died within 6 hours of admission.

All patients in the study were admitted to the hospital within the first 8 days of illness (median: 3 days; range: 1–8). EV-A71 and non-EV-A71 EVs were detected in 86% (74/86) of the enrolled patients, with EV-A71 being the most frequently detected (80%, 59/74). EV-A71 was detected in 82% (45/55) of patients presenting with severe disease, including 8/11, 12/14, 21/26 and 4/4 of those presenting with Grade 2b1, 2b2, 3 and 4 disease, respectively. The remaining 10 patients with severe HFMD were either positive for other EVs (n=5) or PCR negative (Table). While our overall PCR positivity rate was in agreement with previous reports [8], the failure to detect enterovirus in 14% (12/86) of the patients may have been attributed to the low level of viral load in the tested samples and/or the miss-inclusion of patients with other diseases presenting with rash (e.g. chickenpox, measles and rubella) into the study.

Forty partial VP1 sequences were successfully obtained. Sequence analysis revealed that the majority (n=36, 90%) belonged to EV-A71 subgenogroup C4, while subgenogroup B5 was sporadically detected (n=4, 10%), (Figure 2). Both rectal and throat swabs of the fatal case were positive for EV-A71 subgenogroup C4.

Of 15 non-EV-A71 enteroviruses, coxsackievirus A6 (CV-A6), CV-A8 and CV-A10 were detected in two, one and eight patients, respectively.

Discussion and conclusion

HFMD is an emerging infection mostly affecting children ≤5 years old [2]. The disease can be caused by various serotypes of the EV A species, of which EV-A71, CV-A6, CV-A10 and CV-A16 are the most frequently

detected serotypes [2,3,9]. In contrast to HFMD associated with other serotypes, the burden of EV-A71 is attributed to both large numbers of hospitalised cases and a relatively high proportion of cases with severe disease and case fatality. While since 2008 (EV-A71-associated) HFMD causes over 1 million reported cases annually across the Asia-Pacific region [10], outbreaks of EV-A71 associated with neurological complications have been documented worldwide, including in Europe and the US [11-13].

An inactivated EV-A71 C₄ vaccine has been approved in China in December 2015 but has not been implemented anywhere outside of China [14,15]. Currently, there is no clinically proven effective antiviral drug available, while intravenous immunoglobulin (IVIg) is widely used to treat patients with severe illness [16]. Thus, with poliomyelitis near eradication [17], the emergence of EV-A71 is considered a serious global threat posed by enterovirus infections. Indeed, the World Health Organization has recently included EV-A71 infection for consideration of its Blueprint List of Priority Diseases [18].

EV-A71 exists as a single serotype, but is genetically divided into several genogroups (including A, B and C) and subgenogroups (e.g. B₁ to B₅ and C₁ to C₅). While there is no genetic evidence of the difference in virulence between different (sub)genogroups, subgenogroup replacement often accompanies large outbreaks of severe HFMD [19]. Of note, a switch from C₅ to C₄ in 2011 coincided with an explosive outbreak in Vietnam, which resulted in >200,000 hospitalisations and >200 fatal cases between 2011 and 2012 [2]. In contrast, the replacement of C₄ by B₅ in 2013 coincided with a decrease in the numbers of reported cases and the proportion of cases with severe illness (including fatality) in subsequent years [20,21].

While the underlying mechanism for EV-A71 subgenogroup switches in endemic countries remains unsolved, the observed 2 to 3 year cycles of EV-A71-associated HFMD are attributed to accumulation of a sufficient number of susceptible young children [9].

In Vietnam, HFMD and EV-A71 annual activity increases during the wet season (June to November) and the current upswing of cases is likely to continue over the next few months. Collectively, these data suggest that the re-emergence of subgenogroup C₄ after an absence of 6 years causing severe HFMD disease in Vietnam is likely driven by the accumulation of a sufficient number of susceptible young children, and is expected to cause a further increase in monthly hospital admissions. The observed increase of cases, including severe cases, the unprotected birth cohort of children born after 2012 and the known association of EV-A71 C₄ with explosive outbreaks and severe illness are therefore a cause of concern of another impending severe outbreak.

Given the absence of vaccines outside China, the absence of specific antiviral treatment and the potential of EV-A71-associated HFMD to spread to other parts of the world including Europe, improving public awareness and public health measures (e.g. hand washing, cleaning/disinfecting surfaces (including toys) and avoiding close contact (hugging and kissing)), as well as accurate case classification and early identification of cases with high risk of deterioration remain essential to minimise the impact of the ongoing outbreak. Clinical trials to demonstrate the effectiveness of current treatment approaches (including IVIg) are of clinical importance and are urgently needed [16].

Acknowledgements

We would like to thank Ms Le Kim Thanh for her logistic support. We are indebted to the patients and their parents for their participation in this study, and all the nursing and medical staff at the paediatric ICU and infectious diseases wards at Children's Hospital 1 who provided care for the patients and helped collect clinical data.

This work was supported by the Wellcome Trust, United Kingdom (106680/B/14/Z, and 204904/Z/16/Z). The funding body did not have any influence on the study design, study conduct, preparation of the manuscript or decision to publish.

Conflict of interest

None declared.

Authors' contributions

Designed the study: Le Nguyen Thanh Nhan, Hoang Minh Tu Van, Tran Tinh Hien, C Louise Thwaites, H. Rogier van Doorn, Guy Thwaites, Nguyen Van Vinh Chau, Nguyen Thanh Hung and Le Van Tan;

Recruited the patients: Do Duong Kim Han, Le Nguyen Thanh Nhan, Phan Tu Qui, Pham Van Quang, Ngo Ngoc Quang Minh, Truong Huu Khanh, Nguyen Thanh Hung;

Performed laboratory experiments: Nguyen Thi Thu Hong, Le Nguyen Truc Nhu, Lam Anh Nguyet, Nguyen Thi Han Ny, Tran Tan Thanh;

Analyzed the data: Nguyen Thi Thu Hong, Le Nguyen Truc Nhu, Lam Anh Nguyet, Nguyen Thi Han Ny, Tran Tan Thanh, Le Nguyen Thanh Nhan and Le Van Tan;

Drafted the manuscript: Le Nguyen Thanh Nhan, C Louise Thwaites, Guy Thwaites, H. Rogier van Doorn, and Le Van Tan;

All authors have read the final manuscript and agreed with its contents.

References

1. Ministry of Health (Vietnam). General Department of Preventive Medicine. Preventing hand, foot and mouth disease. Hanoi: Ministry of Health; Oct 2018. Vietnamese. Available from: <http://vncdc.gov.vn/vi/tin-tuc-trong-nuoc/2361/chu-dong-phong-benh-tay-chan-mieng>

2. Khanh TH, Sabanathan S, Thanh TT, Thoa PK, Thuong TC, Hang V, et al. Enterovirus 71-associated hand, foot, and mouth disease, Southern Vietnam, 2011. *Emerg Infect Dis.* 2012;18(12):2002-5. <https://doi.org/10.3201/eid1812.120929> PMID: 23194699
3. Anh NT, Nhu LNT, Van HMT, Hong NTT, Thanh TT, Hang VTT, et al. Emerging Coxsackievirus A6 Causing Hand, Foot and Mouth Disease, Vietnam. *Emerg Infect Dis.* 2018;24(4):654-62. <https://doi.org/10.3201/eid2404.171298> PMID: 29553326
4. Thanh TT, Anh NT, Tham NT, Van HM, Sabanathan S, Qui PT, et al. Validation and utilization of an internally controlled multiplex Real-time RT-PCR assay for simultaneous detection of enteroviruses and enterovirus A71 associated with hand foot and mouth disease. *Virology.* 2015;12(1):85. <https://doi.org/10.1186/s12985-015-0316-2> PMID: 26050791
5. Brown BA, Oberste MS, Alexander JP Jr, Kennett ML, Pallansch MA. Molecular epidemiology and evolution of enterovirus 71 strains isolated from 1970 to 1998. *J Virol.* 1999;73(12):9969-75. PMID: 10559310
6. Tan V, Tuyen NT, Thanh TT, Ngan TT, Van HM, Sabanathan S, et al. A generic assay for whole-genome amplification and deep sequencing of enterovirus A71. *J Virol Methods.* 2015;215-216:30-6. <https://doi.org/10.1016/j.jviromet.2015.02.011> PMID: 25704598
7. Kroneman A, Vennema H, Deforche K, v d Avoort H, Peñaranda S, Oberste MS, et al. An automated genotyping tool for enteroviruses and noroviruses. *J Clin Virol.* 2011;51(2):121-5. <https://doi.org/10.1016/j.jcv.2011.03.006> PMID: 21514213
8. Gao L, Zou G, Liao Q, Zhou Y, Liu F, Dai B, et al. Spectrum of Enterovirus Serotypes Causing Uncomplicated Hand, Foot, and Mouth Disease and Enteroviral Diagnostic Yield of Different Clinical Samples. *Clin Infect Dis.* 2018;67(11):1729-35. <https://doi.org/10.1093/cid/ciy341> PMID: 29688329
9. Takahashi S, Metcalf CJE, Arima Y, Fujimoto T, Shimizu H, Rogier van Doorn H, et al. Epidemic dynamics, interactions and predictability of enteroviruses associated with hand, foot and mouth disease in Japan. *J R Soc Interface.* 2018;15(146):20180507. <https://doi.org/10.1098/rsif.2018.0507> PMID: 30209044
10. Xing W, Liao Q, Viboud C, Zhang J, Sun J, Wu JT, et al. Hand, foot, and mouth disease in China, 2008-12: an epidemiological study. *Lancet Infect Dis.* 2014;14(4):308-18. [https://doi.org/10.1016/S1473-3099\(13\)70342-6](https://doi.org/10.1016/S1473-3099(13)70342-6) PMID: 24485991
11. Hassel C, Mirand A, Lukashev A, TerletskaiaLadwig E, Farkas A, Schuffenecker I, et al. Transmission patterns of human enterovirus 71 to, from and among European countries, 2003 to 2013. *Euro Surveill.* 2015;20(34):30005. <https://doi.org/10.2807/1560-7917.ES.2015.20.34.30005> PMID: 26530407
12. Midgley SE, Nielsen AG, Trebbien R, Poulsen MW, Andersen PH, Fischer TK. Co-circulation of multiple subtypes of enterovirus A71 (EV- A71) genotype C, including novel recombinants characterised by use of whole genome sequencing (WGS), Denmark 2016. *Euro Surveill.* 2017;22(26):30565. <https://doi.org/10.2807/1560-7917.ES.2017.22.26.30565> PMID: 28681718
13. Antona D, Kossorotoff M, Schuffenecker I, Mirand A, Leruez-Ville M, Bassi C, et al. Severe paediatric conditions linked with EV-A71 and EV-D68, France, May to October 2016. *Euro Surveill.* 2016;21(46):30402. <https://doi.org/10.2807/1560-7917.ES.2016.21.46.30402> PMID: 27918268
14. World Health Organization (WHO). China produces world's first vaccine against virus that causes Hand, Foot and Mouth Disease. Geneva: WHO; Dec 2015. Available from: <http://www.wpro.who.int/china/mediacentre/releases/2015/20151208/en/>
15. Mao QY, Wang Y, Bian L, Xu M, Liang Z. EV71 vaccine, a new tool to control outbreaks of hand, foot and mouth disease (HFMD). *Expert Rev Vaccines.* 2016;15(5):599-606. <https://doi.org/10.1586/14760584.2016.1138862> PMID: 26732723
16. Chea S, Cheng YB, Chokephaibulkit K, Chotpitayasunondh T, Rogier van Doorn H, Hafy Z, et al. Workshop on use of intravenous immunoglobulin in hand, foot and mouth disease in Southeast Asia. *Emerg Infect Dis.* 2015;21(1). <https://doi.org/10.3201/eid2101.140992> PMID: 25531166
17. World Health Organization (WHO). Poliomyelitis. Geneva: WHO; Mar 2018. Available from: <http://www.who.int/news-room/fact-sheets/detail/poliomyelitis>
18. World Health Organization (WHO). List of Blueprint Priority Diseases. Geneva: WHO. [Accessed 8 Nov 2018]. Available from: <https://www.who.int/blueprint/priority-diseases/en/>
19. Solomon T, Lewthwaite P, Perera D, Cardosa MJ, McMinn P, Ooi MH. Virology, epidemiology, pathogenesis, and control of enterovirus 71. *Lancet Infect Dis.* 2010;10(11):778-90. [https://doi.org/10.1016/S1473-3099\(10\)70194-8](https://doi.org/10.1016/S1473-3099(10)70194-8) PMID: 20961813
20. Geoghegan JL, Tan V, Kühnert D, Halpin RA, Lin X, Simenauer A, et al. Phylogenetics of enterovirus A71-associated hand, foot, and mouth disease in Viet Nam. *J Virol.* 2015;89(17):8871-9. <https://doi.org/10.1128/JVI.00706-15> PMID: 26085170
21. World Health Organization (WHO). Hand, Foot and Mouth Disease Situation Update (Archives 2015, 2016 and 2017). Geneva: WHO. Available from: <http://www.who.int/westernpacific/emergencies/surveillance/archives/hand-foot-and-mouth-disease>

License and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence, and indicate if changes were made.

This article is copyright of the authors or their affiliated institutions, 2018.