

SCIENTIFIC REPORTS



OPEN

Analysis of gene expression of secreted factors associated with breast cancer metastases in breast cancer subtypes

Received: 12 January 2015

Accepted: 02 June 2015

Published: 15 July 2015

Elana J. Fertig^{1,*}, Esak Lee^{2,*}, Niranjan B. Pandey² & Aleksander S. Popel^{1,2}

Breast cancer is a heterogeneous disease, having multiple subtypes with different malignant phenotypes. The triple-negative breast cancer, or basal breast cancer, is highly aggressive, metastatic, and difficult to treat. Previously, we identified that key molecules (*IL6*, *CSF2*, *CCL5*, *VEGFA*, and *VEGFC*) secreted by tumor cells and stromal cells in basal breast cancer can promote metastasis. It remains to assess whether these molecules function similarly in other subtypes of breast cancer. Here, we characterize the relative gene expression of the five secreted molecules and their associated receptors (*GP130*, *GMRA*, *GMRB*, *CCR5*, *VEGFR2*, *NRP1*, *VEGFR3*, *NRP2*) in the basal, HER2 (human epidermal growth factor receptor 2) positive, luminal A, and luminal B subtypes using high throughput data from tumor samples in The Cancer Genome Atlas (TCGA) and Molecular Taxonomy of Breast Cancer International Consortium (METABRIC). *IL6* and *CCL5* gene expression are basal breast cancer specific, whereas high gene expression of *GP130* was observed in luminal A/B. *VEGFA/C* and *CSF2* mRNA are overexpressed in HER2 positive breast cancer, with *VEGFA* and *CSF2* also overexpressed in basal breast cancer. Further study of the specific protein function of these factors within their associated cancer subtypes may yield personalized biomarkers and treatment modalities.

Breast cancer is the most frequently diagnosed cancer among women in the United States¹. Primary breast tumors are divided in four main molecular subtypes: Basal (also known as, triple negative), HER2 (human epidermal growth factor receptor 2) positive, Luminal A, and Luminal B. Each of these subtypes has characteristic traits and expected patient outcome. For example, basal breast cancer is the most aggressive and metastatic subtype. Basal breast tumors do not express typical breast cancer cell receptors, such as the estrogen receptor (ER), the progesterone receptor (PR), and does not overexpress the human epidermal growth factor receptor 2 (HER2) that are activated in the other subtypes². Thus, current hormonal therapies and HER2 inhibition cannot be used to treat basal breast cancer. Moreover, therapeutic resistance is common when treating tumors from other subtypes with hormonal therapies³. Therefore, new therapeutics that target additional molecular factors in breast tumors are needed. Optimal therapeutics would target the factors that promote tumor growth and metastasis resulting from interactions between cancer cells, stromal cells, and extracellular matrix.

Secreted factors from each of the diverse cells in a tumor regulate inter-cellular signaling between tumor cells and the microenvironment to promote breast cancer growth and metastasis⁴. Specifically, the secreted factors cytokines, chemokines, and growth factors contribute to distinct modes of metastasis and subsequent mortality⁵. Cytokines represent soluble proteins secreted by mammalian cells that are important in cell signaling. Among them, cytokines related to inflammatory signals are involved

¹Department of Oncology, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD, USA. ²Department of Biomedical Engineering, Johns Hopkins University School of Medicine, Baltimore, MD, USA. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to A.S.P. (email: apopel@jhu.edu)




Figure 1. Summary of relationship of ligand and receptor pairs previously associated with metastasis (Lee *et al.* 2014) in tumor cells and adjacent BEC and LEC.

in many human diseases. For example, in cancer, inflammation induces tumor growth, tumor drug resistance, and metastasis⁶. Chemokines have been studied in immunology and are known to serve as immune cell-recruiting and cell-trafficking factors⁷. Though chemokines are a subset of cytokines, they are categorized as distinct secreted factors for this study, as they can play a role in tumor cell motility and recruitment, which are critical for metastatic dissemination. Growth factors are essential secreted factors for cancer cell proliferation, maintenance, migration, and adhesion⁸. Angiogenic and lymphangiogenic growth factors can regulate angiogenesis and lymphangiogenesis in both primary tumors and pre-metastatic niches^{9,10}.

Because of their critical role activating the signaling processes responsible for tumor maintenance and progression, tumor secreted factors (“tumor secretome”) can serve as targets to inhibit the primary tumor growth¹¹. The specific secreted factor in an individual tumor would provide the most promising therapeutic target in that individual’s disease. Therapeutics that thus target the secretome have the greatest potential for translation to the clinic when the factor they target are common to multiple tumors from each molecular subtype.

We previously reported that the secreted factors IL6, CSF2, CCL5, VEGFA, and VEGFC are pivotal orchestrators of basal breast cancer growth and metastasis^{4,12}. Specifically, this previous study reported that basal breast cancer cells secrete interleukin 6 (IL6), a cytokine, which conditions (educates, reprograms) lymphatic endothelial cells (LEC) within pre-metastatic organs and primary tumors to secrete the chemokine CC-chemokine ligand 5 (CCL5) and the growth factor, vascular endothelial growth factor A (VEGFA)⁴. This group of secreted factors including a cytokine, a chemokine, and a growth factor make a self-reinforcing paracrine loop to promote basal breast cancer metastasis. LEC-derived CCL5 recruits CCR5-positive cancer cells into the lymphatic vessels and triggers tumor dissemination. LEC-derived VEGFA interacts with blood endothelial cells (BEC) of the vasculature enhancing vascular permeability in the lungs, and promoting angiogenesis in the lymph nodes. These are important steps for tumor cell extravasation and colonization. VEGFC is a lymphangiogenic growth factor, secreted by cancer cells and stromal cells to promote lymphatic vessel growth in primary tumors¹³. We also demonstrated a crosstalk between basal breast cancer cells and BEC/LEC that was important for primary tumor growth¹². We showed that the secretomes of the BEC and LEC are perturbed in distinct ways, influencing primary tumor growth, pericyte infiltration, and angiogenesis in different ways¹².

In addition to having a pivotal role in basal breast cancer growth and metastasis, the secreted factors implicated in our previous study (IL6, CSF2, CCL5, VEGFA, and VEGFC) may also serve critical roles in other subtypes of breast cancers. In this case, inhibitors of these factors and the pathways they regulate could be used to treat a wider array of breast cancers. High throughput genomic data can indicate the molecular profile of these factors to infer such candidate targets from the secretome. Therefore, in this study we identify the relative gene expression of IL6, CSF2, CCL5, VEGFA, and VEGFC and their receptors (GP130, GMRA, GMRB, CCR5, VEGFR2, NRP1, VEGFR3, and NRP2, Fig. 1), in multiple breast cancer subtypes (Basal, HER2+, Luminal A, and Luminal B) using high-throughput genomic data of primary tumors from the Cancer Genome Atlas (TCGA)¹⁴ and Molecular Taxonomy of Breast Cancer International Consortium METABRIC¹⁵. The goals of this study are: (a) to understand how these key pro-metastatic factor genes are expressed in breast cancer subtypes, (b) to examine how their associated receptor genes are expressed in the subtypes, and (c) to evaluate whether these gene expression profiles can predict the survival rates within each breast cancer subtype.

Pro-metastatic factor	TCGA	METABRIC
IL6	IL6	ILMN_1699651
GP130	IL6ST	ILMN_1849013
GMCSF	CSF2	ILMN_1661861
GMRA	CSF2RA	ILMN_2376455
GRMB	CSF2RB	ILMN_1798475
CCL5	CCL5	ILMN_2098126
CCR5	CCR5	
VEGFA	VEGFA	
VEGFR2	KDR	ILMN_1686405
NRP1	NRP1	ILMN_1742547
VEGFC	VEGFC	ILMN_1701204
VEGFR3	FLT4	ILMN_2390427
NRP2	NRP2	ILMN_1787190

Table 1. Pro-metastatic factors analyzed in this study. List of pro-metastatic factors considered in this study (Fig. 1). The TCGA column indicates gene aliases used to match the gene annotations in TCGA. Similarly, the METABRIC column indicates the single array probe used to obtain data for each gene in METABRIC, as described in the methods. A blank entry indicates that no measurements were available for that gene.

Ligand	Receptor	TCGA		METABRIC	
		R	p-value	R	p-value
IL6	GP130 (IL6ST)	-0.18	3×10^{-6}	-0.20	3×10^{-9}
CSF2 (GMCSF)	GMRA (CSF2RA)	0.34	$<2 \times 10^{-16}$	0.23	8×10^{-12}
	GMRB (CSF2RB)	0.42	$<2 \times 10^{-16}$	0.12	4×10^{-4}
CCL5	CCR5	0.86	$<2 \times 10^{-16}$		
VEGFA	VEGFR2 (KDR)	0.11	6×10^{-3}		
	NRP1	0.06	0.13		
VEGFC	VEGFR3 (FLT4)	0.49	$<2 \times 10^{-16}$	0.05	0.1
	NRP2	0.42	$<2 \times 10^{-16}$	0.16	3×10^{-6}

Table 2. Correlation between gene expression of pro-metastatic ligand and receptor pairs. Spearman correlation coefficient (R) and corresponding p-value for mRNA expression of a ligand with its associated receptor(s) (Fig. 1) from RNA sequencing data from primary breast tumors in TCGA and microarray gene expression data from primary tumors in METABRIC. Gene names are consistent with Fig. 1, with aliases used in TCGA or METABRIC in parentheses.

Results

Selection of secreted factors associated with metastasis. Figure 1 shows the ligand and receptor interactions between the factors we previously associated with metastasis⁴ and have selected for analysis in this current study (listed in Table 1). Specifically, this figure summarizes the relationship between these factors and adjacent blood endothelial cells (BEC) and lymphatic endothelial cells (LEC) from our previous study. Primary tumor samples used for genomic profiling contain a mixture of tumor cells and cells from the microenvironment, including adjacent BEC and LEC. We test whether the putative regulatory relationships between ligands and their associated receptor(s) are represented in mRNA expression of such primary tumors by correlating mRNA expression of each ligand with its associated receptor(s). Correlation analyses are run on large cohorts of gene expression data from 638 primary tumors in TCGA (Table 2) and from 897 primary tumors in METABRIC (Table 2), with sample characteristics in Supplemental Tables 1 and 2, respectively.

In TCGA, *CSF2*, *CCL5*, and *VEGFC* are all significantly correlated to their target receptors (*GMRA* and *GMRB*; *CCR5*; and *VEGFR3* and *NRP2*, respectively, Table 2). *VEGFA* is significantly correlated to only one of its receptors (*VEGFR2*), but not to the other target receptor *NRP1*. *IL6* is significantly anti-correlated with its target receptor (*GP130*). The significant anti-correlation between *IL6* and *GP130* and correlation between *VEGFC* and *NRP2* and *CSF2* and targets *GMRA* and *GMRB* are all confirmed




Figure 2. Gene expression of *IL6* in (a) TCGA by tumor subtype and in (b) METABRIC by Pam50 gene expression subtype. (c) and (d) provide corresponding boxplots for expression of the *IL6* ligand target receptor *GP130*.

in gene expression data from METABRIC (Table 2). However, the correlation between *VEGFC* and target receptor *VEGFR3* fails to meet statistical significance. We are also unable to confirm associations with *CCL5* with *CCR5* or *VEGFA* with *VEGFR2* because the array used in METABRIC does not contain probes that measure gene expression of *CCR5* or *VEGFA*.

IL6 is overexpressed in basal breast cancer while its receptor GP130 is overexpressed in luminal breast cancer. We compared expression of the ligand *IL6* and its target receptor *GP130* in each of the breast cancer subtypes. *IL6* is significantly overexpressed in the basal subtype relative to other subtypes in both TCGA (p-value of 5×10^{-8}) and METABRIC (p-value of 1×10^{-9}) data (Fig. 2a,b, respectively). On the other hand, the target receptor *GP130* is significantly overexpressed in both luminal subtypes (Fig. 2c for TCGA and 2d for METABRIC with corresponding one-sided p-values below 2×10^{-16} in both datasets). Survival analyses were run for these genes using only METABRIC, due to the relatively long patient follow-up times in that dataset. We observed a trend towards longer survival times based upon *GP130* (Supplemental Fig. 3) expression in the luminal A subtype (p-value of 0.06) not observed in the other subtypes; no significant trend was observed for *IL6* (Supplemental Fig. 4).

Overexpression of CSF2 target receptors GMRA and GMRB are associated with survival in basal and HER2+ breast cancer. In TCGA, *CSF2* is significantly overexpressed in basal and HER2+ breast cancer relative to luminal subtypes (Fig. 3a, p-values of 1×10^{-8} and 0.03, respectively). This discrepancy in p-values is consistent with a higher log fold change in basal relative to luminal breast cancer (0.7) than HER2+ relative to luminal breast cancer (0.3). A similar trend is confirmed in METABRIC (Fig. 3b), with a p-value of 1×10^{-8} with log fold change of 0.1 in basal breast cancer relative to luminal and p-value 0.006 with log fold change of 0.05 in HER2+ relative to luminal. However, in




Figure 3. Gene expression of *CSF2* in (a) TCGA by tumor subtype and in (b) METABRIC by Pam50 gene expression subtype. (c) and (d) provide corresponding boxplots for expression of the *CSF2* ligand target receptor *GMRA* and (e) and (f) for *GMRB* in TCGA and METABRIC, respectively.

both datasets target receptor *GMRA* was only overexpressed in basal breast cancer relative to all other subtypes (Fig. 3c,d with p-values of 2×10^{-4} for TCGA and 7×10^{-10} for METABRIC). Similar association with basal breast cancer was observed for the other target receptor, *GMRB* (Fig. 3e,f and p-values of 1×10^{-4} for TCGA and 9×10^{-4} for METABRIC).

CSF2 expression was not significantly associated with survival in any subtypes (Supplemental Fig. 5). Nonetheless, higher *GMRA* expression significantly associated with better survival in basal breast cancer (Supplemental Fig. 6, p-value of 0.02) and *GMRB* expression trended towards higher survival in HER2+ breast cancer (Supplemental Figure 7, p-value of 0.08).

***CCL5* overexpression is associated with basal breast cancer and with survival in HER2+ breast cancer.** Similar to *CSF2*, chemokine *CCL5* is significantly overexpressed in basal over luminal breast cancer in both TCGA (Fig. 4a, p-value of 9×10^{-8}) and METABRIC (Fig. 4b, p-value below 2×10^{-16}). *CCL5* is also significantly overexpressed in HER2+ breast cancer relative to luminal in METABRIC (p-value of 9×10^{-9}) with a similar trend that failed to reach statistical significance in TCGA (p-value of 0.09). Likewise, its target receptor *CCR5* is significantly overexpressed only in basal breast cancer in TCGA (Fig. 4c, p-value of 0.02). We are unable to confirm these relationships in METABRIC because there was no associated probe for this gene on the array measuring expression in this study. Nonetheless, increased *CCL5* expression was associated with better survival in HER2+ breast cancer (Supplemental Figure 8, p-value of 0.01).




Figure 4. Gene expression of *CCL5* in (a) TCGA by tumor subtype and in (b) METABRIC by Pam50 gene expression subtype. Corresponding boxplot of expression of *CCL5* target receptor, *CCR5*, in TCGA is in (c).




Figure 5. Gene expression of *VEGFA* in (a) TCGA by tumor subtype and in (b) METABRIC by Pam50 gene expression subtype. (c) and (d) provide corresponding boxplots for expression of the *VEGFA* ligand target receptor *VEGFR2* and (e) and (f) for *NRP1* in TCGA and METABRIC, respectively.




Figure 6. Gene expression of *VEGFC* in (a) TCGA by tumor subtype and in (b) METABRIC by Pam50 gene expression subtype. (c) and (d) provide corresponding boxplots for expression of the *VEGFC* ligand target receptor *VEGFR3* and (e) and (f) for *NRP2* in TCGA and METABRIC, respectively.

VEGFA is associated with HER2+ breast cancer and its target VEGFR2 with survival in basal breast cancer. In TCGA data, *VEGFA* expression is highest in HER2+ breast cancer (Fig. 5a, p-value of 5×10^{-3} relative to other subtypes). It is also overexpressed in basal breast cancer relative to luminal breast cancer (p-value of 5×10^{-9}). These relationships could not be confirmed in METABRIC because no probe measures *VEGFA* gene expression.

Although expression of the target receptor *VEGFR2* was not associated with any subtype in TCGA data (Fig. 5b), it was significantly overexpressed in HER2+ breast cancer in METABRIC (Fig. 5c, p-value of 4×10^{-3}). The other target receptor, *NRP1*, was not differentially expressed in any breast cancer subtypes in TCGA (Fig. 5d), but was overexpressed in basal breast cancer relative to other subtypes in METABRIC (Fig. 5e, p-value of 2×10^{-4}). Moreover, increased *VEGFR2* expression was significantly associated with better survival in basal breast cancer (Supplemental Figure 9, p-value of 0.008). No significant associations with survival were observed in any subtype for *NRP1* (Supplemental Figure 10).

VEGFC is significantly overexpressed in HER2+ breast cancer. *VEGFC* is significantly overexpressed in HER2+ breast cancer relative to basal breast cancer in TCGA (Fig. 6a, p-value of 0.005) and in HER2+ breast cancer relative to all other subtypes in METABRIC data (Fig. 6b, p-value of 4×10^{-6}). It is also overexpressed in both luminal subtypes relative to basal breast cancer in TCGA (p-value of 0.003), not confirmed in the METABRIC data. We observe significant overexpression of target receptor *VEGFR3* in the luminal A subtype in TCGA (Fig. 6c, p-value of 2×10^{-3}), while we observe significant overexpression of *VEGFR3* in basal breast cancer in METABRIC (Fig. 6d, p-value of 9×10^{-3}). *NRP2*, which is not

significantly differentially expressed in any subtype in TCGA (Fig. 6e) but is significantly overexpressed in basal breast cancer in METABRIC (Fig. 6f, p-value of 9×10^{-8}). Moreover, we do not observe any significant survival differences based upon expression of genes in this pathway (Supplemental Figures 11–13 for *VEGFC*, *VEGFR3*, and *NRP2*, respectively).

Discussion

In this study, we characterized gene expression of multiple secreted factors and their receptors, in all the breast cancer subtypes (Basal, Her2, Luminal A, and Luminal B) by using large genomic studies (TCGA and METABRIC). The factors and receptors analyzed were *IL6*, *CSF2*, *CCL5*, *VEGFA*, *VEGFC*, *GP130*, *GMRA*, *GMRB*, *CCR5*, *VEGFR2*, *NRP1*, *VEGFR3*, and *NRP2*, because we found their protein expression to be critical to basal breast cancer metastasis in previous studies^{4,12}. In this study, we found that *IL6* and *CCL5* gene expression are basal breast cancer specific. High gene expression of *GP130* is observed in luminal A/B. *VEGFA/C* and *CSF2* mRNA are overexpressed both basal and HER2+ breast cancer relative to luminal subtypes.

IL6 mRNA expression is higher in basal breast cancer when compared to other subtypes of breast cancer. Basal breast cancer is considered as aggressive and metastatic, and effective therapeutic treatments are very limited. IL6 protein has been studied in breast cancer¹⁶; it promotes formation of cancer stem cells¹⁷. Mesenchymal stem cell derived IL6 protein promotes breast cancer cell migration and invasion¹⁸. IL6 protein is also involved in drug resistance in breast cancer¹⁹. Studies of protein expression in basal breast tumors during lymph node metastasis in mouse models have shown that IL6 was highly expressed in lymph node positive basal breast tumors, compared to lymph node negative basal breast²⁰. IL6 protein can activate a STAT3 (signal transducer and activator of transcription 3) pathway. Binding of IL6 protein to the GP130 receptor triggers STAT3 phosphorylation by JAK2²¹. Recent bioinformatics study showed that STAT3-associated genes can be a prognostic marker in basal breast cancer²². Although we did not observe an association between *IL6* gene expression and survival, the association of *IL6* gene expression with basal breast cancer and its protein function documented in other previous studies suggest that IL6 protein expression may serve as a therapeutic and diagnostic marker for basal breast cancer growth and metastasis.

Surprisingly, we observed that *GP130*, a functional receptor gene for IL6 signal transduction, has lower mRNA expression in basal breast cancer, compared to luminal breast cancer. At the same time, *IL6* mRNA was not highly expressed in luminal breast cancer, but enriched in basal subtype. This was unexpected because autocrine signaling of IL6-GP130-STAT3 in basal cancer cells is well-studied^{23–25}. This discrepancy may be attributed to differences between gene expression and protein expression or function. Specifically, the membrane receptor *GP130* mRNA must be translated into protein and then bind IL6 protein to function as a signal transducer. Thus, functional studies at the protein level are needed. Nonetheless, lower *GP130* gene expression is consistent with reduced GP130 protein expression observed in a recent study by Lee *et al.*²⁶. Specifically, this study showed that multiple cancer cells (e.g., lymphoma, adenocarcinoma, breast and prostate cancer) maintain activated STAT3 persistently via SIP-S1PR1 signaling, without IL6-GP130 signaling²⁶. Sphingosine-1-phosphate receptor-1 (S1PR1), a receptor for the sphingosine-1-phosphate (S1P), is elevated in STAT3-positive tumors. S1P-S1PR1-induced STAT3 activation is persistent, in contrast to transient STAT3 activation by IL6 protein. This may suggest that basal breast tumor with lower mRNA expression of *GP130* may employ other pathways, such as S1P-S1PR1²⁷. Since paracrine roles of IL6 protein are relatively less understood, our data suggest that *IL6* mRNA expressed by basal tumor cells can play a role in paracrine activators for other types of breast cancer cells (e.g., luminal) or stromal cells. It has been reported that HOXB13 protein mediates tamoxifen resistance and invasiveness in luminal breast cancer by suppressing estrogen receptor (ER) and inducing IL6 protein expression²⁸. This study demonstrates that IL6 signaling promotes aggressiveness in luminal breast cancer cells making them more basal-like. We also showed that IL6 protein expressed by basal cancer cells activated other stromal cells (e.g., lymphatic endothelial cells) to promote tumor metastasis. Paracrine roles of the IL6 protein are still less understood, warranting further studies, particularly in the luminal subtype of breast cancer and in other stromal cells in basal breast cancer.

In addition to *IL6*, *CCL5* mRNA expression is highest in basal breast cancer, which is consistent with our experimental study showing that IL6 protein expressed by basal cancer cells conditions LEC to overexpress CCL5 protein⁴. That previous study also reported a significant correlation between *CCL5* and *IL6* gene expression in lymph node positive basal breast cancer samples from TCGA; there was no correlation for lymph node negative samples. We note that our previous study associated CCL5 protein expression in the lungs and lymph nodes with metastatic potential, but not normal LECs or cancer cells. Whereas that study analyzed protein expression in isolated cell types, the present study analyzes gene expression in primary tumors. The primary tumor samples in this study are from not purified cancer cells. As a result, this study cannot quantify the expression of *CCL5* in distinct cell types. Therefore, further study is required to establish the relative expression of *CCL5* in distinct cell types or metastatic sites suggested in our previous study. Nonetheless, the primary samples contain a mixture of cells so that the gene expression profiling may also characterize expression from LECs that are located in the tumor and express CCL5. We therefore hypothesize that *IL6* and *CCL5* gene expression within basal cancer tumor samples may determine their metastatic potential.

In this study higher *CCL5* gene expression is associated with better prognosis in HER2+ breast cancer samples, but not basal breast cancer. The survival data are in contrast to association of *CCL5* protein expression with metastatic potential in basal breast cancer. Nonetheless, the role of *CCL5* in HER2+ breast cancer metastases warrants further study. For example, HER2+ breast cancer shows high rate of brain metastasis, compared to other subtypes^{29,30} and shows severe drug resistance³¹. Our previous study found that both of these phenotypes are consistent with *CCL5* overexpression. Moreover, recently, a STAT3-*CCL5* loop was studied in drug resistance in luminal cancer¹⁹ and distal metastasis in basal breast cancer⁴. As was the case for *GP130*, differences may be attributed to discrepancies between measurements of mRNA expression and protein function. Discrepancy between *CCL5* gene expression and survival in HER2+ breast cancer may also arise from confounding clinical factors in the survival analysis that are independent of either therapeutic resistance or metastatic site. Associations of *CCL5* expression with time to metastasis or metastatic site may be more consistent with the metastatic potential established in previous studies. However, adequate clinical data for these analyses are not available for either METABRIC or TCGA. Future prospective studies are required to establish the link between *CCL5* expression, metastatic potential, and survival in breast cancer subtypes. The *CCL5* protein may alternatively play a different role in HER2+ breast cancer and must be studied in tumor-drug resistance.

VEGFA and VEGFC are angiogenic and lymphangiogenic growth factors, respectively. We showed that mRNA expression for these growth factors is highest in HER2+ breast cancer compared to other subtypes and that *VEGFR2* mRNA expression predicted survival in HER2 patients. It has been shown that HER2 and angiogenesis signaling pathways exhibit molecular crosstalk³². In that study, higher microvascular density (enhanced angiogenesis) in human breast tumor samples predicted higher co-expression of HER2 and ER³³. Angiogenesis impairment in *Id*-deficient mice completely suppressed HER2/neu-dependent breast tumors³⁴, suggesting a role of angiogenic and lymphangiogenic growth factors in supporting tumor growth and hematogenous metastasis. Nonetheless, higher mRNA expression is associated with better prognosis in HER2+ breast cancer, similar to inconsistencies between the metastatic potential and survival of HER2+ breast cancer for *CCL5*. *NRP1* was overexpressed in basal breast cancer relative to other subtypes in METABRIC (Fig. 5e). It has been reported that expression of both VEGF and semaphorin genes are altered in basal breast cancer³⁵. Semaphorin proteins are ligands of NRP proteins and exhibit anti-angiogenic and anti-lymphangiogenic property³⁶. A pattern of high VEGFA expression with low expression of secreted semaphorins was associated with 60% of basal breast tumors³⁵. Though *VEGFC* mRNA expression in HER2+ breast cancer is less well-understood, recent study showed that HER2/neu expression correlates with VEGFC and lymphangiogenesis in lymph node-positive breast cancer³⁷. Molecular crosstalk between *VEGFC* mRNA expression and HER2 / HER2-dependent transcription factors remains to be investigated in future studies.

We also found that *GMRA* and *GMRB*, possible receptors for *CSF2* (*GM-CSF*), to have high gene expression in basal breast cancer. Their high mRNA expression in HER2+ breast cancer was correlated with better survival. Roles of GM-CSF signaling in breast cancer are still controversial. GM-CSF is known as either an anti-tumorigenic host immune booster^{38,39} or anti-angiogenic factor⁴⁰ or pro-metastatic factor⁴¹. These suggest that targeting *CSF2* signaling must be considered carefully, and needs to be further clarified with more mechanistic studies.

In summary, we analyzed expression of pro-metastatic factor genes in breast cancer subtypes and showed correlation between factor/receptor gene expression and patient survival rates using TCGA and METABRIC datasets. From the study, we found that *IL6* and *CCL5* are overexpressed in basal breast cancer, suggesting their potential as therapeutic targets. It remains to be determined if VEGFA and its receptor *VEGFR2* and VEGFC and its receptor *VEGFR3*, and *CSF2* and its receptor *GMRA/GMRB* can also serve as therapeutic targets for HER2+ breast cancer since the associations we found were modest. High levels of gene expression of *IL6* receptor, *GP130*, in luminal A and B warrant further studies of paracrine roles of *IL6* in luminal cancer. Other cell types such as tumor-associated macrophages can contribute to secretion of *IL6* in tumor microenvironment; this has been observed in other tumor types^{42–44}. In our previous study, we showed by immunohistochemistry that *CCL5* protein expression was co-localized with LECs in the lungs in tumor-bearing mice, however, normal mice without tumors did not show *CCL5* expression in LEC⁴. However, the primary tumor samples profiled in both TCGA and METABRIC contain a mixture tumor cells and cells in the microenvironment. Therefore, future screening studies of the protein expression of these factors on microdissected tissues must be performed to assess regulatory relationships within the different subtypes for treatment selection. Future prospective studies of associating *CCL5*, *VEGFR2*, and *CSF2* expression with metastases could mitigate the confounding factors that may be contributing to contradictory associations in the survival analyses.

Methods

Primary breast cancer gene expression data in TCGA. Analyses of TCGA data¹⁴ are performed on primary breast cancer tumor samples with both RNA-sequencing data and clinical annotations. Level 3 normalized gene expression (RNA Seq V2) is obtained from cBioPortal using the CRAN CGDS-R package (version 1.1.30)⁴⁵. Gene expression data is log₂ transformed and subset to the genes of interest in Fig. 1. The following aliases identified from the Bioconductor package org.Hs.eg.db (version 2.14.0) are used to match the gene annotations used in the TCGA alignment and normalization pipeline according to Table 2.

Clinical data for each TCGA sample is downloaded directly from the TCGA Data Portal. ER and PR status are assessed using the consensus of clinical tests and summarized in “breast carcinoma estrogen receptor status” and “breast carcinoma progesterone receptor status”, respectively. HER2 status is obtained from IHC in the variable labeled “lab proc her2 neu immunohistochemistry receptor status.” Samples missing data for any one of these tests are excluded from analysis, leaving a total of 638 samples with RNA-sequencing data. Breast cancer samples are defined as “Basal” if all three markers are negative, “HER2+” if only HER2 is positive, “Luminal A” if either ER or PR are positive but not HER2, and “Luminal B” if HER2 is positive in addition to either ER or PR. Supplementary Table 1 summarizes the clinical attributes of each of the 638 samples by these subtypes.

Primary breast cancer gene expression data in METABRIC. Gene expression data from METABRIC¹⁵ are obtained from the public domain training data. Normal samples are excluded from analysis, and subtypes are defined using the PAM50 class available in the clinical data. Supplementary Table 2 summarizes the clinical attributes of the 897 primary tumors by these subtypes. We compute survival times from the difference between diagnosis and follow-up dates. Patients are considered to have an event if they died of their disease, as indicated with the label “d-d.s” in the “last follow up status” variable.

We link METABRIC gene identifiers of the genes of interest in Fig. 1. In each case, we select probes indicated as having “Perfect” evidence in the annotation. We select the probe that with the lowest p-value for differential expression between subtypes for genes with multiple probes based upon the differential expression analysis described below, listed in Table 2.

Differential expression analysis. For both TCGA and METABRIC, one-sided t-tests are applied to each gene to compare expression of samples in each subtype to samples from all other subtypes. P-values are adjusted using the Benjamini-Hochberg procedure to account for multiple hypothesis testing. In the case of METABRIC, differential expression and survival statistics are reported for the probe that has the lowest p-value in any subtype relative to the other subtypes for each gene. All analyses are performed in R, version 3.1.1.

Survival analysis. Due to the relatively limited follow-up time in TCGA, survival analyses are performed only for METABRIC data. The function “survdif” in the CRAN package survival (2.37.7) applies the *G-rho* family of tests to compare survival curves for samples with high expression to samples with low expression to each gene in Fig. 1 that is also measured in METABRIC. We distinguish samples as having high or low expression of a gene relative to the distribution of expression values in the subtype that has lowest average expression of that gene. Specifically, a sample is defined as having high expression of a gene if its expression is at least one standard deviation above its mean expression in the subtype with lowest average expression. We do not perform analysis on combinations of genes and subtypes that have fewer than 10 samples with high or low expression.

References

- DeSantis, C., Ma, J., Bryan, L. & Jemal, A. Breast cancer statistics, 2013. *CA Cancer J Clin* **64**, 52–62 (2014).
- Lehmann, B. D. *et al.* Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* **121**, 2750–2767 (2011).
- Eroles, P., Bosch, A., Bermejo, B. & Lluch, A. Mechanisms of resistance to hormonal treatment in breast cancer. *Clin Transl Oncol* **12**, 246–252 (2010).
- Lee, E. *et al.* Breast cancer cells condition lymphatic endothelial cells within pre-metastatic niches to promote metastasis. *Nat Commun* **5**, 4715 (2014).
- Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *Cell* **144**, 646–674 (2011).
- Elinav, E. *et al.* Inflammation-induced cancer: crosstalk between tumours, immune cells and microorganisms. *Nat Rev Cancer* **13**, 759–771 (2013).
- Rot, A. & von Andrian, U. H. Chemokines in innate and adaptive host defense: basic chemokine grammar for immune cells. *Annu Rev Immunol* **22**, 891–928 (2004).
- Athale, C. A. & Deisboeck, T. S. The effects of EGF-receptor density on multiscale tumor growth patterns. *J Theor Biol* **238**, 771–779 (2006).
- Li, T., Yang, J., Zhou, Q. & He, Y. Molecular regulation of lymphangiogenesis in development and tumor microenvironment. *Cancer Microenviron* **5**, 249–260 (2012).
- Cao, Y. *et al.* Forty-year journey of angiogenesis translational research. *Sci Transl Med* **3**, 114rv113 (2011).
- Chen, S. T. *et al.* Breast tumor microenvironment: proteomics highlights the treatments targeting secretome. *J Proteome Res* **7**, 1379–1387 (2008).
- Lee, E., Pandey, N. B. & Popel, A. S. Lymphatic endothelial cells support tumor growth in breast cancer. *Sci Rep* **4**, 5853 (2014).
- Stacker, S. A. *et al.* Lymphangiogenesis and lymphatic vessel remodelling in cancer. *Nat Rev Cancer* **14**, 159–172 (2014).
- Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61–70 (2012).
- Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature* **486**, 346–352 (2012).
- Dethlefsen, C., Hojfeldt, G. & Hojman, P. The role of intratumoral and systemic IL-6 in breast cancer. *Breast Cancer Res Treat* **138**, 657–664 (2013).
- Iliopoulos, D., Hirsch, H. A., Wang, G. & Struhl, K. Inducible formation of breast cancer stem cells and their dynamic equilibrium with non-stem cancer cells via IL6 secretion. *Proc Natl Acad Sci U S A* **108**, 1397–1402 (2011).
- De Luca, A., Lamura, L., Gallo, M., Maffia, V. & Normanno, N. Mesenchymal stem cell-derived interleukin-6 and vascular endothelial growth factor promote breast cancer cell migration. *J Cell Biochem* **113**, 3363–3370 (2012).
- Yi, E. H. *et al.* STAT3-RANTES autocrine signaling is essential for tamoxifen resistance in human breast cancer cells. *Mol Cancer Res* **11**, 31–42 (2013).

20. Roberti, M. P. *et al.* Protein expression changes during human triple negative breast cancer cell line progression to lymph node metastasis in a xenografted model in nude mice. *Cancer Biol Ther* **13**, 1123–1140 (2012).
21. Wang, S. W. & Sun, Y. M. The IL-6/JAK/STAT3 pathway: potential therapeutic strategies in treating colorectal cancer (Review). *Int J Oncol* **44**, 1032–1040 (2014).
22. Tell, R. W. & Horvath, C. M. Bioinformatic analysis reveals a pattern of STAT3-associated gene expression specific to basal-like breast cancers in human tumors. *Proc Natl Acad Sci U S A* **111**, 12787–12792 (2014).
23. Ara, T. *et al.* Critical Role of STAT3 in IL-6-Mediated Drug Resistance in Human Neuroblastoma. *Cancer Res* **73**, 3852–3864 (2013).
24. Ashizawa, T. *et al.* Clinical significance of interleukin-6 (IL-6) in the spread of gastric cancer: role of IL-6 as a prognostic factor. *Gastric Cancer* **8**, 124–131 (2005).
25. Xiong, H. *et al.* Roles of STAT3 and ZEB1 proteins in E-cadherin down-regulation and human colorectal cancer epithelial-mesenchymal transition. *J Biol Chem* **287**, 5819–5832 (2012).
26. Lee, H. *et al.* STAT3-induced S1PR1 expression is crucial for persistent STAT3 activation in tumors. *Nat Med* **16**, 1421–1428 (2010).
27. Taga, T. IL6 signalling through IL6 receptor and receptor-associated signal transducer, gp130. *Res Immunol* **143**, 737–739 (1992).
28. Shah, N. *et al.* HOXB13 mediates tamoxifen resistance and invasiveness in human breast cancer by suppressing ERalpha and inducing IL-6 expression. *Cancer Res* **73**, 5449–5458 (2013).
29. Hicks, D. G. *et al.* Breast cancers with brain metastases are more likely to be estrogen receptor negative, express the basal cytokeratin CK5/6, and overexpress HER2 or EGFR. *Am J Surg Pathol* **30**, 1097–1104 (2006).
30. Azim, H. A. & Azim, H. A., Jr. Systemic treatment of brain metastases in HER2-positive breast cancer: current status and future directions. *Future Oncol* **8**, 135–144 (2012).
31. Campone, M., Frenel, J. S., Andre, F., Bachelot, T. & Juin, P. [Tumor resistance to HER2 inhibitors: the drug sedimentation concept]. *Bull Cancer* **99**, 665–672 (2012).
32. Alameddine, R. S., Otrrock, Z. K., Awada, A. & Shamseddine, A. Crosstalk between HER2 signaling and angiogenesis in breast cancer: molecular basis, clinical applications and challenges. *Curr Opin Oncol* **25**, 313–324 (2013).
33. Vamesu, S. Angiogenesis and co-expressed of ER and c-erbB-2 (HER2/neu) protein in primary breast cancer patients: an analysis of 158 needle core biopsies. *Rom J Morphol Embryol* **49**, 469–478 (2008).
34. de Candia, P. *et al.* Angiogenesis impairment in Id-deficient mice cooperates with an Hsp90 inhibitor to completely suppress HER2/neu-dependent breast tumors. *Proc Natl Acad Sci U S A* **100**, 12337–12342 (2003).
35. Bender, R. J. & Mac Gabhann, F. Expression of VEGF and semaphorin genes define subgroups of triple negative breast cancer. *PLoS One* **8**, e61788 (2013).
36. Sakurai, A., Doci, C. L. & Gutkind, J. S. Semaphorin signaling in angiogenesis, lymphangiogenesis and cancer. *Cell Res* **22**, 23–32 (2012).
37. Schoppmann, S. F. *et al.* HER2/neu expression correlates with vascular endothelial growth factor-C and lymphangiogenesis in lymph node-positive breast cancer. *Ann Oncol* **21**, 955–960 (2010).
38. Emens, L. A. GM-CSF-secreting vaccines for solid tumors. *Curr Opin Investig Drugs* **10**, 1315–1324 (2009).
39. Emens, L. A. *et al.* A phase I vaccine safety and chemotherapy dose-finding trial of an allogeneic GM-CSF-secreting breast cancer vaccine given in a specifically timed sequence with immunomodulatory doses of cyclophosphamide and doxorubicin. *Hum Gene Ther* **15**, 313–337 (2004).
40. Eubank, T. D. *et al.* Granulocyte macrophage colony-stimulating factor inhibits breast cancer growth and metastasis by invoking an anti-angiogenic program in tumor-educated macrophages. *Cancer Res* **69**, 2133–2140 (2009).
41. Park, B. K. *et al.* NF-kappaB in breast cancer cells promotes osteolytic bone metastasis by inducing osteoclastogenesis via GM-CSF. *Nat Med* **13**, 62–69 (2007).
42. Wan, S. *et al.* Tumor-associated macrophages produce interleukin 6 and signal via STAT3 to promote expansion of human hepatocellular carcinoma stem cells. *Gastroenterology* **147**, 1393–1404 (2014).
43. Xu, H. *et al.* Tumor-associated macrophage-derived IL-6 and IL-8 enhance invasive activity of LoVo cells induced by PRL-3 in a KCNN4 channel-dependent manner. *BMC Cancer* **14**, 330 (2014).
44. Sung, S. Y. *et al.* Loss of let-7 microRNA upregulates IL-6 in bone marrow-derived mesenchymal stem cells triggering a reactive stromal response to prostate cancer. *PLoS One* **8**, e71637 (2013).
45. Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* **6**, pl1 (2013).

Acknowledgements

This work was supported by the National Institutes of Health grants R01 CA138264 (ASP) and K25CA141053 (EJF).

Author Contributions

E.J.F., E.L., N.B.P. and A.S.P. designed the study. E.J.F. performed all statistical analyses. All authors wrote and revised the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Fertig, E. J. *et al.* Analysis of gene expression of secreted factors associated with breast cancer metastases in breast cancer subtypes. *Sci. Rep.* **5**, 12133; doi: 10.1038/srep12133 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>