


Editorial

Sexual abuse and mental ill health in boys and men: what we do and don't know

Simon M. Rice, Scott D. Easton, Zac E. Seidler and John L. Oliffe



Summary

The spectrum of adverse mental health trajectories caused by sexual abuse, broadly defined as exposure to rape and unwanted physical sexual contact, is well-known. Few studies have systematically appraised the epidemiology and impact of sexual abuse among boys and men. New meta-analytic insights ($k = 44$; $n = 45\,172$) reported by Zarchev and colleagues challenge assumptions that men experiencing mental ill health rarely report sexual abuse exposure. Adult-onset sexual abuse rates of 1–7% are observed in the general population, but for men experiencing mental ill health, adult lifetime prevalence was 14.1% (95% CI 7.3–22.4%), with past-year exposure 5.3% (95% CI 1.6–12.8%). We note that these rates are certainly underestimates, as childhood sexual abuse exposures were excluded. Boys and men with a sexual abuse history experience substantial disclosure and treatment barriers. We draw attention to population health gains that could be achieved via implementation of

gender-sensitive assessment and intervention approaches for this at-risk population.

Keywords

Sexual abuse; gender; help-seeking; masculinity; male.

Copyright and usage

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Simon Rice (pictured) is Head of Gender and Social Psychiatry in Youth Mental Health at Orygen and the Centre for Youth Mental Health at the University of Melbourne, Australia. **Scott Easton** is Associate Professor, Mental Health Department Chair and co-director of the Trauma Integration Initiative at the Boston College School of Social Work, Boston, USA. **Zac Seidler** is a Senior Research Fellow at Orygen and the Centre for Youth Mental Health at the University of Melbourne, Australia and Director of Mental Health Training at Movenber, Victoria, Australia. **John Oliffe** is Professor and Tier 1 Canada Research Chair in Men's Health Promotion at the School of Nursing, University of British Columbia, Vancouver, Canada.

New prevalence data

Social taboos related to male sexual victimisation, coupled with significant disclosure barriers faced by boys and men exposed to sexual abuse, catalyse societal and self-stigma, in addition to perceived inadequacy and shame.¹ Collectively, these factors adversely affect the mental health and well-being of boys and men with sexual abuse histories. Relative to girls and women, boys and men are less likely to: disclose sexual abuse at the time, accurately label experiences as sexual abuse and receive acknowledgement or validation of sexual abuse experiences from clinicians and/or parents.² It is commonplace for men to defer initial disclosure for upwards of three decades.³ Best estimates regarding the latency of sexual abuse disclosure among men reasonably suggest a proportion of these men never experience sufficient safety⁴ to speak of their abuse experience(s), with the psychosocial impacts silently concealed and suffered or self-managed.

Compelling data show that at the population health level, exposure to traumatic events such as childhood and adult-onset sexual abuse can have a powerful impact on the mental ill health trajectories of boys and men. This includes elevated risk for suicide attempts and death by suicide,⁵ as well as engaging in maladaptive male-specific coping behaviours, influenced by gender norms especially among younger males,⁶ including aggression, substance misuse and risk-taking.⁷ The knowledge base related to sexual abuse among boys and men is underdeveloped.⁸

Zarchev and colleagues⁹ present compelling meta-analytic prevalence data on adult-onset sexual abuse (defined as exposure to rape and unwanted physical sexual contact at ≥ 18 years of age) among men with a history of mental ill health. As the authors correctly state, scant research exists on the male experience of sexual abuse⁸ (including epidemiological data), and in their review ($k = 44$; $n = 45\,172$) using Bayesian multi-level models, important contributions are offered to advance the field. First, Zarchev and colleagues differentiated sexual abuse exposure according to abuse perpetrated within familial or intimate relationships versus non-familial/partner perpetrators. Second, analyses examined disorder and comorbidity-specific effects. Third, meta-regression techniques were used to investigate effects according to study characteristics (e.g. age, year of data collection, screening instrument, sampling design, implementation of instrument).

Findings indicated that for men experiencing mental ill health, adult lifetime sexual abuse prevalence was 14.1% (95% CI 7.3–22.4%) and past-year exposure was 5.3% (95% CI 1.6–12.8%). There was little change to the adulthood lifetime rate when studies at higher risk of bias were omitted from analysis. Men with mental illness were not at higher risk of sexual abuse perpetrated by an intimate partner than men in the general population; however, the key finding indicated that the prevalence of adult-onset lifetime abuse for men diagnosed with a mental disorder was potentially three times higher relative to the general population (1–7%). This underscores the need for research to sensitively screen for and assess sexual abuse exposure among men attending clinical settings (including those presenting for care in later life¹⁰) and to develop trauma-informed care mindful of masculine norms, some of which are known barriers to disclosure and treatment engagement.¹ We applaud the data-driven insights provided by Zarchev and colleagues and offer reflections for driving the field forward.

Suggestions for future research

Improved data collection

Over 50% of the included studies used convenience sampling, with a similar proportion failing to report response rates. Nonetheless, the reported adult lifetime and past-year rates reported by Zarchev and colleagues are certain to be underestimates of prevalence among men experiencing mental ill health, owing to the known latency of male sexual abuse disclosure (e.g. including those who potentially deny true exposure in clinical interview) and the exclusion of studies reporting childhood sexual abuse (onset at <18 years of age). Furthermore, all but one of the included studies involved a clinician or trained specialist assigning psychiatric classification. Although this approach is considered gold standard for enhancing validity of coded diagnostic impressions, the lack of anonymity may have resulted in a proportion of men denying true abuse exposure because of shame and other factors.⁴

Zarchev and colleagues⁹ highlight that reported lifetime prevalence rates would be markedly higher if childhood sexual abuse studies were included in the analysis, with the true prevalence of male childhood sexual abuse in men with mental illness yet to be determined. Prior meta-analysis in the general population (<18 years of age) indicates that 8% (95% CI 4–16%) of men have experienced childhood sexual abuse broadly defined,¹¹ and a recent self-report study adopting convenience sampling among 1277 young men (≤30 years of age) seeking mental health help yielded a childhood sexual abuse prevalence rate of 21.9%.⁷ This rate is remarkably similar to the 21.1% self-reported prevalence in the Zarchev review, which was notably 11.5% higher than the 9.6% prevalence reported in face-to-face studies.

There are of course significant challenges in collecting accurate childhood sexual abuse prevalence data among boys and men, and research on trauma history assessment that avoids causing harm is needed. Improved prevalence data matter for a number of important reasons. Recent longitudinal research has shown that, after controlling for sociodemographic, health and background factors, childhood sexual abuse undermines men's mental health across the life course well into men's sixth, seventh and eighth decades of life.¹⁰ Hence, there is a rationale for reducing the duration of untreated (or unacknowledged) sexual abuse among boys and men, with the aim of offsetting subsequent distress or illness. That said, it is acknowledged that humans have extraordinary reservoirs of, and capacity for, resilience in the aftermath of trauma exposures. Formal psychological intervention may not be necessary for many men with a sexual abuse history – indeed sexual abuse exposure need not consign boys and men to lives of misery. Many men articulately reflect on experiences of post-traumatic growth¹² and positive identity development after abuse exposure.

Intervention studies: how and when to intervene


Previous studies have called for an urgent focus on intervention-based research for males with histories of sexual abuse, including the roles of formal and informal support groups, psychoeducation and awareness campaigns, and refinement of clinical treatment modalities.⁸ This may extend to case conceptualisation, assessment and diagnosis, inclusive of older men who may be presenting to mental healthcare services for the first time. Following exposure to traumatic events, including sexual abuse, males are known to be less likely to experience threshold post-traumatic stress disorder (PTSD) compared with females, but report comparatively higher rates of externalising symptoms, including alcohol and drug use, irritability, anger and violent behaviour.¹³ Of note, DSM-5-TR acknowledges that this constellation of externalising symptoms

are particularly likely to co-occur with major depressive disorder in men.¹⁴ Traumatic events may exacerbate underlying socially influenced gender differences in response to distress, resulting in alternative constellations of post-traumatic symptom patterns among boys and men.¹³

Clinicians reticent to ask about men's sexual abuse experiences may inadvertently add to the harms experienced by men seeking care. Poor treatment experiences among male sexual abuse populations can reinforce a sense of shame and secrecy, perpetuate rejection and prohibit the development of therapeutic trust, acceptance, connection and honest disclosure with treating therapists.² Future research should examine differential outcomes for boys and men based on whether disclosure (and treatment access) is provided proximally to exposure, and the types of supportive approaches and interventions that boys and men experience as acceptable, engaging and safe. Emerging evidence suggests that (a) early disclosure (within 1 year) of sexual abuse, (b) the nature of the response to abuse disclosure and (c) having in-depth discussions about abuse experiences are each protective for men's mental distress.⁸ These factors do, however, raise complexities, including safety implications for those less than 18 years of age, the readiness, confidence and willingness of clinicians, peers and family to appropriately listen to and validate sexual abuse disclosure, and the extremely limited availability of appropriately sensitive services for boys and men. These are areas requiring urgent research and practice advances.

Implications for population health

Adequately addressing the mental health needs of boys and men who have experienced sexual abuse is certain to spur major advances in population health. It is increasingly clear from the meta-analysis of Zarchev et al⁹ and others¹¹ that sexual abuse among boys and men is massively underrecognised and undertreated. The associated burden of disease, comorbidity and suffering, both for the individual and that imposed on others, is immense. Attention to this problem would likely reduce rates of suicide among men, which globally eclipse suicide rates among women by a factor of two to four.¹⁵ When extrapolating conservative prevalence rates of undetected and untreated sexual abuse, maladaptive coping strategies (e.g. substance use, aggression, risk-taking) that risk harm to self and others could be reduced in hundreds of millions of boys and men globally. There is growing attention to the ways in which gender norms affect the social and emotional development of boys and men¹⁵ and future tailored interventions and programme development for boys and men affected by sexual abuse must be cognisant and transformative of such norms. This extends to initial disclosure and help-seeking, tailoring of specialised services and clinical trials to improve long-term psychosocial functioning. Too often ignored and hidden in plain sight, boys and men with sexual abuse histories require focused attention to address their unmet mental health needs. Doing so will improve not only the mental health of affected individuals, but extend health benefits to the families and communities where these men live.

Simon M. Rice , PhD, Orygen, Parkville, Melbourne, Victoria, Australia; and Centre for Youth Mental Health, University of Melbourne, Victoria, Australia; **Scott D. Easton**, PhD, School of Social Work, Boston College, Massachusetts, USA; **Zac E. Seidler**, PhD, Orygen, Parkville, Melbourne, Victoria, Australia; and Centre for Youth Mental Health, University of Melbourne, Victoria, Australia; **John L. Oliffe**, PhD, School of Nursing, University of British Columbia, Vancouver, Canada; and Department of Nursing, University of Melbourne, Victoria, Australia

Correspondence: Simon Rice. Email: simon.rice@orygen.org.au

First received 17 Dec 2021, final revision 10 Apr 2022, accepted 12 May 2022

Data availability

Data availability is not applicable to this editorial as no new data were created or analysed in its preparation.

Author contributions

S.M.R. developed the initial manuscript draft. All authors contributed to editing and refining the final manuscript.

Funding

This work received no specific grant from any funding agency, commercial or not-for-profit sectors.

Declaration of interest

S.M.R. was supported by a Career Development Fellowship from the National Health and Medical Research Council of Australia (NHMRC; GNT1158881) and the Dame Kate Campbell Fellowship from the Faculty of Medicine, Dentistry and Health Sciences at the University of Melbourne. In the past 5 years S.M.R. has received funding and research support from the NHMRC, Movember Foundation, International Olympic Committee, Australian Football League, TAL Foundation, McCusker Charitable Foundation, Royal Melbourne Hospital, Foundation for Positive Masculinity, Northwestern Melbourne Primary Health Network and the Marian and EH Flack Trust.

References

- Easton SD, Parchment TM. 'The whole wall fell apart, and I felt free for the first time': men's perceptions of helpful responses during discussion of child sexual abuse. *Child Abuse Negl* 2021; **112**: 104922.
- Rapsey C, Campbell A, Clearwater K, Patterson T. Listening to the therapeutic needs of male survivors of childhood sexual abuse. *J Interpers Violence* 2020; **35**: 2033–54.
- Easton SD. Disclosure of child sexual abuse among adult male survivors. *Clin Soc Work J* 2013; **41**: 344–55.
- Easton SD, Saltzman LY, Willis DG. 'Would you tell under circumstances like that?': barriers to disclosure of child sexual abuse for men. *Psychol Men Mascul* 2014; **15**: 460–9.
- Easton SD, Renner LM, O'Leary P. Suicide attempts among men with histories of child sexual abuse: examining abuse severity, mental health, and masculine norms. *Child Abuse Negl* 2013; **37**: 380–7.
- Rice SM, Purcell R, McGorry PD. Adolescent and young adult male mental health: transforming system failures into proactive models of engagement. *J Adolesc Health* 2018; **62**: S9–17.
- Rice SM, Kealy D, Seidler ZE, Walton CC, Oliffe JL, Ogronczuk JS. Male-type depression symptoms in young men with a history of childhood sexual abuse and current hazardous alcohol use. *Psychiatry Res* 2021; **304**: 114110.
- Easton SD. Childhood disclosure of sexual abuse and mental health outcomes in adulthood: assessing merits of early disclosure and discussion. *Child Abuse Negl* 2019; **93**: 208–14.
- Zarchev M, Roos RE, Mulder C, Kamperman AM. Prevalence of adult sexual abuse in men with mental illness: Bayesian meta-analysis. *BJPsych Open* 2022; **8**(1): e16.
- Easton SD, Kong J, Gregas MC, Shen C, Shafer K. Child sexual abuse and depression in late life for men: a population-based, longitudinal analysis. *J Gerontol* 2019; **74**: 842–52.
- Barth J, Bermetz L, Heim E, Trelle S, Tonia T. The current prevalence of child sexual abuse worldwide: a systematic review and meta-analysis. *Int J Public Health* 2013; **58**: 469–83.
- Easton SD, Coohy C, Rhodes AM, Moorthy MV. Posttraumatic growth among men with histories of child sexual abuse. *Child Maltreat* 2013; **18**: 211–20.
- Tolin DF, Foa EB. Sex differences in trauma and posttraumatic stress disorder: a quantitative review of 25 years of research. *Psychol Bull* 2006; **132**: 959–92.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders* (5th edn, text revision) (DSM-5-TR). American Psychiatric Association, 2022.
- Rice S, Oliffe J, Seidler Z, Borschmann R, Pirkis J, Reavley N, et al. Gender norms and the mental health of boys and young men. *Lancet Public Health* 2021; **6**(8): e541–2.

