

ADOPTED: 15 September 2017

doi: 10.2903/j.efsa.2017.4999

Pest categorisation of *Ips sexdentatus*

EFSA Panel on Plant Health (PLH),

Michael Jeger, Claude Bragard, David Caffier, Thierry Candresse, Elisavet Chatzivassiliou, Katharina Dehnen-Schmutz, Gianni Gilioli, Josep Anton Jaques Miret, Alan MacLeod, Maria Navajas Navarro, Björn Niere, Stephen Parnell, Roel Potting, Trond Rafoss, Vittorio Rossi, Gregor Urek, Ariena Van Bruggen, Wopke Van der Werf, Jonathan West, Stephan Winter, Virág Kertész, Mitesha Aukhojee and Jean-Claude Grégoire

Abstract

The Panel on Plant Health performed a pest categorisation of the six-toothed bark beetle, *Ips sexdentatus* (Börner) (Coleoptera: Curculionidae, Scolytinae), for the EU. *I. sexdentatus* is a well-defined and distinguishable species, native to Eurasia and recognised mainly as a pest of pine (*Pinus* spp., in the pest's whole range) and spruce (mainly *Picea orientalis* in Turkey and Georgia). It also might occasionally attack *Larix* spp. and *Abies* spp. It is distributed throughout the EU (24 Member States). It is a protected zone quarantine pest in Ireland, Cyprus and the United Kingdom (Northern Ireland, Isle of Man), listed in Annex IIB of Council Directive 2000/29/EC. Wood, wood products, bark and wood packaging material are considered as pathways for this pest, which is also able to disperse by flight over tens of kilometres. The adults normally establish on fallen or weakened trees (e.g. after a fire or a drought) and can also mass-attack healthy trees. The males produce aggregation pheromones that attract conspecifics of both sexes. The insects also inoculate pathogenic fungi to their hosts. There are one to five generations per year. The wide current geographical range of *I. sexdentatus* suggests that it is able to establish anywhere in the EU where its hosts are present. Sanitary thinning or clear-felling are the major control methods. Pheromone mass-trapping is also locally implemented. Quarantine measures are implemented to prevent entry into the protected zones. All criteria for consideration as potential protected zone quarantine pest are met. The criteria for considering *I. sexdentatus* as a potential regulated non-quarantine pest are not met since plants for planting are not viewed as a pathway.

© 2017 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

Keywords: Curculionidae, European Union, pest risk, plant health, plant pest, quarantine, six-toothed bark beetle

Requestor: European Commission

Question number: EFSA-Q-2017-00203

Correspondence: alpha@efsa.europa.eu

Panel members: Claude Bragard, David Caffier, Thierry Candresse, Elisavet Chatzivassiliou, Katharina Dehnen-Schmutz, Gianni Gilioli, Jean-Claude Grégoire, Josep Anton Jaques Miret, Michael Jeger, Alan MacLeod, Maria Navajas Navarro, Björn Niere, Stephen Parnell, Roel Potting, Trond Rafoss, Vittorio Rossi, Gregor Urek, Ariena Van Bruggen, Wopke Van der Werf, Jonathan West and Stephan Winter.

Acknowledgements: The Panel wishes to acknowledge all European competent institutions, Member State bodies and other organisations that provided data for this scientific output.

Suggested citation: EFSA PLH Panel (EFSA Panel on Plant Health), Jeger M, Bragard C, Caffier D, Candresse T, Chatzivassiliou E, Dehnen-Schmutz K, Gilioli G, Jaques Miret JA, MacLeod A, Navajas Navarro M, Niere B, Parnell S, Potting R, Rafoss T, Rossi V, Urek G, Van Bruggen A, Van der Werf W, West J, Winter S, Kertész V, Aukhojee M and Grégoire J-C, 2017. Scientific Opinion on the pest categorisation of *Ips sexdentatus*. EFSA Journal 2017;15(11):4999, 28 pp. <https://doi.org/10.2903/j.efsa.2017.4999>

ISSN: 1831-4732

© 2017 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs License](#), which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.

Reproduction of the images listed below is prohibited and permission must be sought directly from the copyright holder:

Figure 1: © EPPO

Figure 2: © European Union, 2017. Reuse is authorised, provided the source is acknowledged



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



Table of contents

Abstract.....	1
1. Introduction.....	4
1.1. Background and Terms of Reference as provided by the requestor.....	4
1.1.1. Background.....	4
1.1.2. Terms of Reference.....	4
1.1.2.1. Terms of Reference: Appendix 1.....	5
1.1.2.2. Terms of Reference: Appendix 2.....	6
1.1.2.3. Terms of Reference: Appendix 3.....	7
1.2. Interpretation of the Terms of Reference.....	8
2. Data and methodologies.....	8
2.1. Data.....	8
2.1.1. Literature search.....	8
2.1.2. Database search.....	8
2.2. Methodologies.....	8
3. Pest categorisation.....	10
3.1. Identity and biology of the pest.....	10
3.1.1. Identity and taxonomy.....	10
3.1.2. Biology of the pest.....	11
3.1.3. Detection and identification of the pest.....	11
3.2. Pest distribution.....	11
3.2.1. Pest distribution outside the EU.....	11
3.2.2. Pest distribution in the EU.....	12
3.3. Regulatory status.....	13
3.3.1. Council Directive 2000/29/EC.....	13
3.3.2. Legislation addressing plants and plant parts on which <i>Ips sexdentatus</i> is regulated.....	13
3.3.3. Legislation addressing the organisms vectored by <i>Ips sexdentatus</i> (Directive 2000/29/EC).....	14
3.4. Entry, establishment and spread in the EU.....	15
3.4.1. Host range.....	15
3.4.2. Entry.....	15
3.4.3. Establishment.....	16
3.4.3.1. EU distribution of main host plants.....	16
3.4.3.2. Climatic conditions affecting establishment.....	18
3.4.4. Spread.....	18
3.5. Impacts.....	18
3.6. Availability and limits of mitigation measures.....	19
3.6.1. Biological or technical factors limiting the feasibility and effectiveness of measures to prevent the entry, establishment and spread of the pest.....	19
3.6.2. Control methods.....	19
3.7. Uncertainty.....	19
4. Conclusions.....	20
References.....	21
Abbreviations.....	24
Appendix A – Methodological notes on Figure 2.....	25

1. Introduction

1.1. Background and Terms of Reference as provided by the requestor

1.1.1. Background

Council Directive 2000/29/EC¹ on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community establishes the present European Union plant health regime. The Directive lays down the phytosanitary provisions and the control checks to be carried out at the place of origin on plants and plant products destined for the Union or to be moved within the Union. In the Directive's 2000/29/EC annexes, the list of harmful organisms (pests) whose introduction into or spread within the Union is prohibited, is detailed together with specific requirements for import or internal movement.

Following the evaluation of the plant health regime, the new basic plant health law, Regulation (EU) 2016/2031² on protective measures against pests of plants, was adopted on 26 October 2016 and will apply from 14 December 2019 onwards, repealing Directive 2000/29/EC. In line with the principles of the above mentioned legislation and the follow-up work of the secondary legislation for the listing of EU regulated pests, EFSA is requested to provide pest categorizations of the harmful organisms included in the annexes of Directive 2000/29/EC, in the cases where recent pest risk assessment/ pest categorisation is not available.

1.1.2. Terms of Reference

EFSA is requested, pursuant to Article 22(5.b) and Article 29(1) of Regulation (EC) No 178/2002³, to provide scientific opinion in the field of plant health.

EFSA is requested to prepare and deliver a pest categorisation (step 1 analysis) for each of the regulated pests included in the appendices of the annex to this mandate. The methodology and template of pest categorisation have already been developed in past mandates for the organisms listed in Annex II Part A Section II of Directive 2000/29/EC. The same methodology and outcome is expected for this work as well.

The list of the harmful organisms included in the annex to this mandate comprises 133 harmful organisms or groups. A pest categorisation is expected for these 133 pests or groups and the delivery of the work would be stepwise at regular intervals through the year as detailed below. First priority covers the harmful organisms included in Appendix 1, comprising pests from Annex II Part A Section I and Annex II Part B of Directive 2000/29/EC. The delivery of all pest categorisations for the pests included in Appendix 1 is June 2018. The second priority is the pests included in Appendix 2, comprising the group of *Cicadellidae* (non-EU) known to be vector of Pierce's disease (caused by *Xylella fastidiosa*), the group of *Tephritidae* (non-EU), the group of potato viruses and virus-like organisms, the group of viruses and virus-like organisms of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L. and the group of *Margarodes* (non-EU species). The delivery of all pest categorisations for the pests included in Appendix 2 is end 2019. The pests included in Appendix 3 cover pests of Annex I part A section I and all pests categorisations should be delivered by end 2020.

For the above mentioned groups, each covering a large number of pests, the pest categorisation will be performed for the group and not the individual harmful organisms listed under "such as" notation in the Annexes of the Directive 2000/29/EC. The criteria to be taken particularly under consideration for these cases, is the analysis of host pest combination, investigation of pathways, the damages occurring and the relevant impact.

Finally, as indicated in the text above, all references to 'non-European' should be avoided and replaced by 'non-EU' and refer to all territories with exception of the Union territories as defined in Article 1 point 3 of Regulation (EU) 2016/2031.

¹ Council Directive 2000/29/EC of 8 May 2000 on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community. OJ L 169/1, 10.7.2000, p. 1–112.

² Regulation (EU) 2016/2031 of the European Parliament of the Council of 26 October 2016 on protective measures against pests of plants. OJ L 317, 23.11.2016, p. 4–104.

³ Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31/1, 1.2.2002, p. 1–24.

1.1.2.1. Terms of Reference: Appendix 1

List of harmful organisms for which pest categorisation is requested. The list below follows the annexes of Directive 2000/29/EC.

Annex IIAI

(a) Insects, mites and nematodes, at all stages of their development

<i>Aleurocantus</i> spp.	<i>Numonia pyrivorella</i> (Matsumura)
<i>Anthonomus bisignifer</i> (Schenkling)	<i>Oligonychus perditus</i> Pritchard and Baker
<i>Anthonomus signatus</i> (Say)	<i>Pissodes</i> spp. (non-EU)
<i>Aschistonyx eppoi</i> Inouye	<i>Scirtothrips aurantii</i> Faure
<i>Carposina niponensis</i> Walsingham	<i>Scirtothrips citri</i> (Moultex)
<i>Enarmonia packardi</i> (Zeller)	<i>Scolytidae</i> spp. (non-EU)
<i>Enarmonia prunivora</i> Walsh	<i>Scrobipalopsis solanivora</i> Povolny
<i>Grapholita inopinata</i> Heinrich	<i>Tachypterellus quadrigibbus</i> Say
<i>Hishomonus phycitis</i>	<i>Toxoptera citricida</i> Kirk.
<i>Leucaspis japonica</i> Ckll.	<i>Unaspis citri</i> Comstock
<i>Listronotus bonariensis</i> (Kuschel)	

(b) Bacteria

Citrus variegated chlorosis	<i>Xanthomonas campestris</i> pv. <i>oryzae</i> (Ishiyama)
<i>Erwinia stewartii</i> (Smith) Dye	Dye and pv. <i>oryzicola</i> (Fang, et al.) Dye

(c) Fungi

<i>Alternaria alternata</i> (Fr.) Keissler (non-EU pathogenic isolates)	<i>Elsinoe</i> spp. Bitanc. and Jenk. Mendes
<i>Anisogramma anomala</i> (Peck) E. Müller	<i>Fusarium oxysporum</i> f. sp. <i>albedinis</i> (Kilian and Maire) Gordon
<i>Apiosporina morbosa</i> (Schwein.) v. Arx	<i>Guignardia piricola</i> (Nosa) Yamamoto
<i>Ceratocystis virescens</i> (Davidson) Moreau	<i>Puccinia pittieriana</i> Hennings
<i>Cercoseptoria pini-densiflorae</i> (Hori and Nambu) Deighton	<i>Stegophora ulmea</i> (Schweinitz: Fries) Sydow & Sydow
<i>Cercospora angolensis</i> Carv. and Mendes	<i>Venturia nashicola</i> Tanaka and Yamamoto

(d) Virus and virus-like organisms

Beet curly top virus (non-EU isolates)	Little cherry pathogen (non- EU isolates)
Black raspberry latent virus	Naturally spreading psorosis
Blight and blight-like	Palm lethal yellowing mycoplasma
Cadang-Cadang viroid	Satsuma dwarf virus
Citrus tristeza virus (non-EU isolates)	Tatter leaf virus
Leprosis	Witches' broom (MLO)

Annex IIB

(a) Insect mites and nematodes, at all stages of their development

<i>Anthonomus grandis</i> (Boh.)	<i>Ips amitinus</i> Eichhof
<i>Cephalcia lariciphila</i> (Klug)	<i>Ips cembrae</i> Heer
<i>Dendroctonus micans</i> Kugelan	<i>Ips duplicatus</i> Sahlberg
<i>Gilpinia hercyniae</i> (Hartig)	<i>Ips sexdentatus</i> Börner
<i>Gonipterus scutellatus</i> Gyll.	<i>Ips typographus</i> Heer
<i>Sternochetus mangiferae</i> Fabricius	

(b) Bacteria

Curtobacterium flaccumfaciens pv. *flaccumfaciens*
(Hedges) Collins and Jones

(c) Fungi

Glomerella gossypii Edgerton

Hypoxylon mammatum (Wahl.) J. Miller

Gremmeniella abietina (Lag.) Morelet

1.1.2.2. Terms of Reference: Appendix 2

List of harmful organisms for which pest categorisation is requested per group. The list below follows the categorisation included in the annexes of Directive 2000/29/EC.

Annex IAI**(a) Insects, mites and nematodes, at all stages of their development**

Group of Cicadellidae (non-EU) known to be vector of Pierce's disease (caused by *Xylella fastidiosa*), such as:

- | | |
|--|---|
| 1) <i>Carneocephala fulgida</i> Nottingham | 3) <i>Graphocephala atropunctata</i> (Signoret) |
| 2) <i>Draeculacephala minerva</i> Ball | |

Group of Tephritidae (non-EU) such as:

- | | |
|--|---|
| 1) <i>Anastrepha fraterculus</i> (Wiedemann) | 12) <i>Pardalaspis cyanescens</i> Bezzi |
| 2) <i>Anastrepha ludens</i> (Loew) | 13) <i>Pardalaspis quinaria</i> Bezzi |
| 3) <i>Anastrepha obliqua</i> Macquart | 14) <i>Pterandrus rosa</i> (Karsch) |
| 4) <i>Anastrepha suspensa</i> (Loew) | 15) <i>Rhacochlaena japonica</i> Ito |
| 5) <i>Dacus ciliatus</i> Loew | 16) <i>Rhagoletis completa</i> Cresson |
| 6) <i>Dacus curcurbitae</i> Coquillett | 17) <i>Rhagoletis fausta</i> (Osten-Sacken) |
| 7) <i>Dacus dorsalis</i> Hendel | 18) <i>Rhagoletis indifferens</i> Curran |
| 8) <i>Dacus tryoni</i> (Froggatt) | 19) <i>Rhagoletis mendax</i> Curran |
| 9) <i>Dacus tsuneonis</i> Miyake | 20) <i>Rhagoletis pomonella</i> Walsh |
| 10) <i>Dacus zonatus</i> Saund. | 21) <i>Rhagoletis suavis</i> (Loew) |
| 11) <i>Epochra canadensis</i> (Loew) | |

(c) Viruses and virus-like organisms

Group of potato viruses and virus-like organisms such as:

- | | |
|----------------------------------|--|
| 1) Andean potato latent virus | 4) Potato black ringspot virus |
| 2) Andean potato mottle virus | 5) Potato virus T |
| 3) Arracacha virus B, oca strain | 6) non-EU isolates of potato viruses A, M, S, V, X and Y (including Yo, Yn and Yc) and Potato leafroll virus |

Group of viruses and virus-like organisms of *Cydonia* Mill., *Fragaria* L., *Malus* Mill., *Prunus* L., *Pyrus* L., *Ribes* L., *Rubus* L. and *Vitis* L., such as:

- | | |
|--------------------------------------|--|
| 1) Blueberry leaf mottle virus | 8) Peach yellows mycoplasma |
| 2) Cherry rasp leaf virus (American) | 9) Plum line pattern virus (American) |
| 3) Peach mosaic virus (American) | 10) Raspberry leaf curl virus (American) |
| 4) Peach phony rickettsia | 11) Strawberry witches' broom mycoplasma |
| 5) Peach rosette mosaic virus | 12) Non-EU viruses and virus-like organisms of <i>Cydonia</i> Mill., <i>Fragaria</i> L., <i>Malus</i> Mill., <i>Prunus</i> L., <i>Pyrus</i> L., <i>Ribes</i> L., <i>Rubus</i> L. and <i>Vitis</i> L. |
| 6) Peach rosette mycoplasma | |
| 7) Peach X-disease mycoplasma | |

Annex IIAI

(a) Insects, mites and nematodes, at all stages of their development

Group of *Margarodes* (non-EU species) such as:

- 1) *Margarodes vitis* (Phillipi)
- 2) *Margarodes vredendalensis* de Klerk
- 3) *Margarodes prieskaensis* Jakubski

1.1.2.3. Terms of Reference: Appendix 3

List of harmful organisms for which pest categorisation is requested. The list below follows the annexes of Directive 2000/29/EC.

Annex IAI

(a) Insects, mites and nematodes, at all stages of their development

<i>Acleris</i> spp. (non-EU)	<i>Longidorus diadecturus</i> Eveleigh and Allen
<i>Amauromyza maculosa</i> (Malloch)	<i>Monochamus</i> spp. (non-EU)
<i>Anomala orientalis</i> Waterhouse	<i>Myndus crudus</i> Van Duzee
<i>Arrhenodes minutus</i> Drury	<i>Nacobbus aberrans</i> (Thorne) Thorne and Allen
<i>Choristoneura</i> spp. (non-EU)	<i>Naupactus leucoloma</i> Boheman
<i>Conotrachelus nenuphar</i> (Herbst)	<i>Premnotrypes</i> spp. (non-EU)
<i>Dendrolimus sibiricus</i> Tschetverikov	<i>Pseudopityophthorus minutissimus</i> (Zimmermann)
<i>Diabrotica barberi</i> Smith and Lawrence	<i>Pseudopityophthorus pruinus</i> (Eichhoff)
<i>Diabrotica undecimpunctata howardi</i> Barber	<i>Scaphoideus luteolus</i> (Van Duzee)
<i>Diabrotica undecimpunctata undecimpunctata</i> Mannerheim	<i>Spodoptera eridania</i> (Cramer)
<i>Diabrotica virgifera zea</i> Krysan & Smith	<i>Spodoptera frugiperda</i> (Smith)
<i>Diaphorina citri</i> Kuway	<i>Spodoptera litura</i> (Fabricus)
<i>Heliothis zea</i> (Boddie)	<i>Thrips palmi</i> Karny
<i>Hirschmanniella</i> spp., other than	<i>Xiphinema americanum</i> Cobb sensu lato (non-EU populations)
<i>Hirschmanniella gracilis</i> (de Man) Luc and Goodey	<i>Xiphinema californicum</i> Lamberti and Bleve-Zacheo
<i>Liriomyza sativae</i> Blanchard	

(b) Fungi

<i>Ceratocystis fagacearum</i> (Bretz) Hunt	<i>Mycosphaerella larici-leptolepis</i> Ito et al.
<i>Chrysomyxa arctostaphyli</i> Dietel	<i>Mycosphaerella populorum</i> G. E. Thompson
<i>Cronartium</i> spp. (non-EU)	<i>Phoma andina</i> Turkensteen
<i>Endocronartium</i> spp. (non-EU)	<i>Phyllosticta solitaria</i> Ell. and Ev.
<i>Guignardia laricina</i> (Saw.) Yamamoto and Ito	<i>Septoria lycopersici</i> Speg. var. <i>malagutii</i> Ciccarone and Boerema
<i>Gymnosporangium</i> spp. (non-EU)	<i>Thecaphora solani</i> Barrus
<i>Inonotus weirii</i> (Murril) Kotlaba and Pouzar	<i>Trechispora brinkmannii</i> (Bresad.) Rogers
<i>Melampsora farlowii</i> (Arthur) Davis	

(c) Viruses and virus-like organisms

Tobacco ringspot virus	Pepper mild tigré virus
Tomato ringspot virus	Squash leaf curl virus
Bean golden mosaic virus	Euphorbia mosaic virus
Cowpea mild mottle virus	Florida tomato virus
Lettuce infectious yellows virus	

(d) Parasitic plants

Arceuthobium spp. (non-EU)

Annex I A I I

(a) Insects, mites and nematodes, at all stages of their development

Meloidogyne fallax Karssen

Rhizoecus hibisci Kawai and Takagi

Popillia japonica Newman

(b) Bacteria

Clavibacter michiganensis (Smith) Davis et al.
ssp. *sepedonicus* (Spieckermann and Kotthoff)
Davis et al.

Ralstonia solanacearum (Smith) Yabuuchi et al.

(c) Fungi

Melampsora medusae Thümen

Synchytrium endobioticum (Schilbersky) Percival

Annex I B

(a) Insects, mites and nematodes, at all stages of their development

Leptinotarsa decemlineata Say

Liriomyza bryoniae (Kaltenbach)

(b) Viruses and virus-like organisms

Beet necrotic yellow vein virus

1.2. Interpretation of the Terms of Reference

Ips sexdentatus is one of a number of pests listed in the Appendices to the Terms of Reference (ToR) to be subject to pest categorisation to determine whether it fulfils the criteria of a quarantine pest or those of a regulated non-quarantine pest (RNQP) for the area of the European Union (EU) excluding Ceuta, Melilla and the outermost regions of Member States (MSs) referred to in Article 355(1) of the Treaty on the Functioning of the European Union (TFEU), other than Madeira and the Azores.

Since *I. sexdentatus* is regulated in the protected zones (PZs) only, the scope of the categorisation is the territory of the PZ (Ireland, Cyprus and the United Kingdom: Northern Ireland, Isle of Man); thus, the criteria refer to the PZ instead of the EU territory.

2. Data and methodologies

2.1. Data

2.1.1. Literature search

A literature search on *I. sexdentatus* was conducted at the beginning of the categorisation in the ISI Web of Science bibliographic database, using the scientific name of the pest as search term. Relevant papers were reviewed and further references and information were obtained from experts as well as from citations within the references and grey literature.

2.1.2. Database search

Pest information, on host(s) and distribution, was retrieved from the European and Mediterranean Plant Protection Organization (EPPO) Global Database (EPPO, 2017) and relevant publications.

Data about import of commodity types that could potentially provide a pathway for the pest to enter the EU were obtained from Eurostat (Statistical Office of the European Communities).

The Europhyt database was consulted for pest-specific notifications on interceptions and outbreaks. Europhyt is a web-based network launched by the Directorate General for Health and Consumers (DG SANCO) and is a subproject of PHYSAN (Phyto-Sanitary Controls) specifically concerned with plant health information. The Europhyt database manages notifications of interceptions of plants or plant products that do not comply with EU legislation as well as notifications of plant pests detected in the territory of the MS and the phytosanitary measures taken to eradicate or avoid their spread.

2.2. Methodologies

The Panel performed the pest categorisation for *I. sexdentatus*, following guiding principles and steps presented in the EFSA guidance on the harmonised framework for pest risk assessment (EFSA

PLH Panel, 2010a) and as defined in the International Standard for Phytosanitary Measures No 11 (FAO,2013) and No 21 (FAO, 2004).

In accordance with the guidance on a harmonised framework for pest risk assessment in the EU (EFSA PLH Panel, 2010a), this work was initiated following an evaluation of the EU's plant health regime. Therefore, to facilitate the decision-making process, in the conclusions of the pest categorisation, the Panel addresses explicitly each criterion for a Union quarantine pest and for a Union RNQP in accordance with Regulation (EU) 2016/2031 on protective measures against pests of plants and includes additional information required in accordance with the specific ToR received by the European Commission. In addition, for each conclusion, the Panel provides a short description of its associated uncertainty.

Table 1 presents the Regulation (EU) 2016/2031 pest categorisation criteria on which the Panel bases its conclusions. All relevant criteria have to be met for the pest to potentially qualify either as a quarantine pest or as a RNQP. If one of the criteria is not met, the pest will not qualify. Note that a pest that does not qualify as a quarantine pest may still qualify as a RNQP that needs to be addressed in the opinion. For the pests regulated in the PZs only, the scope of the categorisation is the territory of the PZ; thus, the criteria refer to the PZ instead of the EU territory.

It should be noted that the Panel's conclusions are formulated respecting its remit and particularly with regard to the principle of separation between risk assessment and risk management (EFSA founding regulation (EU) No 178/2002); therefore, instead of determining whether the pest is likely to have an unacceptable impact, the Panel will present a summary of the observed pest impacts. Economic impacts are expressed in terms of yield and quality losses and not in monetary terms, whereas addressing social impacts is outside the remit of the Panel, in agreement with EFSA guidance on a harmonised framework for pest risk assessment (EFSA PLH Panel, 2010a).

Table 1: Pest categorisation criteria under evaluation, as defined in Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column)

Criterion of pest categorisation	Criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest	Criterion in Regulation (EU) 2016/2031 regarding protected zone quarantine pest (articles 32–35)	Criterion in Regulation (EU) 2016/2031 regarding Union regulated non-quarantine pest
Identity of the pest (Section 3.1)	Is the identity of the pest established, or has it been shown to produce consistent symptoms and to be transmissible?	Is the identity of the pest established, or has it been shown to produce consistent symptoms and to be transmissible?	Is the identity of the pest established, or has it been shown to produce consistent symptoms and to be transmissible?
Absence/ presence of the pest in the EU territory (Section 3.2)	Is the pest present in the EU territory? If present, is the pest widely distributed within the EU? Describe the pest distribution briefly!	Is the pest present in the EU territory? If not, it cannot be a protected zone quarantine organism.	Is the pest present in the EU territory? If not, it cannot be a regulated non-quarantine pest. (A regulated non-quarantine pest must be present in the risk assessment area).
Regulatory status (Section 3.3)	If the pest is present in the EU but not widely distributed in the risk assessment area, it should be under official control or expected to be under official control in the near future.	The protected zone system aligns with the pest-free area system under the International Plant Protection Convention (IPPC). The pest satisfies the IPPC definition of a quarantine pest that is not present in the risk assessment area (i.e. protected zone).	Is the pest regulated as a quarantine pest? If currently regulated as a quarantine pest, are there grounds to consider its status could be revoked?
Pest potential for entry, establishment and spread in the EU territory (Section 3.4)	Is the pest able to enter into, become established in and spread within the EU territory? If yes, briefly list the pathways!	Is the pest able to enter into, become established in and spread within the protected zone areas? Is entry by natural spread from EU areas where the pest is present possible?	Is spread mainly via specific plants for planting, rather than via natural spread or via movement of plant products or other objects? Clearly state if plants for planting is the main pathway!

Criterion of pest categorisation	Criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest	Criterion in Regulation (EU) 2016/2031 regarding protected zone quarantine pest (articles 32–35)	Criterion in Regulation (EU) 2016/2031 regarding Union regulated non-quarantine pest
Potential for consequences in the EU territory (Section 3.5)	Would the pests' introduction have an economic or environmental impact on the EU territory?	Would the pests' introduction have an economic or environmental impact on the protected zone areas?	Does the presence of the pest on plants for planting have an economic impact, as regards the intended use of those plants for planting?
Available measures (Section 3.6)	Are there measures available to prevent the entry into, establishment within or spread of the pest within the EU such that the risk becomes mitigated?	Are there measures available to prevent the entry into, establishment within or spread of the pest within the protected zone areas such that the risk becomes mitigated? Is it possible to eradicate the pest in a restricted area within 24 months (or a period longer than 24 months where the biology of the organism so justifies) after the presence of the pest was confirmed in the protected zone?	Are there measures available to prevent pest presence on plants for planting such that the risk becomes mitigated?
Conclusion of pest categorisation (Section 4)	A statement as to whether (1) all criteria assessed by EFSA above for consideration as a potential quarantine pest were met and (2) if not, which one(s) were not met.	A statement as to whether (1) all criteria assessed by EFSA above for consideration as potential protected zone quarantine pest were met, and (2) if not, which one(s) were not met.	A statement as to whether (1) all criteria assessed by EFSA above for consideration as a potential regulated non-quarantine pest were met, and (2) if not, which one(s) were not met.

The Panel will not indicate in its conclusions of the pest categorisation whether to continue the risk assessment process, but, following the agreed two-step approach, will continue only if requested by the risk managers. However, during the categorisation process, experts may identify key elements and knowledge gaps that could contribute significant uncertainty to a future assessment of risk. It would be useful to identify and highlight such gaps so that potential future requests can specifically target the major elements of uncertainty, perhaps suggesting specific scenarios to examine.

3. Pest categorisation

3.1. Identity and biology of the pest

3.1.1. Identity and taxonomy

Is the identity of the pest established, or has it been shown to produce consistent symptoms and to be transmissible?

Yes, the identity of the pest is established. It can be identified at species level using conventional entomological keys.

I. sexdentatus is an insect of the family Curculionidae, subfamily Scolytinae.⁴

⁴ Although the leading taxonomists in the 2000s (Wood, 1982; Bright and Skidmore, 2002) still considered the Scolytidae to be a family distinct from the Curculionidae according to morphological criteria, modern phylogenetics supports the position of scolytine beetles (subfamily Scolytinae) within the family Curculionidae (Knížek and Beaver, 2004; Hulcr et al., 2015). This is reflected by the growing number of citations in Scopus (2017) referring to Scolytinae (18 in 1990 vs 177 in 2016), as opposed to citations referring to Scolytidae (50 in 1990 vs 15 in 2016). The Scolytinae includes two subcategories, the 'bark beetles' which live in the phloem and the 'ambrosia beetles' which live in the sapwood.

3.1.2. Biology of the pest

A general description of the biology and ecology of *I. sexdentatus* is provided by Chararas (1962), Bakke (1968) and Léviex et al. (1985). The adults overwinter in the bark of their hosts or in the litter and disperse in the spring, flying in search for new hosts, sometimes over large distances. In flight mill experiments, Jactel and Gaillard (1991) observed that, out of a sample of 38 individuals, 98% flew more than 5 km, 50% more than 20 km and 10% more than 45 km. In Europe, *I. sexdentatus* preferentially colonise weakened pines, such as cut logs or wind-felled trees (Samalens et al., 2007; Rossi et al., 2009), trees affected by forest fires (Fernandez and Salgado Costas, 1999; Fernandez, 2006; Lombardero and Ayres, 2011; Santolamazza-Carbone et al., 2011) or drought-stressed trees (Lieutier et al., 1988). However, they can also attack living trees when population levels are high (Rossi et al., 2009; Pineau et al., 2017). In Turkey and Georgia, *I. sexdentatus* is a major primary pest of *Picea orientalis*, attacking living trees (Schimitschek, 1939; Lozovoj, 1966; Ozcan et al., 2011). The males arrive first, start a mating chamber and emit pheromones that attract females as well as other males (Vité et al., 1974; Francke et al., 1986; Kohnle et al., 1992; Etxebeste et al., 2012). After having excavated a nuptial chamber in the phloem, each male is joined by one to five females, which bore each a maternal gallery in the phloem, parallel to the fibres. Single eggs are laid at regular intervals along these galleries. After egg laying, the parent adults often re-emerge and establish sister broods on the same tree or on a new host. Each larva excavates an individual gallery perpendicular to the maternal gallery. Pupation occurs in a small niche in the phloem, at the end of the larval gallery. Productivity varies between 1 and 60 offspring per female and is inversely dependent upon attack density (Pineau et al., 2017). After metamorphosis, the young adults remain under the bark for maturation feeding before they disperse. Upon emergence, the sex ratio is balanced (Pineau et al., 2017). There are possibly one to five generations per year (Léviex et al., 1985). Pathogenic ophiostomatoid fungi are carried by the beetles, some of them in pit mycangia on the body (Léviex et al., 1991) and are inoculated to the host (Kirisits, 2004; Romón et al., 2008; Bueno et al., 2010; Jankowiak, 2012). They cause blue staining of the wood and some of them can contribute to tree death.

3.1.3. Detection and identification of the pest

Are detection and identification methods available for the pest?

Yes, the organism can be detected by visual searching, often after damage symptoms are seen, and by pheromone trapping. The species can be identified by examining morphological features, for which taxonomic keys exist, e.g. Balachowsky (1949); Grüne (1979); Schedl (1981); Wood (1982).

The standing trees attacked by *I. sexdentatus* die during the colonisation process, with an obvious discolouration of their crown, which becomes brown and then grey after the needles have shed. During the attacks, brown sawdust is expelled from the entry holes and, when the broods have metamorphosed and the young adults start feeding on the phloem around the galleries, the bark can flake off. This phenomenon can be amplified by the action of woodpeckers. Within and under the phloem, maternal galleries, parallel to the fibres and up to 50 cm long, and transversal larval galleries can be seen. Pheromone lures and traps are commercially available for *I. sexdentatus* but, because of the large dispersal capacity of the pest, trap catches do not necessarily reflect local establishment. The sapwood shows blue staining due to the fungi introduced by the beetles. The adults are dark brown or black in colour, cylindrical, 7–8 mm long. The larvae are apodous, with a dark amber cephalic capsule.

3.2. Pest distribution

3.2.1. Pest distribution outside the EU

I. sexdentatus is present in Europe and Asia. In non-EU European countries, the insect has been reported from Armenia, Azerbaijan, Belarus, Georgia, Macedonia, Moldova, Norway, Russia, Serbia, Switzerland, Turkey and Ukraine (Figure 1).




Figure 1: Global distribution map for *Ips sexdentatus* (extracted from the EPPO global database accessed on 22 August 2017)

3.2.2. Pest distribution in the EU

Is the pest present in the EU territory? If present, is the pest widely distributed within the EU?

Yes, *I. sexdentatus* is present and widely distributed in the EU, it has been reported from 24 MSs (Table 2). The pest is absent in the protected zones (Ireland, Cyprus and the United Kingdom: Northern Ireland and the Isle of Man).

Table 2: Current distribution of *Ips sexdentatus* in the 28 EU MS based on information from the EPPO Global Database

Country	EPPO Global Database Last updated: 12 July 2017 Date Accessed: 22 August 2017
Austria	Present, no details
Belgium	Present, no details
Bulgaria	Present, widespread
Croatia	Present, restricted distribution
Cyprus	No information ^(a)
Czech Republic	Present, restricted distribution
Denmark	Absent, intercepted only
Estonia	Present, no details
Finland	Present, restricted distribution
France	Present, restricted distribution
Germany	Present, widespread
Greece	Present, no details
Hungary	Present, restricted distribution
Ireland	Absent, confirmed by survey
Italy	Present, widespread Sardinia: Present, no details Sicily: Present, no details
Latvia	Present, no details
Lithuania	Present, restricted distribution

Country	EPPO Global Database Last updated: 12 July 2017 Date Accessed: 22 August 2017
Luxembourg	Present, no details
Malta	No information
Poland	Present, widespread
Portugal	Present, no details
Romania	Present, no details
Slovakia	Present, restricted distribution
Slovenia	Present, restricted distribution
Spain	Present, widespread
Sweden	Present, restricted distribution
The Netherlands	Present, no details
United Kingdom	Present, restricted distribution England: Present, restricted distribution Northern Ireland: Absent, confirmed by survey

(a): Cyprus is a protected zone, and therefore, regular surveys are carried out to confirm absence/presence.

3.3. Regulatory status

3.3.1. Council Directive 2000/29/EC

I. sexdentatus is listed in Council Directive 2000/29/EC. Details are presented in Tables 3 and 4.

Table 3: *Ips sexdentatus* in Council Directive 2000/29/EC

Annex II, Part B	Harmful organisms whose introduction into, and whose spread within, certain protected zones shall be banned if they are present on certain plants or plant products		
(a)	Insects, mites and nematodes, at all stages of their development		
	Species	Subject of contamination	Protected zones
6 (d)	<i>Ips sexdentatus</i>	Plants of <i>Abies</i> Mill., <i>Larix</i> Mill., <i>Picea</i> A. Dietr., <i>Pinus</i> L. over 3 m in height, other than fruit and seeds, wood of conifers (<i>Coniferales</i>) with bark, isolated bark of conifers	IRL, CY, UK (Northern Ireland, Isle of Man)

3.3.2. Legislation addressing plants and plant parts on which *Ips sexdentatus* is regulated

Table 4: Regulated hosts and commodities that may involve *Ips sexdentatus* in Annexes III, IV and V of Council Directive 2000/29/EC

Annex III, Part A	Plants, plant products and other objects the introduction of which shall be prohibited in all Member States		
	Description		Country of origin
1.	Plants of <i>Abies</i> Mill., [...], <i>Larix</i> Mill., <i>Picea</i> A. Dietr., <i>Pinus</i> L., [...], other than fruit and seeds		Non-European Countries
Annex IV, Part B	Special requirements which shall be laid down by all member states for the introduction and movement of plants, plant products and other objects into and within certain protected zones		
	Plants, plant products and other objects	Special requirements	Protected zone(s)

6.	Wood of conifers (<i>Coniferales</i>)	Without prejudice to the requirements applicable to the wood listed in Annex IV(A)(I)(1.1), (1.2), (1.3), (1.4), (1.5), (1.6), (1.7), where appropriate, and Annex IV(B)(1), (2), (3), (4), (5): the wood shall be stripped of its bark; or official statement that the wood originates in areas known to be free from <i>Ips sexdentatus</i> or there shall be evidence by a mark .Kilndried., .KD. or another internationally recognised mark, put on the wood or on its packaging in accordance with current commercial usage, that it has undergone kiln-drying to below 20% moisture content, expressed as a percentage of dry matter, at time of manufacture, achieved through an appropriate time/temperature schedule.	IRL, CY, UK (Northern Ireland, Isle of Man)
12.	Plants of <i>Abies</i> Mill., <i>Larix</i> Mill., <i>Picea</i> A. Dietr. and <i>Pinus</i> L. over 3 m in height, other than fruit and seeds	Without prejudice to the provisions applicable to the plants listed in Annex III(A)(1), Annex IV(A)(I)(8.1), (8.2), (9), (10), Annex IV(A)(II)(4), (5), and Annex IV(B)(7), (8), (9), (10), (11), where appropriate, official statement that the place of production is free from <i>Ips sexdentatus</i>	IRL, CY, UK (Northern Ireland, Isle of Man)
14.5	Isolated bark of conifers (<i>Coniferales</i>)	Without prejudice to the provisions applicable to the bark listed in Annex IV(B)(14.1), (14.2), (14.3), (14.4), official statement that the consignment: has been subjected to fumigation or other appropriate treatments against bark beetles; or originates in areas known to be free from <i>Ips sexdentatus</i>	IRL, CY, UK (Northern Ireland, Isle of Man)
Annex V	Plants, plant products and other objects which must be subject to a plant health inspection (at the place of production if originating in the Community, before being moved within the Community—in the country of origin or the consignor country, if originating outside the Community) before being permitted to enter the Community		
Part A	Plants, plant products and other objects originating in the Community		
Section II	Plants, plant products and other objects produced by producers whose production and sale is authorised to persons professionally engaged in plant production, other than those plants, plant products and other objects which are prepared and ready for sale to the final consumer, and for which it is ensured by the responsible official bodies of the Member States, that the production thereof is clearly separate from that of other products		
2.1	Plants intended for planting other than seeds of the genera <i>Abies</i> Mill., [...], <i>Larix</i> Mill., [...], <i>Picea</i> A. Dietr., <i>Pinus</i> L., [...]		

3.3.3. Legislation addressing the organisms vectored by *Ips sexdentatus* (Directive 2000/29/EC)

Although several phytopathogenic ophiostomatoid fungi are regularly associated with *I. sexdentatus*, (Kirisits, 2004; Romón et al., 2007, 2008; Bueno et al., 2010; Jankowiak, 2012), there is currently no legislation addressing this issue. However, the pest has been also found associated with the pitch canker fungus, *Fusarium circinatum* (*Gibberella circinata*) Nirenberg and O'Donnell (Romón

et al., 2008). *F. circinatum* is a quarantine organism in the EU (Commission Decision 2007/433/EC; EFSA PLH Panel, 2010b).

3.4. Entry, establishment and spread in the EU

3.4.1. Host range

Ips sexdentatus attacks mainly pines. It has been reported on the following species: *Pinus brutia* (Agbaba and Celepirovic, 2008); *Pinus halepensis* (Agbaba and Celepirovic, 2008); *Pinus heldreichii* (Ivojinovi, 1960); *Pinus koraiensis* (Arefin, 1983); *Pinus leucodermis* (Frisullo et al., 2003); *Pinus nigra* (Kondur et al., 2012); *Pinus nigra* subsp. *salzmannii* (Etxebeste and Pajares, 2011); *Pinus pallasiana* (Kondur et al., 2012); *Pinus peuce* (Ivojinovi, 1960); *Pinus pinaster* (Bueno et al., 2010); *Pinus pithyusa* (Lozovoi, 1961); *Pinus radiata* (Cobos Suarez and Ruiz Urrestarazu, 1990); *Pinus sibirica* (Kobzar, 1968); *Pinus strobus* (Beffa, 2006); *Pinus sylvestris* (Croisé et al., 1998); *P. sylvestris* var. *mongolica* (Wang et al., 2011); *Pinus tabulaeformis* (Wang et al., 2011). Spruce (e.g. *P. orientalis* (Besceli and Ekici, 1969); *Picea abies* (Slankis, 1969) and *Picea schrenkiana* (Ismukhambetov, 1964)) are also attacked. Chararas (1962) mentions attacks on *Larix decidua* and *Abies* sp.

The hosts for which *I. sexdentatus* is regulated are comprehensive of the host range: the pest is regulated on four genera: *Abies*, *Larix*, *Picea* and *Pinus*.

3.4.2. Entry

Is the pest able to enter into the protected zone areas of the EU territory? If yes, identify and list the pathways.

Yes, the pest is already established in 24 MSs and can enter the protected zones by human assisted spread or by natural spread from EU areas where the pest is present.

The main pathways of entry are:

- Wood of *Abies*, *Larix*, *Picea* and *Pinus* from countries where the pest occurs
- Wood chips of conifers from countries where the pest occurs
- Bark of conifers from countries where the pest occurs
- Wood packaging material and dunnage from countries where the pest occurs

Ips species are regularly intercepted on wood, wood packaging material and dunnage. During the period 1985–2000, among the 2.740 Scolytinae intercepted at the US ports of entry and identified to species, 157 *I. sexdentatus* were found (Haack, 2001). In the Europhyt database, there are in total 66 records of *Ips* species (1994–2017), all on coniferous wood or packaging material. For *I. sexdentatus*, there are two records of interception, one from Bulgaria on coniferous wood and one from Ukraine on *P. sylvestris* wood. Lopez and Goldarazena (2012) report the introduction of *I. sexdentatus* from France to Spain with transport of wind thrown *P. pinaster* after a major storm in 2009.

There are no records of interception that indicate that plants for planting can be a pathway for *I. sexdentatus*. Plants for planting are not considered a pathway for *I. sexdentatus* since young plants are not attacked by the pest.

According to the Eurostat database, there is trade of wood from third countries and EU countries where the pest is present, into the PZs (Table 5). It should be noted that for the PZs in the United Kingdom (Northern Ireland and Isle of Man), no specific data were available and the actual imports into these PZs are a fraction of the volume imported into the United Kingdom.

Table 5: Import of wood (in tonnes) originating from Third countries and EU countries where the pest is present into protected zones. Eurostat data, period 2011–2015, GN codes: 44032010, 44032011, 44032019, 44032030, 44032031, 4403239, 44032090, 44032091, 44032099.

Protected zone	3rd Countries	EU Countries
Cyprus	28	–
United Kingdom	3.001	1.326.402
Ireland	1.662	836.992
Total	4.690	2.163.394


3.4.3. Establishment

Is the pest able to become established in the protected zone areas of the EU territory?

Yes, the pest is already established in 24 MS. The climate of the EU Protected Zones is similar to that of the MS where *I. sexdentatus* is established, and the pest's main host plants are present (Figure 2)

3.4.3.1. EU distribution of main host plants

The wide distribution of host trees in the EU territory allowed *I. sexdentatus* to establish in most MSs (Table 2) (Figure 2A,B).



A) Distribution map of the genus *Pinus* in the European Union territory (based on data from the species: *P. sylvestris*, *P. pinaster*, *P. halepensis*, *P. nigra*, *P. pinea*, *P. contorta*, *P. cembra*, *P. mugo*, *P. radiata*, *P. canariensis*, *P. strobus*, *P. brutia*, *P. banksiana*, *P. ponderosa*, *P. heldreichii*, *P. leucodermis*, *P. wallichiana*) B) Distribution map of the genus *Picea* in the European Union territory (based on data from the species: *P. abies*, *P. sitchensis*, *P. glauca*, *P. engelmannii*, *P. pungens*, *P. omorika*, *P. orientalis*).

Figure 2: Left panel: Relative probability of presence (RPP) of the genera *Pinus* and *Picea* in Europe, mapped at 100 km² resolution. The underlying data are from European-wide forest monitoring data sets and from national forestry inventories based on standard observation plots measuring in the order of hundreds m². RPP represents the probability of finding at least one individual of the taxon in a standard plot placed randomly within the grid cell. For details, see Appendix A (courtesy of JRC, 2017). Right panel: Trustability of RPP. This metric expresses the strength of the underlying information in each grid cell and varies according to the spatial variability in forestry inventories. The colour scale of the trustability map is obtained by plotting the cumulative probabilities (0–1) of the underlying index (for details see Appendix A)

3.4.3.2. Climatic conditions affecting establishment

Given the current distribution of *I. sexdentatus*, the whole EU area (including PZs) is suitable for establishment.

3.4.4. Spread

Is the pest able to spread within the protected zone areas of the EU territory following establishment? How?

Yes, adults can disperse naturally. They can fly over tens of kilometres or even more (Jactel, 1991; Jactel and Gaillard, 1991). The pest can also spread by human assistance, e.g. with the transportation of wood, wood chips, bark, wood packaging material and dunnage of conifers.

RNQPs: Is spread mainly via specific plants for planting, rather than via natural spread or via movement of plant products or other objects?

No, plants for planting are not a pathway (see Section 3.4.2).

3.5. Impacts

Would the pests' introduction have an economic or environmental impact on the protected zone areas of the EU territory?

Yes, the pest is known to have killed thousands of trees, after triggering events such as storms, forest fires or dry summers.

This species is mostly a secondary pest, which generally uses stumps, fallen trees and large branches as host material. It will attack trees alone or together with other bark beetles such as *Tomicus piniperda* (Bouhot et al., 1988). Specific economic loss is not known for *I. sexdentatus*. However, approximately one million *P. orientalis* trees were lost due to sporadic *I. sexdentatus* infestations in Turkey (Schimitschek, 1939; Besceli and Ekici, 1969; Schönherr et al., 1983).

Kirisits (2004), Romón et al. (2007, 2008), Bueno et al. (2010) and Jankowiak (2012) report finding the following ophiostomatoid species either on the body or in the galleries of *I. sexdentatus*:

Ambrosiella ips; Ambrosiella tingens; Ceratocystiopsis minuta; Graphium pseudormiticum; Graphium sp.; Leptographium cf. truncatum; Leptographium guttulatum; Leptographium procerum; Leptographium sp.; Ophiostoma ainoae (= Ophiostoma brunneo-ciliatum?); Ophiostoma araucariae; Ophiostoma brunneo-ciliatum; Ophiostoma cf. abietinum; Ophiostoma cf. rectangulosporium; Ophiostoma clavatum; Ophiostoma floccosum; Ophiostoma ips; Ophiostoma japonicum (= Ophiostoma arborea?); Ophiostoma minus; Ophiostoma obscura; Ophiostoma olivaceum; Ophiostoma piceae; Ophiostoma piceaperdum; Ophiostoma pluriannulatum; Ophiostoma quercus; Ophiostoma rectangulosporium-like; Ophiostoma stenoceras; Ophiostoma sp.; Pesotum fragrans; Sporothrix 12.

Jankowiak (2012) found that *Leptographium cf. truncatum* and *O. minus* were very virulent and considered them as serious pine pathogens.

It is important to note that Romón et al. (2008) found *F. circinatum* (8.6%; n = 35 beetles) and *Fusarium verticillioides* (2.9%) associated with *I. sexdentatus* in Spain. *F. circinatum* is a quarantine pathogen in the EU.

3.6. Availability and limits of mitigation measures

Are there measures available to prevent the entry into, establishment within or spread of the pest within the protected zone areas such that the risk becomes mitigated?

Yes, in isolated areas (e.g. islands) that cannot be reached by natural spread, measures can be put in place to prevent the introduction with wood and bark. Debarking wood and heat treatment of wood, bark and chips are effective as specified in annex IVB of 2000/29/EC. When such geographical barriers do not exist, the pest will eventually be able to enter new territories by natural dispersal.

Is it possible to eradicate the pest in a restricted area within 24 months after the presence of the pest was confirmed in the PZ?

Yes. Eradication is possible as the pest is mainly attacking fallen or weakened trees in the EU territory. Provided incipient populations are localised very early (i.e. preferably before the new brood has emerged), the attacked material can be removed and destroyed. However, eradication is difficult because all suitable host material (fallen or weakened trees) in the surrounding area within a radius of several kilometres should be localised and removed.

3.6.1. Biological or technical factors limiting the feasibility and effectiveness of measures to prevent the entry, establishment and spread of the pest

- Quarantine measures are not fully effective. Despite quarantine regulations bearing on round wood, wood packaging material and wood products, the pest is regularly intercepted at ports.
- It is difficult to successfully eradicate the pest from forest areas after an introduction. All infested trees and tree parts (including pieces of fallen or broken material) have to be detected and removed within a suitable radius of several kilometres.
- Silvicultural control is not fully effective. In areas where it is established, the pest continues to develop outbreaks whenever climatic conditions are favourable.

3.6.2. Control methods

- Monitoring methods by roadside sampling techniques, focusing on log pile storage areas if any, have been developed and tested by Samalens et al. (2007).
- Silvicultural practices are the usual control methods. They include sanitation thinning and clear-felling with rapid removal of the infested material (Stadelmann et al., 2013; Fettig and Hilszczanski, 2015; Grégoire et al., 2015).
- Pheromone mass-trapping is attempted in several countries, such as in Turkey (Ozcan et al., 2011) and in Spain (Etxebeste et al., 2012), but it is not a generalised control method.
- Verbenone and non-host volatiles have been successfully used experimentally to protect pines or pine logs (Jactel et al., 2001; Etxebeste and Pajares, 2011; Etxebeste et al., 2013).

3.7. Uncertainty

I. sexdentatus is a secondary pest in the EU, causing limited damage on pine. It appears more aggressive on *P. orientalis* in Turkey and Georgia, but does not aggressively attack *P. abies* in the EU. The reasons for these differences in aggressiveness are unknown and could constitute an increased risk if, for example, they correspond to intraspecific variations in aggressiveness or host range within the species.

4. Conclusions

Table 6: The Panel's conclusions on the pest categorisation criteria defined in Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column)

Criterion of pest categorisation	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding protected zone quarantine pest (articles 32–35)	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding Union regulated non-quarantine pest	Key uncertainties
Identity of the pest (Section 3.1)	The identity of the pest is established. It can be identified to the species level using conventional entomological keys.	The identity of the pest is established. It can be identified to the species level using conventional entomological keys.	None
Absence/ presence of the pest in the EU territory (Section 3.2)	<i>I. sexdentatus</i> is present and widely distributed in the EU; it has been reported from 24 EU MS. The protected zones, Ireland, Cyprus ^(a) and the United Kingdom (Northern Ireland and the Isle of Man), are free from the pest.	<i>I. sexdentatus</i> is present and widely distributed in the EU; it has been reported from 24 EU MS. The protected zones, Ireland, Cyprus and the United Kingdom (Northern Ireland and the Isle of Man), are free from the pest.	None
Regulatory status (Section 3.3)	The pest is currently officially regulated by 2000/29/EC on plants of <i>Abies</i> , <i>Larix</i> , <i>Picea</i> and <i>Pinus</i> over 3 m in height, other than fruit and seeds, wood of conifers (Coniferales) with bark, isolated bark of conifers. <i>I. sexdentatus</i> is regulated as a quarantine pest in protected zones (Annex IIB): Ireland, Cyprus and the United Kingdom (Northern Ireland, Isle of Man).	The pest is currently officially regulated by 2000/29/EC on plants of <i>Abies</i> , <i>Larix</i> , <i>Picea</i> , <i>Pinus</i> over 3 m in height, other than fruit and seeds, wood of conifers (Coniferales) with bark, isolated bark of conifers. <i>I. sexdentatus</i> is regulated as a quarantine pest in protected zones (Annex IIB): Ireland, Cyprus and the United Kingdom (Northern Ireland, Isle of Man).	Although the pest is regulated on <i>Abies</i> and <i>Larix</i> spp., there is no scientific evidence in the literature, apart from a brief mention by Chararas (1962), that <i>Abies</i> spp. and <i>Larix</i> spp. are hosts for <i>I. sexdentatus</i> .
Pest potential for entry, establishment and spread in the EU territory (Section 3.4)	<u>Entry</u> : the pest is established in 24 MS. Since entry by natural spread from EU areas where the pest is present is possible, only isolated areas (e.g. islands) can be long-term protected zones. <u>Establishment</u> : the climate of the EU protected zones is similar to that of MSs where <i>I. sexdentatus</i> is established, and the pest's main host plants are present. <u>Spread</u> : adults can disperse naturally. They can fly over tens of kilometres. The pest can also spread by human assistance, e.g. with the transportation of wood, wood chips, bark, wood packaging material and dunnage of conifers.	Plants for planting are not a pathway for the spread of <i>I. sexdentatus</i> .	None

Criterion of pest categorisation	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding protected zone quarantine pest (articles 32–35)	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding Union regulated non-quarantine pest	Key uncertainties
Potential for consequences in the EU territory (Section 3.5)	The pest is secondary, but is known to have killed thousands of trees after triggering events such as storms or dry summers.	Young trees are not attacked by <i>I. sexdentatus</i> ; therefore, impacts in nurseries are not expected.	None This is illustrated by the pest's past history in the EU.
Available measures (Section 3.6)	In isolated areas (e.g. islands) that cannot be reached by natural spread, measures can be put in place to prevent the introduction of the pest. For wood, wood products, wood chips and bark this can be achieved by debarking wood and heat treatment of wood, bark and chips. When such geographical barriers do not exist, there is no possibility to prevent the entry, establishment and spread of <i>I. sexdentatus</i> by natural dispersal.	Young plants are not attacked by <i>I. sexdentatus</i> .	Inspections of large shipments at entry are difficult to perform with complete accuracy.
Conclusion on pest categorisation (Section 4)	All criteria assessed by EFSA above for consideration as potential protected zone quarantine pest were met.	The criteria for considering <i>I. sexdentatus</i> as a potential regulated non-quarantine pest are not met since plants for planting are not a pathway.	See above
Aspects of assessment to focus on/ scenarios to address in future if appropriate	The difference of aggressiveness between the attacks in the EU (mostly secondary, on fallen or weak pines) and the more primary attacks (on standing, healthy Oriental spruce) in Turkey and Georgia raises the issue of possible intraspecific variations within the <i>I. sexdentatus</i> species, with potential quarantine implications.		

(a): Although there is no report on absence of the pest in Cyprus in the EPPO Global Database, it is assumed that Cyprus is free from the pest since it is a protected zone, and therefore, regular surveys are carried out to confirm their absence/presence

References

- Agbaba SN and Celepirovic N, 2008. Health condition of the forest vegetation on the island of Veliki Brijun, National Park Brijuni, Croatia. Radovi Sumarskog Fakulteta Univerziteta u Sarajevu, 38, 35–45.
- Arefin VS, 1983. A method of estimating *Ips sexdentatus* Coleoptera Ipidae density of egg production. Lesovedenie, 1, 56–59.
- Bakke A, 1968. Ecological studies on bark beetles (Coleoptera: Scolytidae) associated with Scots pine (*Pinus sylvestris* L.) in Norway with particular reference to the influence of temperature. Meddelelser fra det Norske Skogsforsoksvesen, 21, 441–602.
- Balachowsky A, 1949. *Faune de France*. 50. *Coleoptères Scolytides*. Lechevalier, Paris, 320 pp.
- Beffa GD, 2006. Attacks by bark beetles on the eastern white pine in Piedmont. Sherwood – Foreste ed Alberi Oggi, 118, 39–42.
- Besceli O and Ekici M, 1969. Control and biology of *Ips sexdentatus* in the *Picea orientalis* region. Ormancilik Arastirma Enstitusu Teknik Bulten, 32, 32–32.
- Bossard M, Feranec J and Otahel J, 2000. CORINE land cover technical guide – Addendum 2000. Tech. Rep. 40, European Environment Agency. Available online: https://www.eea.europa.eu/ds_resolveuid/032TFUPGVR, INRMM-MiD:13106045
- Bouhot L, Lieutier F and Debouzie D, 1988. Spatial and temporal distribution of attacks by *Tomicus piniperda* L. and *Ips sexdentatus* Boern. (Col., Scolytidae) on *Pinus sylvestris*. Journal of Applied Entomology, 106, 356–371.

- Bright DE and Skidmore RE, 2002. *A catalogue of Scolytidae and Platypodidae (Coleoptera), Supplement 2 (1995–1999)*. NRC Research Press, Ottawa, Canada, 523 pp.
- Bueno A, Diez JJ and Fernández MM, 2010. Ophiostomatoid Fungi Transported by *Ips sexdentatus* (Coleoptera, Scolytidae) in *Pinus pinaster* in NW Spain. *Silva Fennica*, 44, 387–397.
- Büttner G, Kosztra B, Maucha G and Pataki R, 2012. Implementation and achievements of CLC2006. Tech. rep., European Environment Agency. Available online: http://www.eea.europa.eu/ds_resolveuid/GQ4JECM8TB, INRMM-MiD:14284151
- Chararas C, 1962. Etude biologique des scolytides des conifères. *Encyclopedie Entomologique*, Ser. A, Nr. 38, Paul Lechevalier, Paris.
- Chirici G, Bertini R, Travaglini D, Puletti N and Chiavetta U, 2011a. The common NFI database. In: Chirici G, Winter S and McRoberts RE (eds.) *National forest inventories: contributions to forest biodiversity assessments*. Springer, Berlin, pp. 99–119.
- Chirici G, McRoberts RE, Winter S, Barbati A, Brändli U-B, Abegg M, Beranova A, Barbati J, Rondeux J, Bertini R, Alberdi Asensio I and Condés S, 2011b. Harmonization tests. In: Chirici G, Winter S and McRoberts RE (eds.) *National forest inventories: contributions to forest biodiversity assessments*. Springer, Berlin, pp. 121–190.
- Cobos Suarez JM and Ruiz Urrestarazu MM, 1990. Phytosanitary problems of the species *Pinus radiata* D. Don in Spain, with special reference to the Basque country. *Boletín de Sanidad Vegetal, Plagas*, 16, 37–53.
- Croisé L, Dreyer E and Lieutier F, 1998. Effects of drought stress and severe pruning on the reaction zone induced by single inoculations with a bark beetle associated fungus (*Ophiostoma ips*) in the phloem of young Scots pines. *Canadian Journal of Forest Research*, 28, 1814–1824.
- EFSA PLH Panel (EFSA Panel on Plant Health), 2010a. PLH Guidance on a harmonised framework for pest risk assessment and the identification and evaluation of pest risk management options by EFSA. *EFSA Journal* 2010;8(2):1495, 66 pp. <https://doi.org/10.2903/j.efsa.2010.1495>
- EFSA PLH Panel (EFSA Panel on Plant Health), 2010b. Risk assessment of *Gibberella circinata* for the EU territory and identification and evaluation of risk management options. *EFSA Journal* 2010;8(6):1620, 93 pp. <https://doi.org/10.2903/j.efsa.2010.1620>. Available online: www.efsa.europa.eu
- EPPO (European and Mediterranean Plant Protection Organization), 2017. EPPO Global Database. Available online: <https://gd.eppo.int> [Accessed: 22 August 2017]
- Etxebeste I and Pajares JA, 2011. Verbenone protects pine trees from colonization by the six-toothed pine bark beetle, *Ips sexdentatus* Boern. (Col.: Scolytinae). *Journal of Applied Entomology*, 135, 258–268.
- Etxebeste I, Alvarez G, Pérez G and Pajares JA, 2012. Field response of the six-toothed pine bark beetle, *Ips sexdentatus* (Col.: Curculionidae, Scolytinae), to pheromonal blend candidates. *Journal of Applied Entomology*, 136, 431–444.
- Etxebeste I, Alvarez G and Pajares JA, 2013. Log colonization by *Ips sexdentatus* prevented by increasing host unsuitability signaled by verbenone. *Entomologia Experimentalis Et Applicata*, 147, 231–240.
- FAO (Food and Agriculture Organization of the United Nations), 2004. ISPM (International Standards for Phytosanitary Measures) 21—Pest risk analysis of regulated non-quarantine pests. FAO, Rome, 30 pp. Available online: https://www.ippc.int/sites/default/files/documents/1323945746_ISPM_21_2004_En_2011-11-29_Refor.pdf
- FAO (Food and Agriculture Organization of the United Nations), 2013. ISPM (International Standards for Phytosanitary Measures) 11—Pest risk analysis for quarantine pests. FAO, Rome, 36 pp. Available online: https://www.ippc.int/sites/default/files/documents/20140512/ispm_11_2013_en_2014-04-30_201405121523-494.65%20KB.pdf
- Fernandez MM, 2006. Colonization of fire-damaged trees by *Ips sexdentatus* (Boerner) as related to the percentage of burnt crown. *Entomologica Fennica*, 17, 381–386.
- Fernandez MM and Salgado Costas JM, 1999. Susceptibility of fire-damaged pine trees (*Pinus pinaster* and *Pinus nigra*) to attacks by *Ips sexdentatus* and *Tomicus piniperda* (Coleoptera: Scolytidae). *Entomologia Generalis*, 24, 105–114.
- Fettig CJ and Hilszczanski J, 2015. Management Strategies for Bark Beetles in Conifer Forests. In: Vega FE, Hofstetter RW (eds.). *Bark Beetles. Biology and Ecology of Native and Invasive Species*. Elsevier, Amsterdam. pp. 555–584.
- Francke W, Pan ML, Bartels J, König WA, Vité JP, Krawielitzki S and Kohnle U, 1986. The odour bouquet of three pine engraver beetles (*Ips* spp.) 1. *Journal of Applied Entomology*, 101, 453–461.
- Frisullo S, et al., 2003. Implication of chromogenous fungi and insects in loricated pine dieback in Pollino National Park. *Informatore Fitopatologico*, 53, 50–52.
- Grégoire JC, Raffa KF and Lindgren BS, 2015. Economics and Politics of Bark Beetles. In: Vega FE, Hofstetter RW (eds.), *Bark Beetles. Biology and Ecology of Native and Invasive Species*. Elsevier, Amsterdam. pp. 585–613.
- Grüne S, 1979. *Brief Illustrated Key to European Bark Beetles*. M&H Schaper, Hannover, 182 pp.
- Haack RA, 2001. Intercepted Scolytidae (Coleoptera) at US ports of entry: 1985–2000. *Integrated Pest Management Reviews*, 6, 253–282.
- Hiederer R, Houston Durrant T, Granke O, Lambotte M, Lorenz M, Mignon B and Mues V, 2007. Forest focus monitoring database system - validation methodology. Vol. EUR 23020 EN of EUR – Scientific and Technical Research. Office for Official Publications of the European Communities. <https://doi.org/10.2788/51364>

- Hiederer R, Houston Durrant T and Micheli E, 2011. Evaluation of BioSoil demonstration project - Soil data analysis. Vol. 24729 of EUR - Scientific and Technical Research. Publications Office of the European Union. <https://doi.org/10.2788/56105>
- Houston Durrant T and Hiederer R, 2009. Applying quality assurance procedures to environmental monitoring data: a case study. *Journal of Environmental Monitoring*, 11, 774–781.
- Houston Durrant T, San-Miguel-Ayanz J, Schulte E and Suarez Meyer A, 2011. Evaluation of BioSoil demonstration project: forest biodiversity - Analysis of biodiversity module. Vol. 24777 of EUR – Scientific and Technical Research. Publications Office of the European Union. <https://doi.org/10.2788/84823>
- Hulcr J, Atkinson T, Cognato AI, Jordal BH and McKenna DD, 2015. Morphology, taxonomy and phylogenetics of bark beetles. In: Vega FE and Hofstetter RW (eds.). *Bark Beetles. Biology and Ecology of Native and Invasive Species*. Elsevier, Amsterdam. pp. 41–84.
- Ismukhambetov Z, 1964. Insect-pests, brought in from the Siberian forest, that are dangerous for Schrenk spruce (*Picea schrenkiana*) From: REF ZH BIOL, 1964, No. 24E231. (Translation). *Tr Nauch Issled Inst Zashch Rast Raz Ssr*, 8, 245–250.
- Ivojinovi S, 1960. A contribution to the knowledge of Scolytidae living on *Pinus heldreichii* Christ, and *Pinus peuce* Griesbach in Serbia English summ. *Plant Protection [Zashtita Bilja]*, 57–58, 21–29.
- Jactel H, 1991. Dispersal and flight behaviour of *Ips sexdentatus* (Coleoptera: Scolytidae) in pine forest. *Official journal of the Institut National de la Recherche Agronomique (INRA)*, 48, 417–428.
- Jactel H and Gaillard J, 1991. A preliminary study of the dispersal potential of *Ips sexdentatus* (Boern) (Col., Scolytidae) with an automatically recording flight mill. *Journal of Applied Entymology*, 112, 138–145.
- Jactel H, Van Halder I, Menassieu P, Zhang QH and Schlyter F, 2001. Non-host volatiles disrupt the response of the stenographer bark beetle, *Ips sexdentatus* (Coleoptera: Scolytidae), to pheromone-baited traps and maritime pine logs. *Integrated Pest Management Reviews*, 6, 197–207.
- Jankowiak R, 2012. Ophiostomatoid fungi associated with *Ips sexdentatus* on *Pinus sylvestris* in Poland. *Dendrobiology*, 68, 43–53.
- Kirisits T, 2004. Fungal associates of european bark beetles with special emphasis on the ophiostomatoid fungi. In: Lieutier F, Day KR, Battisti A, Grégoire JC and Evans HF (eds.). *Bark and wood boring insects in living trees in Europe, a synthesis*. Springer, The Netherlands. pp. 181–236.
- Knížek M and Beaver R, 2004. Taxonomy and systematics of bark and ambrosia beetles. In: Lieutier F, Day K, Battisti A, Grégoire JC, Evans H (eds). *Bark and Wood Boring Insects in Living Trees in Europe, a Synthesis*. Kluwer, Dordrecht. pp. 41–54. https://doi.org/10.1007/978-1-4020-2241-8_11
- Kobzar VF, 1968. Statistical evaluation of methods of counting *Ips sexdentatus* populations. *Nauc. Trud. Leningr. Lsoteh. Akad.*, 115, 32–36.
- Kohnle U, Meyer M and Kluber J, 1992. Formulation of population attractant for the pine bark beetle, *Ips sexdentatus* (Col, Scolytidae). *Allgemeine Forst Und Jagdzeitung*, 163, 81–87.
- Kondur Y, Bilgili BC, Şimsek Z and Öner N, 2012. Monitoring the epidemic of bark beetles determined in Uludag fir stands in Ilgaz (Derbent and Doruk) via Landsat satellite images. *Kastamonu Universitesi Orman Fakultesi Dergisi*, 12, 86–90.
- Lévieux J, Lieutier F and Delplanque A, 1985. Les Scolytes ravageurs du Pin sylvestre. *Revue Forestière Française*, 37, 431–440.
- Lévieux J, Cassier P, Guillaumin D and Roques A, 1991. Structures implicated in the transportation of pathogenic fungi by the European bark beetle, *Ips sexdentatus* Boerner – ultrastructure of a mycangium. *Canadian Entomologist*, 123, 245–254.
- Lieutier F, Faure T and Garcia J, 1988. Les attaques de scolytes et le dépérissement du pin sylvestre dans la region Provence-Côte d'Azur. *Revue Forestière-Française*, 40, 224–232.
- Lombardero MJ and Ayres MP, 2011. Factors influencing bark beetle outbreaks after forest fires on the Iberian Peninsula. *Environmental Entomology*, 40, 1007–1018.
- Lopez S and Goldarazena A, 2012. Flight Dynamics and Abundance of *Ips sexdentatus* (Coleoptera: Curculionidae: Scolytinae) in Different Sawmills from Northern Spain: Differences between Local *Pinus radiata* (Pinales: Pinaceae) and Southern France Incoming *P. pinaster* Timber. *Psyche (Cambridge)*, 2012, 1–6.
- Lozovoi DI, 1961. Koroedy khvoinykh drevesnykh porod v parkovykh i lesnykh nasazhdeniyakh Gruzii [Bark beetles (Ipidae) of coniferous species in the park and forest plantings of Georgia]. *Akademiia Nauk Gruzinskoi SSR, Tbilisskii Botaniche skii Institut, Vestnik*, 67, 91–113.
- Lozovoj DI, 1966. Hozjajstvenno vaznye vidy koroedov hvoinykh (elovyh) nasazdenij Gruzii i mery bor'by s nimi. [Economically important species of barkbeetles in the conifer (Spruce) stands of Soviet Georgia, and their control]. *Hozjajstvenno vaznye vidy koroedov hvoinykh (elovyh) nasazdenij Gruzii i mery bor'by s nimi*. 1966 pp. 90 pp. ref. 5 pp. of refs. From CAB Direct. Available online: <https://www.cabdirect.org/cabdirect/abstract/19660601005> [Accessed: 14 June 2017].
- Ozcan GE, Eroglu M and Akinci HA, 2011. Use of pheromone-baited traps for monitoring *Ips sexdentatus* (Boerner) (Coleoptera: Curculionidae) in oriental spruce stands. *African Journal of Biotechnology*, 10, 16351–16360.
- Pineau X, Bourguignon M, Jactel H, Lieutier F and Sallé A, 2017. Pyrrhic victory for bark beetles: successful standing tree colonization triggers strong intraspecific competition for offspring of *Ips sexdentatus*. *Forest Ecology and Management*, 399, 188–196.

- de Rigo D, 2012. Semantic Array Programming for environmental modelling: application of the Mastrave library. In: Seppelt R, Voinov AA, Lange S and Bankamp D (eds.) International Environmental Modelling and Software Society (iEMSS) 2012 International Congress on Environmental Modelling and Software - Managing Resources of a Limited Planet: Pathways and Visions under Uncertainty, Sixth Biennial Meeting. pp. 1167-1176.
- de Rigo D, Caudullo G, Busetto L and San-Miguel-Ayanz J, 2014. Supporting EFSA assessment of the EU environmental suitability for exotic forestry pests: final report. EFSA Supporting Publications 11, EN-434+. <https://doi.org/10.2903/sp.efsa.2014.en-434>, INRM-MiD:13114000
- de Rigo D, Caudullo G, Houston Durrant T and San-Miguel-Ayanz J, 2016. The European Atlas of Forest Tree Species: modelling, data and information on forest tree species. In: San-Miguel-Ayanz J, De Rigo D, Caudullo G, Houston Durrant T and Mauri A (eds.). European Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg, pp. e01aa69+. Available online: <https://w3id.org/mtv/FISE-Comm/v01/e01aa69>
- de Rigo D, Caudullo G, San-Miguel-Ayanz J and Barredo JI, 2017. Robust modelling of the impacts of climate change on the habitat suitability of forest tree species. Publication Office of the European Union, 58 pp. ISBN:978-92-79-66704-6, <https://doi.org/10.2760/296501>, INRM-MiD:14314400
- Romón P, Zhou X, Iturrondobeitia JC, Wingfield MJ and Goldarazena A, 2007. *Ophiostoma* species (Ascomycetes: Ophiostomatales) associated with bark beetles (Coleoptera: Scolytinae) colonizing *Pinus radiata* in northern Spain. Canadian Journal of Microbiology, 53, 756–767.
- Romón P, Troya M, Fernández de Gamarra ME, Eguzkitza A, Iturrondobeitia JC and Goldarazena A, 2008. Fungal communities associated with pitch canker disease of *Pinus radiata* caused by *Fusarium circinatum* in northern Spain: association with insects and pathogen-saprophyte antagonistic interactions. Canadian Journal of Plant Pathology-Revue Canadienne De Phytopathologie, 30, 241–253.
- Rossi JP, Samalens JC, Guyon D, Van Halder I, Jactel H, Menassieu P and Piou D, 2009. Multiscale spatial variation of the bark beetle *Ips sexdentatus* damage in a pine plantation forest (Landes de Gascogne, Southwestern France). Forest Ecology and Management, 257, 1551–1557.
- Samalens JC, Rossi JP, Guyon D, Van Halder I, Menassieu P, Piou D and Jactel H, 2007. Adaptive roadside sampling for bark beetle damage assessment. Forest Ecology and Management, 253, 177–187.
- San-Miguel-Ayanz J, 2016. The European Union Forest Strategy and the Forest Information System for Europe. In: San-Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A (eds.). European Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg, pp. e012228+.
- San-Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T and Mauri A. (Eds.), 2016. *European Atlas of Forest Tree Species*. Publication Office of the European Union, Luxembourg. ISBN: 978-92-79-36740-3, <https://doi.org/10.2788/4251>, Available online: <https://w3id.org/mtv/FISE-Comm/v01/>
- Santolamazza-Carbone S, Pestana M and Antonio Vega J, 2011. Post-fire attractiveness of maritime pines (*Pinus pinaster* Ait.) to xylophagous insects. Journal of Pest Science, 84, 343–353.
- Schedl KE, 1981. 91. Familie: Scolytidae (Borken- und Ambrosiakäfer). In: Freude H, Harde KW, Lohse GA (eds.). *Die Käfer Mitteleuropas*. Goecke & Evers, Krefeld, pp. 34–99.
- Schimitschek E, 1939. Contributions to the Forest Entomology of Turkey III. The Outbreak of *Ips sexdentatus* in the Region of the Oriental Spruce. First Part. Zeitschrift für Angewandte Entomologie, 26, 545–588.
- Schönherr J, Vité JP and Serez M, 1983. Überwachung von *Ips sexdentatus*-Populationen mit synthetischem Lockstoff. Zeitschrift für Angewandte Entomologie, 95, 51–53.
- Scopus, 2017. Available online: <https://www.scopus.com/home.uri> [Accessed: 28 March 2017].
- Slankis A, 1969. A new species of endoparasitic nematode, *Contortylenchus pseudodiplogaster* n.sp. from the bark beetle, *Ips sexdentatus*. Mater. nauch. Konf. vses. Obshch. Gel'mint. (Part II), pp. 302–305.
- Stadelmann G, Bugmann H, Meier F, Wermelinger B and Bigler C, 2013. Effects of salvage logging and sanitation felling on bark beetle (*Ips typographus* L.) infestations. Forest Ecology and Management, 305, 273–281. <https://doi.org/10.1016/j.foreco.2013.06.003>
- Vité JP, Bakke A and Hughes PR, 1974. Ein populationslockstoff des zwölftzahnigen kiefernborckenkäfers *Ips sexdentatus*. Naturwissenschaften, 61, 365–366.
- Wang X, et al., 2011. Effect of X-ray (9 MeV) irradiation on the development and propagation of *Ips sexdentatus*. Plant Quarantine (Shanghai), 25, 28–31.
- Wood SL, 1982. *The bark and ambrosia beetles of North and Central America (Coleoptera: Scolytidae), a taxonomic monograph*. Brigham Young University, Provo, UT, 1359 pp.

Abbreviations

CLC	Corine Land Cover
EPPO	European and Mediterranean Plant Protection Organization
EUFGIS	European Information System on Forest Genetic Resources
EU MS	European Union Member State
FAO	Food and Agriculture Organization
GD ²	Georeferenced Data on Genetic Diversity
IPPC	International Plant Protection Convention

JRC	Joint Research Centre of the European Commission
PLH	EFSA Panel on Plant Health
PZ	protected zone
RNQP	regulated non-quarantine pest
RPP	relative probability of presence
RRO	risk reduction option
SMFA	spatial multiscale frequency analysis
TFEU	Treaty on the Functioning of the European Union
ToR	Terms of Reference

Appendix A – Methodological notes on Figure 2

The relative probability of presence (RPP) reported here for *Pinus* and *Picea* spp. in Figure 2 and in the European Atlas of Forest Tree Species (de Rigo et al., 2016; San-Miguel-Ayanz et al., 2016) is the probability of that genus to occur in a given spatial unit (de Rigo et al., 2017). In forestry, such a probability for a single taxon is called 'relative'. The maps of RPP are produced by means of the constrained spatial multiscale frequency analysis (C-SMFA) (de Rigo et al., 2014, 2017) of species presence data reported in geolocated plots by different forest inventories.

A.1. Geolocated plot databases

The RPP models rely on five geodatabases that provide presence/absence data for tree species and genera: four European-wide forest monitoring data sets and a harmonised collection of records from national forest inventories (de Rigo et al., 2014, 2016, 2017). The databases report observations made inside geolocalised sample plots positioned in a forested area, but do not provide information about the plot size or consistent quantitative information about the recorded species beyond presence/absence.

The harmonisation of these data sets was performed within the research project at the origin of the European Atlas of Forest Tree Species (de Rigo et al., 2016; San-Miguel-Ayanz, 2016; San-Miguel-Ayanz et al., 2016). Given the heterogeneity of strategies of field sampling design and establishment of sampling plots in the various national forest inventories (Chirici et al. 2011a,b), and also given legal constraints, the information from the original data sources was harmonised to refer to an INSPIRE compliant geospatial grid, with a spatial resolution of 1 km² pixel size, using the ETRS89 Lambert Azimuthal Equal-Area as geospatial projection (EPSG: 3035, <http://spatialreference.org/ref/epsg/etrs89-etrs-laea/>).

A.1.1. European National Forestry Inventories database

This data set was derived from National Forest Inventory data and provides information on the presence/absence of forest tree species in approximately 375,000 sample points with a spatial resolution of 1 km²/pixel, covering 21 European countries (de Rigo et al., 2014, 2016).

A.1.2. Forest Focus/Monitoring data set

This project is a Community scheme for harmonised long-term monitoring of air pollution effects in European forest ecosystems, normed by EC Regulation No 2152/2003⁵. Under this scheme, the monitoring is carried out by participating countries on the basis of a systematic network of observation points (Level I) and a network of observation plots for intensive and continuous monitoring (Level II). For managing the data, the JRC implemented a Forest Focus Monitoring Database System, from which the data used in this project were taken (Hiederer et al., 2007; Houston Durrant and Hiederer, 2009). The complete Forest Focus data set covers 30 European Countries with more than 8,600 sample points.

A.1.3. BioSoil data set

This data set was produced by one of a number of demonstration studies performed in response to the 'Forest Focus' Regulation (EC) No 2152/2003 mentioned above. The aim of the BioSoil project was to provide harmonised soil and forest biodiversity data. It comprised two modules: a Soil Module (Hiederer et al., 2011) and a Biodiversity Module (Houston Durrant et al., 2011). The data set used in the C-SMFA RPP model came from the Biodiversity module, in which plant species from both the tree layer and the ground vegetation layer were recorded for more than 3,300 sample points in 19 European Countries.

⁵ Council of the European Union, 2003. Regulation (EC) No 2152/2003 of the European Parliament and of the Council of 17 November 2003 concerning monitoring of forests and environmental interactions in the Community (Forest Focus). Official Journal of the European Union 46 (L 324), 1–8.

A.1.4. European Information System on Forest Genetic Resources (EUFGIS)

EUFGIS (<http://portal.eufgis.org>) is a smaller geodatabase providing information on tree species composition in over 3,200 forest plots in 34 European countries. The plots are part of a network of forest stands managed for the genetic conservation of one or more target tree species. Hence, the plots represent the natural environment to which the target tree species are adapted.

A.1.5. Georeferenced Data on Genetic Diversity (GD²)

GD² (<http://gd2.pierroton.inra.fr>) provides information about 63 species of interest for genetic conservation. The database covers 6,254 forest plots located in stands of natural populations that are traditionally analysed in genetic surveys. While this database covers fewer species than the others, it covers 66 countries in Europe, North Africa and the Middle East, making it the data set with the largest geographic extent.

A.2. Modelling methodology

For modelling, the data were harmonised in order to have the same spatial resolution (1 km²) and filtered to a study area comprising 36 countries in the European continent. The density of field observations varies greatly throughout the study area and large areas are poorly covered by the plot databases. A low density of field plots is particularly problematic in heterogeneous landscapes, such as mountainous regions and areas with many different land use and cover types, where a plot in one location is not representative of many nearby locations (de Rigo et al., 2014). To account for the spatial variation in plot density, the model used here (C-SMFA) considers multiple spatial scales when estimating RPP. Furthermore, statistical resampling is systematically applied to mitigate the cumulated data-driven uncertainty.

The presence or absence of a given forest tree species then refers to an idealised standard field sample of negligible size compared with the 1 km² pixel size of the harmonised grid. The modelling methodology considered these presence/absence measures as if they were random samples of a binary quantity (the punctual presence/absence, not the pixel one). This binary quantity is a random variable having its own probability distribution which is a function of the unknown average probability of finding the given tree species within a plot of negligible area belonging to the considered 1 km² pixel (de Rigo et al., 2014). This unknown statistic is denoted hereinafter with the name of 'probability of presence'.

C-SMFA performs spatial frequency analysis of the geolocated plot data to create preliminary RPP maps (de Rigo et al., 2014). For each 1 km² grid cell, the model estimates kernel densities over a range of kernel sizes to estimate the probability that a given species is present in that cell. The entire array of multiscale spatial kernels is aggregated with adaptive weights based on the local pattern of data density. Thus, in areas where plot data are scarce or inconsistent, the method tends to put weight on larger kernels. Wherever denser local data are available, they are privileged ensuring a more detailed local RPP estimation. Therefore, a smooth multiscale aggregation of the entire arrays of kernels and data sets is applied instead of selecting a local 'best performing' one and discarding the remaining information. This array-based processing, and the entire data harmonisation procedure, are made possible thanks to the semantic modularisation which defines the Semantic Array Programming modelling paradigm (de Rigo, 2012).

The probability to find a single species (e.g. a particular coniferous tree species) in a 1 km² grid cell cannot be higher than the probability of presence of all the coniferous species combined. The same logical constraints applied to the case of single broadleaved species with respect to the probability of presence of all the broadleaved species combined. Thus, to improve the accuracy of the maps, the preliminary RPP values were constrained so as not to exceed the local forest-type cover fraction with an iterative refinement (de Rigo et al., 2014). The forest-type cover fraction was estimated from the classes of the Corine Land Cover (CLC) maps which contain a component of forest trees (Bossard et al., 2000; Büttner et al., 2012).

The resulting probability of presence is relative to the specific tree taxon, irrespective of the potential co-occurrence of other tree taxa with the measured plots, and should not be confused with the absolute abundance or proportion of each taxon in the plots. RPP represents the probability of finding at least one individual of the taxon in a plot placed randomly within the grid cell, assuming that the plot has

negligible area compared with the cell. As a consequence, the sum of the RPP associated with different taxa in the same area is not constrained to be 100%. For example, in a forest with two co-dominant tree species which are homogeneously mixed, the RPP of both may be 100% (see e.g. the Glossary in San-Miguel-Ayanz et al. (2016), <http://forest.jrc.ec.europa.eu/media/atlas/Glossary.pdf>).

The robustness of RPP maps depends strongly on sample plot density, as areas with few field observations are mapped with greater uncertainty. This uncertainty is shown qualitatively in maps of 'RPP trustability'. RPP trustability is computed on the basis of the aggregated equivalent number of sample plots in each grid cell (equivalent local density of plot data). The trustability map scale is relative, ranging from 0 to 1, as it is based on the quantiles of the local plot density map obtained using all field observations for the species. Thus, trustability maps may vary among species based on the number of databases that report a particular species (de Rigo et al., 2014, 2016).

The RPP and relative trustability range from 0 to 1 and are mapped at a 1 km spatial resolution. To improve visualisation, these maps can be aggregated to coarser scales (i.e. 10×10 pixels or 25×25 pixels, respectively, summarising the information for aggregated spatial cells of 100 and 625 km²) by averaging the values in larger grid cells.