



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short Communication

In vitro evaluation of antiviral activity of single and combined repurposable drugs against SARS-CoV-2

Andrés Pizzorno^a, Blandine Padey^{a,b}, Julia Dubois^a, Thomas Julien^{a,c}, Aurélien Traversier^a, Victoria Dulière^{a,c}, Pauline Brun^{a,c}, Bruno Lina^{a,d}, Manuel Rosa-Calatrava^{a,c,**,1}, Olivier Terrier^{a,*,1}

^a CIRI, Centre International de Recherche en Infectiologie, (Team VirPath), Univ Lyon, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, F-69007, Lyon, France

^b Signia Therapeutics SAS, Lyon, France

^c VirNext, Faculté de Médecine RTH Laennec, Université Claude Bernard Lyon 1, Université de Lyon, Lyon, France

^d Laboratoire de Virologie, Centre National de Référence des Virus Influenza Sud, Institut des Agents Infectieux, Groupement Hospitalier Nord, Hospices Civils de Lyon, Lyon, France



ARTICLE INFO

Keywords:

COVID-19
Antivirals
Drug combination
Remdesivir
Berberine
Diltiazem

ABSTRACT

In response to the current pandemic caused by the novel SARS-CoV-2, identifying and validating effective therapeutic strategies is more than ever necessary. We evaluated the *in vitro* antiviral activities of a shortlist of compounds, known for their cellular broad-spectrum activities, together with drugs that are currently under evaluation in clinical trials for COVID-19 patients. We report the antiviral effect of remdesivir, lopinavir, chloroquine, umifenovir, berberine and cyclosporine A in Vero E6 cells model of SARS-CoV-2 infection, with estimated 50% inhibitory concentrations of 0.99, 5.2, 1.38, 3.5, 10.6 and 3 μ M, respectively. Virus-directed plus host-directed drug combinations were also investigated. We report a strong antagonism between remdesivir and berberine, in contrast with remdesivir/diltiazem, for which we describe high levels of synergy, with mean Loewe synergy scores of 12 and peak values above 50. Combination of host-directed drugs with direct acting antivirals underscore further validation in more physiological models, yet they open up interesting avenues for the treatment of COVID-19.

Despite unprecedented global efforts during the last months in the fight of the COVID-19 pandemic, which include more than 500 randomized clinical trials (“Covid-19 living Data,” n.d.), neither a preventive vaccine nor an effective treatment as a potential complement to standard supportive care have yet been validated. This situation is more problematic in the case of severe patients, provided increasing clinical reports account for a rather biphasic pathophysiology. In fact, the initial phase of the disease is characterized by a typical virus-driven response, which might subsequently (by day 7–10 after symptom onset) shunt into an acute immunopathologic phase with an associated hypercytokinemia (Huang et al., 2020; Mehta et al., 2020; Zhu et al., 2020), hence adding more complexity to the choice of a potentially effective treatment. However, such “cytokine storm” is not independent of viral replication

but frequently an attempt to clear the viral aggression. In that regard, recent data suggest that high virus titers during the first week of infection could be correlated with a higher probability of moderate to severe clinical outcome (Liu et al., 2020b; Zheng et al., 2020). This highlights the value of identifying and validating molecules and/or combination of molecules with a strong SARS-CoV-2 antiviral activity for an early and effective reduction of viral load that could ultimately contribute to a more favorable clinical prognosis.

In this context, we sought to assess the *in vitro* antiviral activity of a shortlist of compounds known for their relative broad-spectrum antiviral activities that in our view have a rationale for a putative inhibitory effect against SARS-CoV-2 and, on top of that, could be readily available for further pre-clinical and clinical evaluation. Favipiravir (T-705; 6-

* Corresponding author.

** Corresponding author. CIRI, Centre International de Recherche en Infectiologie, (Team VirPath), Univ Lyon, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, F-69007, Lyon, France.

E-mail addresses: manuel.rosa-calatrava@univ-lyon1.fr (M. Rosa-Calatrava), olivier.terrier@univ-lyon1.fr (O. Terrier).

¹ These authors contributed equally.

Table 1
Antiviral activity of post-infection treatment with candidate molecules against SARS-CoV-2 *in vitro*.

	Drug class	Drug indication	CAS Number	IC50 (μM)	CC50 (μM)	SI
Remdesivir	Antimetabolites Antiviral agents	Treatment of Ebola virus disease	1809249-37-3	0.987	275	278.62
Lopinavir	Antiviral agents	Treatment of HIV-infection	192725-17-0	5.246	45	8.57
Chloroquine	Antimalarial agents	Treatment of malaria, rheumatoid arthritis, glioma	54-05-7	1.38	238	172.46
Berberine	Alkaloid	Treatment of parasitic and fungal infections	2086-83-1	10.577	>400	>37.84
Cyclosporine A	Enzyme inhibitors Immunosuppressive agents	Prophylaxis of organ rejection, treatment of rheumatoid arthritis & dermatologic diseases	59865-13-3	3.048	>90	>29.53
Arbidol (umifenovir)	Antiviral agents	Treatment of Influenza infection	131707-25-0	3.537	75	21.20
Diltiazem	antihypertensive & vasodilating agents	Management of hypertension	42399-41-7	>45	424	<9.42
Ribavirin	Hepatitis C & antiviral agents	Treatment of HCV infection	36791-04-5	>10	>100	nd
Favipiravir	Antiviral agents	Treatment of Influenza infection	259793-96-9	>100	631	<6.31




Fig. 1. Evaluation of antiviral activity of six drug candidates against SARS-CoV-2 *in vitro*. Vero E6 cells were infected by SARS-CoV-2 at a MOI of 0.01 and treated 1 h post infection (hpi) with serial dilutions of remdesivir, lopinavir, chloroquine, berberine, cyclosporine A and arbidol (umifenovir). Viral production was measured 48 hpi and expressed in relative values compared to the vehicle-treated control. The impact of treatment on cell viability was assessed by MTS assay (CellTiter 96® AQueous One Solution Cell Proliferation Assay, Promega), and expressed in relative values compared to control. Cell viability measurements were performed to ensure that molecular viral quantification was performed within a non-cytotoxic concentration range that could bias the measurement.

fluoro-3-hydroxy-2-pyrazinecarboxamine) is a pyrazine derivative that has demonstrated potent antiviral activity against multiple RNA viruses, being currently approved in Japan for the treatment of influenza infections (Furuta et al., 2013, 2002). The ribonucleoside analog ribavirin (1- β -D-ribofuranosyl-1,2,4-triazole-3-carboxamide) is also known to show broad-spectrum antiviral activity and is used in combination with interferon- α to treat hepatitis C virus infection (Crotty et al., 2000; Sidwell et al., 1972). Umifenovir (Arbidol) is a small indole-derivative molecule, licensed in Russia and China for the prophylaxis and treatment of influenza and other respiratory viral infections (Blaising et al., 2014). Berberine is a plant-derived isoquinoline alkaloid, described as having broad antiviral activity, notably against influenza and alphaviruses (Varghese et al., 2016a; Yan et al., 2018). Cyclosporine A is a widely used immunosuppressant drug also known for its ability to inhibit the replication of several viruses, including several coronaviruses (de Wilde et al., 2011). Lastly, diltiazem is a voltage gated Ca²⁺ channel

blocker commonly used as anti-hypertensive, which we have recently repurposed as a host-directed inhibitor of influenza virus (Pizzorno et al., 2019).

Molecules were obtained from Sigma-Aldrich and 10 mM stock solutions were prepared in DMSO (favipiravir, lopinavir, remdesivir, umifenovir), ethanol (berberine, cyclosporine A) or water (chloroquine, diltiazem, ribavirin). We therefore performed experimental infections in Vero E6 cells at a multiplicity of infection (MOI) of 0.01 using the BetaCoV/France/IDF0571/2020 SARS-CoV-2 strain, previously isolated from one of the first COVID-19 cases in France. One-hour post infection (hpi), the viral inoculum was removed, and cells were treated with serial dilutions of the candidate molecules in infection media. Dilutions of the corresponding solvents in infection media were used for mock-treated controls. Supernatants were collected 48 hpi for subsequent viral titration by RT-qPCR, as described elsewhere (Pizzorno et al., 2020). Noteworthy, we included the active principles of the three drugs evaluated in




Fig. 2. Evaluation of remdesivir-berberine combined treatment against SARS-CoV-2 *in vitro*. A. Vero E6 cells were infected by SARS-CoV-2 at a MOI of 0.01 and treated 1 hpi with serial dilutions of remdesivir in the presence of different fixed concentrations of berberine (A, left panel) or, alternatively, serial dilutions of berberine in the presence of different fixed concentrations of remdesivir (A, right panel). Viral production was measured 48 hpi and expressed in relative values compared to the vehicle-treated control. B. Dose-response percent inhibition matrix of single and combined remdesivir-berberine treatments C. Interaction landscape between remdesivir and berberine as calculated using the Loewe additive model. Areas with synergy scores of -10 or lower (green) represent zones of drug antagonism.

the WHO's COVID-19 Solidarity Trial ("Solidarity 2020" clinical trial for COVID-19 treatments," 2020) as comparators: i) remdesivir, an experimental adenosine nucleotide analog prodrug previously evaluated in the context of the 2018 Kivu Ebola epidemic (Mulangu et al., 2019); ii) lopinavir, a protease inhibitor antiretroviral approved for the treatment of HIV infections (Chu, 2004); iii) chloroquine, an aminoquinolone derivative mainly used for the treatment of malaria but also for certain chronic inflammatory diseases such as rheumatoid arthritis and lupus erythematosus (Al-Bari, 2017).

Table 1 and Fig. 1 recapitulate the main features of the evaluated compounds, including the calculated 50% inhibitory concentration (IC50), 50% cytotoxic concentration (CC50) and selectivity index (SI) values as well as the dose-response curves of the compounds with such an effect. As expected, calculated IC50 values for lopinavir, remdesivir and chloroquine were in the low micromolar range (5.2, 0.99 and 1.38 μM , respectively), with the latter two showing a very favorable SI (278 and 172, respectively), in good agreement with recent reports (Choy et al., 2020; Wang et al., 2020). On the other hand, although their efficacy against influenza viruses was previously validated (data not

shown), neither diltiazem nor ribavirin or favipiravir showed antiviral activity against SARS-CoV-2 in the conditions tested. This is not surprising in the case of diltiazem, considering its recently described antiviral properties are mostly based on the capacity to induce type I and III interferon (IFN) antiviral responses (Pizzorno et al., 2019), which are partially abrogated in the Vero E6 cell line (Prescott et al., 2010). However, we would *a priori* have expected that broad spectrum inhibitors of viral RNA synthesis ribavirin and favipiravir show at least mild antiviral activity against SARS-CoV-2, notably in the context of preliminary clinical data suggesting a putative benefit of favipiravir for the treatment of COVID-19 patients in comparison with umifenovir (Chen et al., 2020a; Liu et al., 2020a). Nonetheless, our results are in line with recent observations by Choy and colleagues (Choy et al., 2020). Interestingly, umifenovir, berberine and cyclosporine A, three compounds previously reported to have broad antiviral activity, displayed dose-dependent antiviral responses against SARS-CoV-2. Umifenovir (IC50: 3.5 μM and SI: 21) was reported to inhibit fusion between the viral envelope and the membrane of the target cell, although the exact mechanism of this inhibition is still not well described. Besides its

Table 2
Inhibitory effect of remdesivir-berberine combined treatment against SARS-CoV-2 *in vitro*. For each experimental condition, IC₅₀, CC₅₀ and the corresponding SI values are listed.

Combination treatment	IC ₅₀ (μM)	CC ₅₀ (μM)	SI
Remdesivir	0.987	275	278.62
Remdesivir + Berberine 12.5 μM	0.409	264	645.47
Berberine	10.577	> 400	> 37.84
Berberine + Remdesivir 0.625 μM	6.95		> 57.55
Berberine + Remdesivir 2.5 μM	8.237		> 48.56

anti-influenza properties, umifenovir has been shown to inhibit some arthropod-borne flaviviruses, including Zika, West Nile and tick-borne encephalitis virus (Haviernik et al., 2018). Umifenovir is currently being evaluated in clinical trials against COVID-19 (Chen et al., 2020a;

Liu et al., 2020a). Berberine (IC₅₀: 10.6 μM and SI: >37) has demonstrated *in vitro* and *in vivo* (mice) antiviral activity against influenza, Chikungunya and enterovirus 71, possibly by downregulating autophagy and the MEK/ERK signaling pathway (Varghese et al., 2016b, 2016b). Cyclosporine A (IC₅₀: 3.0 μM and SI: >29) was shown to be very effective in inhibiting two different coronaviruses *in vitro*, human coronavirus 229E (HCoV-229E) and mouse hepatitis virus (MHV) but not SARS-CoV (de Wilde et al., 2011). Although the exact mechanism by which cyclosporine A inhibits coronavirus replication remains to be established, an early blockage of viral RNA and protein synthesis has been suggested (de Wilde et al., 2011).

There is arguably a consensus among virologists that combining two compounds with antiviral activity and different viral or cellular targets could result in better virological and physiological responses than those of antiviral monotherapy, for which we decided to explore this hypothesis in the case of SARS-CoV-2. Provided recent results of clinical trials failed to show a therapeutic benefit of the use of either the lopinavir-ritonavir association or chloroquine (or its derivative hydroxychloroquine) in COVID-19 patients (Cao et al., 2020; Chen et al., 2020b; Magagnoli et al., 2020), we then selected remdesivir as the main




Fig. 3. Synergistic effect of remdesivir-diltiazem combined treatment against SARS-CoV-2 *in vitro*. A. Vero E6 cells were infected by SARS-CoV-2 at a MOI of 0.01 and treated 1 hpi with serial dilutions of remdesivir in the presence of different fixed concentrations of diltiazem (A, left panel) or, alternatively, serial dilutions of diltiazem in the presence of different fixed concentrations of remdesivir (A, right panel). Viral production was measured 48 hpi and expressed in relative values compared to the vehicle-treated control. B. Dose-response percent inhibition matrix of single and combined remdesivir-diltiazem treatments C. Interaction landscape between remdesivir and diltiazem as calculated using the Loewe additive model. Areas with synergy scores of 10 or higher (red) represent zones of drug synergy.

Table 3
Inhibitory effect of remdesivir-diltiazem combined treatment against SARS-CoV-2 *in vitro*. For each experimental condition, IC50, CC50 and the corresponding SI values are listed.

Combination treatment	IC50 (μM)	CC50 (μM)	SI
Remdesivir	0.987	275	278.62
Remdesivir	0.627	267	425.83
+ Diltiazem 5.62 μM			
Remdesivir	0.116		2301.72
+ Diltiazem 11.5 μM			
Remdesivir	0.228		1171.05
+ Diltiazem 22.5 μM			
Diltiazem	>45	424	<9.42
Diltiazem	5.268	414	78.59
+ Remdesivir 0.625 μM			
Diltiazem	1.892		218.81
+ Remdesivir 1.25 μM			
Diltiazem	0.554		747.29
+ Remdesivir 2.5 μM			

antiviral agent for our drug combination studies. Following the same infection protocol described above, remdesivir dose-response experiments were repeated in the presence of berberine or diltiazem, given their expected host-directed antiviral properties. Alternatively, berberine and diltiazem dose-response curves were performed in the presence of different fixed concentrations of remdesivir. Our results show that the presence of 12.5 μM berberine induces a shift of the remdesivir dose-response towards lower drug concentration values (Fig. 2A, left panel), as evidenced by a >2-fold reduction of remdesivir IC50 (0.41 μM versus 0.99 μM), with the associated increase on the SI (Table 2). A similar trend was observed when we analyzed the berberine dose-response profile in the presence of 0.625 μM remdesivir but not at higher remdesivir concentrations (Fig. 2A, right panel and Table 2). To further explore this combination, we used SynergyFinder (Ianevski et al., 2017) to calculate and visualize synergy scores, using a wider range of remdesivir/berberine concentrations spanning 24 different pairwise combinations. Interestingly, although the dose-response matrix shows a large range of pairwise combination with significant inhibitory effect (Fig. 2B), calculated Loewe synergy scores (Fig. 2C) account for a strong antagonism between remdesivir and berberine, with negative values distributed all throughout the synergy map regardless of drug concentrations (Fig. 2B and C). Following an analogous approach, the presence of diltiazem induced a stronger shift in the dose-response curve of remdesivir (Fig. 3A, left panel), with an up to 8.5-fold reduction of remdesivir IC50 values (0.12 μM versus 0.99 μM in presence of 11.5 μM diltiazem) and the consequent SI increase (Table 3). A similar trend was observed for diltiazem dose-response curves in the presence of remdesivir (Fig. 3A, right panel). SynergyFinder analysis revealed a clear synergistic inhibitory effect of the combination, driven mainly by remdesivir but also by diltiazem, as shown by both the dose-response matrix (Fig. 3B) and the three-dimensional synergy map (Fig. 3C), with mean Loewe synergy scores of 12 and peak values above 50. Similar results were obtained using the Zero Interaction Potency (ZIP), Highest Single Agent (HAS), and Bliss independence models. Interestingly, the Loewe synergy scores obtained were quite high in comparison to the previously published synergistic combination of remdesivir and emetine (Choy et al., 2020).

In conclusion, our study demonstrated the *in vitro* efficacy against SARS-CoV-2 of broad-spectrum antivirals such as Remdesivir, Lopinavir, chloroquine, Arbidol and Ribavirin and repurposable host-targeted drug candidates such as Berberine or cyclosporine A in monotherapy. As

already reported by our group and others for the treatment of severe forms of influenza infection (Dunning et al., 2014; Pizzorno et al., 2019), drug combination highlighted the added value of combining drugs with different modes of action as a strategy to improve antiviral therapy. From this perspective, the remdesivir/diltiazem combination seems particularly interesting for the treatment of COVID-19, with a synergistic antiviral effect against SARS-CoV-2 in a wide dose range that remains below reported remdesivir therapeutic plasma concentrations. Although berberine, cyclosporine A and the remdesivir-diltiazem combination showed interesting potential as SARS-CoV-2 inhibitors, we are nonetheless aware of the inherent limits of the Vero E6 *in vitro* model, which may not accurately reflect the potential efficacy of antiviral molecules in more physiological models. In addition, the *a priori* counter-intuitive antagonistic effect observed for the remdesivir/berberine combination further illustrates the challenge of proposing effective drug combinations based on individual drug-efficacy data. We therefore expect our data constitute a starting point for further validation of selected candidates in more complex and biologically relevant pre-clinical models of SARS-CoV-2 infection before consideration for evaluation in the clinical setting.

Acknowledgments

This work was funded by INSERM REACTING (REsearch & ACTION emergING infectious diseases), CNRS, and Mérieux research grants. AP, BP, TJ, AT, OT and MRC are co-inventors of a patent application filed by INSERM, CNRS, Université Claude Bernard Lyon 1 and Signia Therapeutics for the repurposing of diltiazem for the treatment of SARS-CoV-2 infections (FR 20/02351).

References

- Al-Bari, M.A.A., 2017. Targeting endosomal acidification by chloroquine analogs as a promising strategy for the treatment of emerging viral diseases. *Pharmacol. Res. Perspect.* 5, e00293 <https://doi.org/10.1002/prp2.293>.
- Blaising, J., Polyak, S.J., Pécheur, E.-I., 2014. Arbidol as a broad-spectrum antiviral: an update. *Antivir. Res.* 107, 84–94. <https://doi.org/10.1016/j.antiviral.2014.04.006>.
- Cao, B., Wang, Y., Wen, D., Liu, W., Wang, Jingli, Fan, G., Ruan, L., Song, B., Cai, Y., Wei, M., Li, X., Xia, J., Chen, N., Xiang, J., Yu, T., Bai, T., Xie, X., Zhang, L., Li, C., Yuan, Y., Chen, H., Li, Huadong, Huang, H., Tu, S., Gong, F., Liu, Y., Wei, Y., Dong, C., Zhou, F., Gu, X., Xu, J., Liu, Z., Zhang, Y., Li, Hui, Shang, L., Wang, K., Li, K., Zhou, X., Dong, X., Qu, Z., Lu, S., Hu, X., Ruan, S., Luo, S., Wu, J., Peng, L., Cheng, F., Pan, L., Zou, J., Jia, C., Wang, Juan, Liu, X., Wang, S., Wu, X., Ge, Q., He, J., Zhan, H., Qiu, F., Guo, L., Huang, C., Jaki, T., Hayden, F.G., Horby, P.W., Zhang, D., Wang, C., 2020. A trial of lopinavir–ritonavir in adults hospitalized with severe covid-19. *N. Engl. J. Med.* 382 (19), 1787–1799. <https://doi.org/10.1056/NEJMoa2001282>.
- Chen, C., Zhang, Yi, Huang, J., Yin, P., Cheng, Z., Wu, J., Chen, S., Zhang, Yongxi, Chen, B., Lu, M., Luo, Y., Ju, L., Zhang, J., Wang, X., 2020a. Favipiravir versus arbidol for COVID-19: a randomized clinical trial. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.17.20037432>, 03.17.20037432.
- Chen, Z., Hu, J., Zhang, Zongwei, Jiang, S., Han, S., Yan, D., Zhuang, R., Hu, B., Zhang, Zhan, 2020b. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.22.20040758>, 03.22.20040758.
- Choy, K.-T., Wong, A.Y.-L., Kaewpreedee, P., Sia, S.F., Chen, D., Hui, K.P.Y., Chu, D.K.W., Chan, M.C.W., Cheung, P.P.-H., Huang, X., Peiris, M., Yen, H.-L., 2020. Remdesivir, lopinavir, emetine, and homoharringtonine inhibit SARS-CoV-2 replication *in vitro*. *Antivir. Res.* 178, 104786 <https://doi.org/10.1016/j.antiviral.2020.104786>.
- Chu, C.M., 2004. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax* 59, 252–256. <https://doi.org/10.1136/thorax.2003.012658>.
- [WWW Document] Covid-19 living Data, 2020 (accessed 5.4.20). <https://covid-nma.com/>.
- Crotty, S., Maag, D., Arnold, J.J., Zhong, W., Lau, J.Y.N., Hong, Z., Andino, R., Cameron, C.E., 2000. The broad-spectrum antiviral ribonucleoside ribavirin is an RNA virus mutagen. *Nat. Med.* 6, 1375–1379. <https://doi.org/10.1038/82191>.
- de Wilde, A.H., Zevenhoven-Dobbe, J.C., van der Meer, Y., Thiel, V., Narayanan, K., Makino, S., Snijder, E.J., van Hemert, M.J., 2011. Cyclosporin A inhibits the replication of diverse coronaviruses. *J. Gen. Virol.* 92, 2542–2548. <https://doi.org/10.1099/vir.0.034983-0>.
- Dunning, J., Baillie, J.K., Cao, B., Hayden, F.G., 2014. Antiviral combinations for severe influenza. *Lancet Infect. Dis.* 14, 1259–1270. [https://doi.org/10.1016/S1473-3099\(14\)70821-7](https://doi.org/10.1016/S1473-3099(14)70821-7).

- Furuta, Y., Gowen, B.B., Takahashi, K., Shiraki, K., Smee, D.F., Barnard, D.L., 2013. Favipiravir (T-705), a novel viral RNA polymerase inhibitor. *Antivir. Res.* 100, 446–454. <https://doi.org/10.1016/j.antiviral.2013.09.015>.
- Furuta, Y., Takahashi, K., Fukuda, Y., Kuno, M., Kamiyama, T., Kozaki, K., Nomura, N., Egawa, H., Minami, S., Watanabe, Y., Narita, H., Shiraki, K., 2002. In vitro and in vivo activities of anti-influenza virus compound T-705. *Antimicrob. Agents Chemother.* 46, 977–981. <https://doi.org/10.1128/AAC.46.4.977-981.2002>.
- Haviernik, J., Štefánik, M., Fojtková, M., Kali, S., Tordo, N., Rudolf, I., Hubálek, Z., Eyer, L., Ruzek, D., 2018. Arbidol (umifenovir): a broad-spectrum antiviral drug that inhibits medically important arthropod-borne flaviviruses. *Viruses* 10, 184. <https://doi.org/10.3390/v10040184>.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H., Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., Cao, B., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet Lond. Engl.* 395, 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- Ianevski, A., He, L., Aittokallio, T., Tang, J., 2017. SynergyFinder: a web application for analyzing drug combination dose–response matrix data. *Bioinformatics* 33, 2413–2415. <https://doi.org/10.1093/bioinformatics/btx162>.
- Liu, Q., Fang, X., Tian, L., Chen, X., Chung, U., Wang, K., Li, D., Dai, X., Zhu, Q., Xu, F., Shen, L., Wang, B., Yao, L., Peng, P., 2020a. The effect of Arbidol Hydrochloride on reducing mortality of Covid-19 patients: a retrospective study of real world data from three hospitals in Wuhan. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.11.20056523>, 04.11.20056523.
- Liu, Y., Yan, L.-M., Wan, L., Xiang, T.-X., Le, A., Liu, J.-M., Peiris, M., Poon, L.L.M., Zhang, W., 2020b. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect. Dis.* 20 (6), 656–657. [https://doi.org/10.1016/S1473-3099\(20\)30232-2](https://doi.org/10.1016/S1473-3099(20)30232-2).
- Magagnoli, J., Narendran, S., Pereira, F., Cummings, T., Hardin, J.W., Sutton, S.S., Ambati, J., 2020. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.16.20065920>, 04.16.20065920.
- Mehta, P., McAuley, D.F., Brown, M., Sanchez, E., Tattersall, R.S., Manson, J.J., 2020. COVID-19: consider cytokine storm syndromes and immunosuppression. *The Lancet* 395 (10229), 1033–1034. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](https://doi.org/10.1016/S0140-6736(20)30628-0).
- Mulangu, S., Dodd, L.E., Davey, R.T., Tshiani Mbaya, O., Proschan, M., Mukadi, D., Lusakibanza Manzo, M., Nzolo, D., Tshomba Oloma, A., Ibanda, A., Ali, R., Coulibaly, S., Levine, A.C., Grais, R., Diaz, J., Lane, H.C., Muyembe-Tamfum, J.-J., the Palm Writing Group, 2019. A randomized, controlled trial of Ebola virus disease therapeutics. *N. Engl. J. Med.* 381, 2293–2303. <https://doi.org/10.1056/NEJMoa1910993>.
- Pizzorno, A., Padey, B., Julien, T., Trouillet-Assant, S., Traversier, A., Errazuriz-Cerda, E., Fouret, J., Dubois, J., Gaymard, A., Lescure, F.-X., Dulière, V., Brun, P., Constant, S., Poissy, J., Lina, B., Yazdanpanah, Y., Terrier, O., Rosa-Calatrava, M., 2020. Characterization and treatment of SARS-CoV-2 in nasal and bronchial human airway epithelia. *bioRxiv* 2020. <https://doi.org/10.1101/2020.03.31.017889>, 03.31.017889.
- Pizzorno, A., Terrier, O., Nicolas de Lamballerie, C., Julien, T., Padey, B., Traversier, A., Roche, M., Hamelin, M.-E., Rhéaume, C., Croze, S., Escuret, V., Poissy, J., Lina, B., Legras-Lachuer, C., Textoris, J., Boivin, G., Rosa-Calatrava, M., 2019. Repurposing of drugs as novel influenza inhibitors from clinical gene expression infection signatures. *Front. Immunol.* 10, 60. <https://doi.org/10.3389/fimmu.2019.00060>.
- Prescott, J., Hall, P., Acuna-Retamar, M., Ye, C., Wathelet, M.G., Ebihara, H., Feldmann, H., Hjelle, B., 2010. New world hantaviruses activate IFN λ production in type I IFN-deficient Vero E6 cells. *PloS One* 5, e11159. <https://doi.org/10.1371/journal.pone.0011159>.
- Sidwell, R.W., Huffman, J.H., Khare Lois, G.P., Allen, B., Witkowski Roland, J.T., Robins, K., 1972. Broad-spectrum antiviral activity of virazole: 1- β -D-ribofuranosyl-1,2,4-triazole-3-carboxamide. *Science* 177, 705–706. <https://doi.org/10.1126/science.177.4050.705>.
- Solidarity, 2020. Clinical trial for COVID-19 treatments [WWW Document]. accessed 4.27.20. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/global-research-on-novel-coronavirus-2019-ncov/solidarity-clinical-trial-for-covid-19-treatments>.
- Varghese, F.S., Kaukinen, P., Gläsker, S., Bespalov, M., Hanski, L., Wennerberg, K., Kümmerer, B.M., Ahola, T., 2016a. Discovery of berberine, abamectin and ivermectin as antivirals against chikungunya and other alphaviruses. *Antivir. Res.* 126, 117–124. <https://doi.org/10.1016/j.antiviral.2015.12.012>.
- Varghese, F.S., Thaa, B., Amrun, S.N., Simarmata, D., Rausalu, K., Nyman, T.A., Merits, A., McInerney, G.M., Ng, L.F.P., Ahola, T., 2016b. The antiviral alkaloid berberine reduces chikungunya virus-induced mitogen-activated protein kinase signaling. *J. Virol.* 90, 9743–9757. <https://doi.org/10.1128/JVI.01382-16>.
- Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., Shi, Z., Hu, Z., Zhong, W., Xiao, G., 2020. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 30, 269–271. <https://doi.org/10.1038/s41422-020-0282-0>.
- Yan, Y.-Q., Fu, Y.-J., Wu, S., Qin, H.-Q., Zhen, X., Song, B.-M., Weng, Y.-S., Wang, P.-C., Chen, X.-Y., Jiang, Z.-Y., 2018. Anti-influenza activity of berberine improves prognosis by reducing viral replication in mice. *Phytother. Res.* PTR 32, 2560–2567. <https://doi.org/10.1002/ptr.6196>.
- Zheng, S., Fan, J., Yu, F., Feng, B., Lou, B., Zou, Q., Xie, G., Lin, S., Wang, R., Yang, X., Chen, W., Wang, Q., Zhang, D., Liu, Y., Gong, R., Ma, Z., Lu, S., Xiao, Y., Gu, Y., Zhang, J., Yao, H., Xu, K., Lu, X., Wei, G., Zhou, J., Fang, Q., Cai, H., Qiu, Y., Sheng, J., Chen, Y., Liang, T., 2020. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January–March 2020: retrospective cohort study. *BMJ* 369, m1443. <https://doi.org/10.1136/bmj.m1443>.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G.F., Tan, W., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* 382, 727–733. <https://doi.org/10.1056/NEJMoa2001017>.