

Risk Factors for Suicidal Ideation of the General Population

Seung-Min Bae,¹ Yu Jin Lee,¹
In Hee Cho,¹ Seog Ju Kim,²
Jeong Soo Im,³ and Seong-Jin Cho¹

¹Department of Psychiatry, Gachon University Gil Medical Center, Incheon; ²Department of Medicine, Seoul National University College of Medicine, Seoul; ³Department of Preventive Medicine, Gachon University, Incheon, Korea

Received: 20 October 2012
Accepted: 30 January 2013

Address for Correspondence:

Yu Jin Lee, MD
Department of Psychiatry, Gachon University Gil Medical Center,
21 Namdong-daero 774-beon-gil, Namdong-gu, Incheon
405-760, Korea
Tel: +82.32-468-9932, Fax: +82.32-468-9962
E-mail: leeyj1203@gmail.com

This work was supported by the Gachon University Gil Medical Center (Grant number: 2012-01).

The aim of this study was to investigate risk factors of suicide ideation in general population. A total of 1,116 adults were sampled with randomization in an urban area. After excluding 116 participants due to incomplete answer, 1,000 participants (500 males; mean age 39.6 ± 11.6) completed self-report questionnaire including the Center for Epidemiologic Study Depression Scale (CES-D), the Beck Scale for Suicide Ideation (BSI), the Spielberger State-Trait Anger Expression Inventory (STAXI), the Barratt Impulsiveness Scale (BIS), and questions on weekday sleep duration. Results showed higher scores on the BSI were related with higher scores on the STAXI, CES-D, a family history of psychiatric illness, and short or long sleep duration (≤ 5 or ≥ 10 hr) (adjusted $R^2 = 0.151$, $P < 0.001$, $P < 0.001$, $P = 0.001$, and $P = 0.008$, respectively). The predictive power of the STAXI scores, the presence of psychiatric family history and short or long sleep duration was stronger in the high CES-D group with scores of 16 or higher (adjusted $R^2 = 0.275$, $P < 0.001$, $P < 0.001$, and $P < 0.001$, respectively). These findings suggest that suicide idea in general population may be related with the presence of family history for psychiatric illness, depressive mood, high anger and short or long sleep duration.

Key Words: General Population; Risk Factor; Suicide Idea

INTRODUCTION

Recently, suicide is a major mental health issue in Korea (1). Suicide is the fourth highest cause of death in Korea with an average of 42 incidents a day (2). The identification of the risk factors of suicide has therefore become a crucial target in health policy.

A number of studies have examined suicide risk factors, prevention and the ways of reducing suicide-related behaviors. Previous studies have shown that suicide is the result of a multifaceted range of social, psychiatric and biological factors, and one or more of these variables may affect suicidal behaviors by complex mechanisms (3-5). Although there are many studies of suicidality and suicide risk factors in people with various mental disorders (6, 7), the findings of these researches may not necessarily generalize to the general population.

The problem of suicide is no longer limited to some psychiatric patients, and consequently the assessment of suicidality and preparing for its appropriate management in the community has important clinical implications (8). For the sake of the mental health of the community, it is therefore necessary to conduct a study of risk factors for suicide idea in the general population sample at large rather than just on those psychiatric patients receiving hospital inpatient or outpatient services. This will enable us to plan suicide prevention strategies not only for psychiatric patients, but also for people in the general community,

who do not visit mental health services.

Thus, the main purpose of the present study was to assess suicidal ideation and related factors in a community sample, which was representative of the general population in an urban area of Korea.

MATERIAL AND METHODS

Subjects

The current study was conducted as a part of the community mental health survey in Incheon, a metropolitan city in Korea. The subjects (sample size: 1,116 participants) were sampled with randomization and recruited by a research agency from all eight administrative districts of the city. The number of subjects recruited from each district was proportional to the relative population size of the district. For each district, equal numbers of males and females and equal numbers of subjects in four different age-groups (the 20s, 30s, 40s, and 50s) were sampled. Fifteen trained surveyors made contact with the subjects. They were educated 2-hr about content of self-report questionnaires used in this study and basic psychiatric concepts including depression and suicide before the interview. The final enrollment comprised a representative population sample of 1,000 people aged over 20 yr (Table 1; 500 males and 500 females; mean age of total sample 39.6 ± 11.6 yr ranging 20-59 yr). All participants completed self-report questionnaires including sociodemo-

Table 1. Comparison of sociodemographic, psychological characteristics and sleep duration between low CES-D and high CES-D groups (n = 1,000)

Parameters	No. (%) of subjects	Low CES-D	High CES-D	χ^2/t	P
		(n = 706)	(n = 294)		
		No. (%)	No. (%)		
Mean age (yr)	39.6 ± 11.6	38.7 ± 11.39	41.8 ± 12.0	-3.814	< 0.001
Sex				4.933	0.026
Male	500 (50)	369 (52.3)	131 (44.6)		
Female	500 (50)	337 (47.7)	163 (55.4)		
Family income (US dollars per month)				2.834	0.726
Below 1,800	48 (4.8)	36 (5.1)	12 (4.1)		
1,800-2,700	170 (17.0)	123 (17.4)	47 (16.0)		
2,700-3,150	318 (31.8)	219 (31.0)	99 (33.7)		
3,150-3,600	257 (25.7)	176 (24.9)	81 (27.6)		
3,600-4,500	148 (14.8)	107 (15.2)	41 (13.9)		
More than 4,500	59 (5.9)	45 (6.4)	14 (4.8)		
Presence of FHx* of psychiatric disorder				23.369	< 0.001
Yes	952 (95.2)	19 (2.7)	29 (9.9)		
No	48 (4.8)	687 (97.3)	265 (90.1)		
Education				7.653	0.176
No education	3 (0.3)	2 (0.3)	1 (0.3)		
Elementary school	14 (1.4)	8 (1.1)	6 (2.0)		
Middle school	43 (4.3)	27 (3.8)	16 (5.4)		
High school	576 (57.6)	398 (56.4)	178 (60.5)		
College	358 (35.8)	265 (37.5)	93 (31.6)		
Graduate school	6 (0.6)	6 (0.8)	0 (0.0)		
Marital status				4.196	0.041
Married	269 (26.9)	503 (71.2)	228 (77.7)		
Unmarried, separated, or divorced	731 (73.1)	203 (28.8)	66 (22.4)		
Presence of religion				12.862	< 0.001
Yes	559 (55.9)	369 (52.3)	190 (64.6)		
No	441 (44.1)	337 (47.7)	14 (35.4)		
CES-D (mean ± SD)	11.6 ± 7.3	7.8 ± 4.4	20.6 ± 4.6	-40.720	< 0.001
BSI (mean ± SD)	3.1 ± 3.8	3.4 ± 4.8	6.3 ± 6.5	-7.148	< 0.001
STAXI (mean ± SD)	34.5 ± 8.1	33.0 ± 7.1	38.1 ± 9.0	-8.648	< 0.001
BIS (mean ± SD)	61.1 ± 9.3	59.7 ± 9.2	64.4 ± 8.7	-7.634	< 0.001
Total sleep time (mean ± SD)	7.2 ± 0.9	7.2 ± 0.9	7.2 ± 0.9	0.657	0.511
Abnormal sleep duration (≤ 5 hr or ≥ 10 hr)	164 (16.4)	110 (15.6)	54 (18.4)	1.176	0.278

Low CES-D, CES-D score < 16; High CES-D, CES-D score ≥ 16. *Family History, any diagnosed psychiatric disorder within second degree relatives. FHx, family history; SD, standard deviation; CES-D, Center for Epidemiologic Study-Depression; BSI, Beck Scale for Suicide Ideation; STAXI, Spielberger State-Trait Anger Expression Inventory; BIS, Barratt Impulsiveness Scale.

graphic information, the Center for Epidemiologic Study Depression Scale (CES-D), the Beck Scale for Suicide Ideation (BSI), the Spielberger State-Trait Anger Expression Inventory (STAXI), the Barratt Impulsiveness Scale (BIS) and questions on sleep duration on weekdays.

Measures

Depressive symptoms were assessed with the Korean version of the CES-D (9), which was originally developed for use as a screening instrument to identify people at risk for clinical depression (10). The scale includes 20 items, which measure the presence and severity of depressive symptoms over the preceding seven days. The response options are four-point Likert scales, with scores ranging 0-3 and anchor points in terms of the number days per week each symptom was experienced: 'rarely or none of the time (less than one day)' to 'most or all of the time (5-7 days)'. The total score ranges from 0 to 60, with a higher score indicating a higher level of depressive symptoms. The scale has

been shown to be reliable and valid in both community and psychiatric samples and a score of 16 or more on the CES-D is generally accepted as an indicative of a clinically significant level of depressive symptoms (10, 11). This cut-off score was used to divide the total sample into high and low depression groups in this study.

The BSI was originally to assess the strength of a person's wishes and plans to commit suicide completed by a clinician using a semi-structured interview (12), modified into a self-report scale with 19 items in Korea (13). The Korean version of the BSI showed acceptable reliability and validity (13). The items are rated on a 3-point scale from 0 to 2. The total score can range from 0 to 38, with higher scores indicating higher suicidal ideation.

The Korean version of STAXI is a self-report scale with 44 items designed to evaluate an individual's anger level. The scale was originally developed by Spielberger (14) and translated into Korean in 1997 (15). Each item is essentially a statement people

use to describe their emotions when they feel angry. Each item is assessed by a four point scale (1 = no anger; 4 = maximum anger). The Korean version of the STAXI has been shown to have satisfactory test-retest reliability, internal consistency and validity (15). The score is considered to correlate positively with anger level in subjects.

The BIS was developed by Barrett (16) and is a scale with 23 items selected to measure impulsiveness. This scale has been widely-used as a trait-based, self-report measure of impulsivity. It was translated into the Korean by Lee in 1992, and this version has been found to have adequate reliability and validity (17). Higher scores indicate higher impulsivity.

Statistical analysis

Group differences were tested using either chi-square or Student's *t*, whichever was appropriate. Partial correlation coefficients were calculated for the correlation analysis. In the stepwise linear regression analysis, the dependent variable was the BSI score and the independent variables were factors known on the basis of past research to be associated with increased risk of suicide (18-20). These were age, gender, level of family income, a family history of psychiatric illness, marital status, education, religious belief, and sleep duration. The data were analyzed using SPSS 15.0 (SPSS Inc, Chicago, IL, USA). Statistical significance was defined as $P < 0.05$ (two-tailed).

Ethics statement

The study protocol was approved by the institutional review board of Gachon University of Medicine and Science (GIRBA-1785). All participants were fully informed about the aims and methods of the study prior to starting the interview and informed consent was obtained.

RESULTS

Table 1 summarizes the sociodemographic, psychological characteristics, and sleep durations of total participants. Two hundred ninety four participants (29.4%) showed significant depressive symptom, defined as high CES-D group (≥ 16) as the previous study (9). Participants with the CES-D score less than

16 were classified as low CES-D group (Table 1). Comparisons between the high and low CES-D groups are summarized in Table 1. Compared with the low CES-D group, the high CES-D group was significantly older (mean ages: 41.8 ± 12.0 vs 38.7 ± 11.39 , $t = -3.814$, $P < 0.001$), had a higher ratio of females (55.4% vs 47.7%, $\chi^2 = 4.933$, $P = 0.026$), were more likely to have a family history of psychiatric disorder (9.9% vs 2.7%, $\chi^2 = 23.369$, $P < 0.001$), were more likely to be married (77.7% vs 71.2%, $\chi^2 = 4.196$, $P = 0.041$), and were more religious (64.6% vs 52.3%, $\chi^2 = 12.862$, $P < 0.001$). The high CES-D group also scored higher on the BSI (6.3 ± 6.5 vs 3.4 ± 3.8 , $t = -7.148$, $P < 0.001$), the STAXI (38.1 ± 9.0 vs 33.0 ± 7.1 , $t = -8.648$, $P < 0.001$) and the BIS (64.4 ± 8.7 vs 59.7 ± 9.2 , $t = -7.634$, $P < 0.001$).

The results of the partial correlation analysis are shown in Table 2. After adjusting for age and gender, the scores on the BSI were found to be positively correlated with the CES-D, STAXI and BIS scores ($r = 0.245$; $P < 0.001$, $r = 0.344$; $P < 0.001$, and $r = 0.152$; $P < 0.001$, respectively). The BSI scores showed no significant correlation with sleep duration on weekday ($r = -0.044$, $P = 0.167$).

The International Classification of Sleep Disorders (21) defines a 'short sleeper' as someone who sleeps five hours or less a day and a 'long sleeper' as someone sleeping 10 hours or more a day. In the present study, we classified short and long sleepers into an abnormal sleep duration group. The regression analysis revealed that higher scores on the BSI were related with higher scores on the STAXI and on the CES-D, the presence of a family history of psychiatric illness and abnormal sleep duration (Table 3; $F = 45.449$, adjusted $R^2 = 0.151$, $P < 0.001$, $P < 0.001$, $P = 0.001$, and $P = 0.008$, respectively). The regression model with high CES-D group showed stronger predictive power, in which higher scores on the BSI were associated with higher scores on the STAXI, an abnormal sleep duration and the presence of a family history of psychiatric illness (Table 4; $F = 38.092$, adjusted $R^2 = 0.275$, $P < 0.001$, $P < 0.001$, and $P < 0.001$, respectively).

DISCUSSION

This study investigated the relationships between suicide ideation

Table 2. Partial correlations controlling for age and gender between suicidal ideation and other variables ($n = 1,000$)

Variables					
	CES-D	STAXI	BIS	Sleep duration	BSI
CES-D	1.00				
STAXI	0.331*	1.00			
BIS	0.230*	0.425*	1.00		
Sleep duration	-0.044	0.009	-0.045	1.00	
BSI	0.245*	0.344*	0.152*	-0.044	1.00

* $P < 0.001$. CES-D, Center for Epidemiologic Study-Depression; STAXI, Spielberger State-Trait Anger Expression Inventory; BIS, Barratt Impulsiveness Scale; BSI, Beck Scale for Suicide Ideation.

Table 3. Stepwise linear regression model predicting suicidal ideation in total sample ($n = 1,000$)

Model	Coefficients			t-value	P value
	B	SE	Beta		
STAXI	0.195	0.021	0.286	9.195	< 0.001
CES-D	0.102	0.023	0.136	4.386	< 0.001
Family history of psychiatric disorder	2.495	0.763	0.097	3.270	0.001
Abnormal sleep duration (≤ 5 hr or ≥ 10 hr)	-1.154	0.434	-0.078	-2.659	0.008

$F = 45.449$, adjusted $R^2 = 0.151$. SE, standard error; STAXI, Spielberger State-Trait Anger Expression Inventory; CES-D, Center for Epidemiologic Study-Depression; Abnormal sleep duration, ≤ 5 hr or ≥ 10 hr.

Table 4. Stepwise linear regression model predicting higher suicidal ideation in high CES-D group (n = 294)

Model	Coefficients			t-value	P value
	B	SE	Beta		
STAXI	0.271	0.036	0.376	7.464	< 0.001
Abnormal sleep duration (≤ 5 hr or ≥ 10 hr)	4.106	0.831	0.246	4.942	< 0.001
Family history of psychiatric disorder	2.759	0.773	0.107	3.568	< 0.001

F = 38.092, adjusted $R^2 = 0.275$. High CES-D group, CES-D score ≥ 16 . SE, standard error; CES-D, Center for Epidemiologic Study-Depression; STAXI, Spielberger State-Trait Anger Expression Inventory; abnormal sleep duration, ≤ 5 hr or ≥ 10 hr.

ation and associated factors in a community population in an urban area of Korea. The results showed that subjective depressive mood, high anger, abnormal sleep duration and a family history of psychiatric illness were related to suicide idea in general population. These relationships were more prominent in the high CES-D group.

The mean CES-D score for the total participants was 11.6 ± 7.3 , and 29.4% showed clinically significant levels of depressive symptoms (scores of 16 or higher in CES-D), and this finding is consistent with those reported in previous studies (22). Just as others have done, we proposed the relationship between depression and suicide ideation. From the study done in the past (20), 95% of people who commit suicide have a diagnosis of mental disorder, and 80% of those with a mental disorder are diagnosed as suffering from depression.

In current study, a positive correlation of anger with suicide idea was also found. Previous studies have demonstrated possible relationships between higher anger and suicidality (23, 24). The present study increases our knowledge about the relationship between anger and suicide idea. In Asian culture, the expression of anger tends to be usually discouraged. This represents a major challenge to Koreans who experience anger, but feel pressure to express and manage the emotion in culturally acceptable ways. Lee et al. (23) showed that anger is an important marker for people, who may be at risk of engaging in suicidal behavior. A similar finding was also reported in a Turkish study (24), in which suppressed anger was shown to increase suicidal thoughts. In the present study, intense feelings and tendency to suppress anger might increase suicidal thought.

Previous studies have shown that individuals who suffer from insomnia or sleep disturbance could be at greater risk of depression and committing suicide (25-27). In the current study, we conducted the question on sleep duration which is a feasible additional risk factor for suicide. While many studies have shown that sleep disorders are highly associated with suicide (28-31), no study has been conducted on the association between suicide idea and sleep duration. Looking at the studies of relativity of sleep only commented relation of insomnia or subjective sleep problem and suicide idea done in the past (30-33), it has never used parametric measure of suicidal ideation with

scales. In this study, abnormal sleep duration (five hours or less or 10 hr or more) was related with higher suicidal ideation.

There was a study on the association of hypersomnia with suicidality the past (27), but the previous study only focused on the increasing rate of suicides in depressed patients with hypersomnia. To the best of knowledge, this is the first study which showed the relationship between long sleep duration and suicide idea. One possible explanation of this association between long sleep duration and suicide idea is that hypersomnia may be mainly related to depression with atypical features (20). Another possible reason is that hypersomnia or long sleep can be associated with sleep disorders, such as sleep apnea and narcolepsy, which are known to be related to depression (30, 31). Moreover, the finding that sleep disorders such as nightmare's can be related to increased suicidality after controlling for depression (28) suggests that suicide may be directly related to sleep disturbances and not always mediated by depression.

In current study, the presence of a family history of psychiatric illness (within their second-degree relatives) was related to higher suicide idea. Psychiatric patients with history of psychiatric illness in their biological family are known to respond less well to treatment, leading to an unfavorable prognosis (34). A feasible explanation is that this unfavorable treatment response can be related to higher suicidality, and the other interpretation imply that suicide can be significantly controlled by biological and genetic factors. This common notion is supported by a number of research findings suggesting biological mechanisms of suicide such as genetic factors (35, 36). In addition, Roy and Segal (37) found higher rates of concordance for suicidality in monozygotic than in dizygotic twins.

Impulsiveness measured by BIS scores used in the present study, did not predict high suicide ideation. In a past research, impulsiveness has been considered to be one of the important factors associated with suicide (38). One possible explanation for this inconsistency is that we related impulsiveness to suicidal ideation rather than actual attempted suicide. This finding suggests that impulsiveness may be strongly related to suicidal behaviors rather than suicide thoughts. More research is needed before any firm conclusions can be drawn about the relationships between impulsiveness and suicidality.

The present study has several limitations. Firstly, the methodology involved subjects completing self-report questionnaires without structured or semi-structured interviews. This precluded an objective assessment of subject's psychiatric status. Secondly, we relied entirely on measure of sleep duration, which was also self-reported. This made it impossible to identify secondary symptoms of sleep disorders may have been related to suicidal ideation. Thirdly, the present study design was cross-sectional, and hence limits our ability to assess causal associations between suicidal ideation and associated risk factors. Thus longitudinal studies are needed to clarify causality.

In spite of several limitations, the current study is meaningful, because it assessed risk factors for high suicide idea in a general urban population using representative samples randomly recruited from city districts and matched equally in terms of gender and age group.

In conclusion, our results suggest that depressive mood, high anger, short or long sleep duration and a family history of psychiatric illness might be risk factors for suicide idea in the general population. Therefore, mental health professionals might pay special attention not only to one's subjective mood state (including depressiveness and anger), also sleep duration and family history of psychiatric illness to evaluate suicide risk factors in a community.

ACKNOWLEDGMENTS

The authors have no conflicts of interest to disclose.

REFERENCES

- Jeon HJ, Lee JY, Lee YM, Hong JP, Won SH, Cho SJ, Kim JY, Chang SM, Lee D, Lee HW, et al. *Lifetime prevalence and correlates of suicidal ideation, plan, and single and multiple attempts in a Korean nationwide study*. *J Nerv Ment Dis* 2010; 198: 643-6.
- Organisation for Economic Co-operation and Development. *The OECD's online Library*. Available at <http://titania.Sourceoecd.Org/vl=1337221/cl=19/nw=1/rpsv/health2007/2-7.Htm> [accessed on 11 April 2009].
- Lewinsohn PM, Rohde P, Seeley JR. *Psychosocial risk factors for future adolescent suicide attempts*. *J Consult Clin Psychol* 1994; 62: 297-305.
- Mann JJ, Brent DA, Arango V. *The neurobiology and genetics of suicide and attempted suicide: a focus on the serotonergic system*. *Neuropsychopharmacology* 2001; 24: 467-77.
- Rowe JL, Bruce ML, Conwell Y. *Correlates of suicide among home health care utilizers who died by suicide and community controls*. *Suicide Life Threat Behav* 2006; 36: 65-75.
- Agargun MY, Besiroglu L, Cilli AS, Gulec M, Aydin A, Inci R, Selvi Y. *Nightmares, suicide attempts, and melancholic features in patients with unipolar major depression*. *J Affect Disord* 2007; 98: 267-70.
- Schmidt NB, Woolaway-Bickel K, Bates M. *Evaluating panic-specific factors in the relationship between suicide and panic disorder*. *Behav Res Ther* 2001; 39: 635-49.
- Borges G, Angst J, Nock MK, Ruscio AM, Walters EE, Kessler RC. *A risk index for 12-month suicide attempts in the National Comorbidity Survey Replication (NCS-R)*. *Psychol Med* 2006; 36: 1747-57.
- Cho MJ, Kim KH. *Use of the Center for Epidemiologic Studies Depression (CES-D) Scale in Korea*. *J Nerv Ment Dis* 1998; 186: 304-10.
- Radloff LS. *The CES-D scale: a self-report depression scale for research in the general population*. *Appl Psychol Meas* 1977; 1: 385-401.
- Beekman AT, Deeg DJ, Van Limbeek J, Braam AW, De Vries MZ, Van Tilburg W. *Criterion validity of the Center for Epidemiologic Studies Depression scale (CES-D): results from a community-based sample of older subjects in the Netherlands*. *Psychol Med* 1997; 27: 231-5.
- Beck AT, Kovacs M, Weissman A. *Assessment of suicidal intention: the Scale for Suicide Ideation*. *J Consult Clin Psychol* 1979; 47: 343-52.
- Shin MS, Park KB, Oh KJ, Kim ZS. *A study of suicidal ideation among high school students: the structural relation among depression, hopelessness, and suicidal ideation*. *Korean J Clin Psychol* 1990; 9: 1-19.
- Spielberger CD. *Manual for the State-Trait Anxiety Inventory*. Palo Alto: Consulting Psychologists Press, 1983.
- Chon KK, Hahn DW, Lee CH, Spielberger CD. *Korean adaptation of the state-trait anger expression inventory: anger and blood pressure*. *Korean J Clin Psychol* 1997; 2: 60-78.
- Barratt ES. *Factor analysis of some psychometric measures of impulsiveness and anxiety*. *Psychol Rep* 1965; 16: 547-54.
- Lee H. *Guidance of impulsivity test*. Seoul: Korean Guidance, 1992.
- Foster T, Gillespie K, McClelland R, Patterson C. *Risk factors for suicide independent of DSM-III-R Axis I disorder: case-control psychological autopsy study in Northern Ireland*. *Br J Psychiatry* 1999; 175: 175-9.
- Gould MS, Fisher P, Parides M, Flory M, Shaffer D. *Psychosocial risk factors of child and adolescent completed suicide*. *Arch Gen Psychiatry* 1996; 53: 1155-62.
- Sadock BJ, Kaplan HI, Sadock VA. *Mood disorder*. In: Sadock BJ, Sadock VA, editors. *Kaplan & Sadock's synopsis of psychiatry: behavioral sciences/clinical psychiatry*. 10th ed. Philadelphia: Wolter Kluwer/Lippincott Williams & Wilkins, 2007, p527-8.
- American Academy of Sleep Medicine. *International classification of sleep disorders, revised: diagnostic and coding manual*. Chicago: American Academy of Sleep Medicine, 2001.
- Cho SJ, Jeon HJ, Kim MJ, Kim JK, Kim US, Lyoo IK, Cho MJ. *Prevalence and correlates of depressive symptoms among the adolescents in an urban area in Korea*. *J Korean Neuropsychiatr Assoc* 2001; 40: 627-39.
- Lee J, Choi H, Kim MJ, Park CG, Shin DS. *Anger as a predictor of suicidal ideation in middle-school students in Korea: gender difference in threshold point*. *Adolescence* 2009; 44: 433-46.
- Engin E, Gurkan A, Dulgerler S, Arabaci LB. *University students' suicidal thoughts and influencing factors*. *J Psychiatr Ment Health Nurs* 2009; 16: 343-54.
- Krakov B, Artar A, Warner TD, Melendrez D, Johnston L, Hollifield M, Germain A, Koss M. *Sleep disorder, depression, and suicidality in female sexual assault survivors*. *Crisis* 2000; 21: 163-70.
- Ağargün MY, Kara H, Solmaz M. *Sleep disturbances and suicidal behavior in patients with major depression*. *J Clin Psychiatry* 1997; 58: 249-51.
- Fawcett J, Scheftner WA, Fogg L, Clark DC, Young MA, Hedeker D, Gibbons R. *Time-related predictors of suicide in major affective disorder*. *Am J Psychiatry* 1990; 147: 1189-94.
- Bernert RA, Joiner TE Jr, Cukrowicz KC, Schmidt NB, Krakow B. *Suicidality and sleep disturbances*. *Sleep* 2005; 28: 1135-41.
- Bernert RA, Joiner TE. *Sleep disturbances and suicide risk: a review of the literature*. *Neuropsychiatr Dis Treat* 2007; 3: 735-43.
- Harris M, Glozier N, Ratnavadivel R, Grunstein RR. *Obstructive sleep apnea and depression*. *Sleep Med Rev* 2009; 13: 437-44.
- Mamelak M. *Narcolepsy and depression and the neurobiology of gamma-hydroxybutyrate*. *Prog Neurobiol* 2009; 89: 193-219.
- Sabo E, Reynolds CF 3rd, Kupfer DJ, Berman SR. *Sleep, depression, and suicide*. *Psychiatry Res* 1991; 36: 265-77.
- Ağargün MY, Kara H, Solmaz M. *Subjective sleep quality and suicidality in patients with major depression*. *J Psychiatr Res* 1997; 31: 377-81.
- Sarchiapone M, Carli V, Janiri L, Marchetti M, Cesaro C, Roy A. *Family*

- history of suicide and personality. Arch Suicide Res 2009; 13: 178-84.*
35. Viana MM, De Marco LA, Boson WL, Romano-Silva MA, Corrêa H. *Investigation of A218C tryptophan hydroxylase polymorphism: association with familial suicide behavior and proband's suicide attempt characteristics. Genes Brain Behav 2006; 5: 340-5.*
36. Mahon PB, Stütz AM, Seifuddin F, Huo Y, Goes FS, Jancic D, Judy JT, Depaulo JR Jr, Gershon ES, McMahon FJ, et al. *Case-control association study of TGOLN2 in attempted suicide. Am J Med Genet B Neuropsychiatr Genet 2010; 153B: 1016-23.*
37. Roy A, Segal NL. *Suicidal behavior in twins: a replication. J Affect Disord 2001; 66: 71-4.*
38. Dougherty DM, Mathias CW, Marsh-Richard DM, Prevette KN, Dawes MA, Hatzis ES, Palmes G, Nouvion SO. *Impulsivity and clinical symptoms among adolescents with non-suicidal self-injury with or without attempted suicide. Psychiatry Res 2009; 169: 22-7.*