




# Streptococcus salivarius K12 Alleviates Oral Mucositis in Patients Undergoing Radiotherapy for Malignant Head and Neck Tumors: A Randomized Controlled Trial

Xingchen Peng, MD<sup>1</sup> ; Zixia Li, MDS<sup>2</sup>; Yiyan Pei, MM<sup>1</sup> ; Shuhao Zheng, DDS<sup>2</sup>; Jinchi Liu, MDS<sup>2</sup>; Jingjing Wang, MD<sup>1</sup>; Ruidan Li, MD<sup>1</sup>; and Xin Xu, DDS<sup>2</sup> 

DOI <https://doi.org/10.1200/JCO.23.00837>

## ABSTRACT

**PURPOSE** Oral mucositis (OM) is a common debilitating toxicity associated with radiotherapy (RT) for malignant head and neck tumors. This prospective, randomized, double-blind, placebo-controlled trial aimed to evaluate the efficacy and safety of *Streptococcus salivarius* K12 (SsK12) in reducing the incidence, duration, and severity of severe OM (SOM).

**METHODS** A total of 160 patients with malignant head and neck tumors undergoing definitive or postoperative adjuvant RT were randomly assigned (1:1) to receive SsK12 probiotic (n = 80) or placebo (n = 80) at West China Hospital, Sichuan University, Chengdu, China. Patients were instructed to suck SsK12 or placebo lozenges thrice daily from the initiation to the end of RT. OM was evaluated twice a week during RT and once a week thereafter for up to 8 weeks. The primary end point was the incidence of SOM. Adverse events were assessed according to the National Cancer Institute Common Terminology Criteria for Adverse Events, version 5.0.

**RESULTS** Baseline patient characteristics were similar in the SsK12 and placebo groups. The incidence of SOM was significantly lower in the SsK12 group as compared with the placebo group (36.6% v 54.2%;  $P = .0351$ ). The duration (median, 0.0 days v 7.0 days; mean, 8.9 days v 18.3 days;  $P = .0084$ ) and time to develop SOM (median, not estimable v 42.0 days; hazard ratio, 0.55 [95% CI, 0.34 to 0.89]; log-rank test:  $P = .0123$ ) were also improved in the case of the SsK12 group. Adverse events were similar between the groups, and mild or moderate gastrointestinal reactions (flatulence or dyspepsia) associated with the lozenges were observed in two patients in the SsK12 group. High-throughput sequencing results indicated that SsK12 inhibited opportunistic pathogens and enriched oral commensals during RT.

**CONCLUSION** In this prospective, randomized clinical trial, SsK12 probiotic significantly reduced the incidence, onset, and duration of SOM with a good safety profile.

## ACCOMPANYING CONTENT

 Article, p. 1436

 Data Sharing Statement

 Data Supplement

 Protocol

Accepted November 6, 2023

Published January 12, 2024

J Clin Oncol 42:1426-1435

© 2024 by American Society of Clinical Oncology



[View Online Article](#)

Creative Commons Attribution  
Non-Commercial No Derivatives  
4.0 License

## INTRODUCTION

Radiotherapy (RT) is an important method of treatment for malignant tumors of the head and neck and can be used alone or in combination with chemotherapy as a radical or adjuvant therapy.<sup>1</sup> Despite improvements in RT equipment and techniques, various acute oral complications persist, including oral mucositis (OM), dry mouth, taste dysfunction, and oral infections.<sup>2</sup> OM is one of the most common acute radiation-related toxicities, and approximately 50%–70% of patients experience severe OM (SOM) defined by the WHO scale as grade 3–4.<sup>3–6</sup> The painful inflammation and ulceration associated with OM not only profoundly affect patients' ability

to eat, swallow, and speak but also decrease their tolerance to anticancer treatment, thereby impairing their quality of life (QoL) significantly and causing interruptions in their cancer treatment.<sup>7</sup> Although some clinical strategies for radiation-induced OM have been recommended by the Multinational Association of Supportive Care in Cancer and the International Society of Oral Oncology,<sup>8</sup> their efficacy and safety still need further clinical validation.

Recent evidence suggests the involvement of oral microbiota in radiation-induced OM, and modulation of oral microbiota is promising for the management of OM.<sup>9–11</sup> *Streptococcus salivarius* K12 (SsK12) is a commercially available oral

## CONTEXT

### Key Objective

Is probiotic *Streptococcus salivarius* K12 (SsK12) able to alleviate oral mucositis (OM) in patients undergoing radiotherapy (RT) for malignant head and neck tumors?

### Knowledge Generated

Topical application of SsK12 significantly reduces the incidence of severe OM (SOM) in patients undergoing RT as compared with those who receive placebo. The development of SOM is delayed, and its duration is also shortened after the administration of SsK12.

### Relevance (J.P.S. Knisely)

Head & neck cancer severe acute RT-associated OM was delayed in onset and decreased in incidence and duration by the use of a SsK12 probiotic in a prospective, randomized, double-blind clinical trial.\*

\*Relevance section written by JCO Associate Editor Jonathan P.S. Knisely, MD.

probiotic with strong oral colonization ability, bacteriocin-like inhibitory substance (BLIS)–producing capability, and immunomodulatory properties and has been used to treat oral candidiasis, pharyngitis, tonsillitis, halitosis, and otitis media.<sup>12–17</sup> More importantly, data from our recent animal study demonstrated that topical use of SsK12 ameliorates radiation-induced OM in mice by modulating the oral microbiota, mainly by suppressing oral anaerobes.<sup>18</sup> However, the efficacy of SsK12 for treating radiation-induced OM still requires validation through well-controlled clinical trials. In this study, a prospective randomized clinical trial was designed to evaluate the effect of SsK12 on SOM in patients undergoing RT for malignant head and neck tumors. The effects of SsK12 on other radiation-related oral complications such as dry mouth, gustatory function, and microbial diversity in saliva were also investigated.

## METHODS

### Study Design and Patients

This prospective, randomized, double-blind, placebo-controlled trial was conducted at the West China Hospital, Sichuan University. The local institutional review board and ethics committee approved the study protocol, and the study was registered at ClinicalTrials.gov (identifier: [NCT05918224](https://clinicaltrials.gov/ct2/show/study?term=NCT05918224)). Written informed consent was obtained from each participant and family member before enrollment.

Patients pathologically diagnosed with nonmetastatic head and neck malignant tumors, age 18–80 years, with an Eastern Cooperative Oncology Group performance status of  $\leq 2$ , and planning to receive definitive RT or postoperative adjuvant RT at a dose of 60–72 Gy with or without concurrent chemotherapy were eligible for this study. Exclusion criteria included patients with a history of allergy to probiotics or severe allergic constitution, use of antibiotics/antifungal drugs within 1 month or

antimicrobial mouthwash within 1 week before the study, poor oral hygiene and/or severe periodontal diseases, any previous RT to the head and neck region, and those deemed unsuitable for the study by the investigators (concomitant with any other severe diseases).

### Random Assignment and Marking

All patients were randomly divided in a 1:1 ratio into two groups to receive either the probiotic SsK12 or a placebo via a computer-generated random assignment list using a randomized permutation block design. Methods for random assignment and marking are detailed in the Data Supplement (online only) and Protocol.

### Interventions

Before the start of RT, all patients received instructions for maintaining oral hygiene, oral clinical examination, and treatment including dental filling, endodontic treatment, extraction of nonrestorable teeth, and nonsurgical periodontal treatment, if necessary. All recruited patients underwent image-guided RT at a daily dose of 1.8–2.2 Gy, five times per week (total, 6–6.5 weeks), thus receiving a cumulative tumor dose of 60–72 Gy. Clinical target volumes and organs at risk were delineated in accordance with the consensus guidelines.<sup>19,20</sup> The oral cavity was defined according to the recommendation in the study by Mir et al.<sup>21</sup> Concurrent chemotherapy consisted of cisplatin (100 mg/m<sup>2</sup>) administered once every 3 weeks.

SsK12 lozenges (NOW Foods, USA) contained at least  $1 \times 10^9$  colony-forming units of viable SsK12 cells as the active ingredient. The placebo lozenges contained sugar and starch as excipients in the active formulation. Patients were instructed to suck the SsK12 lozenges or placebo lozenges thrice daily enduring the entire RT period. The patients were

advised to avoid eating, drinking, and conducting any oral hygiene activities for at least 1 hour after using the lozenges.<sup>22</sup> The number of lozenges used each day was recorded in patients' diaries. During RT, supportive care such as normal oral rinses, topical analgesics, and nutritional support was allowed, whereas anti-inflammatory agents, antibiotics/antifungal agents, antimicrobial mouthwash, and other experimental systemic or topical pharmaceuticals or devices were not permitted.

## Assessment

OM was evaluated by trained investigators according to the WHO Oral Toxicity Scale (Data Supplement, Table S1). OM was evaluated twice weekly during RT and once a week thereafter for up to 8 weeks. During RT, patients reported mouth and throat soreness (MTS) scores and the degree of impact of MTS on oral activities via the weekly oral mucositis questionnaire. Measures taken to ensure the consistency and accuracy of clinical data collection are detailed in the Data Supplement and Protocol.

Other clinical parameters, including hyposalivation, gustatory function, QoL, adverse events, weight loss, use of analgesics, use of intravenous hydration or total parenteral nutrition, and RT break fractions, are evaluated as detailed in the Data Supplement and Protocol.

## Saliva Collection and Bioinformatics

Unstimulated whole saliva was collected at baseline (T0), during mid-term RT (T1, week 3), and at the end of RT (T2, week 6 or 6.5). Bacterial genomic DNA was extracted, and the hypervariable regions 3-4 (V3-V4) of the 16S rRNA gene

were amplified for high-throughput sequencing. Bioinformatics were performed using the online Majorbio Cloud Platform (Majorbio, Shanghai, China<sup>23</sup>). The Benjamini-Hochberg method was used to calculate the false-discovery rate for multiple comparisons, and  $P_{FDR} < .05$  was considered significant (see the Data Supplement and Protocol for details).


## Outcome

The primary end point was the incidence of SOM (WHO grade 3-4). The secondary end points included the duration of SOM, time to onset of SOM, incidence and duration of OM, time to onset of OM, number of RT interruptions, and safety. The exploratory assessments included the presence of hyposalivation, changes in gustatory function, average MTS scores, incidence of analgesic use, use of intravenous hydration or total parenteral nutrition, BMI, QoL, and salivary microbiota.

## Sample Size Estimation and Statistical Analysis

On the basis of the literature, the average incidence of SOM observed during head and neck RT is approximately 65%,<sup>4</sup> and we expected a 25% decrease in the incidence of SOM in the SsK12 group. Accordingly, 142 patients (71 patients per group) were needed (two-sided  $\alpha = .05$ ,  $1 - \beta = .8$ , 1:1 ratio), and considering a patient dropout rate of 10%, at least 158 patients (79 patients per group) were targeted for enrollment.

All patients who completed the study (per-protocol set [PPS]) and all randomly assigned patients (full analysis set [FAS]) were included in the efficacy analyses. The safety analyses



**FIG 1.** CONSORT diagram: patient random assignment. <sup>a</sup>Patients who discontinued the acute oral mucositis evaluation phase of the study. ITT, intent to treat; PPS, per-protocol set; SsK12, *Streptococcus salivarius* K12.

included all randomly assigned patients who received the study medication. The duration of SOM/OM was determined as detailed in the Data Supplement and Protocol. Time to develop SOM/OM was analyzed by the Kaplan–Meier method, and log-rank tests were used to compare the curves. Hazard ratio with 95% CI was estimated using a Cox proportional hazard model. Continuous variables of the different groups were analyzed using the t-test; for nonparametric data, the comparison was performed using the Wilcoxon rank-sum test. Categorical variables were compared using the  $\chi^2$  test or Fisher's exact test.  $P < .05$  was considered as significant. The SAS statistical package version 9.3 (SAS Institute, Cary, NC) was used for data analysis.

## RESULTS

A total of 160 eligible patients were enrolled in the study (Fig 1). All patients were included in the safety analysis, among which 143 patients (SsK12,  $n = 71$ ; placebo,  $n = 72$ ) were evaluated for efficacy. Seventeen patients (11%) who discontinued the study were excluded from the PPS analysis for different reasons listed in Figure 1. The characteristics of the 17 excluded patients are presented in the Data Supplement (Table S3). The baseline characteristics of patients, who were fully evaluated, were similar between the two groups (Table 1). A majority of patients had tumors in the nasopharynx (25.87%), followed by the oral cavity (17.48%) and oropharynx (13.99%). The mean dose to oral cavity was 42.1 Gy in the SsK12 group and 42.2 Gy in the placebo group ( $P > .05$ ; Table 1). Patients displayed good adherence in both groups, with a mean adherence rate of 97% in the SsK12 group and 96% in the placebo group.

### Efficacy

The incidence of grade 3–4 SOM, which was the primary efficacy end point, was significantly reduced among patients receiving SsK12 lozenges as compared with those receiving placebo (36.6% v 54.2%;  $P = .0351$ ; Table 2). The distribution of OM during each patient's observation period is shown in Figure 2. The SOM duration in the SsK12 group was significantly shorter than that in the placebo group (median, 0.0 days v 7.0 days; mean, 8.9 days v 18.3 days;  $P = .0084$ ; Table 2). Similarly, the median duration of SOM among the patients who had SOM in the SsK12 group was also significantly shorter than that in the placebo group (19.0 days v 35.0 days;  $P = .0485$ ). Similar results were obtained in the FAS (Data Supplement, Table S4). The time to develop SOM in the SsK12 group was significantly delayed as compared with that in the placebo group ( $P = .0123$ ; Table 2; Data Supplement, Fig S2). The incidence of grade 4 OM was significantly lower in the SsK12 group vis-a-vis the placebo group (2.8% v 15.3%;  $P = .0169$ ; Table 2). The administration of SsK12 effectively reduced the incidence and duration and delayed the time to onset of OM (Table 2). One patient in the SsK12 group and two patients in the placebo group missed five or more consecutive radiation fractions because of SOM (Table 2; Fig 2).

The distribution of the mean WHO OM grade and incidence of SOM during the RT course in the two groups are shown in Figure 3. As the radiation dose increased, the average OM grade of patients in the SsK12 group was consistently lower than that in the placebo group, with an increasing trend in both groups peaking at the sixth week.

### Other Clinical Parameters

The mean weekly MTS and oral activity scores were slightly lower in the SsK12 group as compared with the placebo group (Data Supplement, Fig S3). The MTS index and oral activity index were also comparatively lower after SsK12 treatment versus the placebo treatment (Data Supplement, Fig S4).

The incidence of analgesic use was lower in the SsK12 group; however, the difference was not statistically significant (7% v 11%). Approximately 45% of patients in the SsK12 group and 53% in the placebo group received intravenous hydration or total parenteral nutrition. The decrease in BMI during RT was similar between the two groups (2.18 and 2.47 in the SsK12 and placebo groups, respectively;  $P = .7451$ ).

Furthermore, 55 (77%) and 56 (78%) patients in the SsK12 and placebo groups, respectively, experienced radiation-related grade 2 or higher hyposalivation at the end of RT, and the incidence of hyposalivation was similar between groups during the observation period (Data Supplement, Table S5). Patients' gustatory function decreased in the two groups immediately at mid-RT and was lowest at the end of treatment, with a trend toward recovery at 1 month after RT (Data Supplement, Fig S5A). The detection of all four classic tastes (sweet, bitter, sour, and salty) was affected by RT, with bitter and salty tastes being the most affected ones (Data Supplement, Fig S5B).

### Oral Microbiota

A total of 420 saliva samples were collected for 16S rRNA sequencing, and 406 saliva samples were eligible for further analysis. The rarefaction curves are presented in the Data Supplement (Fig S6). No significant differences in Alpha diversity were detected within or between groups ( $P > .05$ ; Figs 4A and 4B). The principal coordinates analysis plots showed a significant difference in the microbial community at T0 (baseline), T1 (mid-term radiation), and T2 (end of radiation) in both groups (Figs 4C and 4D); however, no significant between-group differences were observed (Data Supplement, Fig S7). The relative abundance of *Firmicutes* in the placebo group exhibited a decreasing trend, whereas it remained relatively stable in the SsK12 group during RT (Data Supplement, Fig S8). Further comparison of the relative abundance of genera at T0 and T2 showed that three of the top 30 dominant genera exhibited different trends of alteration between the two groups (Figs 4E and 4F). The relative abundance of *Streptococcus* significantly decreased

**TABLE 1. Baseline Characteristics of Patients Who Completed This Study**

Characteristic	SsK12 (n = 71)	Placebo (n = 72)	All Patients (N = 143)	P
Sex, No. (%)				.0695
Male	41 (57.75)	52 (72.22)	93 (65.03)	
Female	30 (42.25)	20 (27.78)	50 (34.97)	
Age, years, mean (SD)	52 (13)	53 (12)	52 (12)	.7200
BMI, kg <sup>2</sup> /m, mean (SD)	23 (9)	24 (11)	24 (10)	.5847
ECOG PS, No. (%)				.9976
0	42 (59.15)	43 (59.72)	85 (59.44)	
1	26 (36.62)	26 (36.11)	52 (36.36)	
2	3 (4.23)	3 (4.17)	6 (4.20)	
Tumor site, No. (%)				.9693
Nasopharynx	19 (26.76)	18 (25.00)	37 (25.87)	
Oral cavity	13 (18.31)	12 (16.67)	25 (17.48)	
Oropharynx	8 (11.27)	12 (16.67)	20 (13.99)	
Hypopharynx	7 (9.86)	7 (9.72)	14 (9.79)	
Larynx	10 (14.08)	9 (12.50)	19 (13.29)	
Others	14 (19.72)	14 (19.44)	28 (19.58)	
Tumor pathologic types, No. (%)				.8874
Squamous carcinoma	53 (74.65)	53 (73.61)	106 (74.13)	
Others	18 (25.35)	19 (26.39)	37 (25.87)	
Treatment type, No. (%)				.5389
Definitive	21 (29.58)	18 (25.00)	39 (27.27)	
Postoperative treatment	50 (70.42)	54 (75.00)	104 (72.73)	
TNM stage, No. (%)				.5792
I-II	14 (19.72)	14 (19.44)	28 (19.58)	
III	22 (30.99)	17 (23.61)	39 (27.27)	
Iva	28 (39.44)	29 (40.28)	57 (39.86)	
IVb	7 (9.86)	12 (16.67)	19 (13.29)	
Tumor primary site, No. (%)				.5640
T1	13 (18.31)	8 (11.11)	21 (14.69)	
T2	21 (29.58)	21 (29.17)	42 (29.37)	
T3	19 (26.76)	19 (26.39)	38 (26.57)	
T4	18 (25.35)	24 (33.33)	42 (29.37)	
Nodal involvement, No. (%)				.1333
N0	24 (33.80)	29 (40.28)	53 (37.06)	
N1	14 (19.72)	5 (6.94)	19 (13.29)	
N2	21 (29.58)	21 (29.17)	42 (29.37)	
N3	12 (16.90)	17 (23.61)	29 (20.28)	
Oropharyngeal cancer HPV status, No. (%)				
Positive	2 (25.00)	3 (25.00)	5 (25.00)	
Negative	6 (75.00)	9 (75.00)	15 (75.00)	
Concurrent chemotherapy, No. (%)	35 (49.30)	32 (44.44)	67 (46.85)	.5611
Tobacco use, No. (%)				.1558
Yes	33 (46.48)	42 (58.33)	75 (52.45)	
No	38 (53.52)	30 (41.67)	68 (47.55)	
Alcohol use, No. (%)				.2077
Yes	30 (42.25)	38 (52.78)	68 (47.55)	
No	41 (57.75)	34 (47.22)	75 (52.45)	
Total RT dose, cGy, mean/median	6,609/6,600	6,644/6,600	6,627/6,600	.3733
Dose to oral cavity, cGy, mean/median (range)	4,208/4,083 (1,027-6,630)	4,220/4,184 (1,032-6,585)	4,214/4,131 (1,027-6,630)	.9548

NOTE. BMI calculated as weight in kilograms divided by height in meters squared.

Abbreviations: ECOG PS, Eastern Cooperative Oncology Group performance status; HPV, human papillomavirus; RT, radiotherapy; SD, standard deviation; SsK12, *S. salivarius* K12.



**TABLE 2.** Efficacy Results

Parameter	SsK12 (n = 71)	Placebo (n = 72)	P
SOM (grade 3 or 4) incidence through RT, % <sup>a</sup>	36.6	54.2	.0351
SOM duration, days, median/mean(SD) <sup>b</sup>	0.0/8.9 (15.0)	7.0/18.3 (22.5)	.0084
SOM onset, days, median <sup>c</sup>	NE	42.0	.0123
Grade 4 OM incidence through RT, % <sup>a</sup>	2.8	15.3	.0169
OM incidence through RT, % <sup>a</sup>	62.0	83.3	.0049
OM duration, days, median <sup>b</sup>	10.5	40.3	.0010
OM onset, days, median <sup>c</sup>	28.0	17.5	<.0001
RT treatment breaks >five consecutive fractions, No. (%)	1 (1.4)	2 (2.8)	

Abbreviations: NE, not estimable; OM, oral mucositis; RT, radiotherapy; SD, standard deviation; SOM, severe OM; SsK12, *S. salivarius* K12.


<sup>a</sup> $\chi^2$  test or Fisher's exact test.

<sup>b</sup>Wilcoxon rank-sum test.


<sup>c</sup>Log-rank test.

at the end of radiation in the placebo group ( $P = .0038$ ;  $P_{FDR} = .0357$ ), but it was unchanged in the SsK12 group ( $P = .1343$ ;  $P_{FDR} = .3542$ ). Conversely, the relative abundance of *Selenomonas* and *Acinetobacter* was increased at the end of radiation in the placebo group ( $P = .0003$ ;  $P_{FDR} = .0050$  for *Selenomonas*, and  $P = .0164$ ;  $P_{FDR} = .0898$  for *Acinetobacter*), whereas it was significantly decreased in the SsK12 group ( $P = .0055$ ;  $P_{FDR} = .0386$  for *Selenomonas* and  $P = .0014$ ;  $P_{FDR} = .0135$  for *Acinetobacter*; Fig 4F). The

relationships between the relative abundance of the three abovementioned genera and time were further analyzed, and the relative abundance of *Streptococcus* was negatively correlated with time in the placebo group (coefficient =  $-0.0175$ ;  $P = .0036$ ;  $P_{FDR} = .0177$ ), whereas it was relatively stable in the SsK12 group during RT (coefficient =  $0.0095$ ;  $P = .1312$ ;  $P_{FDR} = .2284$ ; Figs 4G and 4H). We also compared the top 30 dominant genera between the two groups at the end of RT. Although not statistically significant after Benjamini-



**FIG 2.** Swimmer plot of OM score for the subsets of patients in (left) the SsK12 group (n = 44) or (right) the placebo group (n = 60) who experienced OM. Each horizontal strip represents a patient. The top indicates the time at or after RT, whereas the vertical blue line indicates the end of RT. The bottom strip represents patients who experienced RT treatment breaks (>five consecutive fractions). OM was viewed and scored twice weekly during RT and weekly thereafter for up to 8 weeks. Dark spot, break time; mauve, WHO grade 1; OM, oral mucositis; purple, WHO grade 2; red, WHO grade 4; RT, radiotherapy; SsK12, *Streptococcus salivarius* K12; V, view; white, WHO grade 0; X, no RT plan; yellow, WHO grade 3.



**FIG 3.** Mean WHO score and SOM (WHO grade 3 or 4) incidence (%) over time for SsK12 and placebo groups during radiotherapy. The bars represent the incidence of SOM, and the solid/dashed lines represent the mean WHO score. WHO scoring was performed twice per week during RT. For some patients who received 33 fractions of RT, the observation period was 6.5 weeks (7a). RT, radiotherapy; SOM, severe oral mucositis; SsK12, *Streptococcus salivarius* K12.

Hochberg false discovery rate adjustment, the placebo group harbored a relatively higher abundance of *Peptostreptococcus* and *Atopobium* and the SsK12 group had a higher abundance of *Streptococcus* and *Eubacterium\_brachy\_* group (Data Supplement, Fig S9).

## Safety

No severe adverse events (SAEs) related to SsK12 lozenges or placebo administration were observed. In the SsK12 group, 48% of the patients versus 51% in the placebo group reported at least one SAE related to RT (Data Supplement, Table S6). The AEs were evenly distributed between groups and were consistent with the known toxicities of RT (Data Supplement, Table S7). Mild or moderate GI reactions (flatulence or dyspepsia) were considered to be potentially related to the study lozenges (two patients in the SsK12 group and one patient in the placebo group). These two symptoms resolved spontaneously without any other medications.


## DISCUSSION

Microbiota plays a pivotal role in the development and progression of radiation-induced OM.<sup>24,25</sup> Recent studies have shown that alterations in genus abundance, particularly the decrease in oral commensals and enrichment of Gram-negative bacteria throughout RT, are associated with the onset and severity of OM,<sup>9-11,26</sup> suggesting that rescuing or countering the alterations of specific taxa would be promising for the management of this disease. Probiotics are live

microorganisms that modulate microecology and confer health benefits to the host.<sup>27</sup> Previous clinical trials have shown the effectiveness of probiotics, including *Lactobacillus brevis* CD2 and other probiotic combinations (*Bifidobacterium longum*, *Lactobacillus lactis*, and *Enterococcus faecium*) in the management of radiation-induced OM.<sup>28,29</sup> However, the beneficial effects of *Lactobacillus brevis* CD2 in reducing the incidence of SOM were not confirmed in a recent clinical trial.<sup>30</sup> Therefore, clinical trials using alternative probiotics in larger cohorts are needed to extensively evaluate the efficacy of probiotics in the management of radiation-induced OM, with a particular focus on the incidence of SOM. SsK12 can modulate oral microbiota by producing BLISs and has exhibited good immunomodulatory properties.<sup>31,32</sup> Our recent data showed that SsK12 effectively ameliorates radiation-induced OM in mice, suggesting its potential application in this disease.<sup>18</sup>

Here, we conducted a prospective randomized clinical trial that demonstrated the effectiveness of using SsK12, wherein it significantly reduced the incidence and severity of OM in patients with malignant head and neck tumors undergoing RT. Furthermore, trends in the secondary end points, that is, the duration and time to the onset of SOM, supported this observation. Moreover, SsK12 did not increase the RT-associated toxicity.

In this study, we monitored dynamic changes of oral microbiota during the course of RT. During treatment, we found that the relative abundance of *Streptococcus* continuously decreased, whereas that of *Selenomonas* and *Acinetobacter*



**FIG 4.** Effects of SsK12 treatment on salivary microbiota during RT. (A and B) Alpha diversity of oral microbial estimated by Shannon index and Chao index. (C and D) Beta diversity calculated using weighted UniFrac by the PCoA plot and further confirmed by ANOSIM analysis. (E) The heatmap of the relative abundances of the top 30 bacterial genera (on average). (F) Genera showing a divergent change trend at T0 and T2 within each group (Wilcoxon rank-sum test). (G and H) The correlations between the relative abundance of *Streptococcus* and time throughout the treatment regimen. The correlations were calculated using MaAsLin. All *P* values were adjusted for multiple comparisons using BH-FDR, and  $P_{FDR} < .05$  was considered significant. \**P* < .05, \*\**P* < .01, \*\*\**P* < .001. ANOSIM, analysis of similarities; BH-FDR, Benjamini-Hochberg false discovery rate; FDR, false discovery rate; OTU, operational taxonomic unit; PC, principal component; PCoA, principal coordinates analysis; Placebo-T0, before the treatment of RT plus a placebo; Placebo-T1, the middle of the treatment of RT plus a placebo; Placebo-T2, the end of the treatment of RT plus a placebo; SsK12, *Streptococcus salivarius* K12; SsK12-T0, before the treatment of RT plus the probiotic *S. salivarius* K12; SsK12-T1, the middle of the treatment of RT plus the probiotic *S. salivarius* K12; SsK12-T2, the end of the treatment of RT plus the probiotic *S. salivarius* K12.

increased in the placebo group, with the administration of SsK12 appearing to rescue or reverse this trend. Most *Streptococcus* species are oral commensals. A lower abundance of *Streptococcus* was associated with a shorter average time to the onset of SOM.<sup>26</sup> Consistently, patients who subsequently developed  $\geq$ grade 2 OM reported a lower abundance (26%) of *Streptococcus*.<sup>10</sup> Our study found that the application of SsK12 countered the decrease in *Streptococcus* during RT, which partially explained the protective effect of this probiotic. *Selenomonas* is a cluster of Gram-negative bacteria implicated in periodontitis<sup>33</sup> and OM.<sup>9</sup> *Acinetobacter* is a group of Gram-

negative bacteria that are usually associated with opportunistic infections.<sup>34,35</sup> In the current study, the use of SsK12 reversed the increasing trend of *Selenomonas* and *Acinetobacter* during RT, which may also benefit mucosal homeostasis. Hence, we speculated that the use of SsK12 could benefit the management of OM by maintaining the abundance of commensal *Streptococcus* while inhibiting opportunistic pathogens during RT.

In addition to OM, we found that patients who received SsK12 had slightly lower MTS scores, oral activity impairment, additional nutritional intake, and reduced BMI than those in



the placebo group; however, the differences were not statistically significant. This is partly because other complications of RT, such as sore throat, hyposalivation, taste dysfunction, nausea, and vomiting, also affect the QoL of patients with malignant tumors. Most patients in both groups developed hyposalivation and taste dysfunction, which is consistent with the results of previous studies.<sup>36,37</sup> This suggests holistic management of patients undergoing head and neck RT.

To our knowledge, this is the first study to evaluate the effectiveness of the oral probiotic SsK12 in preventing and treating radiation-induced OM. The major strength of this randomized trial is the strict inclusion and exclusion criteria, standardized RT delivery and supportive care, and the use of a placebo and double-blind design with minimized bias. In addition, previous studies have evaluated the efficacy of probiotics in the treatment of radiation-induced OM but without a longitudinal track of oral microbiota. Furthermore, compared with palifermin (keratinocyte growth factor-1), the only agent approved by the US Food and Drug Administration for minimizing OM in patients with hematologic malignancies undergoing hematopoietic cell transplantation and receiving chemotherapy or RT, SsK12 is cost-effective, is easy to use, and has a good safety profile with only mild or

moderate GI reactions, thereby favoring its daily use in patients with malignant tumors undergoing RT.

This study has a few limitations. Some patients undergoing RT/chemotherapy may develop systemic infections because of immune suppression; in such cases, antibiotic intervention is needed, which may compromise the clinical outcome of probiotic treatment for OM. In addition, the placebo group had more T4 and N3 cases (although not statistically significant), which could potentially lead to dietary changes in patients because of dysphagia and sore throat, thereby decreasing the patient's resistance to mucositis. In addition, some patients reported difficulty in sucking lozenges because of dry mouth. An improved design for the form or delivery of probiotics is warranted in future studies. Furthermore, this study was conducted at a single cancer center and therefore requires multi-institutional validation.

In conclusion, the current study has demonstrated that the use of SsK12 can significantly reduce the incidence and severity of OM in patients with malignant head and neck tumors receiving RT with a good safety profile. The use of SsK12 has the potential to become a novel strategy for the oral mucosal care of patients receiving RT.

## AFFILIATIONS

<sup>1</sup>Department of Biotherapy, Cancer Center, West China Hospital & State Key Laboratory of Biotherapy, Sichuan University, Chengdu, China

<sup>2</sup>Department of Cariology and Endodontics, State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu, China

## CORRESPONDING AUTHOR

Xin Xu, DDS, West China Hospital of Stomatology, Sichuan University, No. 14, Third Section Renmin Nan Rd, Chengdu 610041, China; e-mail: xin.xu@scu.edu.cn.

## EQUAL CONTRIBUTION

X.P. and Z.L. contributed equally as the first authors, and X.P. and X.X. contributed equally as senior authors of this work.

## SUPPORT

Supported by the Science & Technology Department of Sichuan Province (2021YFQ0064); the Health Commission of Sichuan Province (21PJ058); the National Key Research and Development Program of China (2021YFE0206600); 1·3·5 project for disciplines of excellence—Clinical Research Incubation Project, West China Hospital, Sichuan University (2022HXFH026); Talent incubation project of General Hospital of Western Theater Command of PLA (2021-XZYG-C49); and the Scientific Research Project of China Baoyuan (CBYI202103).

## REFERENCES

1. Colevas AD, Yom SS, Pfister DG, et al: NCCN guidelines insights: Head and neck cancers, version 1.2018. *J Natl Compr Canc Netw* 16:479-490, 2018
2. Epstein JB, Thariat J, Bensadoun RJ, et al: Oral complications of cancer and cancer therapy: From cancer treatment to survivorship. *CA Cancer J Clin* 62:400-422, 2012
3. Anderson CM, Lee CM, Saunders DP, et al: Phase IIb, randomized, double-blind trial of GC4419 versus placebo to reduce severe oral mucositis due to concurrent radiotherapy and cisplatin for head and neck cancer. *J Clin Oncol* 37:3256-3265, 2019

## CLINICAL TRIAL INFORMATION

ChiCTR2100054689

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.23.00837>.

## DATA SHARING STATEMENT

A data sharing statement provided by the authors is available with this article at DOI <https://doi.org/10.1200/JCO.23.00837>.

## AUTHOR CONTRIBUTIONS

**Conception and design:** Ruidan Li, Xingchen Peng, Xin Xu

**Financial support:** Ruidan Li, Xingchen Peng, Xin Xu

**Administrative support:** Ruidan Li, Xingchen Peng, Xin Xu

**Provision of study materials or patients:** Ruidan Li, Xingchen Peng, Xin Xu

**Collection and assembly of data:** Zixia Li, Yiyan Pei, Shuhao Zheng

**Data analysis and interpretation:** Jinchi Liu, Jingjing Wang, Ruidan Li, Xingchen Peng, Xin Xu

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

4. Henke M, Alfonsi M, Foa P, et al: Palifermin decreases severe oral mucositis of patients undergoing postoperative radiochemotherapy for head and neck cancer: A randomized, placebo-controlled trial. *J Clin Oncol* 29:2815-2820, 2011
5. Le QT, Kim HE, Schneider CJ, et al: Palifermin reduces severe mucositis in definitive chemoradiotherapy of locally advanced head and neck cancer: A randomized, placebo-controlled study. *J Clin Oncol* 29:2808-2814, 2011
6. Maria OM, Eliopoulos N, Muanza T: Radiation-induced oral mucositis. *Front Oncol* 7:89, 2017
7. Mallick S, Benson R, Rath GK: Radiation induced oral mucositis: A review of current literature on prevention and management. *Eur Arch Otorhinolaryngol* 273:2285-2293, 2016
8. Elad S, Cheng KKF, Lalla RV, et al: MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer* 126:4423-4431, 2020
9. Hou J, Zheng H, Li P, et al: Distinct shifts in the oral microbiota are associated with the progression and aggravation of mucositis during radiotherapy. *Radiother Oncol* 129:44-51, 2018
10. Vesty A, Gear K, Biswas K, et al: Oral microbial influences on oral mucositis during radiotherapy treatment of head and neck cancer. *Support Care Cancer* 28:2683-2691, 2020
11. Zhu XX, Yang XJ, Chao YL, et al: The potential effect of oral microbiota in the prediction of mucositis during radiotherapy for nasopharyngeal carcinoma. *EBioMedicine* 18:23-31, 2017
12. Burton JP, Wescombe PA, Moore CJ, et al: Safety assessment of the oral cavity probiotic *Streptococcus salivarius* K12. *Appl Environ Microbiol* 72:3050-3053, 2006
13. Burton JP, Cowley S, Simon RR, et al: Evaluation of safety and human tolerance of the oral probiotic *Streptococcus salivarius* K12: A randomized, placebo-controlled, double-blind study. *Food Chem Toxicol* 49:2356-2364, 2011
14. Cosseau C, Devine DA, Dullaghan E, et al: The commensal *Streptococcus salivarius* K12 downregulates the innate immune responses of human epithelial cells and promotes host-microbe homeostasis. *Infect Immun* 76:4163-4175, 2008
15. Burton JP, Chilcott CN, Moore CJ, et al: A preliminary study of the effect of probiotic *Streptococcus salivarius* K12 on oral malodour parameters. *J Appl Microbiol* 100:754-764, 2006
16. Ross KF, Ronson CW, Tagg JR: Isolation and characterization of the lantibiotic salivaricin A and its structural gene salA from *Streptococcus salivarius* 20P3. *Appl Environ Microbiol* 59:2014-2021, 1993
17. Upton M, Tagg JR, Wescombe P, et al: Intra- and interspecies signaling between *Streptococcus salivarius* and *Streptococcus pyogenes* mediated by SalA and SalA1 lantibiotic peptides. *J Bacteriol* 183:3931-3938, 2001
18. Wang Y, Li J, Zhang H, et al: Probiotic *Streptococcus salivarius* K12 alleviates radiation-induced oral mucositis in mice. *Front Immunol* 12:684824, 2021
19. Brouwer CL, Steenbakkers RJ, Bourhis J, et al: CT-based delineation of organs at risk in the head and neck region: DAHANCA, EORTC, GORTEC, HKNPCSG, NCIC CTG, NCRI, NRG Oncology and TROG consensus guidelines. *Radiother Oncol* 117:83-90, 2015
20. Grégoire V, Evans M, Le QT, et al: Delineation of the primary tumour Clinical Target Volumes (CTV-P) in laryngeal, hypopharyngeal, oropharyngeal and oral cavity squamous cell carcinoma: AIRO, CACA, DAHANCA, EORTC, GEORCC, GORTEC, HKNPCSG, HNCIG, IAG-KHT, LPRHHT, NCIC CTG, NCRI, NRG Oncology, PHNS, SBRT, SOMERA, SRO, SSHNO, TROG consensus guidelines. *Radiother Oncol* 126:3-24, 2018
21. Mir R, Kelly SM, Xiao Y, et al: Organ at risk delineation for radiation therapy clinical trials: Global Harmonization Group consensus guidelines. *Radiother Oncol* 150:30-39, 2020
22. He L, Yang H, Chen Z, et al: The effect of *Streptococcus salivarius* K12 on halitosis: A double-blind, randomized, placebo-controlled trial. *Probiotics Antimicrob Proteins* 12:1321-1329, 2020
23. Ren Y, Yu G, Shi C, et al: Majorbio Cloud: A one-stop, comprehensive bioinformatic platform for multiomics analyses. *iMeta* 1:e12, 2022
24. Vanhoecke B, De Ryck T, Stringer A, et al: Microbiota and their role in the pathogenesis of oral mucositis. *Oral Dis* 21:17-30, 2015
25. Sonis ST: New thoughts on the initiation of mucositis. *Oral Dis* 16:597-600, 2010
26. Reyes-Gibby CC, Wang J, Zhang L, et al: Oral microbiome and onset of oral mucositis in patients with squamous cell carcinoma of the head and neck. *Cancer* 126:5124-5136, 2020
27. Hill C, Guarner F, Reid G, et al: Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol* 11:506-514, 2014
28. Sharma A, Rath GK, Chaudhary SP, et al: *Lactobacillus brevis* CD2 lozenges reduce radiation- and chemotherapy-induced mucositis in patients with head and neck cancer: A randomized double-blind placebo-controlled study. *Eur J Cancer* 48:875-881, 2012
29. Jiang C, Wang H, Xia C, et al: A randomized, double-blind, placebo-controlled trial of probiotics to reduce the severity of oral mucositis induced by chemoradiotherapy for patients with nasopharyngeal carcinoma. *Cancer* 125:1081-1090, 2019
30. Sanctis VDE, Belgioia L, Cante D, et al: *Lactobacillus brevis* CD2 for prevention of oral mucositis in patients with head and neck tumors: A multicentric randomized study. *Anticancer Res* 39:1935-1942, 2019
31. Barbour A, Tagg J, Abou-Zied OK, et al: New insights into the mode of action of the lantibiotic salivaricin B. *Sci Rep* 6:31749, 2016
32. Geng M, Austin F, Shin R, et al: Covalent structure and bioactivity of the type AII lantibiotic salivaricin A2. *Appl Environ Microbiol* 84:e02528, 2018
33. Hawkes CG, Hinson AN, Vashishta A, et al: *Selenomonas sputigena* interactions with gingival epithelial cells that promote inflammation. *Infect Immun* 91:e0031922, 2023
34. Visca P, Seifert H, Towner KJ: Acinetobacter infection—An emerging threat to human health. *IUBMB Life* 63:1048-1054, 2011
35. Lee CR, Lee JH, Park M, et al: Biology of *Acinetobacter baumannii*: Pathogenesis, antibiotic resistance mechanisms, and prospective treatment options. *Front Cell Infect Microbiol* 7:55, 2017
36. Gunn L, Gilbert J, Nenclares P, et al: Taste dysfunction following radiotherapy to the head and neck: A systematic review. *Radiother Oncol* 157:130-140, 2021
37. Mercadante V, Al Hamad A, Lodi G, et al: Interventions for the management of radiotherapy-induced xerostomia and hyposalivation: A systematic review and meta-analysis. *Oral Oncol* 66:64-74, 2017

## **AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**

### **Streptococcus salivarius K12 Alleviates Oral Mucositis in Patients Undergoing Radiotherapy for Malignant Head and Neck Tumors: A Randomized Controlled Trial**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/jco/authors/author-center](http://ascopubs.org/jco/authors/author-center).

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

No potential conflicts of interest were reported.