

ORIGINAL ARTICLE

Resting-state abnormalities in amnesic mild cognitive impairment: a meta-analysis

WKW Lau^{1,2,6}, M-K Leung^{2,3,6}, TMC Lee^{2,3,4,5} and ACK Law^{1,4}

Amnesic mild cognitive impairment (aMCI) is a prodromal stage of Alzheimer's disease (AD). As no effective drug can cure AD, early diagnosis and intervention for aMCI are urgently needed. The standard diagnostic procedure for aMCI primarily relies on subjective neuropsychological examinations that require the judgment of experienced clinicians. The development of other objective and reliable aMCI markers, such as neural markers, is therefore required. Previous neuroimaging findings revealed various abnormalities in resting-state activity in MCI patients, but the findings have been inconsistent. The current study provides an updated activation likelihood estimation meta-analysis of resting-state functional magnetic resonance imaging (fMRI) data on aMCI. The authors searched on the MEDLINE/PubMed databases for whole-brain resting-state fMRI studies on aMCI published until March 2015. We included 21 whole-brain resting-state fMRI studies that reported a total of 156 distinct foci. Significant regional resting-state differences were consistently found in aMCI patients relative to controls, including the posterior cingulate cortex, right angular gyrus, right parahippocampal gyrus, left fusiform gyrus, left supramarginal gyrus and bilateral middle temporal gyri. Our findings support that abnormalities in resting-state activities of these regions may serve as neuroimaging markers for aMCI.

Translational Psychiatry (2016) 6, e790; doi:10.1038/tp.2016.55; published online 26 April 2016

INTRODUCTION

Mild cognitive impairment (MCI) refers to an intermediate state between the cognitive changes of normal aging and the symptomatic pre-dementia stage.¹ The prevalence of MCI is 10–20% in adults aged 65 years or above, and more than 50% of MCI patients progress to dementia within 5 years.^{2,3} These figures draw great attention from clinicians and society at large because there is currently no effective drug that can cure dementia. MCI can be subdivided into amnesic (characterized by primarily memory impairment) and non-amnesic (characterized by executive function impairment) types based on the neuropsychological symptom profile. Amnesic MCI (aMCI) is generally regarded as a prodromal stage of Alzheimer's disease (AD), with an annual conversion rate of up to 25%.^{2,4} This indicates the importance of early diagnosis and intervention for people with aMCI. Indeed, delayed MCI diagnosis in people with high education has been reported to impact mortality.⁵ According to the National Institute on Aging–Alzheimer's Association workgroups,⁶ the criteria for MCI include (1) a change in cognition, in comparison with the person's prior level; (2) lower performance in one or more cognitive domains that is greater than what would be expected for the patient's age and educational background; (3) independence of function in daily life is generally maintained but is less efficient compared with the past, and minimal aids may be required; and (4) the person is not demented. The standard diagnostic procedure of MCI primarily relies on subjective neuropsychological examinations that require the judgment of experienced clinicians. The development of other objectives and

reliable MCI markers, such as neural markers,² is therefore required.

There is a growing interest in the use of resting-state functional magnetic resonance imaging (fMRI) to explore the neurophysiological mechanisms associated with aMCI owing to its noninvasive and task-free nature. The term 'resting state' refers to spontaneous brain activity during a passive (resting) state when one is lying quietly with the eyes closed or passively viewing a stimulus.⁷ In fact, resting-state activity is influenced by self-consciousness, ongoing thoughts, states of alertness and readiness to process stimuli from the outside world.⁸ Previous studies using fMRI have revealed a highly stereotypical pattern of spontaneous activity in a number of brain regions, namely the default mode network that manifests greater activity during passive task states compared with various active task states.^{9–11} Numerous studies have attempted to work out the difference in resting-state activity between MCI patients and healthy age-matched controls. However, the findings have been inconsistent, which might be due to the inclusion of different types of MCI patients, and/or the use of different methods across studies.

A systemic and quantitative analysis is highly warranted to delineate the meanings of the existing findings in the field. To our knowledge, only one meta-analysis has explored the difference in resting-state activity in MCI and controls.¹² In the Supplementary Materials of this article, the authors reported reduced resting-state activity in the middle temporal gyrus, middle frontal gyrus, medial frontal gyrus and precuneus, and increased resting-state activity in a large cluster spanning the temporo-parietal parts of the brain in

¹Neural Dysfunction Research Laboratory, Department of Psychiatry, The University of Hong Kong, Hong Kong; ²Laboratory of Cognitive Affective Neuroscience, The University of Hong Kong, Hong Kong; ³Laboratory of Neuropsychology, The University of Hong Kong, Hong Kong; ⁴The State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong, Hong Kong and ⁵Institute of Clinical Neuropsychology, The University of Hong Kong, Hong Kong. Correspondence: Professor TMC Lee, Laboratory of Neuropsychology, The University of Hong Kong, Room 656, The Jockey Club Tower, The University of Hong Kong, Pokfulam Road, Hong Kong or Dr ACK Law, Neural Dysfunction Research Laboratory, Department of Psychiatry, The University of Hong Kong, Room 219, Block J, 2nd Floor, Queen Mary Hospital, Pokfulam Road, Hong Kong. E-mail: tmclee@hku.hk or acklaw@hku.hk

⁶These authors contributed equally to this work.

Received 19 August 2015; revised 28 February 2016; accepted 5 March 2016

MCI patients from 17 independent studies. The included MCI patients were heterogeneous, including subcortical vascular MCI (svMCI) patients¹³ who might have a different pathophysiology compared with aMCI. Such heterogeneity likely impacted on the consistency of the results, given the small number of included studies. In particular, reduced resting-state activity was previously reported in the posterior cingulate cortex (PCC),¹⁴ hippocampal and parahippocampal regions¹⁵ in aMCI patients, such that the decreased PCC activity during resting state was further amplified in AD compared with aMCI.¹⁴ However, these findings were not confirmed by the previous meta-analysis study.¹² Hence, the present analysis aimed to comprehensively review the abnormalities in resting-state activity in aMCI patients specifically, who have a high risk of conversion to AD, using activation likelihood estimation (ALE). An updated ALE analysis focusing on aMCI patients could provide a reasonable means to resolving discrepancies in previously reported findings.

MATERIALS AND METHODS

A comprehensive online literature search on the MEDLINE/PubMed databases was conducted, focusing on functional neuroimaging studies on MCI. Keyword searches were conducted using the following search terms: (1) 'neuroimaging' <OR> 'fMRI,' (2) 'resting state' OR 'default network' and (3) 'mild cognitive impairment' <OR> 'MCI.' These searches were confined to articles published in English up to March 2015, which yielded 228 original or review articles. We also searched through the reference lists of relevant review articles. From these research articles, we included studies that reported Montreal Neurological Institute (MNI) or Talairach¹⁶ coordinates of whole-brain contrast comparing the amnesic type of MCI and healthy controls. Two of the authors (WKWL and MKL) confirmed the inclusions of the identified studies. For included articles that reported only brain images, we attempted to obtain the MNI or Talairach coordinates from the corresponding author(s) via email (three out of eight studies replied). Studies were excluded if (1) only non-amnesic MCI or only subtypes of aMCI were included; (2) no control group was included; (3) patients had a history of neurological, psychiatric or any systemic disease that could affect cognitive functions (for example, stroke, depression, alcoholism and drug abuse); (4) *a priori* region of interest analysis or a seed-based functional connectivity analysis was conducted; or (5) the effects of medication were tested without reporting fMRI data at baseline. One study contained multiple independent patient samples; the appropriate coordinates were extracted as two separate experiments.¹⁷ We did not intentionally exclude studies that used a modality other than fMRI, nor focused on a particular analytic approach. As a result, the methods used in our included studies covered the following approaches: regional homogeneity, amplitude of low-frequency fluctuations and independent component analysis. Overall, 21 fMRI studies (22 experiments) reporting 156 foci were included for the ALE meta-analysis (Table 1). One of them used the arterial spin labeling perfusion MRI technique to measure resting-state abnormality in cerebral blood flow in aMCI.

GingerALE version 2.3.2 (The BrainMap Database, www.brainmap.org; San Antonio, TX, USA) was used to conduct the coordinate-based ALE analysis, which is a widely used technique for synthesizing neuroimaging data.^{36–38} ALE estimates the convergence of significant effects in terms of foci across different neuroimaging studies. Coordinates in Talairach space were imported into the software. If coordinates were reported in MNI space, the 'icbm2tal' algorithm was used to transform them into Talairach space.³⁹ For coordinates that had been transformed to Talairach space by brett-transformation, they were first 'un-bretted' into MNI space before applying Lancaster's transformation. Imported foci were modeled as three-dimensional Gaussian spatial probability distributions using a full-width-half-maximum kernel estimated based on the corresponding experiment's sample size.³⁸ These probability distributions were combined into a modeled activation map using the 'non-additive' method.³⁷ Next, the union of the modeled activation maps of each experiment was created to form the ALE image that contains the combined probability distribution of finding an activation being located at that particular voxel (that is, ALE scores). The ALE image was then thresholded using uncorrected $P < 0.001$ and a cluster-level inference threshold of $P < 0.05$ with 1000 permutations of simulated random data based on the characteristics of the imported data.³⁶ In the cluster-level inference, contiguous voxels (that is, clusters) that exceed the cluster-forming threshold were compared against the

simulated random clusters. The cluster-inference threshold approach is considered optimal because it is more stringent than uncorrected voxel-level thresholds and less conservative than conventional false discovery rate and family-wise error rate corrections. Clusters contributed by a single study only were not reported even if they exceeded the cluster-inference threshold. A total of two ALE analyses were conducted for each contrast (healthy control > aMCI or healthy control < aMCI) in all aMCI patients.

RESULTS

aMCI patients showed decreases in resting-state activity compared with healthy controls in the right and medial PCC (Brodmann area/BA23 and 31), right angular gyrus (BA39, extending to the right middle temporal gyrus, BA19), right parahippocampal gyrus (BA35) and left fusiform gyrus (BA37) (Figure 1a and Table 2). Furthermore, aMCI patients showed increases in resting-state activity in the left middle temporal gyrus (BA39) and left supramarginal gyrus (BA40) (Figure 1b and Table 2).

DISCUSSION

The results of the current aggregate analyses demonstrated reduced resting-state activity in the middle temporal gyrus in aMCI compared with controls, which is consistent with a previous meta-analysis report.¹² In addition, reduced resting-state activity was also found in the PCC, right angular gyrus, right parahippocampal gyrus and left fusiform gyrus that were not reported in the previous ALE study.¹² Importantly, the absence of reduced resting-state activity in these regions in the previous ALE study included a heterogeneous sample of MCI patients (that is, aMCI and svMCI), suggesting that the two subtypes of MCI may possess distinct pathophysiological characteristics. In fact, increases in resting-state activity in the PCC have been reported in patients with svMCI.¹³ This further supports our hypothesis that reduced resting-state activity in the PCC may be unique to aMCI patients compared with other subtypes such as svMCI.

Episodic memory decline is the most prominent cognitive impairment marker that is used to differentiate aMCI patients from age-matched healthy controls.⁴⁰ Brain regions including the PCC, parahippocampal gyrus, inferior and middle temporal gyri are associated with many cognitive processes, including episodic memory.^{29,41–44}

Decreases in resting-state activity in the PCC were consistently observed in patients with aMCI compared with controls. According to the literature, metabolic reductions in the PCC are present in very early AD, even before a definitive clinical diagnosis,⁴⁵ which corroborates our findings. The PCC is mainly involved in episodic memory processing.^{41,42} Abnormal connectivity was found between the PCC and the hippocampus in early AD.⁴⁶ Furthermore, positive correlations have been reported between the resting-state activity of the PCC and mini-mental state examination scores in MCI and AD, suggesting that resting-state activity changes in this region are associated with altered cognitive performance.^{17,23}

Decreased gray matter volume (GMV) in the PCC has been reported in both MCI and AD compared with controls using MRI volumetric measurements,^{47,48} which could lead to a reduction in resting-state activity. To confirm that the reduction in resting-state activity of the PCC was not driven by brain atrophy, a small subgroup analysis was performed using 7 studies (38 foci) that reported resting-state abnormalities after corrected for GMV or the reported resting-state abnormalities did not overlap with the reported GMV reduction (Supplementary eTable 1). The resting-state hypoactivation in the medial PCC was maintained in aMCI patients in this subgroup analysis (Supplementary eTable 2). This finding suggests that the reduction in resting-state activity of the PCC may be independent of the GMV changes in aMCI patients.

Table 1. List of included studies on resting state in aMCI

Study	Method of analysis	N	Age, mean (s.d.), years	Education, mean (s.d.), years	MMSE, mean (s.d.)	Contrasts	Foci
Xu et al. ¹⁸	ASL	10 MCI	77.0 (4.5)	13.7 (2.7)	27.8 (1.5)	HC > MCI	1
		12 HC	70.0 (3.9)	15.6 (2.1)	29.6 (0.8)		
Sorg et al. ¹⁹	ICA	24 MCI	69.3 (8.1)	13.8 (4.0)	27.7 (1.1)	HC > MCI	21
		16 HC	68.1 (3.8)		29.6 (0.5)		
Bai et al. ²⁰	ReHo	20 MCI	71.3 (3.8)	14.0 (3.1)	27.2 (1.6)	HC > MCI	9
		20 HC	69.4 (3.8)	13.8 (4.0)	28.3 (1.4)	MCI > HC	4
Qi et al. ²¹	ICA	14 MCI	71.8 (7.3)	11.6 (1.0)	26.6 (0.3)	HC > MCI	7
		14 HC	70.4 (5.8)	10.0 (0.9)	28.5 (0.2)	MCI > HC	5
Zhang et al. ²²	ReHo	48 MCI	72.0 (4.4)	15.5 ^a	27.0 ^a	HC > MCI	9
		36 HC	71.6 (3.7)	16.0 ^a	29.0 ^a	MCI > HC	3
Wang et al. ²³	ALFF	16 MCI	69.4 (7.0)	10.9 (3.4)	26.5 (1.0)	HC > MCI	5
		22 HC	66.6 (7.7)	10.0 (3.9)	28.6 (0.6)	MCI > HC	4
Bai et al. ²⁴	ICA	26 MCI	71.4 (4.3)	13.8 (2.8)	27.2 (1.5)	MCI > HC	2
		18 HC	70.3 (4.7)	15.1 (3.1)	28.3 (1.3)		
Xi et al. ¹⁵	ALFF	18 MCI	67.3 (7.9)	12.4 (3.2)	24.8 (3.8)	HC > MCI	3
		20 HC	64.7 (5.6)	12.2 (2.5)	28.2 (1.8)	MCI > HC	2
Han et al. ²⁵	ALFF	17 MCI	69.7 (7.6)	8.8 (4.0)	25.2 (3.5)	HC > MCI	6
		18 HC	66.5 (6.2)	8.4 (5.6)	29.2 (0.7)	MCI > HC	6
Bai et al. ²⁶	ALFF	43 MCI	72.0 (4.8)	13.6 (3.0)	27.1 (1.5)	HC > MCI	1
		30 HC	73.0 (3.5)	14.9 (2.7)	28.2 (1.4)	MCI > HC	1
Wang et al. ²⁷	BOLD	8 MCI	66.4 (11.0)	10.6 (3.5)	25.4 (1.3)	HC > MCI	2
		14 HC	66.1 (5.8)	11.0 (4.5)	28.0 (1.4)	MCI > HC	9
Zhuang et al. ²⁸	ALFF	47 MCI	72.0 (4.8)	15.9 (11.4)	27.0 (1.5)	HC > MCI	1
		33 HC	72.8 (3.4)	14.7 (2.9)	28.2 (1.3)	MCI > HC	1
Song et al. ¹⁴	ICA	18 MCI	70.2 (7.9)	9.4 (4.8)	21.9 (5.0)	HC > MCI	1
		21 HC	65.0 (8.1)	11.0 (4.4)	28.5 (1.4)		
Xi et al. ²⁹	ALFF	18 MCI	67.4 (7.7)	12.3 (3.2)	25.2 (3.4)	HC > MCI	3
		18 HC	65.4 (5.8)	12.3 (2.4)	28.1 (1.8)	MCI > HC	2
Zhou et al. ³⁰	ALFF	17 MCI	67.0 (7.9)	11.1 (3.3)	26.4 (2.1)	HC > MCI	3
		17 HC	63.8 (5.8)	11.7 (3.0)	28.6 (1.1)	MCI > HC	4
Zhao et al. ²	ALFF	20 MCI	65.1 (9.9)	11.8 (3.3)	25.2 (2.2)	HC > MCI	7
		18 HC	66.8 (7.4)	12.0 (2.9)	29.3 (1.2)	MCI > HC	3
Liang et al. ¹⁷	BOLD	24 MCI	72.8 (6.6)	13.8 (2.6) ^a	28.1 (1.5)	HC > MCI	2
		35 HC	74.3 (5.9)		28.9 (1.6)		
Liang et al. ¹⁷	BOLD	29 MCI	73.2 (7.3)	14.2 (2.5) ^a	27.1 (2.3)	HC > MCI	7
		35 HC	74.3 (5.9)		28.9 (1.6)		
Liu et al. ³¹	ALFF	46 MCI	71.9 (4.9)	13.8 (2.6) ^a	27.1 (1.4) ^a	HC > MCI	1
		32 HC	72.8 (3.5)	14.2 (2.5) ^a	28.3 (1.1) ^a	MCI > HC	1
Liu et al. ³²	ReHo	12 MCI	59.3 (3.3)	10.5 (1.8)	26.4 (0.9)	HC > MCI	4
		12 HC	60.6 (5.8)	10.6 (2.1)	29.8 (0.4)	MCI > HC	4
Wang et al. ³³	ReHo	30 MCI	69.1 (5.8)	13.8 (2.6) ^a	26.2 (2.2)	HC > MCI	5
		32 HC	70.1 (5.5)		28.1 (1.5)	MCI > HC	6
Zhou et al. ³⁴	fALFF	17 MCI	76.7 (5.5)	13.8 (2.6) ^a	26.1 (0.7)	HC > MCI	1
		14 HC	76.3 (8.3)		29.2 (0.4)		

Abbreviations: ALFF, amplitude of low-frequency fluctuations; aMCI, amnesic mild cognitive impairment; ASL, arterial spin labeling; BOLD, blood oxygen level dependent; fALFF, fractional ALFF; HC, healthy control; ICA, independent component analysis; MCI, mild cognitive impairment; MMSE, mini-mental state examination; ReHo, regional homogeneity. ^aMean and s.d. were estimated from median or calculated according to the formulas published by Hoza et al.³⁵




Figure 1. (a) Resting-state hypoactivation in patients with amnesic mild cognitive impairment (aMCI) compared with matched control subjects (in blue). (b) Resting-state hyperactivation in patients with aMCI compared with matched control subjects (in red). AG, angular gyrus; FG, fusiform gyrus; MTG, middle temporal gyrus; ParaHG, parahippocampal gyrus; PCC, posterior cingulate cortex; R, right; SupraMG, supramarginal gyrus.

Table 2. Resting-state abnormalities in aMCI patients compared with controls in 21 studies

	Side	Brain region	BA	Coordinates (Talairach)			Volume (mm ³)	Extrema value
				x	y	z		
HC > MCI	Right		23	12	-56	18		0.0150
	Medial	Posterior cingulate	23	0	-52	16	1472	0.0135
	Right		31	4	-52	24		0.0118
		Angular gyrus	39	46	-66	28		0.0189
	Right	Middle temporal gyrus	19	42	-74	22	664	0.0139
	Right	Parahippocampal gyrus	35	22	-20	-26	312	0.0190
	Left	Fusiform gyrus	37	-30	-44	-12	304	0.0155
MCI > HC	Left	Middle temporal gyrus	39	-50	-64	28	1280	0.0260
		Supramarginal gyrus	40	-54	-54	28		0.0192

Abbreviations: aMCI, amnesic mild cognitive impairment; BA, Brodmann area; HC, healthy control; MCI, mild cognitive impairment.

Whether the atrophy of the PCC in MCI or AD patients is driven by an early change in brain functional activity requires further study.

Reduced resting-state activity in fusiform gyrus and parahippocampal gyrus may also contribute to memory deficits in aMCI patients. For instance, the fusiform gyrus connects with the medial temporal lobe, including the parahippocampal gyrus,⁴⁹ which has an accessory role in memory processes in healthy elderly people.^{21,50} In addition, the previously reported positive association between the resting state of the parahippocampal gyrus and mini-mental state examination scores¹⁵ suggests a functional role of the resting-state activity of the parahippocampal gyrus in episodic memory.⁵¹

On the other hand, lower resting-state activity in the angular gyrus in the inferior parietal lobule of aMCI patients may be related to their poorer verbal working memory performance that involves short-term storage and retrieval of phonological representations.^{52,53} Taken together, hypoactivation of the above-mentioned regions is a consistent physiological change in aMCI patients, which may serve as a potential neuroimaging biomarker for aMCI.

Decreases in resting-state activity of the medial frontal gyrus and left middle frontal gyrus have been reported in a previous ALE study that included different types of MCI patients.¹² Indeed, a previous study that reported a similar resting-state hypoactivity in the medial frontal gyrus was conducted in svMCI patients compared with controls,¹³ and the study was also included in the previous ALE meta-analysis. We did not, however, observe any significant hypoactivity in these regions in aMCI. One possible explanation for the discrepancy is that the hypoactivity of these prefrontal regions is not a defining feature of aMCI. Nevertheless, more primary studies are needed to generate a clearer picture.

Increases in resting-state activity were found in the left middle temporal gyrus and left supramarginal gyrus in aMCI patients

compared with controls, which is consistent with the previous ALE study of MCI.¹² One interpretation is that these regions are activated to compensate for the reduction in function of other brain regions. However, whether such hyperactivation of the brain is truly a compensatory response or indicates other pathological changes, such as excitotoxicity that triggers neuronal cell death, requires more empirical studies to confirm.

Five studies that compared the spontaneous brain activity between aMCI and controls were not included in the current meta-analysis due to the unavailability of the resultant coordinates. Consistent with our findings, hypoactivity in the PCC and/or precuneus was commonly reported in aMCI patients compared with controls in four out of the five studies.⁵⁴⁻⁵⁷ In contrast to our findings, one study reported increased resting-state activity in the orbitofrontal gyrus, anterior cingulate cortex, parahippocampal gyrus, hippocampus and fusiform gyrus in multiple-domain aMCI patients, which was interpreted as a compensatory mechanism for the recruitment of cognitive resources in the patients.⁵⁵ It is possible that multiple impairments (in memory and at least one other cognitive domain) in multiple-domain aMCI patients may trigger such compensatory responses for cognitive resources that were not present in our included aMCI patients. Another study reported increased resting-state activity of the left inferior parietal lobule in aMCI patients compared with controls, which was also interpreted as a compensatory recruitment in aMCI patients.⁵⁸ A compensatory response may be sensitive to the stage of an illness, and in fact, the inferior parietal lobule abnormality was regarded as a sensitive marker for the transition from MCI to early stages of AD.⁵⁸ To further understand this difference between their result and our meta-analytic findings, more information relating to the disease severity is needed.

There are several limitations in the current meta-analysis. First, although we tried to include studies that used similar diagnostic

criteria for aMCI,^{4,59} subject heterogeneity, such as subtypes of aMCI that could influence neuroimaging results,⁶⁰ cannot be ruled out. Second, the significance level of the contributing results was not taken into consideration by the current ALE technique. Nonetheless, our study did adjust for the effect of different sample sizes. Third, different analytic approaches and imaging modalities could reveal different aspects of resting-state abnormalities in the brain.⁶¹ For instance, regional homogeneity and amplitude of low-frequency fluctuations measure the local synchrony and power spectrum of low-frequency signals, respectively, whereas independent component analysis separates linearly mixed low-frequency signals, which are all indirect measures of neural activity. In addition, arterial spin labeling perfusion MRI can indirectly reflect the regional brain metabolism and neural activity during resting state by measuring regional cerebral blood flow during task-free condition. Nonetheless, different analytic approaches and modalities could be complementary to each other and provide a more complete picture of the field.⁶¹ Given that the number of studies using each individual method is not sufficient for conducting independent ALE studies, our results represent a common pattern of resting-state abnormalities that could be revealed across imaging methodologies and have high generalizability toward understanding the neuropathology of aMCI. As shown by our findings, different analytic approaches or modalities could generate similar results, such that reduced resting-state activity was found in PCC in most of our included studies regardless of their analytic approaches or imaging modality. Finally, other confounds such as mini-mental state examination scores, age, gender and education levels that could influence resting-state activity in aMCI could not be controlled for in the current ALE analysis due to the limitation of the current analysis technique. Future studies that incorporate these variables as covariates would provide more conclusive findings on resting-state abnormality in aMCI.

CONCLUSIONS

The current meta-analysis supports that abnormalities in resting-state activity in the PCC, right angular gyrus (extending to middle temporal gyrus), right parahippocampal gyrus, left fusiform gyrus, left middle temporal gyrus and left supramarginal gyrus are commonly found in aMCI patients. These regional abnormalities in resting-state activity may serve as neuroimaging markers for the early detection of aMCI.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by May Endowed Professorship in Neuropsychology by The University of Hong Kong and the generous donation from Chow Tai Fook Charity Foundation and Henderson Warmth Foundation.

REFERENCES

- Petersen RC, Negash S. Mild cognitive impairment: an overview. *CNS Spectr* 2008; **13**: 45–53.
- Zhao Z, Lu J, Jia X, Chao W, Han Y, Jia J et al. Selective changes of resting-state brain oscillations in aMCI: an fMRI study using ALFF. *Biomed Res Int* 2014; **2014**: 920902.
- Langa KM, Levine DA. The diagnosis and management of mild cognitive impairment: a clinical review. *JAMA* 2014; **312**: 2551–2561.
- Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV et al. Current concepts in mild cognitive impairment. *Arch Neurol* 2001; **58**: 1985–1992.
- Vassilaki M, Cha RH, Geda YE, Mielke MM, Knopman DS, Petersen RC et al. Mortality in mild cognitive impairment varies by subtype, sex, and lifestyle factors: the mayo clinic study of aging. *J Alzheimers Dis* 2015; **45**: 1237–1245.

- Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011; **7**: 270–279.
- Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci USA* 2001; **98**: 676–682.
- Buckner RL, Vincent JL. Unrest at rest: default activity and spontaneous network correlations. *Neuroimage* 2007; **37**: 1091–1096, discussion 7–9.
- Binder JR, Frost JA, Hammeke TA, Bellgowan PS, Rao SM, Cox RW. Conceptual processing during the conscious resting state. A functional MRI study. *J Cogn Neurosci* 1999; **11**: 80–95.
- Greicius MD, Menon V. Default-mode activity during a passive sensory task: uncoupled from deactivation but impacting activation. *J Cogn Neurosci* 2004; **16**: 1484–1492.
- McKiernan KA, Kaufman JN, Kucera-Thompson J, Binder JR. A parametric manipulation of factors affecting task-induced deactivation in functional neuroimaging. *J Cogn Neurosci* 2003; **15**: 394–408.
- Li HJ, Hou XH, Liu HH, Yue CL, He Y, Zuo XN. Toward systems neuroscience in mild cognitive impairment and Alzheimer's disease: a meta-analysis of 75 fMRI studies. *Hum Brain Mapp* 2015; **36**: 1217–1232.
- Yi L, Wang J, Jia L, Zhao Z, Lu J, Li K et al. Structural and functional changes in subcortical vascular mild cognitive impairment: a combined voxel-based morphometry and resting-state fMRI study. *PLoS One* 2012; **7**: e44758.
- Song J, Qin W, Liu Y, Duan Y, Liu J, He X et al. Aberrant functional organization within and between resting-state networks in AD. *PLoS One* 2013; **8**: e63727.
- Xi Q, Zhao X, Wang P, Guo Q, Jiang H, Cao X et al. Spontaneous brain activity in mild cognitive impairment revealed by amplitude of low-frequency fluctuation analysis: a resting-state fMRI study. *Radiol Med* 2012; **117**: 865–871.
- Talairach J, Tournoux P. *Co-planar Stereotaxic Atlas of the Human Brain: 3-Dimensional Proportional System—An Approach to Cerebral Imaging*. Thieme Medical Publishers: New York, NY, USA, 1988.
- Liang P, Xiang J, Liang H, Qi Z, Li K, Alzheimer's Disease Neuroimaging Initiative. Altered amplitude of low-frequency fluctuations in early and late mild cognitive impairment and Alzheimer's disease. *Curr Alzheimer Res* 2014; **11**: 389–398.
- Xu G, Antuono PG, Jones J, Xu Y, Wu G, Ward D et al. Perfusion fMRI detects deficits in regional CBF during memory-encoding tasks in MCI subjects. *Neurology* 2007; **69**: 1650–1656.
- Sorg C, Riedl V, Muhlau M, Calhoun VD, Eichele T, Laer L et al. Selective changes of resting-state networks in individuals at risk for Alzheimer's disease. *Proc Natl Acad Sci USA* 2007; **104**: 18760–18765.
- Bai F, Zhang Z, Yu H, Shi Y, Yuan Y, Zhu W et al. Default-mode network activity distinguishes amnesic type mild cognitive impairment from healthy aging: a combined structural and resting-state functional MRI study. *Neurosci Lett* 2008; **438**: 111–115.
- Qi Z, Wu X, Wang Z, Zhang N, Dong H, Yao L et al. Impairment and compensation coexist in amnesic MCI default mode network. *Neuroimage* 2010; **50**: 48–55.
- Zhang Z, Deng L, Bai F, Shi Y, Yu H, Yuan Y et al. Alteration of resting brain function by genetic variation in angiotensin converting enzyme in amnesic-type mild cognitive impairment of Chinese Han. *Behav Brain Res* 2010; **208**: 619–625.
- Wang Z, Yan C, Zhao C, Qi Z, Zhou W, Lu J et al. Spatial patterns of intrinsic brain activity in mild cognitive impairment and Alzheimer's disease: a resting-state functional MRI study. *Hum Brain Mapp* 2011; **32**: 1720–1740.
- Bai F, Watson DR, Shi Y, Wang Y, Yue C, Teng Y et al. Specifically progressive deficits of brain functional marker in amnesic type mild cognitive impairment. *PLoS One* 2011; **6**: e24271.
- Han Y, Lui S, Kuang W, Lang Q, Zou L, Jia J. Anatomical and functional deficits in patients with amnesic mild cognitive impairment. *PLoS One* 2012; **7**: e28664.
- Bai F, Shi Y, Yuan Y, Yue C, Zhuang L, Xu X et al. Association of a GSK-3beta polymorphism with brain resting-state function in amnesic-type mild cognitive impairment. *J Alzheimers Dis* 2012; **32**: 387–396.
- Wang Z, Nie B, Li D, Zhao Z, Han Y, Song H et al. Effect of acupuncture in mild cognitive impairment and Alzheimer disease: a functional MRI study. *PLoS One* 2012; **7**: e42730.
- Zhuang L, Liu X, Xu X, Yue C, Shu H, Bai F et al. Association of the interleukin 1 beta gene and brain spontaneous activity in amnesic mild cognitive impairment. *J Neuroinflammation* 2012; **9**: 263.
- Xi Q, Zhao XH, Wang PJ, Guo QH, He Y. Abnormal intrinsic brain activity in amnesic mild cognitive impairment revealed by amplitude of low-frequency fluctuation: a resting-state functional magnetic resonance imaging study. *Chin Med J (Engl)* 2013; **126**: 2912–2917.
- Zhou X, Zhang J, Chen Y, Ma T, Wang Y, Wang J et al. Aggravated cognitive and brain functional impairment in mild cognitive impairment patients with type 2 diabetes: a resting-state functional MRI study. *J Alzheimers Dis* 2014; **41**: 925–935.
- Liu X, Bai F, Yue C, Shi Y, Yu H, Luo B et al. The association between TOMM40 gene polymorphism and spontaneous brain activity in amnesic mild cognitive impairment. *J Neurol* 2014; **261**: 1499–1507.

- 32 Liu Z, Wei W, Bai L, Dai R, You Y, Chen S et al. Exploring the patterns of acupuncture on mild cognitive impairment patients using regional homogeneity. *PLoS One* 2014; **9**: e99335.
- 33 Wang Y, Zhao X, Xu S, Yu L, Wang L, Song M et al. Using regional homogeneity to reveal altered spontaneous activity in patients with mild cognitive impairment. *Biomed Res Int* 2015; **2015**: 807093.
- 34 Zhou Y, Yu F, Duong TQ, Alzheimer's Disease Neuroimaging Initiative. White matter lesion load is associated with resting state functional MRI activity and amyloid PET but not FDG in mild cognitive impairment and early Alzheimer's disease patients. *J Magn Reson Imaging* 2015; **41**: 102–109.
- 35 Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol* 2005; **5**: 13.
- 36 Eickhoff SB, Bzdok D, Laird AR, Kurth F, Fox PT. Activation likelihood estimation meta-analysis revisited. *Neuroimage* 2012; **59**: 2349–2361.
- 37 Turkeltaub PE, Eickhoff SB, Laird AR, Fox M, Wiener M, Fox P. Minimizing within-experiment and within-group effects in Activation Likelihood Estimation meta-analyses. *Hum Brain Mapp* 2012; **33**: 1–13.
- 38 Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT. Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: a random-effects approach based on empirical estimates of spatial uncertainty. *Hum Brain Mapp* 2009; **30**: 2907–2926.
- 39 Lancaster JL, Tordeillas-Gutierrez D, Martinez M, Salinas F, Evans A, Zilles K et al. Bias between MNI and Talairach coordinates analyzed using the ICBM-152 brain template. *Hum Brain Mapp* 2007; **28**: 1194–1205.
- 40 De Jager CA, Hogervorst E, Combrinck M, Budge MM. Sensitivity and specificity of neuropsychological tests for mild cognitive impairment, vascular cognitive impairment and Alzheimer's disease. *Psychol Med* 2003; **33**: 1039–1050.
- 41 Gusnard DA, Raichle ME, Raichle ME. Searching for a baseline: functional imaging and the resting human brain. *Nat Rev Neurosci* 2001; **2**: 685–694.
- 42 Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 2008; **1124**: 1–38.
- 43 Aminoff EM, Kveraga K, Bar M. The role of the parahippocampal cortex in cognition. *Trends Cogn Sci* 2013; **17**: 379–390.
- 44 Fujishima M, Maikusa N, Nakamura K, Nakatsuka M, Matsuda H, Meguro K. Mild cognitive impairment, poor episodic memory, and late-life depression are associated with cerebral cortical thinning and increased white matter hyperintensities. *Front Aging Neurosci* 2014; **6**: 306.
- 45 Minoshima S, Giordani B, Berent S, Frey KA, Foster NL, Kuhl DE. Metabolic reduction in the posterior cingulate cortex in very early Alzheimer's disease. *Ann Neurol* 1997; **42**: 85–94.
- 46 Zhou Y, Dougherty JH, Hubner KF, Bai B, Cannon RL, Hutson RK. Abnormal connectivity in the posterior cingulate and hippocampus in early Alzheimer's disease and mild cognitive impairment. *Alzheimers Dement* 2008; **4**: 265–270.
- 47 Choo IH, Lee DY, Oh JS, Lee JS, Lee DS, Song IC et al. Posterior cingulate cortex atrophy and regional cingulum disruption in mild cognitive impairment and Alzheimer's disease. *Neurobiol Aging* 2010; **31**: 772–779.
- 48 Boccia M, Acierno M, Piccardi L. Neuroanatomy of Alzheimer's disease and late-life depression: a coordinate-based meta-analysis of MRI studies. *J Alzheimers Dis* 2015; **46**: 963–970.
- 49 Powell HW, Guye M, Parker GJ, Symms MR, Boulby P, Koepp MJ et al. Noninvasive in vivo demonstration of the connections of the human parahippocampal gyrus. *Neuroimage* 2004; **22**: 740–747.
- 50 Hamalainen A, Pihlajamaki M, Tanila H, Hanninen T, Niskanen E, Tervo S et al. Increased fMRI responses during encoding in mild cognitive impairment. *Neurobiol Aging* 2007; **28**: 1889–1903.
- 51 Carcaillon L, Amieva H, Auriacombe S, Helmer C, Dartigues JF. A subtest of the MMSE as a valid test of episodic memory? Comparison with the Free and Cued Reminding Test. *Dement Geriatr Cogn Disord* 2009; **27**: 429–438.
- 52 Jonides J, Schumacher EH, Smith EE, Koeppel RA, Awh E, Reuter-Lorenz PA et al. The role of parietal cortex in verbal working memory. *J Neurosci* 1998; **18**: 5026–5034.
- 53 Bokde AL, Karmann M, Born C, Teipel SJ, Omerovic M, Ewers M et al. Altered brain activation during a verbal working memory task in subjects with amnesic mild cognitive impairment. *J Alzheimers Dis* 2010; **21**: 103–118.
- 54 Nishi H, Sawamoto N, Namiki C, Yoshida H, Dinh HD, Ishizu K et al. Correlation between cognitive deficits and glucose hypometabolism in mild cognitive impairment. *J Neuroimaging* 2010; **20**: 29–36.
- 55 Li X, Cao M, Zhang J, Chen K, Chen Y, Ma C et al. Structural and functional brain changes in the default mode network in subtypes of amnesic mild cognitive impairment. *J Geriatr Psychiatry Neurol* 2014; **27**: 188–198.
- 56 Förster S, Buschert VC, Teipel SJ, Frieze U, Buchholz HG, Drzezga A et al. Effects of a 6-month cognitive intervention on brain metabolism in patients with amnesic MCI and mild Alzheimer's disease. *J Alzheimers Dis* 2011; **26**: 337–348.
- 57 Sanabria-Diaz G, Martinez-Montes E, Melie-Garcia L, Alzheimer's Disease Neuroimaging Initiative. Glucose metabolism during resting state reveals abnormal brain networks organization in the Alzheimer's disease and mild cognitive impairment. *PLoS One* 2013; **8**: e68860.
- 58 Zhang Z, Liu Y, Jiang T, Zhou B, An N, Dai H et al. Altered spontaneous activity in Alzheimer's disease and mild cognitive impairment revealed by Regional Homogeneity. *Neuroimage* 2012; **59**: 1429–1440.
- 59 Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med* 2004; **256**: 183–194.
- 60 Li X, Zhang ZJ. Neuropsychological and neuroimaging characteristics of amnesic mild cognitive impairment subjects: a selective overview. *CNS Neurosci Ther* 2015; **21**: 776–783.
- 61 Alonso Bde C, Hidalgo Tobon S, Dies Suarez P, Garcia Flores J, de Celis Carrillo B, Barragan Perez E. A multi-methodological MR resting state network analysis to assess the changes in brain physiology of children with ADHD. *PLoS One* 2014; **9**: e99119.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Supplementary Information accompanies the paper on the Translational Psychiatry website (<http://www.nature.com/tp>)