



Serum proteins differentially expressed in early- and late-onset preeclampsia assessed using iTRAQ proteomics and bioinformatics analyses

Chengcheng Tu^{1,*}, Feng Tao^{1,*}, Ying Qin², Mingzhu Wu¹, Ji Cheng¹, Min Xie², Bing Shen², Junjiao Ren³, Xiaohong Xu⁴, Dayan Huang³ and Hongbo Chen¹

¹ Department of Obstetrics and Gynecology, Maternal and Child Health Hospital Affiliated to Anhui Medical University, Hefei, Anhui, China

² School of Basic Medicine, Anhui Medical University, Hefei, Anhui, China

³ Department of Science and Education, Maternal and Child Health Hospital Affiliated to Anhui Medical University, Hefei, Anhui, China

⁴ Department of Clinical Laboratory, Maternal and Child Health Hospital Affiliated to Anhui Medical University, Hefei, Anhui, China

* These authors contributed equally to this work.

ABSTRACT

Background. Preeclampsia remains a serious disorder that puts at risk the lives of perinatal mothers and infants worldwide. This study assessed potential pathogenic mechanisms underlying preeclampsia by investigating differentially expressed proteins (DEPs) in the serum of patients with early-onset preeclampsia (EOPE) and late-onset preeclampsia (LOPE) compared with healthy pregnant women.

Methods. Blood samples were collected from four women with EOPE, four women with LOPE, and eight women with normal pregnancies, with four women providing control samples for each preeclampsia group. Serum proteins were identified by isobaric tags for relative and absolute quantitation combined with liquid chromatography–tandem mass spectrometry. Serum proteins with differences in their levels compared with control groups of at least 1.2 fold-changes and that were also statistically significantly different between the groups at $P < 0.05$ were further analyzed. Bioinformatics analyses, including gene ontology and Kyoto Encyclopedia of Genes and Genomes signaling pathway analyses, were used to determine the key proteins and signaling pathways associated with the development of PE and to determine those DEPs that differed between women with EOPE and those with LOPE. Key protein identified by mass spectrometry was verified by enzyme linked immunosorbent assay (ELISA).

Results. Compared with serum samples from healthy pregnant women, those from women with EOPE displayed 70 proteins that were differentially expressed with significance. Among them, 51 proteins were significantly upregulated and 19 proteins were significantly downregulated. In serum samples from women with LOPE, 24 DEPs were identified, with 10 proteins significantly upregulated and 14 proteins significantly downregulated compared with healthy pregnant women. Bioinformatics analyses indicated that DEPs in both the EOPE and LOPE groups were associated with abnormalities in the activation of the coagulation cascade and complement system as well as with lipid metabolism. In addition, 19 DEPs in the EOPE group were closely related to placental development or invasion of tumor cells. Downregulation of

Submitted 26 November 2019

Accepted 28 July 2020

Published 1 September 2020

Corresponding author

Hongbo Chen,
chenhongbo@ahmu.edu.cn

Academic editor

Edward Braun

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.9753

© Copyright
2020 Tu et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

pregnancy-specific beta-1-glycoprotein 9 (PSG9) in the LOPE group was confirmed by ELISA.

Conclusion. The pathogenesis of EOPE and LOPE appeared to be associated with coagulation cascade activation, lipid metabolism, and complement activation. However, the pathogenesis of EOPE also involved processes associated with greater placental injury. This study provided several new proteins in the serum which may be valuable for clinical diagnosis of EOPE and LOPE, and offered potential mechanisms underpinning the development of these disorders.

Subjects Bioinformatics, Gynecology and Obstetrics, Women's Health

Keywords Preeclampsia, Early-onset preeclampsia, Late-onset preeclampsia, Proteomics, LC-MS/MS, iTRAQ

INTRODUCTION

Preeclampsia (PE), a hypertensive disorder complicating pregnancy, is the main cause of increased perinatal mortality among mothers and infants worldwide. The rate of PE in the United States is approximately 3.4% (*Ghulmiyyah & Sibai, 2012*), and it is higher in developing countries (*Ananth, Keyes & Wapner, 2013*). PE is marked by new-onset hypertension occurring after 20 weeks of gestation, accompanied by either new-onset proteinuria or systemic multiple organ damage (*Committee on Practice Bulletins—Obstetrics, 2019*). Severe PE may lead to convulsions, coma, cerebral hemorrhage, heart failure, placental abruption, disseminated intravascular coagulation, and even death (*Bibbins-Domingo et al., 2017*). Fetal delivery is the most effective treatment for PE, with other treatments of relieving symptoms used only as an attempt to gain time for enabling further maturity of the fetus. As a result, PE is the leading cause of premature birth and low birth weight, especially severe PE (*Ananth & Vintzileos, 2006*).

The exact pathogenesis of PE remains unclear. However, PE is considered a placenta-derived disease because the syndrome resolves once the placenta is removed. There are two subtypes of PE: early- and late-onset preeclampsia. Their pathogeneses are not identical. Early-onset preeclampsia (EOPE) has a higher degree of placental damage, whereas late-onset preeclampsia (LOPE) may focus on the interaction between the normal senescence of the placenta and a maternal genetic susceptibility to cardiovascular and metabolic disease (*Burton et al., 2019*). However, pathophysiological changes of the placenta further lead to endothelial dysfunction and systemic inflammatory response, which are their common pathogenesis link (*Young, Levine & Karumanchi, 2010*).

The placenta is in direct contact with the maternal circulation. Therefore, any changes in the protein expression of placental tissue are reflected in maternal serum proteins (*Norwitz, 2007*). If the types and levels of differentially expressed proteins (DEPs) in the serum of patients with PE can be researched quantitatively and holistically, specific serum biomarkers may be found for predicting PE and for further study of its pathogenesis. Compared with other proteomics methods, isobaric tags for relative and absolute quantitation (iTRAQ) combined with liquid chromatography–tandem mass spectrometry (LC-MS/MS)

is considered more effective for searching for serum or plasma biomarkers (*Moulder et al., 2018*). Therefore, in the present study, we used iTRAQ and LC-MS/MS methods to identify serum proteins differentially expressed between EOPE and LOPE and between women with PE and healthy pregnant women. We then used bioinformatics analyses to determine the key proteins and related signaling pathways associated with the development of EOPE and LOPE.

MATERIALS & METHODS

Participants and clinical samples

The Medical Research Ethics Committee of Anhui Medical University reviewed and approved our research protocol and an informed patient consent form (Anhui Medical Ethics approval No. 20150192). All patients signed the approved written informed consent form prior to being included in the study.

Blood samples were collected from December 2018 to May 2019 from 16 pregnant women, eight of whom had received a diagnosis of PE, and were in the Maternal and Child Health Hospital Affiliated with Anhui Medical University. A PE diagnosis was made consistent with the 2019 American College of Obstetricians and Gynecologists pregnancy hypertension guidance (*Committee on Practice Bulletins—Obstetrics, 2019*). All included pregnant women were primipara and without a disease that may have affected their serum protein levels, such as infection, multiple pregnancy, or gestational diabetes mellitus. Four serum samples were obtained from and allocated to each of the following four groups: EOPE, the EOPE control, LOPE, and the LOPE control groups.

Protein extraction and quality testing

High-abundance proteins were removed from the serum samples using Pierce Top 12 Abundant Protein Depletion Spin Columns (Thermo Fisher). The protein concentration was determined by the Bradford method. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) was performed to separate the proteins and evaluate the quality of samples.

Trypsin enzymatic hydrolysis and peptide iTRAQ isobaric labeling

After protein quantification, a centrifuge tube containing 60 μg of protein solution was mixed with 5 μL 1 M dithiothreitol at 37 °C for 1 h. Then 20 μL of 1 M iodoacetamide was added and the solution allowed to react for 1 h at room temperature. All samples were pipetted into ultrafiltration tubes, and the filtrate was discarded after centrifugation. UA buffer (8 M urea, 100 mM Tris-HCl, pH 8.0, 100 μL) was added and the sample centrifuged at 14,000 g for 10 min; this step was repeated twice. Then, 50 mM NH_4HCO_3 (50 μL) was added and the filtrate discarded after centrifugation; this step was repeated three times. Trypsin buffer (40 μL) was added and mixed, and the samples were centrifuged at 600 rpm for 1 min. The samples were then subjected to enzymatic hydrolysis at 37 °C for 12–18 h. Transfer the enzymatic hydrolysate to a new centrifuge tube. After labeling with 8plex iTRAQ reagents multiplex kit, the same volume of each sample was mixed together and desalted using a C18 cartridge.




Figure 1 Flowchart of the data analysis procedure.

Full-size DOI: [10.7717/peerj.9753/fig-1](https://doi.org/10.7717/peerj.9753/fig-1)

LC-MS/MS analysis

The labeled samples were redissolved in 40 μ L of 0.1% formic acid aqueous solution. The peptides were loaded onto a C18-reversed phase column (3 μ m C18 resin, 75 μ m \times 15 cm). The mobile phases consisted of 2% methyl cyanide/0.1% formic acid/98% water and 80% methyl cyanide/0.08% formic acid/20% water. The gradient for the B phase increased linearly at 0–68 min from 7% to 36% and at 68–75 min, from 36% to 100%. Each sample was separated by capillary high-performance liquid chromatography and was analyzed by an Orbitrap Fusion Lumos mass spectrometer (Thermo Science). After the data were collected, they were processed according to the flowchart shown in Fig. 1. Protein identification and quantification were done by using the ProteinPilot Software, version 4.2 (Sciex). Human proteome databases containing UniProt sequences were used to perform peptide identification. Serum proteins with fold changes in their levels compared with control groups of at least 1.2 (in two of two replicates) and that were also statistically significantly different between the groups at $P < 0.05$ were further analyzed.

Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) signaling pathway analyses

Metascape, a web-based resource (<http://metascape.org>) for gene and protein annotation, visualization, and integration discovery (Fang *et al.*, 2019; Soonthornvacharin *et al.*, 2017) was used to perform GO analyses. The KOBAS online analysis database (<http://kobas.cbi.pku.edu.cn/kobas3>) was used to performed KEGG pathway analyses (Kanehisa & Goto, 2000). A two-sided $P < 0.05$ was regarded significantly different.

Enzyme linked immunosorbent assay (ELISA)

PSG9, the downregulated protein in the LOPE group, was verified by ELISA kit (specification and catalogue number: 96T, OM626395). The experiment was performed according to the manufacturer instructions. Blood samples were collected from women with LOPE or normal pregnancies.

Statistical analysis

Two-tailed Mann–Whitney U test was performed with SigmaPlot software. Values are expressed as means \pm SEM. A value of $P < 0.05$ was considered statistically significant.

RESULTS

Participants

The mean and standard deviation (SD) age of the patients in the EOPE group was 31.8 ± 5.0 years, and they were at a mean (SD) gestation of 31.9 ± 2.9 weeks; the mean (SD) age of the participants in the EOPE control group was 30.0 ± 1.0 years, with a mean (SD) gestation of 31.5 ± 1.4 weeks. No significant difference was found in age ($P = 0.57$) or gestational weeks ($P = 0.84$) between these two groups. The mean (SD) age of the patients in the LOPE group was 24.8 ± 1.5 years, with 37.8 ± 1.5 weeks' gestation; the mean (SD) age of the participants in the LOPE control group was 27.0 ± 1.3 years, with a mean (SD) of 38.9 ± 1.3 weeks' gestation. No significant difference was found in age ($P = 0.15$) or gestation ($P = 0.37$) between these two groups.

SDS-PAGE

The total proteins in the molecular weight range of 10–220 kDa from 16 samples were effectively separated by SDS-PAGE. The proteins were not degraded, and the high-abundance proteins were obtained (Fig. 2).

MS/MS spectrum analysis and Identification of DEPs

LC-MS/MS is a powerful tool for identifying proteins in serum samples. We identified 413 serum proteins in the EOPE and EOPE healthy control groups, of which 70 were significantly and differentially expressed between the two groups, with 51 upregulated and 19 downregulated (Table 1, Fig. 3). We also identified 470 proteins in the LOPE and LOPE control groups, of which 24 proteins were significantly and differentially expressed between the two groups, with 10 upregulated and 14 downregulated (Table 2, Fig. 3). Clustergrams generated describing the expression of these DEPs indicated that the expression patterns between the patient EOPE or LOPE groups obviously differed from their control groups, but the two patient groups, EOPE and LOPE, clustered together (Figs. 3B and 3D).

GO functional annotation and enrichment analysis

GO analysis is an important method and tool in the field of bioinformatics. It includes three categories: cellular component, molecular function, and biological process. GO functional annotation analysis results show the number of DEPs under each item in the three categories. GO functional enrichment analysis provides significant GO functional




Figure 2 Removal of high-abundance proteins. (A) Lane E is the original sample from the early-onset preeclampsia (EOPE) group. Lane M contains markers. Lanes 1–4 are from serum samples obtained from the EOPE group after removing the highly abundant proteins, whereas lanes 5–8 are from the EOPE control group after removing the highly abundant proteins. (B) Lane L is the original sample from the late-onset preeclampsia (LOPE) group. Lane M contains markers. Lanes –1 to –4 are from the LOPE group after removing the highly abundant proteins, whereas lanes –5 to –8 are from the LOPE control group after removing the high-abundance proteins. The bands are clear and uniform and without protein degradation.

Full-size [DOI: 10.7717/peerj.9753/fig-2](https://doi.org/10.7717/peerj.9753/fig-2)

terms associated with the DEPs, that is, those biological functions significantly correlated with the DEP. In vivo, different proteins interact to generate a biological behavior, and a pathway-based analysis helps to further understand those biological functions. A significant pathway enrichment analysis can determine the most important biochemical and metabolic pathways and signal transduction pathways associated with the DEPs.

The GO functional annotation analysis results for the EOPE group are shown in Fig. 4. For the biological process category, the highest percentages of the proteins were associated with the term *biological regulation* ($n = 43$ proteins; with the top three upregulated proteins in this process being CHL1, LRP1, and CNTN1). For the cellular component category, the highest percentages of the proteins were associated with the term *extracellular region* ($n = 49$ proteins; with the top 2 upregulated proteins in this process being MMP2 and CHL1). For the molecular function category, the highest percentages of the proteins were associated with the term *binding* ($n = 45$ proteins; with the top 3 upregulated proteins in this process being MMP2, CHL1, and PEPD). For the LOPE group vs. their controls, the highest percentages of the proteins in the biological process category were associated with the term *biological regulation* ($n = 17$; with the top 3 upregulated proteins in this process being PAPP2, F7, and vWF). The highest percentages of the proteins in the cellular component category were associated with term *extracellular region* ($n = 18$; with the top three upregulated proteins in this process being PAPP2, CETP, and F7). The highest percentages of the proteins in the molecular function category were associated with the term *binding* ($n = 13$; with the top 3 upregulated proteins in this process being APPA2, CETP, and F7). The terms in the GO functional enrichment analysis for the EOPE group vs. their controls mainly included response to stress, defense response, and negative regulation of catalytic activity, whereas the terms in the GO functional enrichment analysis

Table 1 List of differentially expressed proteins in the early-onset preeclampsia group.

Regulated type	Protein accession	Gene name	Protein description	FC
UP	A0A024R6R4	MMP2	Matrix metalloproteinase 2	21.48
UP	A8K2X4	-	cDNA FLJ75401	12.87
UP	O00533	CHL1	Neural cell adhesion molecule L1-like protein	7.69
UP	J3K000	PEPD	PEPD protein	6.6
UP	A8K3I0	-	cDNA FLJ78437	6.31
UP	Q6UXB8	PI16	Peptidase inhibitor 16	5.76
UP	Q07954	LRP1	Prolow-density lipoprotein receptor-related protein 1	4.82
UP	Q12860	CNTN1	Contactin-1	4.64
UP	V9HWB4	HEL-S-89n	Epididymis secretory sperm binding protein Li 89n	4.63
UP	P35527	KRT9	Keratin, type I cytoskeletal 9	4.51
UP	P14543	NID1	Nidogen-1	3.86
UP	O95236	APOL3	Apolipoprotein L3	3.4
UP	A6XNE2	-	Complement factor D preproprotein	3.34
UP	P41222	PTGDS	Prostaglandin-H2	3.17
UP	V9H1C1	-	Gelsolin exon 4 (Fragment)	2.95
UP	Q9Y5Y7	LYVE1	Lymphatic vessel endothelial hyaluronic acid receptor 1	2.95
UP	Q13201	MMRN1	Multimerin-1	2.83
UP	B3KQF4	-	cDNA FLJ90373	2.74
UP	Q6MZL2	DKFZp686M0562	Uncharacterized protein DKFZp686M0562 (Fragment)	2.71
UP	J3KPA1	CRISP3	Cysteine-rich secretory protein 3	2.65
UP	P01034	CST3	Cystatin-C	2.64
UP	A0A0S2Z4F1	EFEMP1	EGF Containing Fibulin Extracellular Matrix Protein 1	2.62
UP	P23142	FBLN1	Fibulin-1	2.57
UP	A0A384N669	-	Epididymis secretory sperm binding protein	2.48
UP	J3KNB4	CAMP	Cathelicidin antimicrobial peptide	2.47
UP	B7Z544	-	cDNA FLJ51742	2.43
UP	A8K061	-	cDNA FLJ77880	2.42
UP	A0A087WV75	NCAM1	Neural cell adhesion molecule 1	2.37
UP	Q9NQ79	CRTAC1	Cartilage acidic protein 1	2.24
UP	B2RBW9	-	cDNA, FLJ95746	2.23
UP	A0A024RAA7	C1QC	Adiponectin B	2.15
UP	P16070	CD44	CD44 antigen	2.06
UP	Q76LX8	ADAMTS13	Von Willebrand Factor-Cleaving Protease	2.01
UP	Q6EMK4	VASN	Vasorin	2
UP	D6RE86	CP	Ceruloplasmin (Fragment)	1.88
UP	H6VRG1	KRT1	Keratin 1	1.87
UP	Q86U17	SERPINA11	Serpin A11	1.81
UP	Q9UKE5	TNIK	TRAF2 and NCK-interacting protein kinase	1.75
UP	P61626	LYZ	Lysozyme C	1.74
UP	P02743	APCS	Serum amyloid P-component	1.74

(continued on next page)

Table 1 (continued)

Regulated type	Protein accession	Gene name	Protein description	FC
UP	Q16853	AOC3	Membrane primary amine oxidase	1.73
UP	Q9NZP8	C1RL	Complement C1r subcomponent-like protein	1.73
UP	A0A024R035	C9	Complement C9	1.71
UP	A0A0A0MRJ7	F5	Coagulation factor V	1.66
UP	P09871	C1S	Complement C1s subcomponent	1.65
UP	Q6PIL8	IGK@	IGK@ protein	1.63
UP	P49908	SELENOP	Selenoprotein P	1.55
UP	P02654	APOC1	Apolipoprotein C-I	1.55
UP	D3DRR6	ITIH2	Inter-alpha (Globulin) inhibitor H2	1.54
UP	B4DPQ0	C1R	Complement C1r subcomponent	1.52
UP	J3KNP4	SEMA4B	Semaphorin-4B	1.51
DOWN	Q6GMX6	IGH@	IGH@ protein	0.56
DOWN	A0A0X9TD47	-	MS-D1 light chain variable region (Fragment)	0.43
DOWN	Q9UNU2	C4B	Complement protein C4B frameshift mutant (Fragment)	0.43
DOWN	P01591	JCHAIN	Immunoglobulin J chain	0.34
DOWN	Q14213	EBI3	Interleukin-27 subunit beta	0.34
DOWN	P28799	GRN	Granulins	0.33
DOWN	B2R7N9	-	cDNA, FLJ93532	0.32
DOWN	P00709	LALBA	Alpha-lactalbumin	0.29
DOWN	G5E9F7	PSG1	Pregnancy specific beta-1-glycoprotein 1	0.25
DOWN	A0A075B6A0	IGHM	Immunoglobulin heavy constant mu (Fragment)	0.23
DOWN	A0A075B6R9	IGKV2D-24	Immunoglobulin kappa variable 2D-24	0.23
DOWN	O75636	FCN3	Ficolin-3	0.22
DOWN	O43866	CD5L	CD5 antigen-like	0.2
DOWN	P06733	ENO1	Alpha-enolase	0.16
DOWN	P09172	DBH	Dopamine beta-hydroxylase	0.16
DOWN	Q15485	FCN2	Ficolin-2	0.13
DOWN	Q86TT1		Full-length cDNA clone CS0DD006YL02 of Neuroblastoma	0.08
DOWN	P00739	HPR	Haptoglobin-related protein	0.06
DOWN	P00738	HP	Haptoglobin	0.01

for the LOPE group vs. their controls included enzyme inhibitor activity and serine-type endopeptidase inhibitor activity (Fig. 5).

KEGG signaling pathway analysis

The KOBAS online analysis tool was used to identify the functions associated with the DEPs and the KEGG signaling pathways. Results of the KEGG pathway enrichment analysis showed that complement and coagulation cascades, proteoglycans in cancer, and metabolic pathways were the main signaling pathways associated with the EOPE group, whereas complement and coagulation cascades were the main signaling pathways associated with the LOPE group (Fig. 6). Thus, the results indicated that a functional change in coagulation was the main finding for both EOPE and LOPE.




Figure 3 Differential protein expression. (A and C) Volcano plots with red dots on the right-hand side indicating upregulation, green dots on the left-hand side indicating downregulation, and black dots indicating no significant change in protein expression levels based on the criteria of an absolute log₁₀ fold change and $P < 0.05$ between early-onset preeclampsia and its control group (A), and between late-onset preeclampsia and its control group (C). (B and D) Clustergram for the expression of the DEPs between early-onset preeclampsia and control (B) and between late-onset preeclampsia and control (D).

Full-size [DOI: 10.7717/peerj.9753/fig-3](https://doi.org/10.7717/peerj.9753/fig-3)

PSG9 protein level in serum

Serum level of PSG9 was measured by ELISA. The data showed a significant difference between LOPE group and control group (4.54 ± 1.22 vs. 6.32 ± 1.73 , Fig. 7). The result is consistent with the study of mass spectrometry.

DISCUSSION

Proteomics methods have provided some important information regarding PE. For example, *Blumenstein et al. (2009)* used a differential in gel electrophoresis-based approach to identify changes in the plasma proteome of pregnant women who subsequently developed PE. They found that those DEPs are mainly involved in lipoprotein metabolism, the blood coagulation system, and the complement system. The results of the present study were

Table 2 List of differentially expressed proteins in the late-onset preeclampsia group.

Regulated type	Protein accession	GENE NAME	PROTEIN DESCRIPTION	FC
UP	Q9BXP8	PAPPA2	Pappalysin-2	5.27
UP	A0A0S2Z3F6	CETP	Cholesteryl ester transfer protein plasma isoform 1	3.51
UP	P08709	F7	Coagulation factor VII	2.8
UP	Q6NS95	IGL@	IGL@ protein	2.67
UP	P04275	VWF	von Willebrand factor	2.46
UP	C0JYY2	APOB	Apolipoprotein B	1.93
UP	A0A140VK24	-	Testicular secretory protein Li 24 protein 1	1.74
UP	P05546	SERPIND1	Heparin cofactor 2	1.52
UP	O95445	APOM	Apolipoprotein M	1.34
UP	P80108	GPLD1	Phosphatidylinositol-glycan-specific phospholipase D	1.26
DOWN	P05543	SERPINA7	Thyroxine-binding globulin	0.82
DOWN	Q9NZP8	C1RL	Complement C1r subcomponent-like protein	0.63
DOWN	A8K2T7	-	Receptor protein-tyrosine kinase	0.61
DOWN	Q00887	PSG9	Pregnancy-specific beta-1-glycoprotein 9	0.6
DOWN	P07333	CSF1R	Macrophage colony-stimulating factor 1 receptor	0.56
DOWN	Q2L9S7	AAT	Alpha-1-antitrypsin MBrescia variant receptor 1	0.53
DOWN	A0A2S0BDD1	ATIII-R2	Antithrombin III isoform	0.49
DOWN	Q8WW79	SELL	L-selectin	0.48
DOWN	P40189	IL6ST	Interleukin-6 receptor subunit beta (Fragment)	0.47
DOWN	B2R7Y0	-	cDNA, FLJ93654	0.41
DOWN	Q12860	CNTN1	member 2 (SERPINB2)	0.36
DOWN	Q8WWZ8	OIT3	Oncoprotein-induced transcript 3 protein	0.31
DOWN	A0A140TA29	C4B	Complement C4-B	0.21
DOWN	B2R950	-	cDNA, FLJ94213	0.18

consistent with those of *Blumenstein et al. (2009)*, but we also made some new discoveries (discussed below). These results suggest that the pathogenesis of PE is complex and is associated with multiple proteins and signaling pathways.

To maintain pregnancy and fetal growth progress within normal reference ranges, healthy pregnant women develop physiological hyperlipidemia. However, they do not experience angiopathy because high-density lipoprotein (a vascular protection factor) and low-density lipoprotein (LDL, an atherogenic factor) also increase to protect the vascular endothelium from injury (*Jin et al., 2016; Wang et al., 2017*). Compared with healthy pregnant women, pregnant women with PE display significantly increased serum triglycerides and LDLs, which may enhance oxidative stress and ultimately lead to vascular endothelial cell injury (*Huda, Sattar & Freeman, 2009; Pohanka, 2013*). Our study found that there were six DEPs associated with lipid regulation increased in the PE groups, including ApoB, LRP1, ApoL3, ApoC-I, CETP, and ApoM. ApoB, the main carrier protein of LDL, is an atherogenic risk factor. Previous evidence supports that ApoB is significantly upregulated in the plasma of patients with PE (*Lin et al., 2019*). CETP plays an important role in high-density lipoprotein metabolism and reverse cholesterol transport and is upregulated in the third trimester of pregnancy. Another study has reported that the TaqIB




Figure 4 Gene ontology functional annotation. The upregulated and downregulated proteins in the three categories of biological process, cellular component, and molecular function in (A) the early-onset preeclampsia group vs. its control group and (B) the late-onset preeclampsia group vs. its control group. Full-size [DOI: 10.7717/peerj.9753/fig-4](https://doi.org/10.7717/peerj.9753/fig-4)

polymorphism of the *CETP* gene is significantly correlated with the triglyceride and total cholesterol levels in patients with severe PE (Belo *et al.*, 2004). Here, we reported for the first time, to our knowledge, that LRP1, ApoC1, and ApoL3 are increased in the EOPE group. LRP1 and ApoC1 accelerate the development of atherosclerosis through different pathways (Mueller *et al.*, 2018; Westerterp *et al.*, 2007). Endothelial function decreases during PE, and endothelial dysfunction is a characteristic of atherosclerosis. ApoL3, as a regulator of MAPK and FAK signaling in endothelial cells, has been shown to be involved in angiogenesis *in vitro* (Khalil *et al.*, 2018). Increased ApoL3 in PE may be a compensatory response to endothelial dysfunction. Therefore, the current evidence suggests that dyslipidemia may be related to the development of PE, but the specific pathogenesis remains to be explored in future studies.

Physiological hypercoagulability in healthy pregnant women can prevent intrapartum and postpartum hemorrhage. However, abnormalities in coagulation, the anti-coagulation




Figure 5 Gene ontology (GO) functional enrichment. Enriched GO terms for the upregulated and downregulated proteins and the associated protein numbers in (A) early-onset preeclampsia vs. its control and (B) late-onset preeclampsia vs. its control.

Full-size DOI: [10.7717/peerj.9753/fig-5](https://doi.org/10.7717/peerj.9753/fig-5)

system, and the fibrinolytic system lead to pathological hypercoagulability in PE (*Dusse et al., 2011*). In the LOPE group of the present study, coagulation factors VII, vWF, and SRPIND1 were upregulated group, whereas AAT and ATIII were downregulated compared with healthy controls. In the EOPE group, coagulation factor V, plasminogen, and ADAMTS13 were upregulated. These results are consistent with those of many previous trials (discussed below). The level of plasma FVII in patients with severe PE is significantly higher than that in healthy pregnant women; therefore plasma FVII levels may show high sensitivity and specificity in differentiating between PE and normal pregnancy (*Dusse et al., 2016*). The coagulation factor vWF is a specific marker that reflects damage of endothelial cells; thus, damaged microvascular endothelial cells in PE promote the expression of




Figure 6 Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment. KEGG pathways enriched for the upregulated and downregulated proteins and the associated protein numbers in (A) early-onset preeclampsia vs. its control and (B) late-onset preeclampsia vs. its control.

Full-size [DOI: 10.7717/peerj.9753/fig-6](https://doi.org/10.7717/peerj.9753/fig-6)

vWF. Owing to the activation of the intrinsic and exogenous coagulation pathways by damaged endothelial cells, a large number of coagulation factors are activated. This leads to a massive generation of thrombin, which antagonizes a large amount of anti-thrombin III, resulting in a significant decrease in its level (Demir & Dilek, 2010; Pottecher et al., 2009). In response to abnormalities in the coagulation mechanism of PE, evidence-based medical study has also shown that oral administration of low-dose aspirin during early pregnancy can significantly reduce the incidence rate of PE (Rolnik et al., 2017).

A growing number of studies have shown that abnormal expression with the complement system is associated with PE (Agostinis et al., 2010; Derzsy et al., 2010). Our results support this evidence, showing that expression levels of CFD, CIQC, C1RL, C9, C1S, and C1R were




Figure 7 Protein level of pregnancy-specific beta-1-glycoprotein 9 (PSG9) in serum. Concentration of PSG9 in the patient serum in control and LOPE groups. Values are shown as the mean \pm SEM (n = 10); *P < 0.05 for Control vs. LOPE.

Full-size DOI: 10.7717/peerj.9753/fig-7

significantly increased, while expression levels of C4B, FCN2, and FCN3 were significantly decreased in the EOPE group compared with their control group. Moreover, the expression levels of complement C1RL and C4B were decreased in the LOPE group compared with their controls.

Downregulation of PSG1 and PSG9 expressed in the PE groups is an interesting finding in our study. PSG is a pregnancy-specific glycoprotein that is synthesized and secreted into the blood by placental syncytiotrophoblast cells. *Rong et al. (2017)* found that PSG9 significantly promotes the angiogenesis of human umbilical vein endothelial cells. *Chang et al. (2016)* found a significant increase in the number of cases with deletions in the PSG gene locus among patients with PE. We hypothesize that downregulation of the PSG protein family may be involved in the pathogenesis of PE by affecting the proliferation of endothelial cells although no specific experimental evidence currently supports or disproves this hypothesis.

PE is considered a placenta-derived disease. In the classic two-stage model, placental stress leads to dysfunction of maternal peripheral endothelial cells, systemic inflammatory response, and the clinical syndrome of PE (*Staff, 2019*). This model is also reflected in our mass spectrometry results. In both the EOPE and LOPE groups, abnormalities of coagulation cascade activation, lipid metabolism, and complement activation were found. However, the main stressor associated with EOPE is placental hypoperfusion secondary to impaired spiral artery remodeling; by contrast, in LOPE, the cause is more likely attributable to a mismatch between normal maternal perfusion and the metabolic demands of the placenta and fetus (*Burton et al., 2019*). Spiral artery remodeling is completed by the placental trophoblast cells continuously invading and destroying the uterine spiral arterial wall and gradually replacing the endothelial cells. Many studies have focused on the control of this invasion (*Pollheimer et al., 2018; Wagner, Otomo & Christians, 2011; Wu et al., 2016*).

Unexpectedly, we also found 19 DEPs in the EOPE group that were closely related to placental development or invasion of tumor cells. This result was not detected in the LOPE group, which showed an increase only in PAPP2, which affects the invasion and metastasis of placental trophoblasts and leads to a gradual decline in placental function (Wagner, Otomo & Christians, 2011). LYVE1, CST3, and NCAM1 are also reportedly involved in placental vascular remodeling (Pawlak et al., 2019; Song et al., 2010; Zhang, Xu & Han, 2019). GRN, CD5L, ENO1, CHL1, CRISP3, VASN, and TNIK promote the invasive ability of tumor cells (Aran et al., 2018; Bhandari et al., 2019; Buhusi et al., 2003; Chon et al., 2016; Song et al., 2014; Voshtani et al., 2019; Wang et al., 2019). By contrast, SEMA4B, KRT1, KRT9, FBLN1, and PTGDS are involved in the anti-invasive activity of tumor cells (Blanckaert et al., 2015; Jian et al., 2015; Marano et al., 2018; Zhang et al., 2018). MMP2 plays an important role in trophoblast invasion and is generally thought to be downregulated in PE (Wu et al., 2016). We found the opposite result, that is, the expression of MMP2 was upregulated in EOPE; however, expression of PI16, which inhibits MMP2 activity (Hazell et al., 2016), was also upregulated. PEPD is a hydrolase that affects collagen biosynthesis, cell proliferation, and matrix remodeling. Pehlivan et al. (2017) have shown that PEPD activity in the plasma, umbilical cord, and placental tissue of women with pregnancy-induced hypertension is higher than that of healthy pregnant women. Our results also indicated an upregulation of PEPD in the EOPE group. The dysregulation of the expression of these proteins may be an important cause of placental dysfunction in EOPE.

Furthermore, previous studies have shown that soluble form of placenta-derived endoglin (sENG) was upregulated in the serum of EOPE, which is involved in endothelial dysfunction in coordination with soluble fms-like tyrosine kinase (Venkatesha et al., 2006). Our mass spectrometry results were similar to that previous finding showing that endoglin was upregulated in EOPE but not in LOPE group. This finding may also indicate more severe endothelial dysfunction in EOPE.

CONCLUSIONS

In summary, the use of iTRAQ combined with LC-MS/MS was effective for screening serum for DEPs in PE. We used bioinformatics to analyze the DEPs that showed significant changes in their expression levels to provide potential indicators for detecting PE and ideas for the study of pathogenesis in PE. In conclusion, the pathogenesis of EOPE and LOPE appeared to be related to dysfunctions in coagulation cascade activation, lipid metabolism, and complement activation. However, compared with that in LOPE, the pathogenesis of EOPE was associated with greater placental injury. This study provides several new proteins in the serum which may potentially have value in clinical diagnosis of EOPE and LOPE and offers potential mechanisms for the development of EOPE and LOPE, contributing to the future research on prediction, prevention, and pathogenesis of PE.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the Anhui Province Science and Technology Innovation Project Demonstration Project (No. 201707d08050003), and the Anhui Province Key Research and Development Project (No. 201904a07020032). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Anhui Province Science and Technology Innovation Project Demonstration Project: 201707d08050003.

The Anhui Province Key Research and Development Project: 201904a07020032.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Chengcheng Tu, Feng Tao and Ying Qin performed the experiments, prepared figures and/or tables, and approved the final draft.
- Mingzhu Wu and Dayan Huang analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Ji Cheng analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Min Xie performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Bing Shen conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Junjiao Ren analyzed the data, prepared figures and/or tables, and approved the final draft.
- Xiaohong Xu performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Hongbo Chen conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Medical Research Ethics Committee of Anhui Medical University reviewed and approved our research protocol and an informed patient consent form (Anhui Medical Ethics approval No. 20150192).

Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Medical Research Ethics Committee of Anhui Medical University reviewed and approved our research protocol and an informed patient consent form (Anhui Medical Ethics approval No. 20150192).

Data Availability

The following information was supplied regarding data availability:

The raw measurements are available as [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9753#supplemental-information>.

REFERENCES

- Agostinis C, Bulla R, Tripodo C, Gismondi A, Stabile H, Bossi F, Guarnotta C, Garlanda C, De Seta F, Spessotto P, Santoni A, Ghebrehiwet B, Girardi G, Tedesco F. 2010. An alternative role of C1q in cell migration and tissue remodeling: contribution to trophoblast invasion and placental development. *The Journal of Immunology* 185(7):4420–4429 DOI 10.4049/jimmunol.0903215.
- Ananth CV, Keyes KM, Wapner RJ. 2013. Pre-eclampsia rates in the United States, 1980–2010: age-period-cohort analysis. *British Medical Journal* 347:f6564 DOI 10.1136/bmj.f6564.
- Ananth CV, Vintzileos AM. 2006. Maternal-fetal conditions necessitating a medical intervention resulting in preterm birth. *American Journal of Obstetrics and Gynecology* 195(6):1557–1563 DOI 10.1016/j.ajog.2006.05.021.
- Aran G, Sanjurjo L, Barcena C, Simon-Coma M, Tellez E, Vazquez-Vitali M, Garrido M, Guerra L, Diaz E, Ojanguren I, Elortza F, Planas R, Sala M, Armengol C, Sarrias MR. 2018. CD5L is upregulated in hepatocellular carcinoma and promotes liver cancer cell proliferation and antiapoptotic responses by binding to HSPA5 (GRP78). *FASEB Journal* 32(7):3878–3891 DOI 10.1096/fj.201700941RR.
- Belo LS, Gaffney D, Caslake M, Santos-Silva A, Pereira-Leite Ls, Quintanilha A, Rebelo I. 2004. Apolipoprotein E and cholesteryl ester transfer protein polymorphisms in normal and preeclamptic pregnancies. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 112(1):9–15 DOI 10.1016/S0301-2115(03)00240-9.
- Bhandari A, Guan Y, Xia E, Huang Q, Chen Y. 2019. VASN promotes YAP/TAZ and EMT pathway in thyroid carcinogenesis in vitro. *American Journal of Translational Research* 11(6):3589–3599.
- Bibbins-Domingo K, Grossman DC, Curry SJ, Barry MJ, Davidson KW, Doubeni CA, Epling Jr JW, Kemper AR, Krist AH, Kurth AE, Landefeld CS, Mangione CM, Phillips WR, Phipps MG, Silverstein M, Simon MA, Tseng CW. 2017.

- Screening for Preeclampsia: US preventive services task force recommendation statement. *The Journal of American Medical Association* **317**(16):1661–1667 DOI [10.1001/jama.2017.3439](https://doi.org/10.1001/jama.2017.3439).
- Blanckaert V, Kerviel V, Lepinay A, Joubert-Durigneux V, Hondermarck H, Chenais B.** 2015. Docosahexaenoic acid inhibits the invasion of MDA-MB-231 breast cancer cells through upregulation of cytokeratin-1. *International Journal of Oncology* **46**(6):2649–2655 DOI [10.3892/ijco.2015.2936](https://doi.org/10.3892/ijco.2015.2936).
- Blumenstein M, McMaster MT, Black MA, Wu S, Prakash R, Cooney J, McCowan LM, Cooper GJ, North RA.** 2009. A proteomic approach identifies early pregnancy biomarkers for preeclampsia: novel linkages between a predisposition to preeclampsia and cardiovascular disease. *Proteomics* **9**:2929–2945 DOI [10.1002/pmic.200800625](https://doi.org/10.1002/pmic.200800625).
- Buhusi M, Midkiff BR, Gates AM, Richter M, Schachner M, Maness PF.** 2003. Close homolog of L1 is an enhancer of integrin-mediated cell migration. *Journal of Biological Chemistry* **278**(27):25024–25031 DOI [10.1074/jbc.M303084200](https://doi.org/10.1074/jbc.M303084200).
- Burton GJ, Redman CW, Roberts JM, Moffett A.** 2019. Pre-eclampsia: pathophysiology and clinical implications. *British Medical Journal* **366**:l2381.
- Chang CL, Chang CY, Lee DX, Cheng PJ.** 2016. Characterization of human pregnancy specific glycoprotein (PSG) gene copy number variations in pre-eclampsia patients. *Advances in Experimental Medicine and Biology* **924**:63–65 DOI [10.1007/978-3-319-42044-8-12](https://doi.org/10.1007/978-3-319-42044-8-12).
- Chon HJ, Lee Y, Bae KJ, Byun BJ, Kim SA, Kim J.** 2016. Traf2- and Nck-interacting kinase (TNIK) is involved in the anti-cancer mechanism of dovitinib in human multiple myeloma IM-9 cells. *Amino Acids* **48**(7):1591–1599 DOI [10.1007/s00726-016-2214-3](https://doi.org/10.1007/s00726-016-2214-3).
- Committee on Practice Bulletins—Obstetrics.** 2019. ACOG Practice Bulletin No. 202: gestational hypertension and preeclampsia. *Obstetricians and Gynecologists* **133**:e1–e25.
- Demir C, Dilek I.** 2010. Natural coagulation inhibitors and active protein c resistance in preeclampsia. *Clinics (Sao Paulo)* **65**(11):1119–1122 DOI [10.1590/S1807-59322010001100011](https://doi.org/10.1590/S1807-59322010001100011).
- Derzsy Z, Prohaszka Z, Rigo Jr J, Fust G, Molvarec A.** 2010. Activation of the complement system in normal pregnancy and preeclampsia. *Molecular Immunology* **47**(7–8):1500–1506 DOI [10.1016/j.molimm.2010.01.021](https://doi.org/10.1016/j.molimm.2010.01.021).
- Dusse LM, Godoi LC, Gomes KB, Carvalho M, Lwaleed BA.** 2016. Tissue factor-dependent pathway in severe preeclampsia revisited: a Brazilian cohort study. *Blood Coagul Fibrinolysis* **27**(4):436–440 DOI [10.1097/MBC.0000000000000507](https://doi.org/10.1097/MBC.0000000000000507).
- Dusse LM, Rios DR, Pinheiro MB, Cooper AJ, Lwaleed BA.** 2011. Pre-eclampsia: relationship between coagulation, fibrinolysis and inflammation. *Clinica Chimica Acta* **412**(1–2):17–21 DOI [10.1016/j.cca.2010.09.030](https://doi.org/10.1016/j.cca.2010.09.030).
- Fang Y, Wang P, Xia L, Bai S, Shen Y, Li Q, Wang Y, Zhu J, Du J, Shen B.** 2019. Aberrantly hydroxymethylated differentially expressed genes and the associated protein pathways in osteoarthritis. *PeerJ* **7**:e6425 DOI [10.7717/peerj.6425](https://doi.org/10.7717/peerj.6425).

- Ghulmiyyah L, Sibai B. 2012.** Maternal mortality from preeclampsia/eclampsia. *Seminars in Perinatology* **36**(1):56–59 DOI [10.1053/j.semperi.2011.09.011](https://doi.org/10.1053/j.semperi.2011.09.011).
- Hazell GGJ, Peachey AMG, Teasdale JE, Sala-Newby GB, Angelini GD, Newby AC, White SJ. 2016.** PI16 is a shear stress and inflammation-regulated inhibitor of MMP2. *Scientific Reports* **6**:39553 DOI [10.1038/srep39663](https://doi.org/10.1038/srep39663).
- Huda SS, Sattar N, Freeman DJ. 2009.** Lipoprotein metabolism and vascular complications in pregnancy. *Clinical Lipidology* **4**(1):91–102 DOI [10.2217/17584299.4.1.91](https://doi.org/10.2217/17584299.4.1.91).
- Jian H, Zhao Y, Liu B, Lu S. 2015.** SEMA4B inhibits growth of non-small cell lung cancer in vitro and in vivo. *Cellular Signalling* **27**(6):1208–1213 DOI [10.1016/j.cellsig.2015.02.027](https://doi.org/10.1016/j.cellsig.2015.02.027).
- Jin WY, Lin SL, Hou RL, Chen XY, Han T, Jin Y, Tang L, Zhu ZW, Zhao ZY. 2016.** Associations between maternal lipid profile and pregnancy complications and perinatal outcomes: a population-based study from China. *BMC Pregnancy and Childbirth* **16**:60 DOI [10.1186/s12884-016-0852-9](https://doi.org/10.1186/s12884-016-0852-9).
- Kanehisa M, Goto S. 2000.** KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Research* **28**(1):27–30 DOI [10.1093/nar/28.1.27](https://doi.org/10.1093/nar/28.1.27).
- Khalil A, Poelvoorde P, Fayyad-Kazan M, Rousseau A, Nuyens V, Uzureau S, Biston P, El-Makhour Y, Badran B, Antwerpen P, Van, Boudjeltia KZ, Vanhamme L. 2018.** Apolipoprotein L3 interferes with endothelial tube formation via regulation of ERK1/2, FAK and Akt signaling pathway. *Atherosclerosis* **279**:73–87 DOI [10.1016/j.atherosclerosis.2018.10.023](https://doi.org/10.1016/j.atherosclerosis.2018.10.023).
- Lin N, Gu W, Hua RY, Hou YY. 2019.** Correlation between blood lipid levels in early pregnancy and preeclampsia. *Journal of Shanghai University (Natural Science)* **25**(1):18–23.
- Marano F, Zunino V, Frairia R, Arvat E, Castellano I, Bosco O, Catalano MG, Fortunati N. 2018.** Fibulin-1 interacts with sex hormone binding globulin and is linked to less aggressive estrogen-dependent breast cancers. *Life Sciences* **207**:372–380 DOI [10.1016/j.lfs.2018.06.024](https://doi.org/10.1016/j.lfs.2018.06.024).
- Moulder R, Bhosale SD, Goodlett DR, Lahesmaa R. 2018.** Analysis of the plasma proteome using iTRAQ and TMT-based Isobaric labeling. *Mass Spectrometry Reviews* **37**(5):583–606 DOI [10.1002/mas.21550](https://doi.org/10.1002/mas.21550).
- Mueller PA, Zhu L, Tavori H, Huynh K, Giunzioni I, Stafford JM, Linton MF, Fazio S. 2018.** Deletion of Macrophage Low-Density Lipoprotein Receptor-Related Protein 1 (LRP1) Accelerates Atherosclerosis Regression and Increases CCR7 Expression in Plaque Macrophages. *Circulation* **138**(17):1850–1863 DOI [10.1161/CIRCULATIONAHA.117.031702](https://doi.org/10.1161/CIRCULATIONAHA.117.031702).
- Norwitz E. 2007.** Defective implantation and placentation laying the blueprint for pregnancy complications. *Reproductive BioMedicine Online* **13**(4):591–599.
- Pawlak JB, Balint L, Lim L, Ma W, Davis RB, Benyo Z, Soares MJ, Oliver G, Kahn ML, Jakus Z, Caron KM. 2019.** Lymphatic mimicry in maternal endothelial cells promotes placental spiral artery remodeling. *Journal of Clinical Investigation* **129**(11):4912–4921 DOI [10.1172/JCI120446](https://doi.org/10.1172/JCI120446).

- Pehlivan M, Ozun Ozbay P, Temur M, Yilmaz O, Verit FF, Aksoy N, Korkmazer E, Ustunyurt E. 2017. Is there any role of prolidase enzyme activity in the etiology of preeclampsia? *Journal of Maternal-Fetal & Neonatal Medicine* **30**:1108–1113 DOI [10.1080/14767058.2016.1205023](https://doi.org/10.1080/14767058.2016.1205023).
- Pohanka M. 2013. Role of oxidative stress in infectious diseases. A review. *Folia Microbiologica* **58**(6):503–513 DOI [10.1007/s12223-013-0239-5](https://doi.org/10.1007/s12223-013-0239-5).
- Pollheimer J, Vondra S, Baltayeva J, Beristain AG, Knofler M. 2018. Regulation of placental extravillous trophoblasts by the maternal uterine environment. *Frontiers in Immunology* **9**:2597 DOI [10.3389/fimmu.2018.02597](https://doi.org/10.3389/fimmu.2018.02597).
- Pottecher J, Huet O, Degos V, Bonnet MP, Gaussem P, Duranteau J, Ozier Y, Mignon A, Tsatsaris V. 2009. In vitro plasma-induced endothelial oxidative stress and circulating markers of endothelial dysfunction in preeclampsia: an observational study. *Hypertens Pregnancy* **28**(2):212–223 DOI [10.1080/10641950802601195](https://doi.org/10.1080/10641950802601195).
- Rolnik DL, Wright D, Poon LC, O’Gorman N, Syngelaki A, De Paco Matallana C, Akolekar R, Cicero S, Janga D, Singh M, Molina FS, Persico N, Jani JC, Plasencia W, Papaioannou G, Tenenbaum-Gavish K, Meiri H, Gizurason S, Maclagan K, Nicolaides KH. 2017. Aspirin versus placebo in pregnancies at high risk for preterm preeclampsia. *New England Journal of Medicine* **377**(7):613–622 DOI [10.1056/NEJMoa1704559](https://doi.org/10.1056/NEJMoa1704559).
- Rong W, Yang L, Yin L, Gao Y, Xiao T, Cheng S. 2017. PSG9 promotes angiogenesis by stimulating VEGFA production and is associated with poor prognosis in hepatocellular carcinoma. *Science China Life Sciences* **60**:528–535 DOI [10.1007/s11427-016-0226-7](https://doi.org/10.1007/s11427-016-0226-7).
- Song G, Bailey DW, Dunlap KA, Burghardt RC, Spencer TE, Bazer FW, Johnson GA. 2010. Cathepsin B, cathepsin L, and cystatin C in the porcine uterus and placenta: potential roles in endometrial/placental remodeling and in fluid-phase transport of proteins secreted by uterine epithelia across placental areolae. *Biology of Reproduction* **82**(5):854–864 DOI [10.1095/biolreprod.109.080929](https://doi.org/10.1095/biolreprod.109.080929).
- Song Y, Luo Q, Long H, Hu Z, Que T, Zhang X, Li Z, Wang G, Yi L, Liu Z, Fang W, Qi S. 2014. Alpha-enolase as a potential cancer prognostic marker promotes cell growth, migration, and invasion in glioma. *Molecular Cancer* **13**:65 DOI [10.1186/1476-4598-13-65](https://doi.org/10.1186/1476-4598-13-65).
- Soonthornvacharin S, Rodriguez-Frandsen A, Zhou Y, Galvez F, Huffmaster NJ, Tripathi S, Balasubramaniam VR, Inoue A, de Castro E, Moulton H, Stein DA, Sanchez-Aparicio MT, De Jesus PD, Nguyen Q, Konig R, Krogan NJ, Garcia-Sastre A, Yoh SM, Chanda SK. 2017. Systems-based analysis of RIG-I-dependent signalling identifies KHSRP as an inhibitor of RIG-I receptor activation. *Nature Microbiology* **2**:17022 DOI [10.1038/nmicrobiol.2017.22](https://doi.org/10.1038/nmicrobiol.2017.22).
- Staff AC. 2019. The two-stage placental model of preeclampsia: an update. *Journal of Reproductive Immunology* **134–135**:1–10.
- Venkatesha S, Toporsian M, Lam C, Hanai J, Mammoto T, Kim Y, Bdolah Y, Lim K, Yuan H, Libermann T, Stillman I, Roberts D, D’Amore P, Epstein FH, Sellke F, Romero R, Sukhatme V, Letarte M, Karumanchi S. 2006. Soluble endoglin

- contributes to the pathogenesis of preeclampsia. *Nature Medicine* **12**(6):642–649 DOI [10.1038/nm1429](https://doi.org/10.1038/nm1429).
- Voshtani R, Song M, Wang H, Li X, Zhang W, Tavallaie MS, Yan W, Sun J, Wei F, Ma X. 2019.** Progranulin promotes melanoma progression by inhibiting natural killer cell recruitment to the tumor microenvironment. *Cancer Letters* **465**:24–35 DOI [10.1016/j.canlet.2019.08.018](https://doi.org/10.1016/j.canlet.2019.08.018).
- Wagner PK, Otomo A, Christians JK. 2011.** Regulation of pregnancy-associated plasma protein A2 (PAPPA2) in a human placental trophoblast cell line (BeWo). *Reproductive Biology and Endocrinology* **9**(1):48 DOI [10.1186/1477-7827-9-48](https://doi.org/10.1186/1477-7827-9-48).
- Wang C, Zhu W, Wei Y, Su R, Feng H, Hadar E, Hod M, Yang H. 2017.** The associations between early pregnancy lipid profiles and pregnancy outcomes. *Journal of Perinatology* **37**(2):127–133 DOI [10.1038/jp.2016.191](https://doi.org/10.1038/jp.2016.191).
- Wang Y, Sheng N, Xie Y, Chen S, Lu J, Zhang Z, Shan Q, Wu D, Zheng G, Li M, Zheng Y, Fan S. 2019.** Low expression of CRISP3 predicts a favorable prognosis in patients with mammary carcinoma. *Journal of Cellular Physiology* **234**(8):13629–13638 DOI [10.1002/jcp.28043](https://doi.org/10.1002/jcp.28043).
- Westerterp M, Berbee JF, Pires NM, Van Mierlo GJ, Kleemann R, Romijn JA, Havekes LM, Rensen PC. 2007.** Apolipoprotein C-I is crucially involved in lipopolysaccharide-induced atherosclerosis development in apolipoprotein E-knockout mice. *Circulation* **116**(19):2173–2181 DOI [10.1161/CIRCULATIONAHA.107.693382](https://doi.org/10.1161/CIRCULATIONAHA.107.693382).
- Wu D, Hong H, Huang X, Huang L, He Z, Fang Q, Luo Y. 2016.** CXCR2 is decreased in preeclamptic placentas and promotes human trophoblast invasion through the Akt signaling pathway. *Placenta* **43**:17–25 DOI [10.1016/j.placenta.2016.04.016](https://doi.org/10.1016/j.placenta.2016.04.016).
- Young BC, Levine RJ, Karumanchi SA. 2010.** Pathogenesis of preeclampsia. *Annual Review of Pathology* **5**:173–192 DOI [10.1146/annurev-pathol-121808-102149](https://doi.org/10.1146/annurev-pathol-121808-102149).
- Zhang B, Bie Q, Wu P, Zhang J, You B, Shi H, Qian H, Xu W. 2018.** PGD2/PTGDR2 signaling restricts the self-renewal and tumorigenesis of gastric cancer. *Stem Cells* **36**(7):990–1003 DOI [10.1002/stem.2821](https://doi.org/10.1002/stem.2821).
- Zhang XL, Xu FX, Han XY. 2019.** siRNA-mediated NCAM1 gene silencing suppresses oxidative stress in pre-eclampsia by inhibiting the p38MAPK signaling pathway. *Journal of Cellular Biochemistry* **120**(11):18608–18617 DOI [10.1002/jcb.28778](https://doi.org/10.1002/jcb.28778).