



REVIEW

Open Access

The CNS microvascular pericyte: pericyte-astrocyte crosstalk in the regulation of tissue survival

Drew Bonkowski^{1,2}, Vladimir Katyshev^{1,3}, Roumen D Balabanov⁴, Andre Borisov⁵, Paula Dore-Duffy^{1*}

Abstract

The French scientist Charles Benjamin Rouget identified the pericyte nearly 140 years ago. Since that time the role of the pericyte in vascular function has been difficult to elucidate. It was not until the development of techniques to isolate and culture pericytes that scientists have begun to understand the true impact of this unique cell in the maintenance of tissue homeostasis. In the brain the pericyte is an integral cellular component of the blood-brain barrier and, together with other cells of the neurovascular unit (endothelial cells, astrocytes and neurons) the pericyte makes fine-tuned regulatory adjustments and adaptations to promote tissue survival. These regulatory changes involve trans-cellular communication networks between cells. In this review we consider evidence for cell-to-cell crosstalk between pericytes and astrocytes during development and in adult brain.

Review

Introduction

The blood-brain barrier (BBB) regulates the passage of nutrients, essential components, proteins, chemical substances and microscopic organisms between the bloodstream and the parenchymal tissue. The anatomical constituents of the BBB are the endothelial cells (EC), pericytes, and basal lamina (matrix proteins) that with astrocytes, neurons, and possibly other glial cells, comprise the neurovascular unit [1]. Together, the cells of the neurovascular unit adapt to environmental changes and make fine-tuned regulatory decisions that maintain homeostasis and promote tissue survival [1-4]. Nowhere is such tight regulation more important than in the brain where bioenergetic and metabolic homeostasis is integral for neuronal survival [1,2]. Dysregulation at the neurovascular level is linked to many common human CNS pathologies, making the unit a potential target for therapeutic intervention [1-4].

The role of the CNS pericyte in the neurovascular unit is still unclear although with improved culturing techniques and the use of genetically engineered animals it has become increasingly clear that pericytes are integral to BBB function [3,4]. It is known that pericytes are highly complex regulatory cells that communicate with

ECs and other cells of the neurovascular unit such as neurons by direct physical contact and through autocrine and paracrine signaling pathways [3-9]. However, while there is abundant indirect evidence there is little direct evidence of pericyte-astrocyte communication. In this review we will summarize what is known about pericyte-astrocyte transcellular communication. We will discuss vascular development, BBB function and regulation of the neurovascular unit. The potential role of pericyte-astrocyte crosstalk in development of disease pathology will also be covered.

1. Pericytes

Pericytes were described over 100 years ago in 1873 by the French scientist Charles-Marie Benjamin Rouget and dubbed the Rouget cell [10]. The Rouget cell was renamed in the early 1900s to reflect its anatomical location with in the microvasculature abluminal to the EC and luminal to parenchymal cells (reviewed in [11]). In the brain, pericytes are located in pre-capillary arterioles, capillaries and post capillary venules. They synthesize and deposit elements of the basal lamina and are totally surrounded by this vascular compartment [3-7]. Pericytes are local regulatory cells that are important for the maintenance of vascular homeostasis and hemostasis, and are a source of adult pluripotent stem cells [12,13]. Pericytes are important cellular constituents of the BBB and actively communicate with other cells of the neurovascular unit such as EC, astrocytes, and neurons. While the communication networks involving

* Correspondence: pd Duffy@med.wayne.edu

¹Department of Neurology, Wayne State University School of Medicine, 421 East Canfield Road, Detroit, Michigan 48201, USA

Full list of author information is available at the end of the article

pericytes and endothelial cells have been considered in a number of publications, less is known of the cell-to-cell communication involving astrocytes.

Since its original discovery there has been much confusion and controversy about the pericyte as reflected by the numerous conflicting publications and definitions found in the literature. The pericyte has been referred to as: a) A contractile cell that surrounds the capillary in a tunic-like fashion [10]. b) A branching contractile cell on the abluminal wall of a capillary. c) A peculiar elongated, contractile cell wrapped around pre-capillary arterioles ‘outside’ the basement membrane [14]. d) A relatively undifferentiated connective tissue cell in the capillaries or other small blood vessels also called the adventitial cell [15]. e) A smooth muscle/pericyte or smooth muscle cell of the capillaries [16,17]. f) A broad flat cell with slender projections that wraps around the capillaries [17]. g) A mesenchymal stem-like cell, associated with the walls of small blood vessels. As a relatively undifferentiated cell, it serves to support these vessels, but it can differentiate into a fibroblast, smooth muscle cell, or macrophage as well other cell populations if required [3,5-7,12,13,18,19]. h) A pluripotent or pericyte progenitor cell [13].

In the mature CNS capillary, the pericyte is located between the EC and parenchymal astrocytes and neurons and is surrounded by the basal lamina [3-7]. Pericytes have a prominent round nucleus that clearly differs in shape from the elongated cigar shaped nucleus of the EC. The pericyte extends long processes that extend over the vessel wall. The morphological pattern of projections appears to be somewhat heterogeneous [13]. Pericyte projections can extend around the capillary as originally described by Rouget [10]. The classic wrapping pattern is also somewhat heterogeneous. The most common association of the pericyte with the capillary is one in which the pericyte processes are large and broad and span a continuous surface of the vessel. Alternatively these processes may form finger-like projections that are more confined and ensheathe a more finite portion of the vessel surface. A third pattern of pericyte orientation in the microvessel involves a retraction of projections with protrusion of the cells away from the capillary surface during angiogenesis, in response to stress stimuli [20-24] and possibly under other conditions [13]. In normal capillaries the wrapping pattern predominates but under pathological conditions the migrating patterns increase [24] and are associated with upregulation of cell surface proteases [20,21,25].

The CNS pericyte is surrounded by the basal lamina on all sides. During development and during angiogenesis the pericyte, assisted by nearby astrocytes discussed below, deposits basal lamina components [3,4,7,13,25-32]. Even extended pericyte projections, observed using electron microscopy, have a thin layer of basal lamina. The basal lamina has been shown to become thicker or thinner in response to stress stimuli [24,33-36]. Changes in the basal lamina can be directly associated with pericyte expression of proteases [20,24,25,37-41] and ultimate migration from its vascular location [20,33,34,42].

The intact basal lamina may provide anchoring and structural integrity to the capillary but it may also be involved in regulation of pericyte function and differentiation. It seems intuitive that there must be a reason why the pericyte is surrounded by laminin proteins. Av β 8 integrin is important in neurovascular cell adhesion [27,43]. Pericytes encased in the basal lamina or exposed to laminin proteins do not usually differentiate (Dore-Duffy, unpublished observations). Thus migration through the basal lamina is necessary before pericytes can function in their stem cell capacity [13]. Regulation at the level of the basal lamina may also be integral to vascular adaptability to an ever-changing environment and to pericyte signaling mechanisms [24].

In its pericapillary location, the pericyte may signal nearby ECs [3,4,13,40,44,45], astrocytes [46,47], neurons, smooth muscle cells and perhaps other pericytes [13]. Pericyte-EC contacts include peg and socket arrangements [8,48] and gap junctions [49-52]. Gap junctions allow pericytes to communicate with ECs through the exchange of ions and small molecules. Peg-and-socket contacts enable pericytes to penetrate through the basal lamina and make contact with other cells and nearby vessels [8,48,53]. Junction complexes including adhesion plaques also support transmission of contractile forces from pericytes to other cells. Pericyte gap junctions contain N-cadherin, a variety of adhesion molecules, β -catenin, extracellular matrix (ECM) molecules such as fibronectin, and a number of integrins [50,51]. Thus, pericytes are involved in highly complex signaling cascades that enable this cell to respond to changes in the microenvironment. However, it is unclear whether gap junctions and peg and socket contacts are naturally present or whether they are initiated during changes in functional activity. For example, it is known that pericytes interdigitate with ECs during the early phases of angiogenesis and with neurons during the maturation of newly-formed vessels [54]. These sites of communication are altered under pathological conditions. During cerebral edema or diabetes, gap junctions are substantially decreased or disrupted in retinal pericytes [54-57]. Diabetes-induced changes in gap junctions may be regulated by high

glucose [56–58]. Pericyte-EC communication via gap junctions is fundamental to the adaptive responses to compromised bioenergetic homeostasis [58]. Crosstalk between ECs, pericytes, as well as astrocytes is involved in regulation of insulin transport [47]. Pericyte/EC crosstalk is also integral to physiological angiogenesis [59], and is likely to be important in adaptation to hypoxic injury and focal capillary contractility.

2. Pericytes and astrocytes during vasculogenesis and angiogenesis

Vasculogenesis is the formation of new blood-vessels by differentiation of vascular precursor cells during development. During retinal development, a role for astrocyte-pericyte communication has been established. Retinal vascularization begins in the inner retinal layer and sprouts radially from the optic nerve to reach the periphery of the retina [60]. Subsequently, retinal vessels sprout into the deep retinal layer to vascularize three parallel nerve fiber layers and two plexiform layers. Experiments using retinal models have tested the hypothesis that astrocytes and pericytes influence this process by affecting the composition of the extracellular matrix. Fibronectin, but not laminin, is expressed in zones of vasculogenesis immediately prior to vessel formation. At this time astrocytes and pericytes spread into the tissue and may be involved in the initiation of vasculogenesis. In fact, it is unclear whether platelet-derived growth factor receptors PDGF β R+/PDGF α R+ pericytes and/or PDGF α R+ immature astrocytes are the regulating cell type [60]. Increased amounts of fibronectin mRNA suggest that fibronectin is synthesized by cells within this same region. Pericytes are known to synthesize most extracellular matrix protein components [61]. Differentiation of endothelial cells is correlated with the appearance of pericytes in the vessel wall and laminin in the vascular basement membrane. Astrocyte-conditioned medium stimulates fibronectin expression by both primary endothelial cells and pericytes [62]. Astrocyte-induced pericyte synthesis of fibronectin and perhaps laminin is an essential step in the initiation of retinal vasculogenesis [63]. It is, however, unclear whether a similar mechanism functions in the brain.


Vascularization in the brain during development is derived from a preformed perineuronal vascular plexus and occurs almost exclusively through sprouting angiogenesis that starts at embryonic day 9 in vertebrates [63]. Angiogenesis is the formation of new vessels from existing vessels. Sprouting angiogenesis is an invasive process that involves proteolytic activities required for degradation of the basal lamina, with pericyte and endothelial migration through tissue matrix [63]. During vessel formation, the recruitment of pericytes and astrocytes to newly-forming vascular tubes is closely associated with the formation of tight junctions. At P4,

invading vessels are in direct contact with pericytes, but not with astrocytes. With the progression of development, foot processes of astrocytes are gathered around retinal vessels and the maturation of tight junction ZO-1 in endothelial cells is more clearly defined [64]. Tight junctions can be formed with contact from pericytes without the ensheathment of the astrocytic foot processes. From *in vitro* studies it is clear that pericytes induce synthesis of both occludin and claudin through the release of angiopoietin-1 [65]. Thus tight junction formation in the BBB could be developed, in part, by cellular interactions between EC and pericytes followed by astrocytes (as reviewed in [66]). The expression of ZO-1 is augmented by astrocytes [67]. Moreover, ZO-1 as well as occludin or claudin together may be biological indicators of barrier maturation [67]. The role of the pericyte in BBB function will be discussed below.

Studies using glial fibrillary acidic protein (GFAP)-knockout mice (GFAP $^{-/-}$) indicate that, although astrocyte foot interactions with the microvasculature are missing, neovascularization and the formation of tight junctions as well as other CNS morphology appears to be natural [68–70]. Astrocyte neuronal interactions are disrupted [70]. These data suggest that astrocyte interaction with the capillaries or neurons is not essential for tight junction formation. Of interest is that GFAP $^{-/-}$ mice are prone to hemorrhagic injury [71]. Similar observations were observed by McCarthy and colleagues [44] in studies using animals lacking alphav integrins. In contrast, administration of the gliotoxin 6-aminonicotamide (6-AN) to chick embryos during both early and late embryonic development showed a good correlation between perivascular glial depletion and BBB impairment suggesting that astroglia do in fact play a role in BBB prenatal differentiation and must in some way contribute to vascular function [72]. Taken together the results suggest that, while astrocyte formation of end-feet is not essential for tight junction formation, astrocytes are an important component contributing to normal vessel structural integrity. These results may also indicate that pericytes or other CNS cellular components can compensate for the lack of direct astrocyte signaling.

Using *in vitro* model systems to study vasculogenesis, scientists have learned that both pericytes and astrocytes are involved in formation of capillary structures. Both cell types form endothelial connections with newly forming vessels (Figure 1) realigning themselves according to the same arrangement seen *in vivo*. Pericytes in co-culture associate with EC more rapidly than astrocytes. In multi-cellular systems involving pericytes, astrocytes, and EC newly forming tubes have the three dimensional structure of a vessel and exhibit both a lumen and tight junctions. Tube formation in triple

A



B




Figure 1 Induction of angiogenesis in triple co-cultures of ECs, pericytes and astrocytes. Primary cells were co-cultured at a ratio of one pericyte to five ECs and to five astrocytes. Cultures were exposed to hypoxia (10%) to induce angiogenesis (A). Tube formation in triple cultures involved crosstalk between astrocytes and pericytes. White arrows point to pericytes and black arrows point to astrocytes. Photograph was taken using phase contrast at 10X. In (B) we show a close up of an astrocyte making contact with both elongating EC and pericytes. The black arrows point to astrocyte contacts with an elongating EC (left) and one of the two contacts made with a pericyte (right). Pericytes have fewer projections and are more spread out than the astrocytes.

cultures is more rapid than that observed in EC/pericyte or EC/astrocyte, co-cultures. The triple co-culture of pericytes and astrocytes with ECs is thought by most investigators to represent a better model system to study the BBB [73-76]. This may be in part due to the role of agrin, aquaporin 4, and astrocyte polarity in the BBB [77].

The process of vasculogenesis and differentiation is a complex system that involves crosstalk between numerous cells and is regulated by a number of signaling pathways.

Cell surface proteases expressed on both EC and pericytes exert additional subtle functions in sprouting angiogenesis. These functions involve membrane type-1 matrix metalloproteinase (MT1-MMP), other MMPs, and ADAMs (a disintegrin and metalloproteinase domain) [78]. Proteases modulate the balance between pro- and anti-angiogenic factors by activation of growth factors and chemokines, shedding of chemokines and cytokines from membrane-bound precursors [78], and generation of (matrix) protein fragments that inhibit or activate angiogenesis. Furthermore, they participate in the recruitment of leukocytes and progenitor cells, which contribute to the initiation and progression of angiogenesis. Pericytes are involved in the initiation as well as termination of angiogenesis (as reviewed in [59]). At the initiation of angiogenesis, pericytes are involved in induction of endothelial activation accompanied by augmentation of a variety of proteases [40], adhesion

molecules and proteoglycans [79]. Astrocyte expression of tumor necrosis factor converting enzyme (TACE/ADAM-17) may facilitate pericyte PDGF β R signaling mechanisms [80]. Alternatively, at maturation the recruitment of pericytes to the newly-formed endothelial tubes is accompanied by silencing of MMP activities [81]. Recruitment to the newly-formed vessel with termination of angiogenesis involves PDGF β [82] as well as transforming growth factor beta (TGF β) [83]. PDGF β is thought to be essential for the retention of pericytes in the newly-formed vessels [84,85]. Ablation of PDGF β results in an embryonic lethal mutation associated with multiple vascular abnormalities [82,86,87].

3. Pericytes, astrocytes and BBB function

3.1 Contractility

The concept that pericytes regulate blood flow at the capillary level was originally proposed by Steinach and Kahn in 1903 [88] and Ni in 1922 [89]. Both scientific groups studied the effects of electrical stimulation or exposure to toxic stimuli on capillary diameter. Doré reviewed this area in 1923 [11]. As stated by Doré, [11]: "Until a few years ago the capillaries were regarded as elastic tubes undergoing passive distension in accordance with the general blood pressure, the state of contraction or dilatation of the supplying arterioles, and the nutrition of the vascular walls". The concepts put forward in this 1923 review are on target with exciting more-recently published data [90-92].

CNS pericytes have receptors for a large number of vasoactive signaling molecules [3,93-95] suggesting that they have the capacity to be involved in cerebrovascular autoregulation. Expression of alpha muscle actin (α SMA) and the intermediate filament desmin, two proteins found in smooth muscle cells, as well as their adherence to the endovascular cells make them potential candidates for the regulation of capillary diameter and focal capillary blood flow [3,8,89,91,96-99]. Electrical stimulation of retinal and cerebellar pericytes is reported to evoke a localized capillary constriction [91,100]. ATP in the retina or noradrenaline in the cerebellum also results in constriction of capillaries by pericytes. Glutamate reverses the constriction produced by noradrenaline [100].

Following simulated ischemia and traumatic brain injury (TBI), capillary pericytes are induced to express α SMA and upregulation of pericyte α SMA is correlated with a focal decrease in capillary diameter [100]. Upregulation of muscle actin was mediated by endothelin-1. Other investigators have shown that capillary contraction can be directly linked to metabolic need [101,102]. Exposure to substances that increase pericyte calcium level induces vessel contraction with a concomitant decrease in the capillary lumen diameter [101-103]. The contractile response appears to involve a cascade of events resulting in the inhibition of $\text{Na}^+/\text{Ca}^{2+}$ exchangers on the EC [103]. Hypoxia, which closes gap junctions, switches the effect of lactate from contraction to relaxation. Thus, pericyte function may be linked with local vascular adaptation to changes in bioenergetic requirements and is intimately linked to astrocyte function.

The calcium-dependent transcription factor NFATc3 is a member of the nuclear factor of activated T cells (NFAT)-family of transcription factors. NFATc3 is critical for embryonic vascular development and differentiation. Filosa and colleagues investigated the role of glutamate in control of NFATc3 regulation in pericytes [104]. Coronal cortical slices from neonatal rats were subjected to electrical field stimulation or were treated with glutamate receptor agonist (+/-)-1-aminocyclopentane-trans-1,3-dicarboxylic acid (t-ACPD). Electrical field stimulation induced NFATc3 nuclear accumulation in pericytes and astrocytes. The response in pericytes was dependent on metabotropic glutamate receptor (mGluR) activation. NFATc3 nuclear accumulation in pericytes was prevented when astrocytic function was abolished with the gliotoxin L-alpha-amino adipate. Results suggest that astrocyte glutamate, via mGluR activation, may regulate gene transcription in pluripotent vascular pericytes [104].

3.2 BBB Permeability

It has been known for decades that the CNS tissue microenvironment provides the cues for BBB induction

and differentiation (as reviewed in [3,7,90,105]). Development of the BBB (barrier genesis) is the result of coordinated molecular signaling at the neurovascular interface [106]. Only recently has the canonical Wnt/beta-catenin pathway and the Wnt7a/7b growth factors been implicated in CNS angiogenesis and in BBB induction [86,105-108]. This pathway interacts with other pathways that are crucial for vascular development such as VEGF. Wnt/beta-catenin pathways enhance pericyte mesenchymal differentiation in the presence of TGF-beta3, as demonstrated by increased Sox-9 expression and glycosaminoglycan release into the extracellular matrix. In contrast, transduction of pericytes with a recombinant adenovirus encoding dominant-negative T-cell factor-4 blocked Wnt/beta-catenin signaling and inhibited pericyte differentiation to chondrocytes, leading to reduced Sox-9, reduced type II collagen expression and reduced glycosaminoglycan accumulation [105,107]. These data demonstrate that TGF-beta3 induces the chondrogenic differentiation of pericytes by inducing Wnt/beta-catenin signaling and T-cell factor-induced gene transcription. In response to injury, the Wnt/beta catenin signaling pathway is enhanced in proliferating NG2 chondroitin sulphate proteoglycan positive cells undergoing differentiation to NG2 glia [109]. It is unclear whether induction of Wnt/beta-catenin signaling in pericytes is responsible for the regulation of the BBB by augmenting the number of NG2⁺ astrocyte subsets.

Pericytes play a key role in vascular remodeling during development and in the adult. Considerable insight into pericyte biology during development has arisen from studies employing genetically-manipulated mice with disrupted PDGF β /platelet PDGF β R signaling [84-87,110]. During development, EC-secreted PDGF β binds to pericyte PDGF β R receptors located on the pericyte plasma membrane resulting in dimerization of the receptor [111]. Pericytes themselves produce PDGF β and thus may signal in an autocrine fashion [13]. PDGF stimulates the proliferation, migration, and recruitment of pericytes to the vascular wall of newly-formed blood vessels [10,14,15,83-87,112]. PDGF β and PDGF β R knockout mice have diminished pericyte numbers and a lack of pericyte coverage [87]. The mutation is embryonic lethal with abnormal vasculogenesis [87,112]. Loss of pericytes also results in loss of BBB integrity during embryogenesis [113] clearly showing that pericytes are essential to the differentiation of the BBB during development and have an important role in regulating the BBB [114]. The exact mechanisms are still somewhat unclear as pericytes have been shown to both induce permeability and to inhibit permeability during development [113-115].

PDGF β and PDGF β R interactions are also involved in pericyte-mediated regulation of vascular function in the

adult brain and have a pivotal role in the regulation of the cerebral microcirculation. However, the functional roles of CNS pericytes in the adult and aging brain are less well understood. In the adult animal, it is generally accepted that pericyte association with the microvessel is essential to vascular integrity and that loss of pericytes leads to an increase in vascular leakage and altered vascular function. However, loss of pericytes does not always result in a permanent loss of function and does not always result in disease. Pericytes migrate naturally during the early phases of physiological angiogenesis to make way for growing sprouts [116-119], or in response to stress or injury [20]. Migration from the microvessels following TBI for example, is thought to promote survival as pericytes remaining in their vascular location show signs of degenerative activity [20]. Migration from the vessel involves upregulation of cell surface proteases [20]. Newly-forming vessels both in adult angiogenesis and neovascularization are characterized by increased leakiness. This is transient under normal circumstances but is abnormal in conditions such as pathological angiogenesis. During angiogenesis there is co-migration of pericytes with EC sprouts [116-120] and concomitant proliferation of migrated cells before renewed pericyte coverage and termination of angiogenesis [59]. Pericytes may also be recruited from the bone marrow in adult mice and may have a role in initiation as well as termination of angiogenesis [84,120]. Renewed pericyte coverage restores vascular integrity. These observations are supported by data derived using *in vitro* models of angiogenesis [121-123]. Lack of pericyte termination of angiogenesis alters vascular function and may promote vascular leakiness in part by structural problems with tight junctions [113,114]. Pericytes may also be essential to regulation of transport mechanisms [86,108]. Taken together it is easy to speculate that loss of pericyte function may result in the induction of edema and inflammation [3-7,113,114].

Is there a role for astrocytes in maintenance of BBB function? Certainly the astrocytes provide a structural barrier that helps promote vascular integrity. As discussed, microvessels from the GFAP knockout mouse that have no astrocyte end-feet have a tendency to form micro-hemorrhages despite the presence of normal tight junctions [68,69]. The interaction of astrocytes with the capillary appears to have an ordered structure as seen in Figure 2a in wild type capillaries. GFAP staining seems to have a fish net orientation that may assist in promoting vascular integrity. Heterozygous animals display a very disordered GFAP staining (Figure 2b). Capillaries from GFAP^{-/-} mice display no staining pattern (Figure 2c). We have found that, to compensate for the loss of astrocyte end-feet, pericytes proliferate. Pericyte to EC ratios are much higher than those observed for wild

type mice and for GFAP⁺⁻ heterozygotes (Figure 3; Table 1). This is in contrast to data by others [84] using PDGF deficient animals. The presence of astrogliosis may also alter pericyte function [87].

The integral association of PDGF/PDGFR signaling in maintenance of pericyte function, makes it easy to speculate that astrocytes may be able to regulate pericyte function by the modulation of PDGF β signaling. Controlled synthesis of astrocyte H₂O₂ alters PDGF signaling and is neuroprotective in conditions of oxidative stress [124-128]. The expression of truncated PDGF β R inhibits PDGF signaling [129]. Inhibition is also seen with upregulation of PDGF α R [130]. Heparan sulphate controls the diffusion, and thus the availability of PDGF β [131]. Perlecan, a heparin sulfate proteoglycan, has been reported to modulate the BBB [132] and is produced by astrocytes during the stress response [133]. Astrocyte modulation of pericyte function is seen in the injured retina [60,64].

4. Pericyte-astrocyte communication and disease

Under normal conditions, the pericyte is relatively quiescent and is essential for vascular stability. Under conditions of stress or injury, the pericyte undergoes phenotypic and functional changes that may include migration, proliferation or differentiation. How these events that include pericyte reprogramming are coordinated at the molecular level needs to be determined. However, it is clear that pericyte dysfunction or the loss of pericytes is likely to play an important role in the pathogenesis of disease.

Pericyte loss or a reduced pericyte-to-EC ratio may be achieved through: 1) migration of pericytes from their microvascular location under pathological or physiological conditions, 2) pericyte death, 3) reduced pericyte turnover or maintenance, and 4) selective alteration of pericyte recruitment to EC that may be associated with dysregulation of angiogenesis and abnormal PDGF β signaling. Loss of pericytes with reduced pericyte-to-EC ratios results in a focal increase in permeability. This can be normal as seen in the transient decrease in pericyte-to-EC ratios due to migration of pericytes during endogenous adaptation to chronic mild hypoxia [59,134] or may be abnormal as observed in a number of pathophysiological diseases. Migration may also play a pathogenic role such as that observed in diabetic retinopathy [135]. Decreased pericyte-to-EC ratios have been observed following TBI (20) and stroke [136], multiple sclerosis [137-140], brain tumors [141-143], diabetic retinopathy [135,144], aging [145,146], and in a variety of angiopathies [147]. Pericyte loss may also play a role in Alzheimer's disease, however; enhanced pericyte coverage of some vessels suggests that increased proliferation of pericytes is an adaptation to focal loss of




Figure 2 Capillaries [2-6 micron diameter determined by Image J software] were isolated from wild type C57BL/6 mice (A), GFAP $^{+/-}$ mice (B), and GFAP $^{-/-}$ mice (C) (Jackson Laboratory). Freshly isolated capillaries were allowed to adhere to coverslips fixed and stained for the expression of GFAP (Santa Cruz Biotechnology Inc). Capillaries were visualized on a Leitz fluorescent microscope at 40x and 100x.

bioenergetic homeostasis [14,140,148,149]. Pericyte loss due to cellular degeneration/apoptosis has been shown in hypertrophic scars, keloids [150,151], early diabetic retinopathy [150,152], brain tumors [153-156], liver cancer [157] hyperglycemia [158], and during development [159]. Premature infants have decreased pericyte coverage [160]. A decreased pericyte-to-EC ratio also observed during vascular regression [160].

Increased pericyte coverage may also be an indicator of vascular dysfunction. Pericyte proliferation has been associated with development of muscularization during pulmonary hypertension and is thought to be due to platelet activating factor [161]. Cochlear pericytes in the *stria vascularis* are markedly affected by acoustic trauma [162]. Levels of the pericyte structural protein, desmin, substantially increase after noise exposure with a corresponding increase in pericyte coverage of vessels. Increased expression levels of desmin were associated with the induction of hypoxia-inducible factor (HIF)-1alpha and the upregulation of vascular endothelial growth factor (VEGF). Inhibition of HIF-1alpha activity decreased VEGF expression levels. Blockade of VEGF activity with SU1498, a VEGF receptor inhibitor, significantly attenuated the expression of desmin in pericytes

and may have altered the adaptive process to acoustic stress. Increased pericyte-to-EC ratios are also observed in response to over expression of PDGF β [163]. The percentage of vessels covered by pericytes is nearly doubled in myelofibrotic bone marrow [164].

Direct evidence for astrocyte augmentation of pericyte coverage is lacking. However, indirect observations suggest that astrocytes may help augment pericyte proliferation through augmentation of PDGF β levels [165] or release of inflammatory mediators [166] that enhance gliosis. In patients with glaucoma, enhanced proliferation is modest but restricted to NG2+ pericytes [167]. Astrocytes and pericytes both express estrogen receptors [167-169]. Astrocytes have been shown to be a source of estrogen following neuronal injury due to an upregulation of aromatase [167]. Estrogens attenuate PDGF β signaling in vascular smooth muscle cells [170,171].

On a more subtle level pericyte stem cell activity may be a target for astrocyte modulation. Pericytes have been shown to be multipotent adult stem cells in a number of tissues [12,13]. Primary CNS pericytes are capable of differentiating along both the mesenchymal and neural lineages depending on the culture conditions [[12], as reviewed [13]]. In the absence of serum in the presence




Figure 3 Pericyte-endothelial cell ratios were determined by isolation of capillaries [2-4 micron diameter determined by Image J software] from wild type B6 mice (A); heterozygous GFAP $^{+/-}$ mice (B) and knockout GFAP $^{-/-}$ mice (C). Freshly isolated capillaries were stained with the fluorescent nuclear dye 4',6-diamidino-2-phenylindole [DAPI]. Round nuclei (pericytes) and elongated nuclei (EC) were counted. Numbers are shown in Table 1. Arrows point to round pericyte nuclei.

Table 1 Pericyte and endothelial cell numbers in freshly-isolated capillaries from wild type and GFAP knockout mice

mouse	#Pericytes*	# EC*	Fragments counted/exp
Wild type	3 ± 2	13 ± 2	40
GFAP -/-	9 ± 3	3 ± 1	55
GFAP +/-	7 ± 2	10 ± 3	42

* Mean +/- SD, N = 2 experiments.

of basic fibroblast growth factor (bFGF) cultured adult pericytes form immature NG2+ astrocytes/radial glial cells, NG2+ oligodendrocyte progenitor-like cells and neurons [12]. Thus in response to injury pericytes may undergo neurogenesis. Astrocyte-generated signaling molecules have been shown to augment neurogenesis and differentiation of oligodendrocyte progenitor cells (OPC) [172]. Radial glial cells also retain neurogenic potential [173,174]. During injury responses in the kindled animal, astrocyte activation and induction of neurogenesis is due in part to augmentation of nestin expression in pericytes [175,176]. Following TBI, Wnt/beta-catenin signaling occurs in quiescent NG2 progenitors in the cortex, in subcallosal zone (SCZ) progenitors, and in subependymal cells surrounding the central canal. Initially after TBI, beta-catenin signaling was predominantly increased in a subset of NG2+ progenitors in the cortex [110]. One week following injury, the majority of beta-catenin signaling appeared in reactive astrocytes but not other glial cells. Interestingly, cells with beta-catenin signaling were not generated following spinal cord injury. This suggests that although augmentation of pericyte generated stem cell activity and differentiation may be essential to tissue plasticity in adult animals responses may be tissue specific. Further, differentiations within microvascular locations, as well as other abnormal pericyte responses to injury, are likely to be highly deleterious.

5. Pericytes here and now

After its identification by Rouget in the late 1800s, relatively little was published about the pericyte until 1902 when the presence of this intriguing cell was confirmed [11]. The development of tissue culture techniques, as well as genetically-manipulated transgenic models in which pericyte coverage is absent or diminished has stimulated considerable interest and provided invaluable tools needed to study pericyte biology. However, while genetically-modified animals can be an important tool some caution should be used when attributing functional roles from observations derived in knockout animals. Adaptive responses to a genetic modification may also confound the interpretation. One must also consider that the phenotype is a function both of the planned genetic manipulation and of potential adaptive responses to the change.

There are a number of examples in the literature that uniquely illustrated this point [177]. Pericytes are integral to the regulation of vascular function as shown in PDGF β knockouts. However, PDGF α knockout is also embryonic lethal and displays a similar phenotype [178,179], and together with PDGF β is also integral to postnatal development [180]. In the future more targeted manipulations as well as fate mapping will likely clarify the functional role of pericytes in many tissues.

Conclusions

In this review we discussed evidence that pericytes and astrocytes undergo direct cell-to-cell communication. We also discuss indirect evidence that suggests that astrocytes and pericytes coordinate BBB function. It is clear that this communication is needed for the maintenance of vascular function and must be integral to endogenous adaptation to injury in the adult neurovascular unit. Loss of pericyte function and/or loss of proper astrocyte contact with the BBB can result in pericyte dysfunction and development of disease. A better understanding of the mechanisms by which pericytes communicate with other cells and how altered communication may result in disease pathology is likely to yield exciting new insights as well as the development of a new therapeutic target in CNS disorders.

Acknowledgements

The work discussed in this manuscript was supported by grants from the National Institutes of Health NINDS, (NS47672) and the National Multiple Sclerosis Society (CA1042A8, PP1517).

Author details

¹Department of Neurology, Wayne State University School of Medicine, 421 East Canfield Road, Detroit, Michigan 48201, USA. ²Department of Neuroscience, University of Michigan, Ann Arbor, Michigan, USA.

³Department of Neurology Wayne State University School of Medicine, Detroit, Michigan 48201, USA. ⁴Department of Neurology, Rush University Medical Center, Chicago, Illinois, 60612, USA. ⁵Department of Biology, Wayne State University, Detroit, Michigan 48201, USA.

Authors' contributions

PDD had primary responsibility for the design of experiments discussed within the review and for the organization and conception of the article. RB performed experiments detailed in the figures and tables as well as development of concepts discussed within. VK conducted experiments to provide some of the data within the manuscript as well as helped with its preparation. DB collected and secured data and interpreted material utilized for the review. AB discussed the organization of the review and possible concepts to be included. All authors have reviewed and approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 10 December 2010 Accepted: 18 January 2011

Published: 18 January 2011

References

1. del Zoppo GJ: The neurovascular unit in the setting of stroke. *J Intern Med* 2010, 267:156-171.

2. Nag S, Kapadia A, Stewart DJ: Molecular Pathogenesis of Blood-Brain Barrier Breakdown in Acute Brain Injury. *Neuropathol Appl Neurobiol* 2010.
3. Balabanov R, Dore-Duffy P: Role of the CNS microvascular pericyte in the blood-brain barrier. *J Neurosci Res* 1998, **53**:637-644.
4. Fisher M: Pericyte signaling in the neurovascular unit. *Stroke* 2009, **40**: S13-15.
5. Sims DE: Recent advances in pericyte biology-implications for health and disease. *Can J Cardiol* 1991, **7**:431-443.
6. Tilton RG: Capillary pericytes: perspectives and future trends. *J Electron Microsc Tech* 1991, **19**:327-344.
7. Shepro D, Morel NM: Pericyte physiology. *FASEB J* 1993, **7**:1031-1038.
8. Nakae R, Verma S, Niwa M, Doghu S, Banks WA: Glucose regulated blood-brain barrier transport of insulin: Pericyte-astrocyte-endothelial cell cross talk. *IJNN* 2007, **3**:195-200.
9. Charbord P, Oostendorp R, Pang W, Herault O, Noel F, Tsuji T, Dzierzak E, Peault B: Comparative study of stromal cell lines derived from embryonic, fetal, and postnatal mouse blood-forming tissues. *Exp Hematol* 2002, **30**:1202-1210.
10. Rouget C: Note sur le développement de la tunique contractile des vaisseaux. *Compt Rend Acad Sci* 1874, **59**:559-562.
11. Doré SE: On the contractility and nervous supply of the capillaries. *Brit J Derma* 1923, **35**:398-404.
12. Dore-Duffy P, Katchev A, Wang X, Van Buren E: CNS microvascular pericytes exhibit multipotential stem cell activity. *J Cereb Blood Flow Metab* 2006, **26**:613-624.
13. Dore-Duffy P: Pericytes: Pluripotent cells of the blood brain barrier. *Curr Pharm Des* 2008, **14**:1581-1593.
14. Blood DC, Studdert VP: *Baillière's Comprehensive Veterinary Dictionary* London: Baillière Tindall; 1988.
15. Stedman TL: *Stedman's Medical Dictionary* Baltimore: Williams & Wilkins; 1995.
16. Fabry Z, Fitzsimmons KM, Herlein JA, Moninger TO, Dobbs MB, Hart MN: Production of the cytokines interleukin 1 and 6 by murine brain microvessel endothelium and smooth muscle pericytes. *J Neuroimmunol* 1993, **47**:23-34.
17. Ding R, Darland DC, Parmacek MS, D'Amore PA: Endothelial-mesenchymal interactions in vitro reveal molecular mechanisms of smooth muscle/pericyte differentiation. *Stem Cells Dev* 2004, **13**:509-520.
18. Pericytes. 2009 [http://en.wikipedia.org/wiki/Pericytes].
19. Krüger M, Bechmann I: Pericytes. In *Central Nervous System Diseases and Inflammation*. Edited by: Lane TE, Bergmann C, Carson M, Wyss-Coray T. Berlin Heidelberg: Springer-Verlag; 2008:33-43.
20. Dore-Duffy P, Owen C, Balabanov R, Murphy S, Beaumont T, Rafols JA: Pericyte migration from the vascular wall in response to traumatic brain injury. *Microvasc Res* 2000, **60**:55-69.
21. Du R, Petritsch C, Lu K, Liu P, Haller A, Ganss R, Song H, Vandenberg S, Bergers G: Matrix metalloproteinase-2 regulates vascular patterning and growth affecting tumor cell survival and invasion in GBM. *Neuro Oncol* 2008, **10**:254-264.
22. Diaz-Flores L, Gutierrez R, Varela H: Angiogenesis: an update. *Histol Histopathol* 1994, **9**:807-843.
23. Nehls V, Herrmann R, Huhnken M: Guide migration as a novel mechanism of capillary network remodeling is regulated by basic fibroblast growth factor. *Histochem Cell Biol* 1998, **109**:319-329.
24. Nehls V, Schuchardt E, Drenckhahn D: The effect of fibroblasts, vascular smooth muscle cells, and pericytes on sprout formation of endothelial cells in a fibrin gel angiogenesis system. *Microvasc Res* 1994, **48**:349-363.
25. del Zoppo GJ: The neurovascular unit, matrix proteases, and innate inflammation. *Ann N Y Acad Sci* 2000, **1207**:46-49.
26. Wolburg H, Noell S, Mask A, Wolburg-Buchholz K, Fallier-Becker P: Brain endothelial cells and the glio-vascular complex. *Cell Tissue Res* 2009, **335**:75-96.
27. del Zoppo GJ, Milner R: Integrin-matrix interactions in the cerebral microvasculature. *Arterioscler Thromb Vasc Biol* 2006, **26**:1966-1975.
28. Stratman AN, Schwindt AE, Malotte KM, Davis GE: Endothelial-derived PDGF-BB and HB-EGF coordinately regulate pericyte recruitment during vasculogenic tube assembly and stabilization. *Blood* 2010, **116**:4720-4730.
29. Kalimo H, Lehto M, Nanto-Salonen K, Jalkanen M, Risteli L, Risteli J, Narva EV: Characterization of the perivascular reticulin network in a case of primary brain lymphoma. Immunohistochemical demonstration of collagen types I, III, IV, and V; laminin; and fibronectin. *Acta Neuropathol* 1985, **66**:299-305.
30. Canfield AE, Schor AM, Loskutoff DJ, Schor SL, Grant ME: Plasminogen activator inhibitor-type I is a major biosynthetic product of retinal microvascular endothelial cells and pericytes in culture. *Biochem J* 1989, **259**:529-535.
31. Mandarino LJ, Sundarrajan N, Finlayson J, Hassel HR: Regulation of fibronectin and laminin synthesis by retinal capillary endothelial cells and pericytes in vitro. *Exp Eye Res* 1993, **57**:609-621.
32. Jeon H, Ono M, Kumagai C, Miki K, Morita A, Kitagawa Y: Pericytes from microvessels fragment produce type IV collagen and multiple laminin isoforms. *Biosci Biotechnol Biochem* 1996, **60**:856-861.
33. Díaz-Flores L, Gutiérrez R, Madrid JF, Varela H, Valladares F, Acosta E, Martín-Vasallo P, Díaz-Flores L Jr: Pericytes. Morphofunction, interactions and pathology in a quiescent and activated mesenchymal cell niche. *Histol Histopathol* 2009, **24**:909-969.
34. Gonul E, Duz B, Kahraman S, Kayali H, Kubar A, Timurkaynak E: Early pericyte response to brain hypoxia in cats: an ultrastructural study. *Microvasc Res* 2002, **64**:116-119.
35. Arismendi-Morillo G, Castellano A: Tumoral micro-blood vessels and vascular microenvironment in human astrocytic tumors. A transmission electron microscopy study. *J Neurooncol* 2005, **73**:211-217.
36. Wiley LA, Rupp GR, Steinle JJ: Sympathetic innervation regulates basement membrane thickening and pericyte number in rat retina. *Invest Ophthalmol Vis Sci* 2005, **46**:744-748.
37. Nielson BS, Sehested M, Kjeldsen L, Borregaard N, Rygaard J, Dan K: Expression of matrix metalloprotease-9 in vascular pericytes in human breast cancer. *Lab Invest* 1997, **77**:345-355.
38. Arihiro S, Ohtani H, Hiwatashi N, Torii A, Sorsa T, Nagura H: Vascular smooth muscle cells and pericytes express MMP-1, MMP-9, TIMP-1 and type I procollagen in inflammatory bowel disease. *Histopathology* 2001, **39**:50-59.
39. Zozulya A, Weidenfeller C, Galla HJ: Pericyte-endothelial cell interaction increases MMP-9 secretion at the blood brain barrier in vitro. *Brain Res* 2008, **1189**:1-11.
40. Xing C, Arai K, Park KP, Lo EH: Induction of vascular endothelial growth factor and matrix metalloproteinase-9 via CD47 signaling in neurovascular cells. *Neurochem Res* 2010, **35**:1092-1097.
41. Thanabalasundaram G, Pieper C, Lischper M, Galla HJ: Regulation of the blood-brain barrier integrity by pericytes via matrix metalloproteinases mediated activation of vascular endothelial growth factor in vitro. *Brain Res* 2010, **1347**:1-10.
42. Hughes SJ, Wall N, Scholfield CN, McGeown JG, Gardiner TA, Stitt AW, Curtis TM: Advanced glycation end product modified basement membrane attenuates endothelin-1 induced $[Ca^{2+}]_i$ signaling and contraction in retinal microvascular pericytes. *Mol Vis* 2004, **10**:996-1004.
43. McCarty JH, Monahan-Earley RA, Brown LF, Keller M, Gerhardt H, Rubin K, Shani M, Dvorak HF, Wolburg H, Bader BL, Dvorak AM, Hynes RO: Defective associations between blood vessels and brain parenchyma lead to cerebral hemorrhage in mice lacking alphav integrins. *Mol Cell Biol* 2002, **22**:7667-7677.
44. McCarty JH: Cell biology of the neurovascular unit: implications for drug delivery across the blood-brain barrier. *Assay Drug Dev Technol* 2005, **3**:89-95.
45. Hayden MR, Sowers JR, Tyagi SC: The central role of vascular extracellular matrix and basement membrane remodeling in metabolic syndrome and type 2 diabetes: the matrix preloaded. *Cardiovasc Diabetol* 2005, **4**:9.
46. Ryan US, Ryan JW, Whitaker C: How do kinins affect vascular tone? *Adv Exp Med Biol* 1979, **120A**:375-391.
47. Filipovich-Rimon T, Fleisher-Berkovich S: Protein synthesis dependent effects of kinins on astrocyte prostaglandin synthesis. *Peptides* 2010, **31**:651-656.
48. Rucker HK, Wynder HJ, Thomas WE: Cellular mechanisms of CNS pericytes. *Brain Res Bull* 2000, **51**:363-369.
49. Carlson EC: Fenestrated subendothelial basement membranes in human retinal capillaries. *Invest Ophthalmol Vis Sci* 1989, **30**:1923-1932.
50. Larson DM, Haudenschild CC, Beyer EC: Gap junction messenger RNA expression by vascular wall cells. *Circ Res* 1990, **66**:1074-1080.
51. Cuevas P, Gutierrez-Diaz JA, Reimers D, Dujovny M, Diaz FG, Ausman JI: Pericyte-endothelial gap junctions in human cerebral capillaries. *Anat Embryol (Berl)* 1984, **170**:155-159.
52. Nakamura K, Kamouchi M, Kitazono T, Kuroda J, Matsuo R, Hagiwara N, Ishikawa E, Ooboshi H, Ibayashi S, Iida M: Role of NHE1 in calcium

- signaling and cell proliferation in human CNS pericytes. *Am J Physiol Heart Circ Physiol* 2008, **294**:H1700-1707.
- 53. Allsopp G, Gamble HJ: An electron microscopic study of the pericytes of the developing capillaries in human fetal brain and muscle. *J Anat* 1979, **128**:155-68.
 - 54. Jójárt I, Joó F, Siklós L, László FA: Immunoelectron histochemical evidence for innervations of brain microvessels by vasopressin-immunoreactive neurons in the rat. *Neurosci Lett* 1984, **51**:259-264.
 - 55. Castejón OJ: Electron microscopic study of capillary wall in human cerebral edema. *J Neuropathol Exp Neurol* 1980, **39**:296-328.
 - 56. Oku H, Kodama T, Sakagami K, Puro DG: Diabetes-induced disruption of gap junction pathways within the retinal microvasculature. *Invest Ophthalmol Vis Sci* 2001, **42**:1915-1920.
 - 57. Kawamura H, Oku H, Li Q, Sakagami K, Puro DG: Endothelin-induced changes in the physiology of retinal pericytes. *Invest Ophthalmol Vis Sci* 2002, **43**:882-888.
 - 58. Li AF, Sato T, Haimovici R, Okamoto T, Roy S: High glucose alters connexin 43 expression and gap junction intercellular communication activity in retinal pericytes. *Invest Ophthalmol Vis Sci* 2003, **44**:5376-5382.
 - 59. Dore-Duffy P, LaManna JC: Physiologic angiodynamics in the brain. *Antioxid Redox Signal* 2007, **9**:1363-1371.
 - 60. Fruttiger M: Development of the mouse retinal vasculature: angiogenesis versus vasculogenesis. *Invest Ophthalmol Vis Sci* 2002, **43**:522-527.
 - 61. Canfield AE, Allen TD, Grant ME, Schor SL, Schor AM: Modulation of extracellular matrix biosynthesis by bovine retinal pericytes in vitro: effects of the substratum and cell density. *J Cell Sci* 1990, **96**:159-169.
 - 62. Jiang B, Liou GI, Behzadian MA, Caldwell RB: Astrocytes modulate retinal vasculogenesis: effects on fibronectin expression. *J Cell Sci* 1994, **107**:2499-2508.
 - 63. Plate KH: Mechanisms of Angiogenesis in the Brain. *J Neuropath Exp Neurol* 1999, **58**:313-320.
 - 64. Gardner TW, Lieth E, Khin SA, Barber AJ, Bonsall DJ, Lesher T, Rice K, Brennan WA Jr: Astrocytes increase barrier properties and ZO-1 expression in retinal vascular endothelial cells. *Arterioscler Thromb Vasc Biol* 2006, **55**:2401-2411.
 - 65. Hori S, Ohtsuki S, Hosoya K, Nakashima E, Terasaki TA: Pericyte-derived angiopoietin-1 multimeric complex induces occludin gene expression in brain capillary endothelial cells through Tie-2 activation in vitro. *J Neurochem* 2007, **89**:503-513.
 - 66. Haseloff RF, Blasig IE, Bauer HC, Bauer H: In search of the astrocytic factor(s) modulating blood-brain barrier functions in brain capillary endothelial cells in vitro. *Cell Mol Neurobiol* 2005, **25**:25-39.
 - 67. Kim JH, Kim JH, Yu YS, Kim DH, Kim KWJ: Recruitment of pericytes and astrocytes is closely related to the formation of tight junction in developing retinal vessels. *Neurosci Res* 2009, **87**:653-659.
 - 68. Benner M, Kisselberth WS, Su Y, Besnard K, Messing A: GFAP promoter direct astrocyte-specific expression in transgenic mice. *J Neurosci* 1994, **14**:1030-1037.
 - 69. Delaney CL, Brenner M, Messing A: Conditional ablation of cerebellar astrocytes in postnatal transgenic mice. *J Neurosci* 1996, **16**:6908-6918.
 - 70. McCall MA, Gregg RG, Behringer RR, Brenner M, Delaney CL, Galbreath EJ, Zhang CL, Pearce RA, Chiu SY, Messing A: Targeted deletion in astrocyte intermediate filament (Gfap) alters neuronal physiology. *Proc Natl Acad Sci USA* 1996, **93**:6361-6366.
 - 71. Nawashiro H, Messing A, Azzam N, Brenner M: Mice lacking GFAP are hypersensitive to traumatic cerebrospinal injury. *Neuroreport* 1998, **9**:1691-1696.
 - 72. Bertossi M, Girolamo F, Errede M, Virgintino D, Roncali L: Effects of 6-aminonicotinamide gliotoxin on blood-brain barrier differentiation in the chick embryo cerebellum. *Anat Embryol* 2003, **207**:209-219.
 - 73. Nakagawa S, Deli MA, Kawaguchi H, Shimizudani T, Shimono T, Kittel A, Tanaka K, Niwa M: A new blood-brain barrier model using primary rat brain endothelial cells, pericytes and astrocytes. *Neurochem Int* 2009, **54**:253-263.
 - 74. Al Ahmad A, Gassmann M, Ogunshola OO: Maintaining blood-brain barrier integrity: pericytes perform better than astrocytes during prolonged oxygen deprivation. *J Cell Physiol* 2009, **218**:612-622.
 - 75. Nakagawa S, Deli MA, Nakao S, Honda M, Hayashi K, Nakaoke R, Kataoka Y, Niwa M: Pericytes from brain microvessels strengthen the barrier integrity in primary cultures of rat brain endothelial cells. *Cell Mol Neurobiol* 2007, **27**:687-694.
 - 76. Bryan BA, D'Amore PA: Pericytes isolation and use in endothelial/pericyte co-culture models. *Methods Enzymol* 2008, **443**:315-331.
 - 77. Wolburg H, Noell S, Wolburg-Buchholz K, Mack A, Fallier-Becker P: Agrin, aquaporin-4, and astrocyte polarity as an important feature of the blood-brain barrier. *Neuroscientist* 2009, **15**:180-193.
 - 78. Mendelson K, Swendeman S, Saftig P, Blobel CP: Stimulation of platelet-derived growth factor receptor beta (PDGFRbeta) activates ADAM17 and promotes metalloproteinase-dependent cross-talk between the PDGFRbeta and epidermal growth factor receptor (EGFR) signaling pathways. *J Biol Chem* 2010, **285**:25024-25032.
 - 79. Fukushi J, Makagiansar IT, Stallcup WB: NG2 proteoglycan promotes endothelial cell motility and angiogenesis via engagement of galectin-3 and alpha3beta1 integrin. *Mol Biol Cell* 2004, **15**:3580-3590.
 - 80. Hurtado O, Lizasoain I, Fernández-Tomé P, Alvarez-Barrientos A, Leza JC, Lorenzo P, Moro MA: TACE/ADAM17-TNF-alpha pathway in rat cortical cultures after exposure to oxygen-glucose deprivation or glutamate. *J Cereb Blood Flow Metab* 2002, **22**:576-585.
 - 81. Van Hinsbergh VW, Koolwijk P: Endothelial sprouting and angiogenesis: matrix metalloproteinases in the lead. *Cardiovasc Res* 2008, **78**:203-212.
 - 82. Risau W, Drexler H, Mironov V, Smits A, Siegbahn A, Funa K, Heldin CH: Platelet-derived growth factor is angiogenic in vivo. *Growth Factors* 1992, **7**:261-266.
 - 83. Darland DC, D'Amore PA: TGF beta is required for the formation of capillary-like structures in three-dimensional cocultures of 10T1/2 and endothelial cells. *Angiogenesis* 2001, **4**:11-20.
 - 84. Hellstrom M, Kalen M, Lindahl P, Abramsson A, Betsholtz C: Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during embryonic blood vessels formation in the mouse. *Development* 1999, **126**:3047-3055.
 - 85. Lindblom P, Gerhardt H, Leibner S, Abramsson A, Enge M, Hellstrom M, Backstrom G, Fredriksson S, Landegren U, Nyström HC, Bergstrom G, Dejana e, Ostman A, Lindahl P, Betsholtz C: Endothelial PDGF-B retention is required for proper investment of pericytes in the microvessel wall. *Genes Dev* 2003, **17**:1835-1840.
 - 86. Armulik A, Abramsson A, Betsholtz C: Endothelial/pericyte interactions. *Circ Res* 2005, **97**:512-523.
 - 87. Bjarnegård M, Enge M, Norlin j, Gustafsdottir S, Fredriksson S, Abramsson A, Takemoto M, Gustafsson E, Fassler R, Betsholtz C: Endothelium-specific ablation of PDGFB leads to pericyte loss and glomerular, cardiac and placental abnormalities. *Development* 2004, **313**:1847-1857.
 - 88. Steinach E, Kahn BH: Echte Contractilität und motorische Innervation der Blutcapillaren. *Pflügers Archiv* 1903, **97**:195, [German].
 - 89. Ni TG: The active response of capillaries of frogs, tadpoles, fish, bats, and men to various forms of excitation. *Amer J Phys* 1922, **62**:282-309.
 - 90. Peppiatt CM, Howarth C, Mobbs P, Attwell D: Bidirectional control of CNS capillary diameter by pericytes. *Nature* 2006, **443**:700-704.
 - 91. Hamilton NB, Attwell D, Hall CN: Pericyte-mediated regulation of capillary diameter: a component of neurovascular coupling in health and disease. *Front Neuroenergetics* 2010, **2**:5.
 - 92. Bell RD, Winkler EA, Sagare AP, Singh I, Larue B, Deane R, Zlokovic BV: Pericytes Control Key Neurovascular Functions and Neuronal Phenotype in the Adult Brain and during Brain Aging. *Neuron* 2010, **68**:409-412.
 - 93. Quignard JF, Harley EA, Duhaud J, Vanhoutte PM, Félix M: K⁺ channels in cultured bovine retinal pericytes: effects of beta-adrenergic stimulation. *J Cardiovasc Pharmacol* 2003, **42**:379-388.
 - 94. Cao C, Goo JH, Lee-Kwon W, Pallone TL: Vasa recta pericytes express a strong inward rectifier K⁺ conductance. *Am J Physiol Regul Integr Comp Physiol* 2006, **290**:R1601-1607.
 - 95. Jackson WF: Potassium channels in the peripheral microcirculation. *Microcirculation* 2005, **12**:113-127.
 - 96. Bondjers C, He L, Takemoto M, Norlin J, Asker N, Hellström M, Lindahl P, Betsholtz C: Microarray analysis of blood microvessels from PDGF-B and PDGF-Rbeta mutant mice identifies novel markers for brain pericytes. *FASEB J* 2006, **20**:1703-1705.
 - 97. Kelley C, D'Amore P, Hechtman HB, Shepro D: Vasoactive hormones and cAMP affect pericyte contraction and stress fibres in vitro. *J Muscle Res Cell Motil* 1988, **9**:184-194.
 - 98. Das A, Frank RN, Weber ML, Kennedy A, Reidy CA, Mancini MA: ATP causes retinal pericytes to contract in vitro. *Exp Eye Res* 1988, **46**:349-362.
 - 99. Matthew KE, Herman IM: The pericyte: Cellular regulation of microvascular blood flow. *Microvasc Res* 2009, **77**:235-246.

100. Anderson DR: Glaucoma, capillaries and pericytes. 1. Blood flow regulation. *Ophthalmologica* 1996, **210**:257-262.
101. Edelman DA, Jiang Y, Tyburski J, Wilson RF, Steffes C: Pericytes and their role in microvasculature homeostasis. *J Surg Res* 135:305-311.
102. Hughes S, Gardiner T, Hu P, Baxter L, Rosinova E, Chan-Ling T: Altered pericyte-endothelial relations in the rat retina during aging: implications for vessel stability. *Neurobiol Aging* 2006, **27**:1838-1847.
103. Kelley C, D'Amore P, Hechtman HB, Shepro D: Vasoactive hormones and cAMP affect pericyte contraction and stress fibres *in vitro*. *J Muscle Res Cell Motil* 1988, **9**:184-194.
104. Filosa JA, Nelson MT, Gonzales Bosc LV: Activity-dependent NFATc3 nuclear accumulation in pericytes from cortical parenchymal microvessels. *J Physiol Cell Physiol* 2007, **293**:1797-1805.
105. Lieber S, Plate KH: Differentiation of the brain vasculature: the answer came blowing by the Wnt. *J Angiogenes Res* 2010, **2**:1.
106. Lee HS, Han J, Bai HJ, Kim KW: Brain angiogenesis in developmental and pathological processes: regulation, molecular and cellular communication at the neurovascular interface. *FEBS J* 2009, **276**:4622-4635.
107. Stenman JM, Rajagopal J, Carroll TJ, Ishibashi M, McMahon J, McMahon AP: Canonical WNT Signaling regulates organ-specific assembly and differentiation of CNS vasculature. *Science* 2008, **322**:1247-1250.
108. Daneman R, Agalliu D, Zhou L, Kuhnert F, Kuo CJ, Barres BA: Wnt/beta-catenin signaling is required for CNS, but not-CNS, angiogenesis. *Proc Natl Acad Sci USA* 2009, **106**:641-646.
109. White BD, Nathe RJ, Maris DO, Nguyen NK, Goodson JM, Moon RT, Horner PJ: Beta-catenin signaling increases in proliferating NG2+ progenitors and astrocytes during post-traumatic gliogenesis in the adult brain. *Stem Cells* 2010, **28**:297-307.
110. Leveen P, Pekny M, Gebre-Medhin S, Swolin B, Larsson E, Betsholtz C: Mice deficient for PDGF B show renal, cardiovascular, and hematological abnormalities. *Genes Dev* 1994, **8**:1875-1887.
111. Siegfried G, Basak A, Prichett-Pejic W, Scamuffa N, Ma L, Benjannet S, Veinot JP, Calvo F, Seidah N, Khatib AM: Regulation of the stepwise proteolytic cleavage and secretion of PDGF-B by the proprotein convertases. *Oncogene* 2005, **24**:6925-6935.
112. Zlokovic BV: The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron* 2008, **57**:178-201.
113. Danema R, Zhou L, Amanuel A, Kebede , Barres BA: Pericytes are required for blood-brain barrier integrity during embryogenesis. *Nature* 2010, **468**:223-231.
114. Armulik A, Genov G, Mae M, Niclancioglu MN, Wallgard E, Niaudet C, Liqun H, Norlin J, Lindblom P, Strittmatter K, Johansson BR, Betsholtz C: Pericytes regulate the blood-brain barrier. *Nature* 2010, **468**:557-561.
115. Girolamo F, Virgintino D, Errede M, Capobianco C, Bernandini N, Bertossi M, Roncali L: Involvement of metalloprotease-2 in the development of human brain microvessels. *Histochem Cell Biol* 2004, **122**:261-270.
116. Diaz-Flores L, Gutierrez R, Valladares F, Varela H, Perez M: Intense vascular sprouting from rat femoral vein induced by prostaglandins E1 and E2. *Anat Rec* 1994, **238**:68-76.
117. Nehls V, Denzer K, Drenckhahn D: Pericyte involvement in capillary sprouting during angiogenesis *in situ*. *Cell Tissue Res* 1992, **270**:469-474.
118. Virgintino D, Ozerdem U, Girolamo F, Roncali L, Stallcup WB, Perris R: Reversal of cellular roles in angiogenesis: implications for anti-angiogenic therapy. *J Vasc Res* 2008, **45**:129-31.
119. Diaz-Flores L, Gutierrez R, Madrid JF, Varela H, Valladares F, Acosta E, Martin-Vasallo P, Diaz-Flores L Jr: Pericytes. Morphofunction, interactions and pathology in a quiescent and activated mesenchymal cell niche. *Histol Histopathol* 2009, **24**:909-969.
120. Kokovay E, Li L, Cunningham LA: Angiogenetic recruitment of pericytes from bone marrow after stroke. *J Cereb Blood Flow Metab* 2006, **26**:545-555.
121. Minakawa T, Bready J, Berliner J, Fisher M, Cancilla PA: In vitro interaction of astrocytes and pericytes with capillary-like structures of brain microvessels oendothelium. *Lab Invest* 1991, **65**:32-40.
122. Lai CH, Kuo KH: The critical component to establish *in vitro* BBB model: Pericyte. *Brain Res Brain Res Rev* 2005, **50**:258-265.
123. Ahmad AA, Taboada CB, Gassmann M, Ogunshola OO: Astrocytes and pericytes differentially modulate blood-brain barrier characteristics during development and hypoxic insult. *J Cereb Blood Flow Metab* 2010, **10**:1038-1045.
124. Haskew RE, Mongin AA, Kimelberg HK: Peroxynitrite enhances astrocytic volume-sensitive excitatory amino acid release via a src tyrosine kinase-dependent mechanism. *J Neurochem* 2002, **82**:903-912.
125. Hede SM, Hansson I, Afink GB, Eriksson A, Nazarenko I, Andrae J, Genove G, Westermark B, Nister M: GFAP promoter driven transgenic expression of PDGFB in the mouse brain leads to glioblastoma in a Trp53 null background. *Glia* 2009, **57**:1143-1153.
126. Tagami M, Yamagata K, Fujino H, Kubota A, Nara Y, Yamory Y: Morphological differentiation of endothelial cells co-cultured with astrocytes on type-I or type IV collagen. *Cell Tissue Res* 1992, **268**:225-232.
127. Choi MH, Lee IK, Kim GW, Kim BU, Han YH, Yu DY, Park HS, Kim KY, Lee JS, Choi C, Bae YS, Lee BI, Rhee SG, Kang SW: Regulation of PDGF signaling and vascular remodeling by peroxiredoxin II. *Nature* 2005, **435**:347-353.
128. Haskew-Layton RE, Payapilly JB, Smirnova NA, Ma TC, Chan KK, Murphy TH, Guo H, Langley B, Sultana B, Butterfield DA, Santagata S, Alldred MJ, Gazaryan IG, Bell GW, Ginsberg SD, Ratan RR: Controlled enzymatic production of astrocytic hydrogen peroxide protects neurons from oxidative stress via an Nrf2-independent pathway. *Proc Natl Acad Sci USA* 2010, **107**:17385-17390.
129. Ueno H, Colbert H, Escobedo JA, Williams LT: Inhibition of PDGF beta receptor signal transduction by coexpression of a truncated receptor. *Science* 1991, **252**:844-848.
130. Zhou T, Liu H, Zhang P, Luo C: Effect of truncated PDGF-Alpha receptor on cellular cycle and proliferation for vascular cells of pulmonary artery. *Sichuan Da Xue Xue Bao Yi Xue Ban* 2003, **34**:31-3.
131. Kurup S, Abramsson A, Li JP, Lindahl U, Kjellen L, Betsholtz C, Genhardt H, Spillmann D: Heparin Sulphate requirement in platelet-derived growth factor B-mediated pericyte recruitment. *Biochem Soc Trans* 2006, **34**:454-455.
132. Deguchi Y, Okutsu H, Okura T, Yamada S, Kimura R, Yuge T, Furukawa A, Morimoto K, Tachikawa m, Ohtsuki S, Hosoya K, Terasaki T: Internalization of basic fibroblast growth factor at the mouse blood-brain barrier involves perlecan, a heparan sulfate proteoglycan. *J Neurochem* 2002, **83**:381-389.
133. Santos-Silva A, Fairless R, Frame MC, Montague P, Smith GM, Toft A, Riddell JS, Barnett SC: FGF/heparin differentially regulates Schwann cell and olfactory ensheathing cell interactions with astrocytes: a role in astrocytosis. *J Neurosci* 2007, **27**:7154-7167.
134. Al Ahmad A, Taboada CB, Gassmann M, Ogunshola OO: Astrocytes and pericytes differentially modulate blood-brain barrier characteristics during development and hypoxic insult. *J Cereb Blood Flow Metab* 2010.
135. Pfister F, Feng Y, vom Hagen F, Hoffmann S, Molema G, Hillebrands JL, Shani M, Deutsch U, Hammes HP: Pericyte migration: a novel mechanism of pericyte loss in experimental diabetic retinopathy. *Diabetes* 2008, **57**:2495-2502.
136. Duz B, Oztas E, Erginay T, Erdogan E, Gonul E: The effect of moderate hypothermia in acute ischemic stroke on pericyte migration: an ultrastructural study. *Cryobiology* 2007, **55**:279-284.
137. Claudio L, Brosnan CF: Effects of prazosin on the blood-brain barrier during experimental autoimmune encephalomyelitis. *Brain Res* 1992, **594**:233-243.
138. Kunz J, Krause D, Gehrmann J, Dermietzel R: Changes in the expression pattern of blood-brain barrier-associated pericytic aminopeptidase N (pAP N) in the course of acute experimental autoimmune encephalomyelitis. *J Neuroimmunol* 1995, **59**:41-55.
139. Dore-Duffy P, Balabanov R, Rafols J, Swborg RH: Recovery phase of acute experimental autoimmune encephalomyelitis in rats corresponds to development of endothelial cell unresponsiveness to interferon gamma activation. *J Neurosci Res* 1996, **44**:223-234.
140. Bolton C: Neurovascular damage in experimental allergic encephalomyelitis: a target for pharmacological control. *Mediators Inflamm* 1997, **6**:295-302.
141. Zlokovic BV: The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron* 2008, **57**:178-201.
142. Ho KL: Ultrastructure of cerebellar capillary hemangioblastoma. IV. Pericytes and their relationship to endothelial cells. *Acta Neuropathol* 1985, **67**:254-264.
143. Winkler F, Kozin SV, Tong RT, Chae SS, Booth MF, Garkavtsev I, Xu L, Hicklin DJ, Fukumura D, di Tomaso E, Munn LL, Jain RK: Kinetics of vascular normalization by VEGFR2 blockade governs brain tumor response to radiation: role of oxygenation, angiopoietin-1, and matrix metalloproteinases. *Cancer Cell* 2004, **6**:553-563.

144. Hammes HP, Lin J, Renner O, Shani M, Lundqvist A, Betsholtz C, Brownlee M, Deutsch U: **Pericytes and the pathogenesis of diabetic retinopathy.** *Diabetes* 2002, **51**:3107-3112.
145. Frank RN, Turczyn TJ, Das A: **Pericyte coverage of retinal and cerebral capillaries.** *Invest Ophthalmol Vis Sci* 1990, **31**:999-1007.
146. Feng Y, Pfister F, Schreiter K, Wang Y, Stock O, Vom Hagen F, Wolburg H, Hoffmann S, Deutsch U, Hammes HP: **Angiopoietin-2 deficiency decelerates age-dependent vascular changes in the mouse retina.** *Cell Physiol Biochem* 2008, **21**:129-136.
147. Szpak GM, Lewandowska E, Wierba-Bobrowicz T, Bertrand E, Pasenik E, Mendel T, Stepien T, Leszczynska A, Rafalowska J: **Small cerebral vessel disease in familial amyloid and non-amyloid angiopathies: FAD-PS-1 (P117L) mutation and CADASIL. Immunohistochemical and ultrastructural studies.** *Folia Neuropathol* 2007, **45**:192-204.
148. Wiegli J, Wisniewski HM: **Tubuloreticular structures in microglial cells, pericytes and endothelial cells in Alzheimer's disease.** *Acta Neuropathol* 1992, **83**:653-658.
149. Stewart PA, Hayakawa K, Akers MA, Vinters HV: **A morphometric study of the blood-brain barrier in Alzheimer's disease.** *Lab Invest* 1992, **67**:734-742.
150. Yamagishi S, Imaizumi T: **Pericyte biology and diseases.** *Int J Tissue React* 2005, **27**:125-135.
151. Kischer CW: **The microvessels in hypertrophic scars, keloids and related lesions: a review.** *J Submicrosc Cytol Pathol* 1992, **24**:281-296.
152. Li W, Yanoff M, Liu X, Ye X: **Retinal capillary pericyte apoptosis in early human diabetic retinopathy.** *Chin Med J (Engl)* 1997, **110**:659-663.
153. Shojaee N, Patton WF, Hechtman HB, Shepro D: **Myosin translocation in retinal pericytes during free-radical induced apoptosis.** *J Cell Biochem* 1999, **75**:118-129.
154. Zagzag D, Amirnovin R, Greco MA, Yee H, Holash J, Wiegand SJ, Zabitski S, Yancopoulos GD, Grumet M: **Vascular apoptosis and involution in gliomas precede neovascularization: a novel concept for glioma growth and angiogenesis.** *Lab Invest* 2000, **80**:837-849.
155. Machein MR, Knedla A, Knoth R, Wagner S, Neuschl E, Plate KH: **Angiopoietin-1 promotes tumor angiogenesis in a rat glioma model.** *Am J Pathol* 2004, **165**:1557-1570.
156. Bexell D, Gunnarsson S, Tormin A, Darabi A, Gisselsson D, Roybon L, Scheding S, Bengzon J: **Bone marrow multipotent mesenchymal stroma cells act as pericyte-like migratory vehicles in experimental gliomas.** *Mol Ther* 2009, **17**:183-190.
157. Vermeulen PB, Colpaert C, Salgado R, Royers R, Hellermans H, Van Den Heuvel E, Goovaerts G, Dirix LY, Van ME: **Liver metastases from colorectal adenocarcinomas grow in three patterns with different angiogenesis and desmoplasia.** *J Pathol* 2001, **195**:336-342.
158. Li W, Liu X, Yanoff M, Cohen S, Ye X: **Cultured retinal capillary pericytes die by apoptosis after an abrupt fluctuation from high to low glucose levels: a comparative study with retinal capillary endothelial cells.** *Diabetologia* 1996, **39**:537-547.
159. Braun A, Xu H, Hu F, Kocherlakota P, Siegel D, Chander P, Ungvari Z, Csizsar A, Nedergaard M, Ballabh P: **Paucity of pericytes in germinal matrix vasculature of premature infants.** *J Neurosci* 2007, **27**:12012-12024.
160. Zhu M, Madigan MC, van Driel D, Maslim J, Billson FA, Provis JM, Penfold PL: **The human hyaloid system: cell death and vascular regression.** *Exp Eye Res* 2000, **70**:767-776.
161. Khouri J, Langleben D: **Platelet-activating factor stimulates lung pericyte growth in vitro.** *Am J Physiol* 1996, **270**:L298-304.
162. Shi X: **Cochlear pericyte responses to acoustic trauma and involvement of hypoxia-inducible factor-1 alpha and vascular endothelial growth factor.** *Am J Pathol* 2009, **174**:1692-1704.
163. Song N, Huang Y, Shi H, Yuan S, Ding Y, Song X, Fu Y, Luo Y: **Overexpression of platelet-derived growth factor-BB increases tumor pericyte content via stromal-derived factor-1 alpha/CXCR4 axis.** *Cancer Res* 2009, **69**:6057-64.
164. Zetterberg E, Alessandro MV, Migliaccio AR, Vainchenker W, Tulliez M, Dickie R, Hasselbach H, Rogers R, Palmlad J: **Pericyte coverage of abnormal blood vessels in myelofibrotic bone marrows.** *Haematologica* 2007, **92**:597-604.
165. Vinukonda G, Dummula K, Malik S, Hu F, Thompson CI, Csizsar A, Ungvari Z, Ballabh : **Effect of prenatal glucocorticoids on cerebral vasculature of the developing brain.** *Stroke* 2010, **41**:1766-1773.
166. Fioravanzo I, Venturini M, Di Liddo R, Marchi F, Grandi C, Parnigotto PP, Folin M: **Involvement of Rat Hippocampal Astrocytes in β -Amyloid - Induced Angiogenesis and Neuroinflammation.** *Curr Alzheimer Res* 2010.
167. Sortino MA, Platania P, Chisari M, Merlo S, Copani A, Catania MV: **A major role for astrocytes in the neuroprotective effect of estrogen.** *Drug Dev Res* 2006, **66**:126-135.
168. Inman DM, Horner PJ: **Reactive nonproliferative gliosis predominates in a chronic mouse model of glaucoma.** *Glia* 2007, **55**:942-953.
169. Bukovsky A, Cekanova M, Caudle MR, Wimalasena J, Foster JS, Henley DC, Elder RF: **Expression and localization of estrogen receptor-alpha protein in normal and abnormal term placentae and stimulation of trophoblast differentiation by estradiol.** *Reprod Biol Endocrinol* 2003, **1**:13.
170. Kappert k, Caglayan E, Huntgeburth M, Baumer AT, Sparwel J, Uebel M, Rosenkranz S: **17Beta-estradiol attenuates PDGF signaling in vascular smooth muscle cells at the postreceptor level.** *Am J Physiol Heart Circ Physiol* 2006, **290**:H538-546.
171. Bonnie J, Korach D, Korach KS: **Estrogen receptors and human disease.** *J Clin Invest* 2006, **116**:561-570.
172. Gaughwin PM, Caldwell MA, Anderson JM, Schwiening CJ, Fawcett JW, Compston DAS, Chandran S: **Astrocytes promote neurogenesis from oligodendrocyte precursor cells.** *Eur J Neurosci* 2006, **23**:945-956.
173. Mori T, Buffo A, Gotz M: **The novel roles of glial cells revisited: the contribution of radial glia and astrocytes to neurogenesis.** *Curr Top Dev Biol* 2005, **69**:67-99.
174. Zheng T, Marshal LL, Laywell ED, Steindler DA: **Neurogenic astrocytes transplanted into the adult mouse lateral ventricle contribute to olfactory neurogenesis, and reveal a novel intrinsic subependymal neuron.** *J Neuroscience* 2006, **142**:175-185.
175. Kraev IV, Godukhin OV, Patrushev IV, Davies HA, Popov VI, Stewart MG: **Partial kindling induces neurogenesis, activates astrocytes and alters synaptic morphology in the dentate gyrus of freely moving adult rats.** *J Neurosci* 2009, **162**:254-267.
176. Nakagawa T, Miyamoto O, Janjua NA, Auer RN, Nagao S, Itano T: **Localization of nestin in amygdaloid kindled rat; an immuno-electron microscopic study.** *Can J Neurol Sci* 2004, **31**:514-519.
177. Williams S, Wagner PD: **Transgenic animals in integrative biology: approaches and interpretations of outcome.** *J Appl Physiol* 2000, **88**:1119-1126.
178. Lindahl P, Karlsson L, Hellstrom M, Gebre-Medhin S, Willets K, Health JK, Betsholtz C: **Alveogenesis failure in PDGF-A-deficient mice is coupled to lack of distal spreading of alveolar smooth muscle cell progenitors during lung development.** *Development* 1997, **124**:3943-3953.
179. Soriano P: **The PDGF alpha receptor is required for neural crest cell development and for normal patterning of the somites.** *Development* 1997, **124**:2691-2700.
180. Betsholtz C: **Role of platelet-derived growth factors in mouse development.** *Int J Dev Biol* 1995, **39**:817-825.

doi:10.1186/2045-8118-8-8

Cite this article as: Bonkowski et al.: The CNS microvascular pericyte: pericyte-astrocyte crosstalk in the regulation of tissue survival. *Fluids and Barriers of the CNS* 2011 **8**:8.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

