




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to Editors

Smoking and COVID-19, the paradox to discover: An Italian retrospective, observational study in hospitalized and non-hospitalized patients



ARTICLE INFO

Keywords

Smoking
SARS-CoV2
COVID-19 disease

We have read with great interest the article published on Medical Hypotheses by de Bernardis & Busà about the putative role for the tobacco mosaic virus in smokers' resistance to COVID-19 [1]. We agree with the Authors and we also believe that smoking has a complex and not yet clear link with COVID-19. We considered 2 cohorts of 601 patients from 2 Northern Italian cities (Brescia and Bergamo) most affected by the SARS-CoV-2 infection: one concerning patients admitted to our hospital in Brescia, and another one concerning non-hospitalized patients evaluated at the territorial medicine level in Bergamo. The aim of the study was to assess whether or not there was a correlation between current active smoking and severity of COVID-19 disease. 299 consecutive hospitalized COVID-19 patients (M/F = 208 (70%)/ 91(30%) were included and among these 46 (15%) were current active smokers. The mean age was 68.8 years (± 11.7). The patients who died as a consequence of COVID-19 disease were 87 (29%). Table 1 shows that active smoking and COVID-19 are independent variables (p -value = 0.99) in our hospitalized patients. Garufi et al. claim that the current available data on the relationship between COVID-19 and smoking only concern hospitalized patients [2]. In our study 302 consecutive non-hospitalized COVID-19 patients (M/F = 176 (58.3%) /126 (41.7%) were also included and among these 92 (30%) were current active smokers. The non-hospitalized patients who died as a result of COVID-19 disease were 28 (9.3%). The table I shows that current active smoking and COVID-19 were independent variables (p -value = 0.66) also for non-hospitalized patients. These data are in line with the results of the study performed by Meini et al. and Lippi et al. that reported an unexpectedly low prevalence of current smokers among COVID-19 hospitalized patients [3,4]. We agree with the Authors and we also found no significant correlation between active smoking and severity of COVID-19 disease. We believe that our study can add important information about the link between smoking and COVID-19 in Italian hospitalized and non-hospitalized patients too.

Probably the link between smoking and COVID-19 is not simple and does not concern a single factor.

If it is true that smoking increases the expression of ACE2 receptors to which the SARS-Cov-2 attaches, it is equally true that up regulation of this enzyme can decrease the risk of developing serious systemic complications of coronavirus infection [5,6]. This is due to the fact that

ACE2 converts angiotensin II (ANG II) to Angiotensin 1–7 (ANG 1–7) thus decreasing the powerful pro inflammatory effect of ANG II [7,8]. Other Authors have confirmed that ACE2 mitigates pro inflammatory effects by decreasing the circulating amount of ANG II [9]. Moreover, studies in mice have shown that in smokers there is an upregulation of ACE2 which protects from the development of the Acute Respiratory Distress Syndrome (ARDS) [10,11]. Consequently, studies have hypothesized that the up-regulation of this enzyme may protect patients from severe lung damage and that this also occurs in patients with COVID-19 [11].

Although it seems like a paradox, the latest scientific evidences are in favour of the hypothesis that smoking is not associated with the severity of COVID-19 patients. Our case series is interesting because it reports the data of hospitalized and non-hospitalized patients, confirming the non-association between smoking and COVID-19 disease. Future studies will be need to confirm this hypothesis and to discover the underlying pathophysiological mechanisms.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] De Bernardis E, Busà L. A putative role for the tobacco mosaic virus in smokers' resistance to COVID-19. *Med Hypotheses* 2020;143:110153.
- [2] Garufi G, Carbognin L, Orlandi A, Tortora GP, Bria E. Smoking habit and hospitalization for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-related pneumonia: the unsolved paradox behind the evidence. *Eur J Intern Med* 2020;77:121–2. Published online 2020 Apr 23.
- [3] Meini S, Fortini A, Andreini R, Sechi LA, Tascini C. The paradox of the low prevalence of current smokers among COVID-19 patients hospitalized in non-intensive care wards: results from an Italian multicentre case-control study. *Nicotine Tob Res* 2020. <https://doi.org/10.1093/ntr/ntaa18>.
- [4] Lippi G, Henry BM. Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19). *Eur J Intern Med* 2020;75:107–8. Published online 2020 Mar 16.
- [5] Leung JM, Yang CX, Tam A, Shaipanich T, Hackett T-L, Singhera GK, et al. ACE-2 expression in the small airway epithelia of smokers and COPD patients:

<https://doi.org/10.1016/j.mehy.2020.110391>

Received 29 October 2020; Accepted 8 November 2020

Available online 12 November 2020

0306-9877/© 2020 Elsevier Ltd. All rights reserved.

Table 1

See text.

Patients	Variable	Deaths	OR	95% CI OR	P-value
Hospitalized	Smoking	No 0.280	1.01	(0.45;2.18)	0.99
		Yes 0.283			
Not-hospitalized	Smoking	No 0.310	0.74	(0.26;1.89)	0.66
		Yes 0.250			

implications for COVID-19. *Eur Respir J.* 2020;55(5):2000688. Published online 2020 May 14.

- [6] Smith JC, Sheltzer JM. Cigarette smoke triggers the expansion of a subpopulation of respiratory epithelial cells that express the SARS-CoV-2 receptor ACE2. *BioRxiv*2020. 03.28.013672.
- [7] Sanchis-Gomar F, Lavie CJ, Perez-Quilis C, Henry BM, Lippi G. Angiotensin-Converting Enzyme 2 and Antihypertensives (Angiotensin Receptor Blockers and Angiotensin-Converting Enzyme Inhibitors) in Coronavirus Disease 2019. *Mayo Clin Proc.* 2020 Jun; 95(6): 1222–1230. Published online 2020 Apr 4.
- [8] Khan A, Benthin C, Zeno B, Albertson TE, Boyd J, Christie JD, et al. A pilot clinical trial of recombinant human angiotensin-converting enzyme 2 in acute respiratory distress syndrome. *Crit Care* 2017;21:234. Published online 2017 Sep 7.

- [9] Imai Y, Kuba K, Rao S, Huan Y, Guo F, Guan B, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature* 2005;436(7047):112–6.
- [10] Imai Y, Kuba K, Penninger JM. The discovery of angiotensin-converting enzyme 2 and its role in acute lung injury in mice. *Exp Physiol* 2008;93(5):543–8.
- [11] Gurwitz D. Angiotensin receptor blockers as tentative SARS-CoV-2 therapeutics. *Drug Dev Res* 2020;81(5):537–40. Epub 2020 Mar 4.

Carlo Lombardi^{a,*}, Elena Roca^b, Laura Ventura^c, Marcello Cottini^d

^a COVID-19 Unit & Departmental Unit of Pneumology & Allergology, Istituto Ospedaliero Fondazione Poliambulanza, Brescia, Italy

^b COVID-19 Unit & Neurosurgery Departmental Unit, Istituto Ospedaliero Fondazione Poliambulanza, Brescia, Italy

^c Department of Statistical Sciences, University of Padova, Padova, Italy

^d Allergy and Pneumology Outpatient Clinic, Bergamo, Italy

* Corresponding author.

E-mail address: carlo.lombardi@poliambulanza.it (C. Lombardi).