


# Unilateral Acoustic Degradation Delays Attentional Separation of Competing Speech

Trends in Hearing  
Volume 25: 1–13  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/23312165211013242  
journals.sagepub.com/home/tia



Frauke Kraus<sup>1,2</sup>, Sarah Tune<sup>1,2</sup>, Anna Ruhe<sup>1,2</sup>, Jonas Obleser<sup>1,2</sup>, and Malte Wöstmann<sup>1,2</sup> 

## Abstract

Hearing loss is often asymmetric such that hearing thresholds differ substantially between the two ears. The extreme case of such asymmetric hearing is single-sided deafness. A unilateral cochlear implant (CI) on the more severely impaired ear is an effective treatment to restore hearing. The interactive effects of unilateral acoustic degradation and spatial attention to one sound source in multitalker situations are at present unclear. Here, we simulated some features of listening with a unilateral CI in young, normal-hearing listeners ( $N = 22$ ) who were presented with 8-band noise-vocoded speech to one ear and intact speech to the other ear. Neural responses were recorded in the electroencephalogram to obtain the spectrotemporal response function to speech. Listeners made more mistakes when answering questions about vocoded (vs. intact) attended speech. At the neural level, we asked how unilateral acoustic degradation would impact the attention-induced amplification of tracking target versus distracting speech. Interestingly, unilateral degradation did not per se reduce the attention-induced amplification but instead delayed it in time: Speech encoding accuracy, modelled on the basis of the spectrotemporal response function, was significantly enhanced for attended versus ignored intact speech at earlier neural response latencies ( $< \sim 250$  ms). This attentional enhancement was not absent but delayed for vocoded speech. These findings suggest that attentional selection of unilateral, degraded speech is feasible but induces delayed neural separation of competing speech, which might explain listening challenges experienced by unilateral CI users.

## Keywords

speech tracking, envelope, unilateral vocoding, cochlear implant, attention, spectrotemporal response function

Received 23 November 2020; Revised 11 February 2021; accepted 7 April 2021

Selecting the relevant auditory signal from a mixture of sounds is a challenging task. Selective attention enables the listener to prioritize target information against concurrent distraction. In recent years, methods have been developed to quantify the phase locking of neural responses in the magneto/electroencephalogram to the envelope of individual speech signals in the mixture of sounds (for reviews, see Brodbeck & Simon, 2020; Ding & Simon, 2014; Peelle & Davis, 2012; Wöstmann, Fiedler, et al., 2017). A number of studies have convergently shown that normal-hearing listeners' magneto/electroencephalogram responses exhibit stronger neural phase locking to the envelope of attended versus ignored speech (e.g., Ding & Simon, 2014; Horton et al., 2013;

O'Sullivan et al., 2015; Zion Golumbic et al., 2013). Furthermore, there is evidence that the degree of neural phase locking to the envelope of speech correlates with speech intelligibility (e.g., Peelle et al., 2013) and with behavioral indices of speech comprehension (e.g., Etard & Reichenbach, 2019). The neural phase locking

<sup>1</sup>Department of Psychology, University of Lübeck, Lübeck, Germany

<sup>2</sup>Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, Lübeck, Germany

### Corresponding author:

Malte Wöstmann, Department of Psychology, University of Lübeck, Maria-Goeppert-Straße 9a, 23562 Lübeck, Germany.  
Email: malte.woestmann@uni-luebeck.de



to the speech envelope—also referred to as neural speech tracking (Obleser & Kayser, 2019)—constitutes an objective measure of the attentional enhancement of target speech against distraction.

Important for the present study, research has associated auditory processing deficits with particular changes in neural speech tracking. For instance, the differential cortical tracking of attended versus ignored speech was found to be reduced in listeners with stronger hearing loss in some studies (Petersen et al., 2017), although results of other studies suggest stronger tracking of target speech with increasing hearing loss (Decruy et al., 2020; Fuglsang et al., 2020). Furthermore, noise vocoding to simulate hearing with a cochlear implant (CI) reduced the attentional enhancement of target speech (Ding et al., 2013; Kong et al., 2015; Rimmele et al., 2015). A recent study found that bilateral CI users showed smaller early neural separation of attended versus ignored speech (< 200 ms) but larger late neural separation (>200 ms) compared with normal-hearing controls (Paul et al., 2020). These findings demonstrate the potential of neural speech tracking to understand interactive effects of degraded acoustics and attention.

A so-far neglected test case for the impact of acoustic degradation on neural speech tracking is asymmetric hearing. There are at least three reasons why asymmetric hearing poses a relevant test case for neural speech tracking. First, asymmetric hearing provides a well-controlled scenario to study the simultaneous neural tracking of degraded versus intact acoustic input in the same neural system. In contrast, comparing speech tracking in hearing-impaired versus normal-hearing listeners poses the challenge to control for between-subject differences. In such cases, it often remains somewhat unclear to what extent changes in speech tracking reflect differences in the perceived acoustic input versus differences in neural processing capabilities. Second, the prevalence of asymmetric hearing loss in the population should not be disregarded (e.g., Golub et al., 2018; Wilson et al., 1993), which emphasizes the importance to study auditory selective attention in such cases. Third, although unilateral CI implantation can partially restore hearing (for review, see Vlastarakos et al., 2014), effects of the CI-induced unilateral spectral degradation on the attentional selection of target speech against concurrent acoustic masking are largely unknown.

Noise vocoding is a common technique used to simulate some aspects of hearing with a CI. In essence, noise vocoding degrades the spectral information of the acoustic input but leaves the temporal envelope largely intact (Rosen et al., 1999). When applied to target speech, noise vocoding compromises speech comprehension in silence (Shannon et al., 1995) and particularly in background noise (for review, see Moore, 2008). Noise vocoding of speech distractors has been shown to

reduce the extent of distraction, evidenced by better target speech reception (e.g., Westermann & Buchholz, 2017) and better memory for target speech (e.g., Ellermeier et al., 2015; Senan et al., 2018; Wöstmann & Obleser, 2016).

Noise vocoding affects the type of masking elicited by speech distractors: Energetic (acoustic) masking is complemented by informational masking for vocoded speech signals that are still intelligible (e.g., Brungart et al., 2001; Mattys et al., 2009), for instance, through noise vocoding with a larger number of vocoder bands. Furthermore, it has been proposed that informational masking additionally increases with higher similarity of target and distracting speech (e.g., Rosen et al., 2013). In consequence, when competing speech signals are presented dichotically, a unilateral CI user should experience a relatively higher degree of informational masking when the masking speech distractor is presented to the better ear and is thus more intelligible.

Different modulations of neural speech tracking are conceivable that might accompany the attentional selection of a unilateral speech signal that is acoustically degraded versus intact. It has been proposed that auditory selective attention depends on two closely intertwined processes, referred to as *auditory object formation* (grouping of sounds that originate from the same physical source) and *auditory object selection* (selecting an object for prioritized processing at the expense of distraction; Shinn-Cunningham, 2008). The neural tracking of speech, which reflects both bottom-up acoustic information such as the syllabic structure as well as top-down attention, might be a neural correlate of both, auditory object formation and selection (Shinn-Cunningham et al., 2017). We have previously observed that earlier components of the speech tracking response are mainly sensitive to acoustic features, whereas later response components reflect the attention focus and the effort of attentional selection (Fiedler et al., 2019). In agreement with this, a recent study proposed a model wherein earlier neural tracking responses (< ~100 ms) reflect filtering and restoration of acoustic signals, whereas later responses correspond to active attentional streaming (Brodbeck et al., 2020). Thus, the latency at which unilateral acoustic degradation impacts neural speech tracking in the present study allows to draw conclusions about the underlying perceptual and cognitive mechanisms.

The present study implemented a dichotic listening task to present an intact speech signal to one ear and a different, noise-vocoded speech signal to the other ear. This way, the present study deliberately avoided simulation of listening challenges associated with the integration of acoustic hearing (with the nonimplanted ear) and electric hearing (with the implanted ear) of sound emitted from a single physical source (see, e.g., Litovsky

et al., 2019; Ma et al., 2016). Spatial cues guided listeners' attention either to the intact or noise-vocoded target speech, while the respective other speech signal served as a to-be-ignored distractor. Because noise vocoding affects the spectral integrity of acoustic signals, we used electroencephalogram (EEG) recordings to derive temporally- but also spectrally resolved neural response functions. We predicted an attention-induced increase in the neural response to attended compared with ignored speech, which would decrease for attending-versus-ignoring vocoded compared with intact speech. It was, however, an open question at which neural response latency unilateral acoustic degradation would affect the tracking of the speech envelope.

## Materials and Methods

### Participants

We tested  $N=22$  young, normal-hearing adults (age: 19–33 years; mean = 24.6 years;  $SD=3.76$  years; 12 females; all right-handed). Pure tone audiometry ensured that the pure tone average (PTA; average hearing threshold across frequencies 500, 1000, 2000, and 4000 Hz) per ear was below 20 dB HL (hearing level). The mean PTA on the right side was 5.18 HL ( $SD=5.97$  dB) and on the left side 2.68 HL ( $SD=3.59$  dB). Two participants had a moderate hearing loss of up to 40 dB HL at the frequencies 500, 750, and 1000 Hz on the right side. Due to the overall good behavioral performance of those participants and PTA smaller than 20 dB HL, we decided to not exclude the data of these participants. Participants were recruited via the participant database of the Department of Psychology at the University of Lübeck. They were native speakers of German and reported no history of neural disorders.

Participants gave their written informed consent and were financially compensated with €10/hour or received course credit. The study was approved by the local ethics committee of the University of Lübeck.

### Stimuli

Two German audiobooks were used as continuous speech streams ("Eine kurze Geschichte der Menschheit" by Yuval Noah Harari, spoken by a male narrator; "Nero Corleone kehrt zurück - Es ist immer genug Liebe da" by Elke Heidenreich, spoken by a female narrator). Of the first 64 min of each audiobook, we used the first half for the to-be-attended stream and the second half for the to-be-ignored stream. We generated individual trials by cutting the respective portions of the audiobook into segments of 4 min each.

For noise vocoding, we used a gamma tone filterbank (implemented in the Auditory Toolbox for MATLAB; Slaney, 1993) with eight logarithmically spaced filters between 0 and 8 kHz. After filtering the speech stimuli with these filters, we multiplied the envelope (magnitude of the Hilbert transform) in each frequency band with white noise filtered with the same filterbank and applied a root mean square normalization. Finally, the eight bands were summed and again normalized to the root mean square value of the intact audiobook parts (Shannon et al., 1995). We used vocoding with eight frequency bands to balance task difficulty in a way that participants could perform the task well above chance level but that listening to vocoded speech was more challenging than listening to intact speech.

### Procedure

Participants performed the continuous speech task sitting in a sound-attenuating chamber. The audiobooks were presented dichotically through in-ear headphones (EARTONE 3 A, 3 M). Participants could set the overall volume to a comfortable level before the experiment started. During the whole experiment, the input to one ear was vocoded speech, while the input to the other was intact speech. Therefore, there were two experimental settings. In Setting 1, participants attended to the intact speech and ignored the vocoded distractor stream. By contrast, in Setting 2, they attended to the vocoded speech and ignored the intact speech stream. The side of vocoding was counterbalanced across participants.

In total, the experiment consisted of 16 trials with 4 min duration each. Each of the two experimental settings was presented in eight trials of the experiment. Participants started with two attention-to-intact speech trials, followed by two attention-to-vocoded speech trials and so on. The to-be-attended audiobook (male vs. female talker) alternated from each trial to the next. Before each trial, a spatial cue and the text "Listen to the left/right" signalled the to-be-attended side and participants started the trial with a button press on a four-button response box (The Black Box Toolkit, Sheffield, UK). After each trial, participants answered four multiple-choice questions, each one with four alternative response options about the content of the to-be-attended audiobook. During the whole task, we measured participants' EEG.

### EEG Recording and Preprocessing

The EEG was recorded from 64 electrodes (ActiChamp, Brain Products, München) at a sampling rate of 1000 Hz, referenced to electrode Fz. After recording, the EEG data were preprocessed with the FieldTrip

toolbox (version 2019-09-20; Oostenveld et al., 2011) for MATLAB (MathWorks, Inc.). The data were rereferenced to the average of all electrodes and filtered between 1 Hz (high-pass) and 100 Hz (low-pass). Following independent component analysis, components corresponding to eye blinks, lateral eye movements, muscle activity, and heartbeat were rejected upon visual inspection. The cleaned data were low-pass filtered at 10 Hz, cut into trials of 4 min starting at the onset of the auditory stimulus and resampled to 125 Hz. For the subsequent analysis, these trials were cut into four consecutive 1-min segments and  $z$ -transformed (Crosse et al., 2016).

### Extraction of Stimulus Features

The experiment was designed to simulate some features of listening with a unilateral CI. We used the onset envelopes of the intact speech signal to estimate the spectrotemporal response function (sTRF, see later). The Neural Systems Laboratory (NSL) toolbox (Chi et al., 2005) for MATLAB was used to compute a spectrally resolved representation of the intact speech stimuli. We extracted an auditory spectrogram consisting of 128 logarithmically spaced frequency bands. To extract the onset envelope in each frequency band, we computed the envelope (magnitude of the Hilbert transform), took the first derivative, and applied half-wave rectification. After downsampling in the time and frequency domain, the envelopes of the auditory stimuli included 16 frequency bands (logarithmically spaced between 0 and 8 kHz) and had the same sampling frequency  $f_s = 125$  Hz as the EEG data. A spectral resolution of 16 frequency bands for the sTRF analysis was used to yield a spectral resolution higher than the number of vocoder bands but still low enough to avoid redundancy.

The major reason for using the onset envelope instead of the conventional (unprocessed) envelope of the speech signal for sTRF modelling was that previous studies (Drennan & Lalor, 2019; Fiedler et al., 2017) found that emphasizing the onsets in the envelope increased the neural encoding response. These observations speak to the view that the brain tracks local changes in amplitude (*edges*) rather than the absolute amplitude of the speech envelope (Ogania & Chang, 2019). Therefore, we used the onset envelope to induce a larger and more robust speech tracking response, which should benefit the statistical power to observe modulatory effects of acoustic detail and attention.

### sTRF Estimation

The sTRF was estimated using the mTRF toolbox (Crosse et al., 2016) for MATLAB using regularized linear regression. First, for every 1-min segment, a

forward (*encoding*) model  $m$  was estimated with  $m = (S^T S + \lambda I)^{-1} S^T R$ .  $S$  is a matrix containing the onset envelopes of the attended and ignored speech stream and their time-lagged versions. We used time lags in the range from  $-150$  to  $+450$  ms. The matrix  $R$  contains the corresponding EEG signal of all 64 electrodes, and  $I$  is the identity matrix. Leave-one-out cross-validation was used to determine the optimal  $\lambda$  parameter, which maximized the Pearson correlation between the predicted and measured neural responses. To allow for an unbiased comparison between the different conditions, the output data of this estimation were  $z$ -scored. Models of attended and ignored speech that belonged to the same experimental trial were jointly estimated. That is, models of to-be-attended intact speech and to-be-ignored vocoded speech were trained together as were the models of to-be-attended vocoded speech and to-be-ignored intact speech. In total, 32 individual models were trained per participant and condition and averaged. For illustration purposes only, the sTRF was also averaged over nine central electrodes (FC1, FCz, FC2, C1, Cz, C2, CP1, CPz, and CP2).

### Neural Encoding Accuracy

Further, the sTRF was used to predict the EEG response to a given speech segment. To this end, using a leave-one-out approach, we convolved the envelopes with the trained sTRF models averaged across all but the current segment. Afterward, we compared each predicted EEG response to the empirical EEG response using Pearson correlation to assess how strongly a speech signal is encoded by the neural response. The resultant  $r$  values were averaged per participant and condition and reflect a continuous, fine-grained measure of neural tracking encoding accuracy (Fiedler et al., 2017, 2019; Fuglsang et al., 2020).

To analyze the temporal unfolding of neural encoding accuracy across sTRF time lags, we predicted the EEG response based on a 48-ms sliding window (24 ms overlap) of the sTRF (Fiedler et al., 2019; Jessen et al., 2019). In essence, we used each 48-ms time window of the sTRF separately to predict the EEG signal. This approach enabled us to test which time windows of the sTRF contribute strongest to EEG prediction. As before, the calculation was carried out using a leave-one-out procedure. The obtained Pearson correlation coefficients were averaged per time window, condition, and individual participants.

Furthermore, for illustration purpose, we analyzed the spectral unfolding of the neural encoding accuracy across the frequency bands of the sTRF. For each frequency band of the sTRF, we used sliding 48-ms time windows to predict the EEG signal. This resulted in

temporal and spectral unfolding of the neural encoding accuracy.

### Statistical Analysis

For behavioral data analysis, we used dependent-samples and independent-samples nonparametric permutation tests on the proportion of correctly answered multiple-choice questions. In detail, a dependent-samples permutation test was used to contrast the proportion of correctly answered content questions about to-be-attended intact versus vocoded speech across all participants ( $N=22$ ). Two independent-samples permutation tests were used to contrast proportion correct scores for to-be-attended intact speech presented on the left ( $n=11$ ) versus right side ( $n=11$ ), and similarly for to-be-attended vocoded speech presented on the left ( $n=11$ ) versus right side ( $n=11$ ). Reported  $p$  values ( $p_{\text{permutation}}$ ) correspond to the proportion of absolute values of 5,000  $t$ -statistics computed on data with permuted condition labels that exceed the absolute empirical  $t$  value for the original (unpermuted) data (Wöstmann et al., 2019). Note that dependent-samples nonparametric permutation tests were furthermore used to contrast neural encoding accuracy derived from the entire sTRF across conditions.

To analyze possible differences between the sTRF in the four conditions of the 2 (acoustic detail: intact vs. vocoded)  $\times$  2 (attention: attended vs. ignored) design, we used a cluster permutation test implemented in the FieldTrip toolbox (Maris & Oostenveld, 2007) for MATLAB. For this statistical analysis, we used the data of all 64 electrodes. First, we calculated two difference-sTRFs for each participant through subtraction of the ignored from the attended sTRF, separately for intact and vocoded speech. The resulting difference-sTRFs [intact (attended–ignored), vocoded (attended–ignored)] were contrasted using multiple dependent-samples  $t$ -tests for all time-frequency-electrode triplets of the sTRF. Resulting  $t$  values were then used to identify clusters by clustering neighboring time-frequency-samples with  $p < .05$  for at least three neighboring electrodes. In each observed cluster, the  $t$  values were summed (reported as  $t$ -sum) over all bins belonging to the cluster (reported as *No. of cluster bins*) and compared with a permutation distribution. The permutation distribution was generated by randomly assigning the sTRF data to conditions, followed by summation of their  $t$  values, through 5,000 iterations. To visually resolve the interaction, we calculated multiple dependent-samples  $t$ -tests for all time-frequency-electrode triplets of the sTRF to contrast the attended with the ignored sTRF (i.e., main effect attention), separately for intact and vocoded speech. Significant clusters for the interaction

(Acoustics  $\times$  Attention) were then overlaid on top of the resulting  $t$ -maps.

For statistical analysis of temporally resolved encoding accuracy, we used cluster permutation tests as well. First, two cluster permutation tests were used to contrast encoding accuracy for attended versus ignored speech, separately for intact and vocoded speech. To test the Acoustics  $\times$  Attention interaction, we used a cluster permutation test to contrast the attended–ignored difference of encoding accuracy for intact versus vocoded speech.

## Results


### Unilateral Degradation Decreases Recall of Attended Speech Content

Participants performed a spatially cued dichotic listening task (Figure 1A). The assignment of vocoded speech to the left or right ear was counterbalanced across participants. On a given trial, participants were cued to attend to the speech signal on the left or right side. At the end of each trial, participants answered four multiple-choice questions about the content of the to-be-attended audiobook.

Individual participants' answers to content questions (Figure 1B) were correct well above chance level (here: proportion correct of 0.25). As expected, the proportion of correct responses to content questions was significantly higher when the to-be-attended audiobook was spectrally intact (mean proportion correct = 0.92;  $SEM=0.001$ ) compared with vocoded (mean proportion correct = 0.82;  $SEM=0.0018$ ;  $p_{\text{permutation}}=0.0002$ ; Cohen's  $d=1.195$ ; for comparable behavioral performance modulation in a small group of unilateral CI users in a pilot experiment, see Supplementary Materials and Supplementary Figure S1). Of note, this effect was very consistent across participants (i.e., present in 19 of 22 participants). The side of speech presentation (left for  $n=11$  and right for  $n=11$ ) did not significantly affect the proportion of correct responses to content questions about intact speech (left vs. right:  $p_{\text{permutation}}=.1722$ ) or vocoded speech (left vs. right:  $p_{\text{permutation}}=.2899$ ).

### Spectrotemporal Responses to Unilateral Intact and Vocoded Speech

In the sTRF that models the neural impulse response to the speech envelope extracted from different frequency bands of the acoustic signal, we observed the typical morphology, including earlier P1, N1, and P2 response components, as well as a later N2 response component (Figure 2). Notably, these response components were strongest for speech envelopes extracted from auditory



**Figure 1.** Experimental Design and Behavioral Results. (A) Design of continuous speech task. On each trial, a visual spatial cue and the text “Listen to the left/right” guided participants’ attention to the left or right side. During the entire experiment, participants listened to intact speech on one ear and to vocoded speech on the other ear (balanced across participants). After each trial, participants had to answer four closed four-choice questions about the content of the to-be-attended audiobook. (B) The 45-degree plot contrasts single-subject proportions of correct answers to content questions for attention to intact versus vocoded speech for participants with vocoded speech on the left side (circles) or right side (crosses). Note that some data points fall on top of each other. Bars and error bars in the inset show mean proportion correct  $\pm$  1 SEM, respectively. \*\*\*  $p_{\text{permutation}} < .001$ .

signals below  $\sim 1000$  Hz and weaker for higher frequencies up to 8000 Hz. Because we had no prior hypotheses regarding effects of the side of vocoding (left vs. right) and because there are no obvious differences between sides of vocoding in Figure 2, we collapsed across these two conditions and focused all subsequent analyses on the factors acoustic detail (intact vs. vocoded) and attention focus (attended vs. ignored).

### Interactive Effects of Unilateral Degradation and Attention on Neural Speech Tracking


Figure 3 shows interactive effects of unilateral vocoding and attention on the sTRF (Panels A and B), as well as on the marginalized temporal response function (TRF; Panels C and D). The corresponding statistical analysis of neural response functions was divided into two steps. First, we analyzed the interactive effect of acoustic detail (intact vs. vocoded)  $\times$  attention focus (attended vs. ignored) on the sTRF (Figure 3A). The interaction was significant in two frontocentral electrode clusters for frequencies below  $\sim 2000$  Hz in time intervals overlapping mainly with the N1 response component (negative cluster with *cluster p* value = .0014, 40–152 ms, 199–1850 Hz; *t*-sum =  $-2,104$ ; No. of cluster bins = 703) and with the N2 response component (positive cluster with *cluster p* value = .0002, 176–328 ms, 199–1850 Hz; *t*-sum = 2,972; No. of cluster bins = 991).

Second, to resolve the significant Acoustic Detail  $\times$  Attention Focus interaction, we analyzed the simple main effect of attention focus on the sTRF, separately for intact and vocoded speech signals (Figure 3B). Interestingly, the earlier interaction cluster was driven by a larger (more negative) N1 component for attended than ignored intact speech, while the same effect was virtually absent for vocoded speech. By contrast, the later interaction cluster was mainly driven by a larger (more negative) N2 component for attended than ignored vocoded speech, while this effect was reduced for intact speech (see also Figure 3C and D).


In addition to interactive effects of vocoding and attention, it is obvious from Figure 3C and D that vocoded speech elicited a generally larger early P1 response component compared with spectrally intact speech (see also Supplementary Figure S2 for a grouping of the TRF according to conditions occurring at the same time in the experiment: attend intact, ignore vocoded speech; attend vocoded, ignore intact speech).

### Acoustic Degradation Delays Attentional Selection of Target Speech

In a final analysis, we focused on a more fine-grained temporal representation of the attentional enhancement of target versus distracting speech that goes beyond individual components of the sTRF. To this end, we applied a sliding-window approach to predict the EEG signal at individual electrodes from the sTRF and speech envelopes (Figure 4). A more positive correlation of the predicted with the measured EEG signal points to a stronger neural representation of the respective speech signal, referred to as higher *neural encoding accuracy*. We asked in which time windows attention would enhance



**Figure 2.** Spectrotemporal Response Functions for All Experimental Conditions. Time-frequency panels show average spectrotemporal response functions (sTRF); solid black lines and shaded areas show the temporal response function (TRF; marginalized across frequencies of the sTRF) and 95% confidence intervals, respectively. The four left-side panels show the response functions for speech materials presented to the left ear split by acoustics and attention condition, the four right-side panels show the same for input to the right ear. Within each participant, neural responses were averaged across nine central electrode positions (see Materials and Methods section for details). Note that A and D show data of  $n = 11$  participants with vocoded speech presented to the left ear and intact speech to the right ear. C and B show data of  $n = 11$  participants with vocoded speech presented to the right ear and intact speech to the left ear.




**Figure 3.** Effects of Acoustics and Attention on the Spectrotemporal Response Function. (A) The time-frequency plot shows  $t$  values for the interaction effect of acoustic detail (intact vs. vocoded)  $\times$  attention (attended vs. ignored) on the spectrotemporal response function, averaged for illustrative purposes across nine central electrodes. White and black outlines indicate a significant negative ( $p = .0014^{**}$ ) and positive cluster ( $p = .0002^{***}$ ), respectively. Topographic maps show average cluster  $t$  values; electrodes belonging to the earlier and later significant cluster are highlighted in white and black, respectively. (B) Time-frequency plots shows  $t$  values for the main effect of attention (attended vs. ignored) for intact speech (left) and vocoded speech (right), overlaid with outlines of significant Acoustics  $\times$  Attention clusters from (A). (C) For illustration purposes, we averaged the sTRF across frequencies, nine central electrodes, and distinct time periods corresponding to individual response components (P1: 16–56 ms, N1: 80–120 ms, P2: 176–224 ms, N2: 312–352 ms). Dots and error bars show the average across participants  $\pm 1$  SEM, respectively. (D) Solid lines and shaded areas show average temporal response functions (TRF; marginalized across frequencies of the sTRF) and 95% confidence intervals.

the encoding accuracy of target versus distracting speech signals in case these were intact versus vocoded.

First, when we used predicted EEG signals derived from the entire sTRF (Figure 4A;  $-150$  to  $+450$  ms), encoding accuracy was significantly increased for attended compared with ignored intact speech ( $p_{\text{permutation}} = .0002$ ) and vocoded speech ( $p_{\text{permutation}} = .0008$ ). However, the Acoustic Detail  $\times$  Attention Focus interaction was not significant (statistical comparison of the attended-ignored difference for intact versus vocoded speech;  $p_{\text{permutation}} = .197$ ) such that acoustic detail (intact vs. vocoded) had no significant effect on encoding accuracy for attended speech ( $p_{\text{permutation}} = .66$ ) or ignored speech ( $p_{\text{permutation}} = .36$ ). In other words, when predictions are based on the entire duration of

the sTRF, attention enhances the encoding accuracy of intact and vocoded target speech to similar extents.

Next, we investigated the temporally resolved dynamics of the attentional enhancement of intact versus vocoded speech (Figure 4B). The attentional enhancement of encoding target versus distracting speech was signified by one cluster for intact speech (48–288 ms; *cluster p* value = .0004; *t*-sum = 1,132; No. of cluster bins = 288) and by two clusters for vocoded speech (first cluster: 96–144 ms; *cluster p* value = .039; *t*-sum = 64; No. of cluster bins = 24; second cluster: 192–408 ms; *cluster p* value = .0002; *t*-sum = 521; No. of cluster bins = 168). The most important finding of this study resulted from the contrast of the attentional enhancement of encoding accuracy for intact versus vocoded



**Figure 4.** Acoustic Degradation Delays Attentional Enhancement of Encoding Accuracy. Encoding accuracy refers to the Pearson correlation of actual EEG data with the EEG data predicted by speech envelope models. (A) Encoding accuracy obtained when the entire sTRF ( $-150$  to  $+450$  ms) was used to predict EEG signals. Circles, dots, and error bars show single-subject, average encoding accuracy  $\pm 1$  SEM, respectively. (B) Encoding accuracy obtained for sliding windows along the sTRF (window length: 48 ms; values on the x axis correspond to centers of time windows). Solid lines and shaded areas indicate mean encoding accuracy and 95% confidence intervals, respectively. Horizontal black lines indicate significant clusters for the difference of encoding accuracy for attended versus ignored speech. Time-frequency spectra on top show encoding accuracy for the effect of attention ( $\Delta$  attention: attend-ignored). (C) To illustrate the attentional enhancement of encoding accuracy, data from (B) were contrasted for attended versus ignored speech ( $\Delta$  attention), separately for intact (red) and vocoded speech (purple). Horizontal black lines indicate significant clusters for the difference in the attentional enhancement for intact versus vocoded speech. Topographic maps show  $t$  values for significant clusters; electrodes of significant clusters are highlighted in black. \*\*\* $p < .001$ ; \*\* $p < .01$ ; \* $p < .05$ ; n.s. = not significant.



speech (Figure 4C): While the attentional enhancement was stronger for intact than vocoded speech in a relatively early cluster (144–192 ms, *cluster p* value = .0054; *t*-sum = 94; No. of cluster bins = 34), the reverse pattern—stronger attentional enhancement for vocoded than intact speech—was observed in a later cluster (312–336 ms; *cluster p* value = .0156; *t*-sum = -68; No. of cluster bins = 23).

It must be noted that cluster-based permutation tests do not establish significance of latencies of the observed clusters (Sassenhagen & Draschkow, 2019). In this sense, it is not warranted to draw inference about when in time the two observed Acoustics  $\times$  Attention effects on encoding accuracy (in Clusters C1 and C2 in Figure 4C) started and ended being significant. However, because the two clusters observed here show significant effects in opposite directions, it can be excluded that the two underlying effects overlap in time (given that the topographical distribution of clusters is very similar). Thus, the attentional enhancement of encoding accuracy for vocoded speech is delayed compared with intact speech (see Supplementary Figure S3 for an additional statistical analysis that confirms the latency difference).

## Discussion

The present study was designed to explore the neural processing constraints on selective attention in challenging listening situations imposed by unilateral spectral degradation, as is common in unilateral CI users. To this end, we studied neural tracking of the envelopes of two continuous, spatially separated speech signals. The most important results can be summarized as follows. First, there was a detrimental effect of unilateral spectral degradation on behavior: Although high overall accuracy (~87%) indicates that participants mastered the listening task quite well, the proportion of correctly answered content questions about to-be-attended speech was about 10% reduced when it was spectrally degraded versus intact. Second, there was an effect of unilateral spectral degradation on neural responses: The preferential neural tracking of attended versus ignored speech was more pronounced for intact speech during earlier neural response components but stronger for degraded speech for later components. These findings suggest that unilateral acoustic degradation impairs mainly the attentional selection of target speech against distraction by delaying it.

### *Effects of Unilateral Acoustic Degradation on Tracking Target Versus Distracting Speech*

Participants' behavioral responses in the present study show a clear benefit of attending to intact compared with spectrally degraded speech on one side.

The reason for this is likely a superposition of two effects. First, target speech comprehension is superior for speech that is spectrally less degraded (e.g., Obleser & Weisz, 2012; Shannon et al., 1995). Second, irrelevant speech distractors that are spectrally degraded interfere less with target speech reception (Westermann & Buchholz, 2017) and memory for target speech (e.g., Ellermeier et al., 2015; Senan et al., 2018; Wöstmann & Obleser, 2016). In this sense, the present study implemented one listening condition that was favorable on two accounts (attend to intact speech, ignore vocoded speech) and another listening condition that was unfavorable on two accounts (attend to vocoded speech, ignore intact speech; Supplementary Figure S2 shows the TRF grouped according to these two listening conditions). Note that these listening conditions correspond to rather extreme cases on the continuum of two-talker situations experienced in case of unilateral vocoding, such as they occur for unilateral CI users.

Although effects of bilateral spectral degradation of speech in multitalker situations on neural envelope tracking have been investigated before (Ding et al., 2013; Kong et al., 2015; Rimmele et al., 2015), less is known about possible effects of unilateral spectral degradation. Here, we show that the typical components of the neural speech tracking response (P1, N1, P2) are present for unilateral intact as well as noise-vocoded target speech but that N1 and P2 were reduced for ignored speech distractors (Figures 2 and 3). The general increase of the early P1 response component for noise-vocoded compared with intact speech (largely irrespective of attention) is most likely due to the fact that noise-vocoded speech is spectrally broader and thus elicits a more pronounced early neural response in auditory regions. This might also be related to the observation that noise-vocoded speech with fewer frequency channels is perceived as being louder (Wöstmann, Lim, et al., 2017).

Interestingly, a late negative N2 component signified the tracking of the attended speech envelope, particularly when it was noise-vocoded. This finding somewhat differs from a previous study that presented listeners with spatially mixed competing speech signals (Fiedler et al., 2019), where we found a larger N2 component in the neural tracking of ignored speech (although with a shorter latency). Despite attempts to assign specific perceptual and cognitive processes to the different components of the neural speech tracking response (for review, see Brodbeck & Simon, 2020), the underlying mechanisms, particularly of the longer-latency components, are still unclear.

Importantly, the effect of unilateral vocoding on the neural tracking of attended versus ignored speech in the present study was best described as a temporal delay, instead of a reduced magnitude of attentional selection:

Early neural tracking responses to intact unilateral speech, mainly during the N1 and P2 components, were larger for attended versus ignored speech. Attending-versus-ignoring vocoded speech increased the magnitude of later neural tracking responses, mainly during the P2 and N2 components. This finding is generally in line with a recent study that found delayed attentional modulation of neural speech tracking responses in bilateral CI users compared with normal-hearing controls (Paul et al., 2020).

What are the possible perceptual and cognitive processes associated with the observed temporally delayed attentional separation of attended versus ignored speech in case of unilateral acoustic degradation? One obvious interpretation is that in case of intact speech, the attentional selection is accomplished early, and largely in parallel with the formation of distinct auditory objects (Shinn-Cunningham, 2008). However, in the case of unilateral acoustic degradation, attentional selection happens later in time, potentially after the formation of competing auditory objects has been accomplished. Of note, neural responses to one speech signal might be significantly affected by the acoustic properties of the masking distractor in the present study. In this sense, an intact speech distractor that is highly intelligible increases the degree of informational masking, which might eventually result in delayed stream segregation (Ezzatian et al., 2015).

The late negative N2 TRF component observed in the present study is somewhat reminiscent of the Nd (negative difference) event-related potential component that refers to a frontocentral scalp negativity in response to attended versus ignored tone stimuli (Hansen & Hillyard, 1980). The Nd component has also been observed in response to auditory probes in the attended versus ignored speech stream in multitalker situations with competing speech signals presented at different spatial locations (Lambrecht et al., 2011; Münte et al., 2010; Nager et al., 2008). Although we can only speculate about the underlying cognitive mechanisms, we consider two processes that might give rise to the delayed attentional selection for vocoded speech in the time interval of the N2 component. First, it might be that perceptual restoration of vocoded target speech is more time consuming such that the attentional separation of target and distractor is delayed in time. Second, it could be that participants use different strategies for the attentional selection of unilateral vocoded versus intact speech. For instance, it might be that the attentional selection of intact target speech exploits mainly the frequency separation of the competing speech signals, whereas the attentional selection of vocoded target speech is based on the additional use of spatial cues (for use of space and frequency cues in speech segregation, see Bonacci et al., 2020).

Of note, results of the present study emphasize the importance of a time-resolved analysis of the neural speech tracking response. In other words, a non-time-resolved analysis on its own (such as the analysis of encoding accuracy in Figure 4A) would have overlooked the temporally delayed attentional separation of unilateral vocoded target versus distractor speech. However, the spectrally resolved analysis of speech tracking responses (sTRF), on the basis of speech envelopes extracted from individual frequency bands of the acoustic signal, did not reveal additional insights in the present study. That is, acoustics- and attention-induced modulations of the sTRF mostly covered a broad frequency range (1–8000 Hz). It might be that the present study, which presented speech materials with envelopes correlated across neighboring frequency bands and recorded the net EEG signal originating from large regions of underlying cortical tissue, lacked the required frequency specificity. Future studies might thus use higher frequency specificity on both, the level of the acoustic stimulus (e.g., by using synthesized stimuli with uncorrelated envelopes across frequency bands) and on the level of the neural response (e.g., through high-density cortical surface recordings in auditory regions; Hullett et al., 2016).

### *What Are the Implications for Listeners With a Unilateral CI?*

Although the present study was designed to simulate some aspects of hearing with a unilateral CI, the results should be transferred to actual unilateral CI user with caution. We presented normal-hearing listeners with unilateral vocoded versus intact speech without prior adaptation or speech comprehension training of spectrally degraded materials. It is thus conceivable that the observed delayed attentional selection of vocoded target speech against distraction is subject to change such that its latency decreases in the course of rehabilitation of CI users (for a recent review on auditory cortical plasticity in CI users, see Glennon et al., 2020). Furthermore, the present study simulated hearing with a CI on one side with normal hearing on the contralateral side. While such a scenario mimics hearing in people with single-sided deafness after CI implantation, a significant amount of unilateral CI users suffers from hearing loss also on the contralateral side, which is treated with a hearing aid in case of bimodal listeners (for comparison of spatial speech-in-noise performance of single-sided deafness and bimodal listeners, see, e.g., Williges et al., 2019).

Although dichotic listening is a useful experimental paradigm that achieves maximal spatial separation of competing sound sources, it must be noted that listening situations with a target signal present at 90° to one side under distraction from 90° to the other side are rare.

Furthermore, the dichotic sound presentation over headphones in the present study did not allow for a partial mixture of the two competing signals before entering either ear, which would happen in real-life listening situations with free field sound presentation. However, a pilot study with free field presentation of competing sound sources (see Supplementary Materials and Supplementary Figure S1) revealed that unilateral vocoding in normal-hearing listeners and listening with a unilateral CI in a small group of CI users produced comparable speech comprehension detriments resulting from spectral degradation. Thus, it is conceivable that the present experimental paradigm simulates some challenges faced when listening with a unilateral CI. Nevertheless, to achieve higher ecological validity, future studies should use free field presentation and locate one sound source in the front and another one on the left or right side, as we have done recently in normal-hearing listeners (Wöstmann et al., 2019). In theory, smaller spatial separation of sound sources should challenge the attentional separation of competing speech signals. Whether systematic variation in spatial separation of sound sources interacts with effects of spectral degradation on speech tracking is a question for future research.

Mechanistically, it is conceivable that the delayed attentional separation of unilateral vocoded target versus distractor speech constitutes a reason for compromised speech-in-noise comprehension in CI users. An ensuing question to be answered in future studies that implement single-trial speech comprehension measures is whether the temporal delay of attentional selection in the speech tracking response relates to target speech comprehension. Alternatively, the delayed attentional separation might signify larger listening effort, which has been shown to increase with decreasing spectral resolution of speech (e.g., Pals et al., 2013).

## Conclusion

The present study presents evidence that unilateral acoustic degradation does not preclude the attention-induced neural selection of target speech from one side against concurrent distraction from the other. Instead, unilateral acoustic degradation delays this neural selection process. These findings improve our understanding of the neurophysiological basis of selective auditory attention to ongoing speech. Furthermore, we here identify a candidate neural signature of aggravated speech-in-noise comprehension under spectral degradation of the acoustic input, which has a potency in better explaining some of the listening challenges experienced by unilateral CI users.

## Acknowledgments

The authors thank Sebastian Puschmann and Martin Orf for their help with the mTRF toolbox.


## Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The authors acknowledge the financial support of Cochlear Inc. (Grant to J. O. and M. W.), the European Research Council (Grant ERC-CoG-2014-646696 to J. O.), and by Land Schleswig-Holstein within the funding programme Open Access Publikationsfonds.

## ORCID iD

Malte Wöstmann  <https://orcid.org/0000-0001-8612-8205>

## Supplemental Material

Supplemental material for this article is available online.

## References

- Bonacci, L. M., Bressi, S., & Shinn-Cunningham, B. (2020). Non-spatial features reduce the reliance on sustained spatial auditory attention. *Ear and Hearing, 41*(6), 1635–1647. <https://doi.org/10.1097/AUD.0000000000000879>
- Brodbeck, C., Jiao, A., Hong, L. E., & Simon, J. Z. (2020). Neural speech restoration at the cocktail party: Auditory cortex recovers masked speech of both attended and ignored speakers. *PLoS Biology, 18*. <https://doi.org/doi.org/10.1371/journal.pbio.3000883>
- Brodbeck, C., & Simon, J. Z. (2020). Continuous speech processing. *Current Opinion in Physiology, 18*, 25–31. <https://doi.org/doi.org/10.1016/j.cophys.2020.07.014>
- Brungart, D. S., Simpson, B. D., Ericson, M. A., & Scott, K. R. (2001). Informational and energetic masking effects in the perception of multiple simultaneous talkers. *Journal of the Acoustical Society of America, 110*(5 Pt 1), 2527–2538. doi: 10.1121/1.1408946
- Chi, T., Ru, P. W., & Shamma, S. A. (2005). Multiresolution spectrotemporal analysis of complex sounds. *Journal of the Acoustical Society of America, 118*(2), 887–906. <https://doi.org/doi.org/10.1121/1.1945807>
- Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The Multivariate Temporal Response Function (mTRF) toolbox: A MATLAB toolbox for relating neural signals to continuous stimuli. *Frontiers in Human Neuroscience, 10*, 604. <https://doi.org/10.3389/fnhum.2016.00604>
- Decruy, L., Vanthornhout, J., & Francart, T. (2020). Hearing impairment is associated with enhanced neural tracking of the speech envelope. *Hearing Research, 393*, 1–13. <https://doi.org/ARTN 107961>

- Ding, N., Chatterjee, M., & Simon, J. Z. (2013). Robust cortical entrainment to the speech envelope relies on the spectro-temporal fine structure. *NeuroImage*, *88C*, 41–46. <https://doi.org/10.1016/j.neuroimage.2013.10.054>
- Ding, N., & Simon, J. Z. (2014). Cortical entrainment to continuous speech: Functional roles and interpretations. *Frontiers in Human Neuroscience*, *8*, 311. <https://doi.org/10.3389/fnhum.2014.00311>
- Drennan, D. P., & Lalor, E. C. (2019). Cortical tracking of complex sound envelopes: Modeling the changes in response with intensity. *eNeuro*, *6*(3), 1–11. <https://doi.org/Artn Eneuro.0082-19.2019>
- Ellermeier, W., Kattner, F., Ueda, K., Doumoto, K., & Nakajima, Y. (2015). Memory disruption by irrelevant noise-vocoded speech: Effects of native language and the number of frequency bands. *Journal of the Acoustical Society of America*, *138*(3), 1561–1569. <https://doi.org/10.1121/1.4928954>
- Etard, O., & Reichenbach, T. (2019). Neural speech tracking in the theta and in the delta frequency band differentially encode clarity and comprehension of speech in noise. *Journal of Neuroscience*, *39*(29), 5750–5759. <https://doi.org/10.1523/Jneurosci.1828-18.2019>
- Ezzatian, P., Li, L., Pichora-Fuller, K., & Schneider, B. A. (2015). Delayed stream segregation in older adults: More than just informational masking. *Ear and Hearing*, *36*(4), 482–484. <https://doi.org/10.1097/Aud.000000000000139>
- Fiedler, L., Wöstmann, M., Graversen, C., Brandmeyer, A., Lunner, T., & Obleser, J. (2017). Single-channel in-ear-EEG detects the focus of auditory attention to concurrent tone streams and mixed speech. *Journal of Neural Engineering*, *14*(3), 036020. <https://doi.org/10.1088/1741-2552/aa66dd>
- Fiedler, L., Wöstmann, M., Herbst, S. K., & Obleser, J. (2019). Late cortical tracking of ignored speech facilitates neural selectivity in acoustically challenging conditions. *NeuroImage*, *186*, 33–42. <https://doi.org/10.1016/j.neuroimage.2018.10.057>
- Fuglsang, S. A., Marcher-Rorsted, J., Dau, T., & Hjortkjaer, J. (2020). Effects of sensorineural hearing loss on cortical synchronization to competing speech during selective attention. *Journal of Neuroscience*, *40*(12), 2562–2572. <https://doi.org/10.1523/Jneurosci.1936-19.2020>
- Glennon, E., Svirsky, M. A., & Froemke, R. C. (2020). Auditory cortical plasticity in cochlear implant users. *Current Opinion in Neurobiology*, *60*, 108–114. <https://doi.org/10.1016/j.conb.2019.11.003>
- Golub, J. S., Lin, F. R., Lustig, L. R., & Lalwani, A. K. (2018). Prevalence of adult unilateral hearing loss and hearing aid use in the United States. *The Laryngoscope*, *128*(7), 1681–1686. <https://doi.org/10.1002/lary.27017>
- Hansen, J. C., & Hillyard, S. A. (1980). Endogenous brain potentials associated with selective auditory attention. *Electroencephalography and Clinical Neurophysiology*, *49*(3–4), 277–290. [https://doi.org/10.1016/0013-4694\(80\)90222-9](https://doi.org/10.1016/0013-4694(80)90222-9)
- Horton, C., D’Zmura, M., & Srinivasan, R. (2013). Suppression of competing speech through entrainment of cortical oscillations. *Journal of Neurophysiology*, *109*(12), 3082–3093. <https://doi.org/10.1152/jn.01026.2012>
- Hullett, P. W., Hamilton, L. S., Mesgarani, N., Schreiner, C. E., & Chang, E. F. (2016). Human superior temporal gyrus organization of spectrotemporal modulation tuning derived from speech stimuli. *Journal of Neuroscience*, *36*(6), 2014–2026. <https://doi.org/10.1523/Jneurosci.1779-15.2016>
- Jessen, S., Fiedler, L., Munte, T. F., & Obleser, J. (2019). Quantifying the individual auditory and visual brain response in 7-month-old infants watching a brief cartoon movie. *NeuroImage*, *202*, 116060. <https://doi.org/10.1016/j.neuroimage.2019.116060>
- Kong, Y. Y., Somarowthu, A., & Ding, N. (2015). Effects of spectral degradation on attentional modulation of cortical auditory responses to continuous speech. *Journal of the Association for Research in Otolaryngology*, *16*(6), 783–796. <https://doi.org/10.1007/s10162-015-0540-x>
- Lambrecht, J., Spring, D. K., & Münte, T. F. (2011). The focus of attention at the virtual cocktail party—electrophysiological evidence. *Neuroscience Letters*, *489*(1), 53–56. <https://doi.org/10.1016/j.neulet.2010.11.066>
- Litovsky, R. Y., Moua, K., Godar, S., Kan, A., Misurelli, S. M., & Lee, D. J. (2019). Restoration of spatial hearing in adult cochlear implant users with single-sided deafness. *Hearing Research*, *372*, 69–79. <https://doi.org/10.1016/j.heares.2018.04.004>
- Ma, N., Morris, S., & Kitterick, P. T. (2016). Benefits to speech perception in noise from the binaural integration of electric and acoustic signals in simulated unilateral deafness. *Ear and Hearing*, *37*(3), 248–259. <https://doi.org/10.1097/Aud.0000000000000252>
- Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. <https://doi.org/10.1016/j.jneumeth.2007.03.024>
- Mattys, S. L., Brooks, J., & Cooke, M. (2009). Recognizing speech under a processing load: Dissociating energetic from informational factors. *Cognitive Psychology*, *59*(3), 203–243. <https://doi.org/10.1016/j.cogpsych.2009.04.001>
- Moore, B. C. (2008). The role of temporal fine structure processing in pitch perception, masking, and speech perception for normal-hearing and hearing-impaired people. *Journal of the Association for Research in Otolaryngology*, *9*(4), 399–406. <https://doi.org/10.1007/s10162-008-0143-x>
- Münte, T. F., Spring, D. K., Szyck, G. R., & Noesselt, T. (2010). Electrophysiological attention effects in a virtual cocktail-party setting. *Brain Research*, *1307*, 78–88. <https://doi.org/10.1016/j.brainres.2009.10.044>
- Nager, W., Dethlefsen, C., & Münte, T. F. (2008). Attention to human speakers in a virtual auditory environment: Brain potential evidence. *Brain Research*, *1220*, 164–170. <https://doi.org/10.1016/j.brainres.2008.02.058>
- Obleser, J., & Kayser, C. (2019). Neural entrainment and attentional selection in the listening brain. *Trends in Cognitive Sciences*, *23*(11), 913–926. <https://doi.org/10.1016/j.tics.2019.08.004>
- Obleser, J., & Weisz, N. (2012). Suppressed alpha oscillations predict intelligibility of speech and its acoustic details. *Cerebral Cortex*, *22*(11), 2466–2477. <https://doi.org/10.1093/cercor/bhr325>

- Oganian, Y., & Chang, E. F. (2019). A speech envelope landmark for syllable encoding in human superior temporal gyrus. *Science Advances*, 5(11), 1–13. <https://doi.org/ARTN eaay6279>
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011, 156869. <https://doi.org/10.1155/2011/156869>
- O’Sullivan, J. A., Power, A. J., Mesgarani, N., Rajaram, S., Foxe, J. J., Shinn-Cunningham, B. G., Slaney, M., Shamma, S. A., & Lalor, E. C. (2015). Attentional selection in a cocktail party environment can be decoded from single-trial EEG. *Cerebral Cortex*, 25(7), 1697–1706. <https://doi.org/10.1093/cercor/bht355>
- Pals, C., Sarampalis, A., & Baskent, D. (2013). Listening effort with cochlear implant simulations. *Journal of Speech Language and Hearing Research*, 56(4), 1075–1084. [https://doi.org/10.1044/1092-4388\(2012\)12-0074](https://doi.org/10.1044/1092-4388(2012)12-0074)
- Paul, B. T., Uzelac, M., Chan, E., & Dimitrijevic, A. (2020). Poor early cortical differentiation of speech predicts perceptual difficulties of severely hearing-impaired listeners in multi-talker environments. *Scientific Reports*, 10(1), 1–12. <https://doi.org/ARTN 6141>
- Peelle, J. E., & Davis, M. H. (2012). Neural oscillations carry speech rhythm through to comprehension. *Frontiers in Psychology*, 3, 320. <https://doi.org/10.3389/fpsyg.2012.00320>
- Peelle, J. E., Gross, J., & Davis, M. H. (2013). Phase-locked responses to speech in human auditory cortex are enhanced during comprehension. *Cerebral Cortex*, 23(6), 1378–1387. <https://doi.org/10.1093/cercor/bhs118>
- Petersen, E. B., Wöstmann, M., Obleser, J., & Lunner, T. (2017). Neural tracking of attended versus ignored speech is differentially affected by hearing loss. *Journal of Neurophysiology*, 117(1), 18–27. <https://doi.org/10.1152/jn.00527.2016>
- Rimmele, J. M., Zion Golumbic, E., Schroger, E., & Poeppel, D. (2015). The effects of selective attention and speech acoustics on neural speech-tracking in a multi-talker scene. *Cortex*, 68, 144–154. <https://doi.org/10.1016/j.cortex.2014.12.014>
- Rosen, S., Faulkner, A., & Wilkinson, L. (1999). Adaptation by normal listeners to upward spectral shifts of speech: Implications for cochlear implants. *Journal of the Acoustical Society of America*, 106(6), 3629–3636.
- Rosen, S., Souza, P., Ekelund, C., & Majeed, A. A. (2013). Listening to speech in a background of other talkers: Effects of talker number and noise vocoding. *Journal of the Acoustical Society of America*, 133(4), 2431–2443. <https://doi.org/10.1121/1.4794379>
- Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, 56(6), 1–8. <https://doi.org/ARTN e13335>
- Senan, T. U., Jelfs, S., & Kohlrausch, A. (2018). Cognitive disruption by noise-vocoded speech stimuli: Effects of spectral variation. *Journal of the Acoustical Society of America*, 143(3), 1407–1416. <https://doi.org/10.1121/1.5026619>
- Shannon, R. V., Zeng, F. G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition with primarily temporal cues. *Science*, 270(5234), 303–304.
- Shinn-Cunningham, B., Best, V., & Lee, A. K. C. (2017). Auditory object formation and selection. *Auditory System at the Cocktail Party*, 60, 7–40. [https://doi.org/10.1007/978-3-319-51662-2\\_2](https://doi.org/10.1007/978-3-319-51662-2_2)
- Shinn-Cunningham, B. G. (2008). Object-based auditory and visual attention. *Trends in Cognitive Sciences*, 12(5), 182–186. <https://doi.org/10.1016/j.tics.2008.02.003>
- Slaney, M. (1993). *An efficient implementation of the Patterson-Holdsworth auditory filter bank*. I. Apple Computer.
- Vlastarakos, P. V., Nazos, K., Tavoulari, E. F., & Nikolopoulos, T. P. (2014). Cochlear implantation for single-sided deafness: The outcomes. An evidence-based approach. *European Archives of Oto-Rhino-Laryngology*, 271(8), 2119–2126. <https://doi.org/10.1007/s00405-013-2746-z>
- Westermann, A., & Buchholz, J. M. (2017). The effect of nearby maskers on speech intelligibility in reverberant, multi-talker environments. *Journal of the Acoustical Society of America*, 141(3), 2214–2223. <https://doi.org/10.1121/1.4979000>
- Williges, B., Wesarg, T., Jung, L., Geven, L. I., Radeloff, A., & Jürgens, T. (2019). Spatial speech-in-noise performance in bimodal and single-sided deaf cochlear implant users. *Trends in Hearing*, 23, 1–16. <https://doi.org/ArtN 2331216519858311>
- Wilson, P. S., Fleming, D. M., & Donaldson, I. (1993). Prevalence of hearing-loss among people aged 65 years and over - Screening and hearing-aid provision. *British Journal of General Practice*, 43(375), 406–409.
- Wöstmann, M., Alavash, M., & Obleser, J. (2019). Alpha oscillations in the human brain implement distractor suppression independent of target selection. *Journal of Neuroscience*, 39(49), 9797–9805. <https://doi.org/10.1523/JNEUROSCI.1954-19.2019>
- Wöstmann, M., Fiedler, L., & Obleser, J. (2017). Tracking the signal, cracking the code: Speech and speech comprehension in non-invasive human electrophysiology. *Language, Cognition and Neuroscience*, 32(7), 1–15. <https://doi.org/10.1080/23273798.2016.1262051>
- Wöstmann, M., Lim, S. J., & Obleser, J. (2017). The human neural alpha response to speech is a proxy of attentional control. *Cerebral Cortex*, 27(6), 3307–3317. <https://doi.org/10.1093/cercor/bhx074>
- Wöstmann, M., & Obleser, J. (2016). Acoustic detail but not predictability of task-irrelevant speech disrupts working memory. *Frontiers in Human Neuroscience*, 10(538), 1–9. <https://doi.org/10.3389/fnhum.2016.00538>
- Zion Golumbic, E. M., Ding, N., Bickel, S., Lakatos, P., Schevon, C. A., McKhann, G. M., Goodman, R. R., Emerson, R., Mehta, A. D., Simon, J. Z., Poeppel, D., & Schroeder, C. E. (2013). Mechanisms underlying selective neuronal tracking of attended speech at a “cocktail party”. *Neuron*, 77(5), 980–991. <https://doi.org/10.1016/j.neuron.2012.12.037>