

RESEARCH ARTICLE

Individual- and country-level correlates of female permanent contraception use in sub-Saharan Africa

Babayemi O. Olakunde^{1,2*}, Jennifer R. Pharr¹, Lung-Chang Chien³, Rebecca D. Benfield⁴, Francisco S. Sy¹

1 Department of Environmental and Occupational Health, School of Public Health, University of Nevada, Las Vegas, NV, United States of America, **2** Department of Community Prevention and Care Services, National Agency for the Control of AIDS, Abuja, Nigeria, **3** Department of Epidemiology and Biostatistics, School of Public Health, University of Nevada, Las Vegas, NV, United States of America, **4** School of Nursing, University of Nevada, Las Vegas, NV, United States of America

* olakunde@unlv.nevada.edu



Abstract

Background

Female permanent contraception is a cost-effective contraceptive method that can help clients with the desire to limit childbearing achieve their reproductive intention. However, despite its benefits, the use of FPC remains low in sub-Saharan Africa (SSA), and limited studies have examined the correlates of its uptake. In this study, we assessed the individual- and country-level factors associated with the use of FPC among married or in-union women using modern contraceptive methods to limit childbearing in SSA.

Methods

This study was a secondary data analysis of individual- and country-level data obtained from the Demographic and Health Surveys (DHS) Program and three open data repositories. The study included 29,777 married or in-union women aged 15–49 years using modern contraceptive methods to limit childbearing from DHS conducted in 33 sub-Saharan African countries between 2010 and 2018. We performed descriptive statistics and fitted multilevel logistic regression models to determine the predisposing, enabling, and need factors associated with the use of FPC.

Results

Approximately 13% of the women used FPC. About 20% of the variance in the odds of using FPC was attributable to between-country differences. In the full model, the significant individual-level factors associated with the use of FPC compared with other modern contraceptive methods were: age (odds ratio [OR] = 1.10; 95%CI = 1.08–1.12), living children (OR = 1.11, 95%CI = 1.04–1.16), high household wealth (OR = 1.39, 95%CI = 1.18–1.64), rural residence (OR = 0.83, 95% CI = 0.71–0.97), joint contraceptive decision with partner (OR = 1.68, 95% = 1.43–1.99), contraceptive decision by partner and others (OR = 2.46, 95% =

OPEN ACCESS

Citation: Olakunde BO, Pharr JR, Chien L-C, Benfield RD, Sy FS (2020) Individual- and country-level correlates of female permanent contraception use in sub-Saharan Africa. *PLoS ONE* 15(12): e0243316. <https://doi.org/10.1371/journal.pone.0243316>

Editor: Mohammad Rifat Haider, Ohio University College of Health Sciences and Professions, UNITED STATES

Received: August 20, 2020

Accepted: November 18, 2020

Published: December 15, 2020

Copyright: © 2020 Olakunde et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data used in the study are publicly available from the following: Demographic Health Survey (<https://dhsprogram.com/>); World Bank (<https://www.worldbank.org/>); World Health Organization (<https://www.who.int/>); and United Nations Educational, Scientific and Cultural Organization (<https://en.unesco.org/>).

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

1.97–3.07), and the number of living children less than the ideal number of children (OR = 1.40, 95%CI = 1.21–1.62). The significantly associated country-level factors were births attended by skilled health providers (OR = 1.03, 95%CI = 1.00–1.05) and density of medical doctors (OR = 1.37, 95%CI = 1.01–1.85).

Conclusions

Our results suggest that both individual- and country-level factors affect uptake of FPC in SSA. Increasing geographic, economic, and psychosocial access to FPC may improve its uptake in SSA.

Background

Despite the increasing proportion of women with demand to limit childbearing in sub-Saharan Africa (SSA) [1, 2], the uptake of female permanent contraception (FPC) which offers a cost-effective, convenient, and safe method to limit births, remains low [3, 4]. The majority of these women do not use contraceptives or rely on reversible methods that are not as effective as FPC [4], thereby increasing their risk of having unintended pregnancies [5, 6]. Estimates indicate that approximately one in three pregnancies in SSA is unintended [7]. Unintended pregnancies are responsible for nearly 30% of maternal deaths and 26% of newborn deaths in SSA [7]. They also have negative social and economic implications [8]. Higher proportion of unintended pregnancies have been found among women who have a desire to limit compared to space childbirths [9].

Globally, FPC is the most widely used contraceptive method [10, 11], representing 24% of the method mix among women using contraceptives [11]. It is the most commonly used method in regions like Central and Southern Asia, and Latin America and the Caribbean [10]. Compared to other regions, SSA has the lowest uptake of FPC in the world, where it accounts for less than 3.9% of the method mix [11].

While women face multilevel barriers in accessing modern contraceptive methods in SSA, how these factors affect uptake of FPC have not been comprehensively examined [12]. A number of quantitative studies that have assessed the use of FPC in SSA conflated FPC with other long-acting reversible methods. Thus, findings reported in these available studies are not specific to FPC. Furthermore, a majority of the studies considered only individual-level factors, leaving out contextual factors which may also affect the uptake of FPC [13, 14].

A better understanding of the factors that influence the uptake of FPC is needed to improve its voluntary and informed utilization among clients who want to limit childbearing in SSA [15]. Accordingly, this study aimed to add to the sparse body of evidence on the use of FPC in SSA by providing insight into its correlates. The objective was to assess the individual- and country-level factors associated with the use of FPC among married or in-union women using modern contraceptive methods to limit childbearing in SSA.

Methods

Conceptual framework

This study drew upon the supply-demand framework for the determinants of fertility and contraceptive use [16, 17] and the behavioral model of health services use [18, 19] to assess the correlates of the use of FPC among married and in-union women in SSA. According to the




Fig 1. Conceptual framework.

<https://doi.org/10.1371/journal.pone.0243316.g001>

supply-demand framework for the determinants of fertility and contraceptive use, the motivation or incentive for fertility regulation is influenced by both the demand for and supply of children [16, 17]. If supply is lower than demand, there would not be a desire to limit fertility and vice versa [16, 17]. However, the framework holds that the use of fertility control when there is excess supply would depend on the costs (subjective and objective) of obtaining and using a contraceptive method relative to the motivation to limit fertility [16, 17]. The behavioral model of health services use (also known as the Andersen model) posits that “people’s use of health services is a function of their predisposition to use services, factors which enable or impede use, and their need for care” [19, p1]. Recognizing the significance of community, structure, and process of service delivery, Andersen later added contextual characteristics as important determinants of health behaviors in the revised version of the model [20]. Andersen noted that, “understanding of utilization of health services can be best achieved by focusing on contextual and individual determinants” [20, p652]. The components of contextual characteristics are similar to individual characteristics (predisposing, enabling, and need factors) [20], but they are measured at an aggregate level (e.g., community or country). Contextual factors can affect individual factors, which will in turn influence health behaviors and outcomes, or they can directly influence health behaviors and outcomes. The addition of contextual characteristics to the model has been described as one of its strengths, and makes it appropriate for multilevel models to assess the determinants of health care utilization [21]. The predisposing, enabling, and need variables considered in this study are factors that have been empirically found to influence the use of contraception or are theoretically plausible (Fig 1).

Study design and sample

This study was a secondary data analysis of data obtained from the Demographic and Health Surveys (DHS) Program and three open data repositories. The study included 29,777 married

or in-union women of reproductive age (15–49 years) who were using modern contraceptive methods (i.e., condom, diaphragm, emergency contraception, female condom, female permanent contraception, foam and jelly, implants, intrauterine device, injections, lactational amenorrhea, male permanent contraception, pill, or standard day method) to limit childbearing.

Data sources

The Individual-level data were obtained from the DHS Program (<https://dhsprogram.com/>). DHS are nationally-representative household surveys that gather data on a number of health-related topics. The DHS program utilizes standardized methodologies and procedures, making the surveys comparable across different countries. The DHS program adopts a stratified two-stage probabilistic sampling design. More details about the survey procedure can be found elsewhere [22]. SSA countries with a standard DHS conducted in the last 10 years (between 2010 and 2019) were considered for inclusion in this study. Based on these inclusion criteria, 33 countries were included, with the years of the surveys ranging from 2010 to 2018 (Table 1). The country-level data were publicly available data obtained from three open data repositories: the World Bank (<https://www.worldbank.org/>), World Health Organization (WHO) (<https://www.who.int/>), and United Nations Educational, Scientific and Cultural Organization (UNESCO) (<https://en.unesco.org/>). We used the most recent available data corresponding or closest to the DHS survey year for each of the countries.

Measures

Dependent and explanatory variables. A binary dependent variable was created from married or in-union women (15–49 years) using modern contraceptive methods to limit childbearing. Married or in-union women using FPC were coded '1', while those using other modern contraceptive methods were coded '0'. The explanatory variables were categorized into two levels: individual and country (Fig 1). Under each level, the variables were grouped into predisposing, enabling, and need. See S1 Table for operational definitions of the variables.

Data analysis

Crude and weighted descriptive statistics were performed to summarize the data. For the weighted percentages, we de-normalized the weights for each of the countries [23], using the population of women of reproductive age (15–49 years) (<https://population.un.org/wpp/Download/Standard/Population/>) corresponding to the year of survey. We performed a 2-level multilevel logistic regression analysis, with individual-level factors at level one and country-level factors at level two. Four random intercept models were fitted as follows: Model 1 (an empty model with no variable); Model 2: (individual-level variables with survey year as a control variable); Model 3 (country-level variables with survey year as a control variable); and Model 4 (individual- and country-level variables with survey year as a control variable). Multicollinearity among the explanatory variables was examined using the variance inflation factor, with a value exceeding 10 used as the cut-off [24]. Multicollinearity was not detected among explanatory variables. Fixed effects (measures of association) were reported as odds ratios (ORs), with their 95% confidence intervals (CIs). Random effects (measures of variation) were measured by the intra-cluster (i.e. intra-country) correlation coefficient (ICC), proportional change in variance (PCV), and median odds ratio (MOR) [25–28]. Responses such as 'don't know' and nonnumeric responses to questions that required numeric answers were treated as missing data. The analysis was conducted using listwise deletion (complete-case analysis). P-value of less than 0.05 was considered statistically significant. The data analysis was performed using Stata Statistical Software: Release 15, College Station, TX, StataCorp LLC.

Table 1. Summary of included DHS.

S/ N	Country	Survey Year	Married or in-union women using modern contraceptive methods to limit childbearing
1	Angola	2015–16	243
2	Benin	2017–18	532
3	Burkina Faso	2010	694
4	Burundi	2016–17	1073
5	Cameroun	2011	498
6	Chad	2014–15	131
7	Comoros	2012	129
8	Congo	2011–12	211
9	Cote d'Ivoire	2011–12	209
10	Democratic Republic of Congo	2013–14	308
11	Ethiopia	2016	1073
12	Gabon	2012	196
13	Gambia	2013	158
14	Ghana	2014	516
15	Guinea	2018	173
16	Kenya	2014	2418
17	Lesotho	2014	1332
18	Liberia	2013	401
19	Malawi	2015–16	5010
20	Mali	2018	335
21	Mozambique	2011	581
22	Namibia	2013	1142
23	Niger	2012	162
24	Nigeria	2018	1471
25	Rwanda	2014–15	1656
26	Senegal	2017	794
27	Sierra Leone	2013	592
28	South Africa	2016	1058
29	Tanzania	2015–16	911
30	Togo	2013–14	471
31	Ugandan	2016	1749
32	Zambia	2013–14	1686
33	Zimbabwe	2015	1864

<https://doi.org/10.1371/journal.pone.0243316.t001>

Ethical consideration

This study was a secondary analysis of publicly available data. All the data used were fully anonymized. The DHS program obtains ethical clearance from appropriate National Ethics Committees in the respective countries before conducting the surveys. Access and permission to use the datasets was granted by ICF. This study was deemed exempt by the University of Nevada, Las Vegas Institutional Review Board.

Results

Descriptive statistics

The individual-level characteristics are summarized in [Table 2](#). The mean age was 36.4 years. About 38% of the respondents had at least secondary education. Approximately 51% of the

Table 2. Descriptive statistics of individual-level characteristics.

Characteristics	Total		FPC		Other modern methods	
	N (%)	Weighted %	N (%)	Weighted %	N (%)	Weighted %
All	29777 (100)	100	4022 (13.5)	12.7	25755 (86.5)	87.3
Age (N = 29777); Mean (SD) ^a	36.3 (6.8)	36.4 (6.8)	40.3 (5.6)	40.6 (5.5)	35.8 (6.8)	35.8 (6.7)
Education (N = 29777)						
None	6539 (22.0)	24.6	829 (12.7)	9.8	5710 (87.3)	90.2
Primary	13419 (45.1)	37.0	2069 (11.4)	14.0	11350 (84.6)	86.0
Secondary or higher	9818 (33.0)	38.3	1124 (13.5)	13.2	8694 (88.6)	86.8
Household wealth (N = 29777)						
Poor	9321 (31.3)	28.8	1083 (11.6)	11.1	8238 (88.4)	88.9
Middle	5926 (19.9)	20.0	774 (13.1)	11.7	5152 (86.9)	88.3
Rich	14530 (48.8)	51.2	2165 (14.9)	13.9	12365 (85.1)	86.1
Area of residence (N = 29777)						
Urban	11471 (38.5)	42.7	1386 (12.1)	12.9	10085 (87.9)	87.1
Rural	18306 (61.5)	57.3	2636 (14.4)	12.5	15670 (85.6)	87.5
Living children (N = 29777); Mean (SD)	4.5 (1.9)	4.4 (1.9)	4.9 (2.0)	4.7 (2.1)	4.4 (1.9)	4.4 (1.9)
Ideal and living children (N = 28870)						
Living equal or greater than ideal	19748 (68.4)	66.9	2517 (12.7)	12.2	17231 (87.3)	87.8
Living less than ideal	9122 (31.6)	33.1	1321 (14.5)	13.7	7801 (85.5)	86.3
Number of living sons (N = 29777); Mean (SD)	2.3 (1.4)	2.2 (1.4)	2.5 (1.5)	2.4 (1.5)	2.2 (1.4)	2.2 (1.4)
Media exposure (N = 29771)						
Yes	16363 (55.0)	53.9	2311 (14.1)	13.6	14052 (85.9)	86.4
No	13408 (45.0)	46.1	1711 (12.8)	11.5	11697 (87.2)	88.5
Decision maker (N = 29651)						
Joint decision	19139 (64.5)	62.5	2717 (14.2)	13.3	16422 (85.8)	86.7
Mainly respondent	7570 (25.5)	28.2	688 (9.1)	9.0	6882 (90.9)	91.0
Mainly husband/partner or Others	2942 (9.9)	9.2	590 (20.1)	18.9	2352 (79.9)	81.1
Husband/Partner's age (N = 29653); Mean (SD)	43.2 (9.5)	43.5 (9.4)	46.8 (8.5)	47.1 (8.7)	42.6 (9.5)	43.0 (9.4)
Husband/Partner's education (N = 29212)						
None	5023 (17.2)	17.0	520 (10.4)	8.2	4503 (89.6)	91.8
Primary	11655 (39.9)	36.5	1908 (16.4)	14.5	9747 (83.6)	85.5
Secondary or higher	12534 (42.9)	36.5	1539 (12.3)	12.9	10995 (87.7)	87.1
Union (N = 29047)						
Monogynous	24074 (82.9)	84.9	3279 (13.6)	12.7	20795 (86.4)	87.3
Polygynous	4973 (17.1)	15.1	655 (13.2)	12.9	4319 (86.8)	87.1
Wantedness (N = 18798)						
Wanted then	10712 (57.0)	58.9	738 (6.9)	6.9	9974 (93.1)	93.1
Wanted later	3975 (21.1)	19.2	250 (6.3)	6.1	3725 (93.7)	93.9
Wanted no more	4111 (21.9)	21.9	454 (11.0)	10.2	3657 (89.0)	89.8

^a SD = Standard deviation.

<https://doi.org/10.1371/journal.pone.0243316.t002>

respondents were from rich households and 57% resided in a rural area. Of the 29,777 married or in-union women using modern contraceptive methods to limit childbearing, 12.7% used FPC (Table 2). The use of FPC varied by individual-level characteristics. It was higher than 12.7% among women who: were ≥ 40 years, had at least a primary education, were from rich households, were urban dwellers, had ≥ 3 living children, whose living children were less than ideal their number of children, had ≥ 3 living sons, were exposed to family planning messages,

Table 3. Descriptive statistics of country-level characteristics.

Characteristics	Mean (SD) ^a	Range
Poverty rate (%)	40.5 (19.0)	3.4–76.6
Literacy rate (%)	53.6 (23.0)	14–88
Births attended by skilled health providers (%)	66.2 (19.3)	20–97
Density of medical doctors (per 10,000 population)	1.4 (1.5)	0.2–8.0
Gross national income (USD)	1617 (1919)	270–9080
Rural population (%)	60.8 (17.2)	13–88
Total fertility rate (per woman)	4.9 (1.0)	2.5–7.4
Out-of-pocket expenditure (%)	37.2 (19.9)	8–78
Under-five mortality rate (per 1,000 live births)	79.8 (27.4)	36.6–136.7

^a SD = Standard deviation.

<https://doi.org/10.1371/journal.pone.0243316.t003>

made joint contraceptive decision with husband/partner, whose husband/partner or others made contraceptive decision, whose husband/partner was ≥ 40 years, whose husband/partner had at least primary education, were in a polygynous union, and wanted no more children at last birth. Table 3 presents the summary of the country-level characteristics. The mean total fertility rate was 4.9 while the mean percentage of births attended by skilled health providers 66.2%. The mean density of medical doctors was 1.4 per 10,000 population.

Multilevel logistic regression analyses

Measures of variations (random effects). From the fixed intercept of the empty model, the odds of using FPC in a typical country was 0.09 (not shown). However, the odds of using FPC varied considerably across the countries. As shown in Model 1 (empty model), there was a significant variation in the odds of using FPC across the 33 countries ($\sigma^2 = 0.82$, 95% CI = 0.49–1.36) (Table 4). The ICC indicated that approximately 20% of the variance in the odds of using FPC to limit childbearing was accounted for by the countries in the study, while 80% of the variance was accounted for by the individual or other unknown factors. The MOR of 2.36 in the empty model also indicated considerable heterogeneity between the countries (Table 4). If a woman moved to another country with a higher probability of FPC use, their odds of using FPC would (in median) increase 2.36 times. In the full model (Model 4), after adjusting for the individual- and country-level factors and survey year, the variation in the odds of using FPC across the countries remained significant ($\sigma^2 = 0.42$, 95% CI = 0.23–0.76). The ICC decreased to 11% and there was a reduction in MOR to 1.85 (Table 4). The full model showed that 49% of the variance in the odds of FPC in the empty model was attributable to the individual- and country-level factors considered in this study (Table 4).

Measures of associations (fixed effects). The fixed effects are shown in Table 4. In Model 2, age, living children, household wealth, area of residence, decision maker, and ideal versus living children were statistically significant. Births attended by skilled health providers and total fertility rate were the only significant factors in Model 3. In the full model (Model 4) that included all the individual- and country-level factors, after adjusting for survey year, all the individual-level factors that were significant in Model 2 (age, living children, household wealth, area of residence, decision maker, and ideal versus living children) remained statistically significant. While for country-level factors, births attended by skilled health providers remained significant as in Model 3, but total fertility rate was no longer statistically significant. However, density of medical doctors which was not significant in Model 3 became significant in the full model. For every one unit increase in age, the odds of using FPC increased by 1.10

Table 4. Multilevel logistic regression models of factors associated with the use of FPC compared with other modern contraceptive methods among married or in-union with desire to limit childbearing.

Variables	Model 1 OR ^b (95% CI ^c)	Model 2 ^a OR (95% CI)	Model 3 ^a OR (95% CI)	Model 4 ^a OR (95% CI)
Fixed effects				
Individual-level factors				
Age (P) ^d		1.10 (1.08–1.12)***		1.10 (1.08–1.12)***
Education (P)				
None		Reference		Reference
Primary		1.00 (0.84–1.20)		0.99 (0.83–1.19)
Secondary or higher		1.03 (0.81–1.30)		1.00 (0.79–1.27)
Husband/partner's age (P)		1.01 (1.00–1.01)		1.01 (1.00–1.01)
Husband/Partner's education (P)				
None		Reference		Reference
Primary		1.12 (0.91–1.38)		1.11 (0.90–1.37)
Secondary or higher		1.13 (0.91–1.45)		1.14 (0.90–1.44)
Union (P)				
Monogynous		Reference		Reference
Polygynous		0.93 (0.78–1.10)		0.93 (0.79–1.11)
Household wealth (E) ^e				
Poor		Reference		Reference
Middle		0.97 (0.81–1.12)		0.97 (0.82–1.16)
Rich		1.38 (1.17–1.63)***		1.39 (1.18–1.64)***
Area of residence (E)				
Urban		Reference		Reference
Rural		0.83 (0.71–0.97)*		0.83 (0.71–0.97)*
Decision maker (E)				
Mainly respondent		Reference		Reference
Joint decision		1.68 (1.43–1.99)***		1.68 (1.43–1.99)***
Mainly husband/partner or others		2.45 (1.96–3.07)***		2.46 (1.97–3.07)***
Media exposure (E)				
No		Reference		Reference
Yes		1.00 (0.88–1.14)		1.00 (0.88–1.14)
Living children (N) ^f		1.10 (1.04–1.16)**		1.11 (1.04–1.16)***
Number of sons (N)		1.03 (0.97–1.08)		1.03 (0.97–1.08)
Ideal and living children (N)				
Living equal or greater than ideal		Reference		Reference
Living less than ideal		1.39 (1.20–1.61)***		1.40 (1.21–1.62)***
Wantedness (N)				
Wanted then		Reference		Reference
Wanted later		0.97 (0.82–1.15)		0.97 (0.82–1.14)
Wanted no more		1.13 (0.98–1.31)		1.12 (0.97–1.30)
Country-level factors				
Literacy rate (P)			1.01 (0.99–1.03)	1.01 (0.99–1.04)
Poverty rate (E)			0.98 (0.96–1.00)	0.98 (0.96–1.11)
Births attended by skilled health providers (E)			1.03 (1.01–1.05)*	1.03 (1.00–1.05)*
Density of medical doctors (E)			1.30 (0.98–1.72)	1.37 (1.01–1.85)*
Gross national income (E)			1.00 (1.00–1.00)	1.00 (0.96–1.01)
Out-of-pocket expenditure (E)			0.99 (0.97–1.01)	0.99 (0.96–1.01)
Rural population (E)			1.01 (0.98–1.04)	1.01 (0.98–1.04)

(Continued)

Table 4. (Continued)

Variables	Model 1 OR ^b (95% CI ^c)	Model 2 ^a OR (95% CI)	Model 3 ^a OR (95% CI)	Model 4 ^a OR (95% CI)
Total fertility rate (N)			1.71 (1.11–2.66)*	1.31 (0.81–2.14)
Under-five mortality rate (N)			1.01 (0.99–1.03)	1.00 (0.98–1.03)
Random effects				
Country level				
Variance (SE) ^g	0.82 (0.21)***	0.67 (0.21)***	0.43 (0.12)***	0.42 (0.19)***
95%CI	(0.49–1.36)	(0.38–1.22)	(0.25–0.73)	(0.23–0.76)
PCV (%) ^h	Reference	18.29	47.56	48.78
ICC (%) ⁱ	19.86 (12.92–29.27)	16.89 (10.03–27.03)	11.50 (7.11–18.06)	11.25 (6.51–18.78)
MOR ^j	2.36	2.17	1.86	1.85

^a Models 2–4 adjusted for survey year.

^b OR = Odds ratio.

^c Confidence interval.

^d P = Predisposing; ^e E = Enabling; ^f N = Need.

^g SE = Standard error. Standard error was used to calculate one-tail p-value for the variance.

^h PCV = Proportional change in variance; ⁱ ICC = Intra-country correlation coefficient. ^j MOR = Median odds ratio.

***p < 0.001

**p < 0.01

*p < 0.05.

<https://doi.org/10.1371/journal.pone.0243316.t004>

(95% CI = 1.08–1.12). Also, for every one unit increase in the number of living children, the odds of using FPC increased by 1.11 (95% CI = 1.04–1.16). Compared to women from poor households, those from rich households had higher odds of using FPC (OR = 1.39, 95% CI = 1.18–1.64). Women residing in rural areas had lower odds of using FPC compared to those who resided in urban areas (OR = 0.89, 95% CI = 0.71–0.97). Women whose partners or others made contraceptive decisions (OR = 2.46, 95% CI = 1.97–3.07) or who made joint contraceptive decisions with their partners (OR = 1.68, 95% CI = 1.43–1.99) had higher odds of using FPC compared to women who made the decision by themselves. The odds of using FPC was significantly higher among women whose number of living children was less than their ideal number of children (OR = 1.40, 95% CI = 1.21–1.62). For every one unit increase in births attended by skilled health providers, the odd of using FPC increased by 1.03 (95% CI = 1.00–1.05). Similarly, for every one unit increase in the density of medical doctors, the odd of using FPC increased by 1.37 (95% CI = 1.01–1.85).

Discussion

Using data from DHS program and three open data repositories, we investigated the individual- and country-level predisposing, enabling, and need factors associated with FPC use among married or in-union women aged 15–49 years who used modern contraceptive methods to limit childbearing in 33 countries in SSA. Approximately 13% of the women used FPC to limit childbearing. Individual-level predisposing, enabling, and needs factors and country-level enabling factors were found to be associated with the use of FPC among married or in-union women who were using modern contraceptive methods to limit childbearing.

FPC is a cost-effective contraceptive method that can help clients with the desire to limit childbearing achieve their reproductive intention. However, our study reaffirms that the majority of women using modern contraceptive to limit childbearing in SSA do not use FPC [4]. Although the classification of modern contraceptive methods may vary across studies [29],

our finding was similar to a previous study which showed that from 2000–2005, FPC accounted for 11% of the modern contraceptive method mix among married women in SSA [30]. Our result is also comparable to the findings from 17 European countries. Using data from the Generations and Gender Survey and DHS from 2004–2011, Dereuddre and colleagues reported that FPC accounted for 10% of the modern contraceptive method mix among women of reproductive age who had a male partner and no childbearing intention in Western, Central, and Eastern Europe [31]. Indeed, the prevalence of FPC in some African countries matches or exceeds the prevalence in some European countries [11]. Our finding on the use of FPC in SSA is, however, low compared to Latin American/Caribbean and Asian countries where FPC accounted for 50% and 47% of the modern contraceptive method mix among married women, respectively from 2000–2005 [30].

There is a dearth of studies on correlates of FPC conducted at regional levels for comparison. However, some of the findings in this study are consistent with similar studies at country levels that have reported positive association between age [14, 32–39], number of living children [35], economic status [37, 40], and urban residence [14, 37] and the use of FPC. Being a permanent contraceptive method, FPC may be less appealing to younger women whose fertility preference may change overtime [41]. Provider bias towards younger women may also account for their lower odds of using FPC to limit childbearing [42]. Owing to the high demand for children in SSA, women with fewer living children may not be willing to permanently limit birth, as the need for children or more children may arise following life circumstances such as the death of a child or remarriage. Policies and laws may also require women to have a specified number of living children before they can have FPC [43]. In settings where FPC requires a high out-of-pocket payment, clients with low income may not be able to afford it. Even in countries where FPC services are free, clients may still incur high out-of-pocket expenditure [44]. In developing countries, FPC services are less likely to be available and accessible in rural areas [45]. However, in developed countries like the U.S., rural dwellers have been found to have higher odds of using FPC [33, 36]. While the reasons for the higher prevalence in rural areas of developed countries are not clear, Lunde et al. opined that it might be due to less access to reversible methods [36].

Contrary to our expectation, the results showed that women who had a lower number of living children than their ideal number of children were more likely to use FPC. This observation cannot be easily explained, and it needs to be further explored in future studies. Nonetheless, “unrealized fertility” is common in SSA [46, 47], and factors such as socioeconomic constraints, competing alternatives, or health issues may make clients opt for permanent contraception despite not having achieved their desired number of children [47].

These results also established the influential role male partners and others, such as peers, relatives, or healthcare providers, play in decision making with regard to contraceptive use and choice in Africa [12, 48–51]. Power imbalances as a result of traditional socio-cultural norms, economic differences, and age disparities allow male partners to influence reproductive health decision making in patriarchal societies that exist in many Africa countries [52, 53]. Given the permanent nature of the procedure, married or in-union women may be reluctant to make the decision alone to avoid consequences such as intimate partner violence, separation, divorce, and extramarital affairs [54–56].

Although the cadre of personnel that constitute skilled birth attendants differ by country [57], a positive relationship between births attended by skilled birth attendants and contraceptive use has been previously reported in studies conducted at the individual level [58, 59]. As a contextual enabling factor in this study, this may reflect access to FPC services in the countries we assessed. Also, with the positive correlation between antenatal care visits (particularly at least four visits) and births attended by skilled birth attendants in SSA [60–62], it is possible

that in countries where a high proportion of births are attended by skilled birth attendants, women may have received information about FPC as an effective contraceptive option to limit childbearing during antenatal care, thereby increasing their preference for it [63]. In Brazil, receiving antenatal care was found to be positively associated with the use of FPC [64]. The higher odds of using FPC in countries with more doctors also illustrates the importance of access to the method, which is largely obtained through surgical procedures [65]. Although task sharing to non-doctors such as clinical officers and other mid-level providers as recommended by the World Health Organization (WHO) [66] is increasing [67–69], there are still restrictions on health care cadres that can perform FPC in many countries with shortages of doctors [3].

Our study is not without limitations. Considering the cross-sectional nature of the survey, causality cannot be inferred from the findings. There is also possibility of underreporting of FPC among some of the women due to its sensitivity [70]. The surveys included in the study were conducted at different time points. However, this limitation was minimized by controlling for the year of survey in the regression models. Also, for some of the country-level variables, the most recent available data did not correspond with the DHS survey year. Although, initially included as some of the important individual-level factors to be explored, religion and insurance coverage were eventually excluded from the analysis because they were not reported in all the countries. These findings are also not generalizable to all women. The study was restricted to married or in-union women because of the assumption that they are at increased risk of pregnancy and may be more interested in limiting childbearing compared to unmarried women. Cultural sensitivity in some African settings may also result in reporting bias on contraceptive use among unmarried women which was another reason for their exclusion [71].

Conclusions

Our results show that individual- and country- level factors are associated with the uptake of FPC in SSA. These factors suggest that increasing geographic, economic, and psychosocial access to FPC may improve its use among clients who desire to limit childbearing in SSA. Interventions such as mobile outreach to rural areas, reduced cost of FPC services, task shifting to lower cadre providers, male partner engagement, and joint family planning counseling may help in addressing some of these factors. Individual-level and other contextual factors specific to each country need to be examined. Qualitative studies are needed to understand the decision-making process among women and their male partners who choose to use FPC.

Supporting information

S1 Table. Description of explanatory variables.
(DOCX)

Author Contributions

Conceptualization: Babayemi O. Olakunde.

Formal analysis: Babayemi O. Olakunde.

Methodology: Babayemi O. Olakunde, Jennifer R. Pharr, Lung-Chang Chien.

Supervision: Jennifer R. Pharr, Lung-Chang Chien, Rebecca D. Benfield, Francisco S. Sy.

Writing – original draft: Babayemi O. Olakunde.

Writing – review & editing: Jennifer R. Pharr, Lung-Chang Chien, Rebecca D. Benfield, Francisco S. Sy.

References

1. Westoff CF. Unmet need for modern contraceptive methods. DHS analytical studies No. 28. Calverton, Maryland: ICF International; 2012.
2. Westoff CF, Bankole A. Trends in the demand for family limitation in developing countries. *Int Fam Plan Perspect.* 2000; 26: 56–62.
3. The RESPOND Project. A matter of fact, a matter of choice: The case for investing in permanent contraceptive methods. EngenderHealth White Paper. New York: EngenderHealth, The RESPOND Project; 2014.
4. Van Lith LM, Yahner M, Bakamjian L. Women's growing desire to limit births in sub-Saharan Africa: Meeting the challenge. *Glob Health Sci Pract.* 2013; 1: 97–107. <https://doi.org/10.9745/GHSP-D-12-00036> PMID: 25276520
5. Guttmacher Institute. Adding it up: Investing in contraception and maternal and newborn health, 2017-Fact sheet. New York: Guttmacher Institute; 2017.
6. Bradley SEK, Croft TN, Rutstein SO. The impact of contraceptive failure on unintended births and induced abortions: Estimates and strategies for reduction. DHS analytical studies No. 22. Calverton, Maryland: ICF International; 2011.
7. Darroch JE, Elizabeth S, Biddlecom A. Adding it up: Investing in contraception and maternal and newborn health, 2017-Supplementary tables. New York: Guttmacher Institute; 2017.
8. Sonfield A, Hasstedt K, Kavanaugh ML, Anderson R. The social and economic benefits of women's ability to determine whether and when to have children. New York: Guttmacher Institute; 2013.
9. Speizer IS, Lance P. Fertility desires, family planning use and pregnancy experience: longitudinal examination of urban areas in three African countries. *BMC Pregnancy Childbirth.* 2015; 15: 294. <https://doi.org/10.1186/s12884-015-0729-3> PMID: 26559486
10. United Nations, Department of Economic and Social Affairs, Population Division. Trends in contraceptive use worldwide 2015. New York: United Nations; 2015.
11. United Nations, Department of Economic and Social Affairs, Population Division. Contraceptive use by method 2019: Data booklet. New York: United Nations; 2019.
12. Olakunde BO, Sam-Agudu NA, Patel TY, Hunt AT, Buffington AM, Phebus TD, et al. Uptake of permanent contraception among women in sub-Saharan Africa: A literature review of barriers and facilitators. *Contraception.* 2019; 99: 205–211. <https://doi.org/10.1016/j.contraception.2018.12.007> PMID: 30685286
13. Ghosh S, Siddiqui MZ. Role of community and context in contraceptive behaviour in rural west bengal, India: A multilevel multinomial approach. *J Biosoc Sci.* 2017; 49: 48–68. <https://doi.org/10.1017/S0021932016000080> PMID: 27088672
14. Stephenson R. District-level religious composition and adoption of sterilization in India. *J Heal Popul Nutr.* 2006; 24: 100–106. PMID: 16796156
15. Marie Stopes International EngenderHealth, International IntraHealth, International Planned Parenthood Federation, Population Council, Jhpiego, et al. Increasing access to high-quality voluntary permanent methods of contraception in low-resource settings. 2014. Available: <https://www.engenderhealth.org/wp-content/uploads/imports/files/external/SIFPO-Symposium-Consensus-Statement-and-Addendum.pdf>
16. Easterlin RA. An economic framework for fertility analysis. *Stud Fam Plann.* 1975; 6: 54–63. PMID: 1118873
17. Easterlin RA, Wongboonsin K, Ahmed MA. The demand for family planning: A new approach. *Stud Fam Plann.* 1988; 19: 257–269. PMID: 3188132
18. Andersen RM. A behavioral model of families' use of health services. Research Series No. 25. Chicago: Center for Health Administration Studies, University of Chicago; 1968.
19. Andersen RM. Revisiting the behavioral model and access to medical care: Does it matter? *J Health Soc Behav.* 1995; 36: 1–10. PMID: 7738325
20. Andersen RM. National health surveys and the behavioral model of health services use. *Med Care.* 2008; 46: 647–653. <https://doi.org/10.1097/MLR.0b013e31817a835d> PMID: 18580382
21. von Lengerke T, Gohl D, Babitsch B. Re-revisiting the behavioral model of health care utilization by Andersen: A review on theoretical advances and perspectives. In: Janssen C, Swart E, von Lengerke T, editors. *Health Care Utilization in Germany.* New York, NY: Springer; 2014. pp. 11–28.

22. Croft TN, Marshall AMJ, Allen CK. Guide to DHS Statistics. Rockville, Maryland: ICF; 2018.
23. Ren R. Note on DHS standard weight de-normalization. [cited 25 Jan 2020]. Available: <http://userforum.measuredhs.com/index.php?t=getfile&id=4&>. Accessed January 12, 2020.
24. Midi H, Sarkar SK, Rana S. Collinearity diagnostics of binary logistic regression model. *J Interdiscip Math*. 2010; 13: 253–267.
25. Austin PC, Merlo J. Intermediate and advanced topics in multilevel logistic regression analysis. *Stat Med*. 2017; 36: 3257–3277. <https://doi.org/10.1002/sim.7336> PMID: 28543517
26. Merlo J, Chaix B, Ohlsson H, Beckman A, Johnell K, Hjerpe P, et al. A brief conceptual tutorial of multi-level analysis in social epidemiology: using measures of clustering in multilevel logistic regression to investigate contextual phenomena. *J Epidemiol Community Health*. 2006; 60: 290–297. <https://doi.org/10.1136/jech.2004.029454> PMID: 16537344
27. Merlo J, Yang M, Chaix B, Lynch J, Rastam L. A brief conceptual tutorial on multilevel analysis in social epidemiology: investigating contextual phenomena in different groups of people. *J Epidemiol Community Health*. 2005; 59: 729–736. <https://doi.org/10.1136/jech.2004.023929> PMID: 16100308
28. Larsen K, Merlo J. Appropriate assessment of neighborhood effects on individual health: integrating random and fixed effects in multilevel logistic regression. *Am J Epidemiol*. 2005; 161: 81–88. <https://doi.org/10.1093/aje/kwi017> PMID: 15615918
29. Festin MPR, Kiarie J, Solo J, Spieler J, Malarcher S, Van Look PFA, et al. Moving towards the goals of FP2020—classifying contraceptives. *Contraception*. 2016; 94: 289–294. <https://doi.org/10.1016/j.contraception.2016.05.015> PMID: 27287693
30. Seiber EE, Bertrand JT, Sullivan TM. Changes in contraceptive method mix in developing countries. *Int Fam Plan Perspect*. 2007; 33: 117–123. <https://doi.org/10.1363/3311707> PMID: 17938094
31. Dereuddre R, Van de Velde S, Bracke P. Gender inequality and the “East-West” divide in contraception: An analysis at the individual, the couple, and the country level. *Soc Sci Med*. 2016; 161: 1–12. <https://doi.org/10.1016/j.socscimed.2016.05.030> PMID: 27239702
32. Anderson JE, Jamieson DJ, Warner L, Kissin DM, Nangia AK, MacAluso M. Contraceptive sterilization among married adults: National data on who chooses vasectomy and tubal sterilization. *Contraception*. 2012; 85: 552–557. <https://doi.org/10.1016/j.contraception.2011.10.009> PMID: 22133657
33. Bass LE. Living in the American South and the likelihood of having a tubal sterilization. *Sociol Focus*. 2013; 46: 47–67.
34. Borrero S, Schwarz EB, Reeves MF, Bost JE, Creinin MD, Ibrahim SA. Race, insurance status, and tubal sterilization. *Obstet Gynecol*. 2007; 109: 94–100. <https://doi.org/10.1097/01.AOG.0000249604.78234.d3> PMID: 17197593
35. de Oliveira IT, Dias JG, Padmadas SS. Dominance of sterilization and alternative choices of contraception in India: an appraisal of the socioeconomic impact. *PLoS One*. 2014; 9: e86654. <https://doi.org/10.1371/journal.pone.0086654> PMID: 24489759
36. Lunde B, Rankin K, Harwood B, Chavez N. Sterilization of rural and urban women in the United States. *Obstet Gynecol*. 2013; 122: 304–311. <https://doi.org/10.1097/AOG.0b013e31829b5a11> PMID: 23969799
37. Perpetuo HIO, Wajnman S. Socioeconomic correlates of female sterilization in Brazil. In: Cosio-Zavala ME, editor. *Poverty, Fertility and Family Planning: Seminar held in Mexico City, June 2–4, 1998*. Paris: CICRED; 2003. pp. 311–333.
38. Thind A. Female sterilisation in rural Bihar: What are the acceptor characteristics? *J Fam Plan Reprod Heal care*. 2005; 31: 34–36. <https://doi.org/10.1783/0000000052972762> PMID: 15720847
39. White K, Potter JE, Zite N. Geographic variation in characteristics of postpartum women using female sterilization. *Women Health Issues*. 2015; 25: 628–633. <https://doi.org/10.1016/j.whi.2015.06.011> PMID: 26232310
40. Edmeades J, Pande RP, Falle T, Krishnan S. Son preference and sterilisation use among young married women in two slums in Bengaluru city, India. *Glob Public Health*. 2011; 6: 407–420. <https://doi.org/10.1080/17441692.2010.533686> PMID: 21218299
41. Sennott C, Yeatman S. Stability and change in fertility preferences among young women in Malawi. *Int Perspect Sex Reprod Health*. 2012; 38: 34–42. <https://doi.org/10.1363/3803412> PMID: 22481147
42. Tumlinson K, Okigbo CC, Speizer IS. Provider barriers to family planning access in urban Kenya. *Contraception*. 2015; 92: 143–151. <https://doi.org/10.1016/j.contraception.2015.04.002> PMID: 25869629
43. EngenderHealth. *Contraceptive sterilization: Global issues and trends*. New York: EngenderHealth; 2002.

44. Mohanty SK, Mishra S, Chatterjee S, Saggurti N. Pattern and correlates of out-of-pocket payment (OOP) on female sterilization in India, 1990–2014. *BMC Womens Health*. 2020; 20: 13. <https://doi.org/10.1186/s12905-020-0884-1> PMID: 31969139
45. Speizer IS, Hotchkiss DR, Magnani RJ, Hubbard B, Nelson K. Do service providers in tanzania unnecessarily restrict clients' access to contraceptive methods? *Int Fam Plan Perspect*. 2000; 26: 13–20&42.
46. Channon MD, Sarah H. Educational differentials in the realisation of fertility intentions: Is sub-Saharan Africa different? *PLoS One*. 2019; 14: e0219736. <https://doi.org/10.1371/journal.pone.0219736> PMID: 31318943
47. Casterline JB, Han S. Unrealized fertility: Fertility desires at the end of the reproductive career. *Demogr Res*. 2017; 36: 427–454.
48. Shattuck D, Kerner B, Gilles K, Hartmann M, Ng'ombe T, Guest G. Encouraging contraceptive uptake by motivating men to communicate about family planning: The Malawi Male Motivator Project. *Am J Public Health*. 2011; 101: 1089–1095. <https://doi.org/10.2105/AJPH.2010.300091> PMID: 21493931
49. Daniele MAS, Ganaba R, Sarrassat S, Cousens S, Rossier C, Drabo S, et al. Involving male partners in maternity care in Burkina Faso: A randomized controlled trial. *Bull World Health Organ*. 2018; 96: 450–461. <https://doi.org/10.2471/BLT.17.206466> PMID: 29962548
50. Musalia JM. Gender, social networks, and contraceptive use in Kenya. *Sex Roles*. 2005; 53: 835–846.
51. Blackstone SR, Nwaozuru U, Iwelunmor J. Factors influencing contraceptive use in sub-Saharan Africa: A systematic review. *Int Q Community Health Educ*. 2017; 37: 79–91. <https://doi.org/10.1177/0272684X16685254> PMID: 28056643
52. Blanc AK. The effect of power in sexual relationships on sexual and reproductive health: an examination of the evidence. *Stud Fam Plann*. 2001; 32: 189–213. <https://doi.org/10.1111/j.1728-4465.2001.00189.x> PMID: 11677692
53. Kriel Y, Milford C, Cordero J, Suleman F, Bekinska M, Steyn P, et al. Male partner influence on family planning and contraceptive use: Perspectives from community members and healthcare providers in KwaZulu-Natal, South Africa. *Reprod Health*. 2019; 16: 89. <https://doi.org/10.1186/s12978-019-0749-y> PMID: 31238960
54. Alemayehu M, Belachew T, Tilahun T. Factors associated with utilization of long acting and permanent contraceptive methods among married women of reproductive age in Mekelle town, Tigray region, north Ethiopia. *BMC Pregnancy Childbirth*. 2012; 12: 6. <https://doi.org/10.1186/1471-2393-12-6> PMID: 22280163
55. Lutala PM, Hugo JF, Lahiriri LN. Psychosocial implications of tubal ligation in a rural health district: A phenomenological study. *Reprod Health*. 2011; 8: 38. <https://doi.org/10.1186/1742-4755-8-38> PMID: 22176816
56. Chibalonza K, Chirhamolekwa C, Bertrand JT. Attitudes toward tubal ligation among acceptors, potential candidates, and husbands in Zaire. *Stud Fam Plann*. 1989; 20: 273–280. PMID: 2815201
57. Adegoke A, Utz B, Msuya SE, van den Broek N. Skilled birth attendants: Who is who? a descriptive study of definitions and roles from nine sub-Saharan African countries. *PLoS One*. 2012; 7: e40220. <https://doi.org/10.1371/journal.pone.0040220> PMID: 22808121
58. Mengesha ZB, Worku AG, Feleke SA. Contraceptive adoption in the extended postpartum period is low in Northwest Ethiopia. *BMC Pregnancy Childbirth*. 2015; 15: 160. <https://doi.org/10.1186/s12884-015-0598-9> PMID: 26231369
59. Rutaremwa G, Kabagenyi A, Wandera SO, Jhamba T, Akiror E, Nviiri HL. Predictors of modern contraceptive use during the postpartum period among women in Uganda: A population-based cross sectional study. *BMC Public Health*. 2015; 15: 262. <https://doi.org/10.1186/s12889-015-1611-y> PMID: 25885372
60. Nyongesa C, Xu X, Hall JJ, Macharia WM, Yego F, Hall B. Factors influencing choice of skilled birth attendance at ANC: Evidence from the Kenya demographic health survey. *BMC Pregnancy Childbirth*. 2018; 18: 88. <https://doi.org/10.1186/s12884-018-1727-z> PMID: 29631549
61. Jacobs C, Moshabela M, Maswenyeho S, Lambo N, Michelo C. Predictors of antenatal care, skilled birth attendance, and postnatal care utilization among the remote and poorest rural communities of Zambia: A multilevel analysis. *Front Public Health*. 2017; 5: 11. <https://doi.org/10.3389/fpubh.2017.00011> PMID: 28239604
62. Baatiema L, Ameyaw EK, Moomin A, Zankawah MM, Koramah D. Does antenatal care translate into skilled birth attendance? Analysis of 2014 Ghana Demographic and Health Survey. *Adv Public Health*. 2019; 6716938: 1–7.
63. Balogun OR, Adewole A, Adeniran AS, Adegboye R. Effect of counseling on contraceptive uptake in Nigeria. *J Med Biomed Sci*. 2017; 6: 1–6.

64. Rodrigues J, Moji K. Factors affecting choice of sterilisation among low income women in Paraiba, Brazil. *J Biosoc Sci.* 1995; 27: 339–345. <https://doi.org/10.1017/s0021932000022860> PMID: 7650051
65. Micks EA, Jensen JT. Permanent contraception for women. *Womens Health.* 2015; 11: 769–777. <https://doi.org/10.2217/whe.15.69> PMID: 26626698
66. World Health Organization. WHO recommendations: Optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting. Geneva: World Health Organization; 2012.
67. Gordon-Maclean C, Nantayi LK, Quinn H, Ngo TD. Safety and acceptability of tubal ligation procedures performed by trained clinical officers in rural Uganda. *Int J Gynaecol Obstet.* 2014; 124: 34–37. <https://doi.org/10.1016/j.ijgo.2013.07.016> PMID: 24156987
68. Nuccio O, Sendek B, Park MH, Mesele T, Okello FO, Gordon-Maclean C. Optimizing tubal ligation service delivery: a prospective cohort study to measure the task-sharing experience of Marie Stopes International Ethiopia. *Health Policy Plan.* 2016; 32: 163–165.
69. Jacobstein R. Lessons from the recent rise in use of female sterilization in Malawi. *Stud Fam Plann.* 2013; 44: 85–95. <https://doi.org/10.1111/j.1728-4465.2013.00345.x> PMID: 23512875
70. Choi Y, Khanna A, Zimmerman L, Radloff S, Zachary B, Ahmad D. Reporting sterilization as a current contraceptive method among sterilized women: lessons learned from a population with high sterilization rates, Rajasthan, India. *Contraception.* 2019; 99: 131–136. <https://doi.org/10.1016/j.contraception.2018.10.006> PMID: 30391289
71. Fabic MS, Jadhav A. Standardizing measurement of contraceptive use among unmarried women. *Glob Health Sci Pract.* 2019; 7: 564–574. <https://doi.org/10.9745/GHSP-D-19-00298> PMID: 31874938