

(*N*-*sec*-Butyl-*N*-*n*-propyldithio-carbamato- κ^2 S, S')triphenyltin(IV)

Normah Awang,^a Ibrahim Baba,^a Bohari M. Yamin,^a Seik Weng Ng^b and Edward R. T. Tiekink^{b*}

^aSchool of Chemical Sciences and Food Technology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 16 August 2010; accepted 17 August 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.032; wR factor = 0.083; data-to-parameter ratio = 21.9.

The Sn atom in the title compound, $[\text{Sn}(\text{C}_6\text{H}_5)_3(\text{C}_8\text{H}_{16}\text{NS}_2)]$, is pentacoordinated by two S atoms, derived from an asymmetrically coordinating dithiocarbamate ligand, and three *ipso*-C atoms. The coordination geometry is intermediate between square-pyramidal and trigonal-bipyramidal, with a leaning towards the latter. The presence of close intramolecular C—H...S contacts preclude the S atoms from forming significant intermolecular interactions. Rather, molecules are consolidated in the crystal structure by C—H... π interactions.

Related literature

For a review of the applications and structural chemistry of tin dithiocarbamates, see: Tiekink (2008). For a related organotin structure having the same dithiocarbamate ligand, see: Abdul Muthalib *et al.* (2010). For additional structural analysis, see: Addison *et al.* (1984).



Experimental

Crystal data

$[\text{Sn}(\text{C}_6\text{H}_5)_3(\text{C}_8\text{H}_{16}\text{NS}_2)]$

$M_r = 540.33$

Monoclinic, $C2/c$
 $a = 14.7997$ (5) Å
 $b = 12.1844$ (5) Å
 $c = 28.8891$ (11) Å
 $\beta = 97.348$ (1)°
 $V = 5166.7$ (3) Å³

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 1.16$ mm⁻¹
 $T = 293$ K
 $0.40 \times 0.30 \times 0.20$ mm

Data collection

Bruker SMART diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.653$, $T_{\max} = 0.801$

17218 measured reflections
5923 independent reflections
5179 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.083$
 $S = 1.02$
5923 reflections

271 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.70$ e Å⁻³
 $\Delta\rho_{\min} = -0.30$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

*Cg*1 and *Cg*2 are the centroids of the C1–C6 and C7–C12 rings, respectively.

<i>D</i> — <i>H</i> ... <i>A</i>	<i>D</i> — <i>H</i>	<i>H</i> ... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> — <i>H</i> ... <i>A</i>
C14—H14...S2	0.93	2.75	3.433 (3)	131
C20—H20...S2	0.98	2.49	3.059 (3)	117
C24—H24a...S1	0.97	2.58	2.938 (3)	102
C25—H25b...S1	0.97	2.84	3.360 (4)	115
C16—H16... <i>Cg</i> 1 ⁱ	0.93	2.78	3.618 (3)	151
C23—H23a... <i>Cg</i> 2 ⁱⁱ	0.96	2.91	3.773 (4)	150

Symmetry codes: (i) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank UKM (UKM-GUP-NBT-08-27-111 and UKM-ST-06-FRGS0092-2010) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2701).

References

- Abdul Muthalib, A. F., Baba, I., Mohamed Tahir, M. I., Ng, S. W. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, m1087.
Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2002). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Tiekink, E. R. T. (2008). *Appl. Organomet. Chem.* **22**, 533–550.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2010). E66, m1144 [doi:10.1107/S1600536810033039]

(*N*-sec-Butyl-*N*-*n*-propyldithiocarbamato- κ^2 S,S')triphenyltin(IV)

N. Awang, I. Baba, B. M. Yamin, S. W. Ng and E. R. T. Tiekink

Comment

Organotin dithiocarbamates display properties that suggest their use as anti-cancer agents, anti-microbial agents and as insecticides (Tiekink, 2008). Such interest motivates on-going structural characterization of such compounds (Abdul Muthalib *et al.*, 2010) and led to the investigation of the title compound, (I).

The Sn atom in (I) is penta-coordinated by two S atoms derived from an asymmetrically coordinating dithiocarbamate ligand and three *ipso*-C atoms from the phenyl substituents, Fig. 1. The resulting C₃S₂ coordination geometry is intermediate between square pyramidal and trigonal bi-pyramidal with a leaning towards the latter. Thus, compared to the ideal values for τ of 0.0 and 1.0 for ideal square pyramidal and trigonal bi-pyramidal geometries, respectively (Addison *et al.*, 1984), the value for τ in (I) computes to 0.55. The asymmetric mode of coordination of the dithiocarbamate ligand is reflected in significant differences in the associated C–S bond distances with that formed by the S1 atom, involved in the shorter of the Sn–S bonds, being considerably longer [S1–C19 = 1.755 (3) Å] than that formed by the S2 atom [S2–C19 = 1.682 (3) Å]. The observed molecular structure is entirely consistent with literature precedents (Tiekink, 2008).

Each of the S atoms is involved in two intramolecular C–H \cdots S contacts and these do not participate in intermolecular interactions, Table 2. The presence of C–H \cdots π contacts are noted, Table 1, and occur between benzene- and methyl-H atoms with two of the Sn-bound benzene rings functioning as the acceptors. These serve to consolidate the molecules into the crystal structure, Fig. 2.

Experimental

Carbon disulfide (30 mmol) was dropped into an ethanol solution (100 ml) of *N*-sec-butyl-*N*-*n*-propylamine (30 mmol). The solution was kept at 273 K for an hour. Triphenyltin chloride (30 mmol) dissolved in ethanol (100 ml) was added to give a white precipitate. This was collected and colourless crystals were obtained by recrystallization from its chloroform/ethanol (1/1) mixture.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 to 0.98 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2 to 1.5 $U_{\text{equiv}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. A view in projection down the *a* axis of (I) showing the unit-cell contents. The C–H... π contacts are shown as purple dashed lines.

(*N*-*sec*-Butyl-*N*-*n*-propyldithiocarbamato- κ^2 *S,S'*)triphenyltin(IV)

Crystal data

[Sn(C₆H₅)₃(C₈H₁₆NS₂)]

M_r = 540.33

Monoclinic, *C2/c*

Hall symbol: -*C* 2yc

a = 14.7997 (5) Å

b = 12.1844 (5) Å

c = 28.8891 (11) Å

β = 97.348 (1)°

V = 5166.7 (3) Å³

Z = 8

F(000) = 2208

D_x = 1.389 Mg m⁻³

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 7436 reflections

θ = 2.1–27.2°

μ = 1.16 mm⁻¹

T = 293 K

Block, colourless

0.40 × 0.30 × 0.20 mm

Data collection

Bruker SMART
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

T_{min} = 0.653, *T_{max}* = 0.801

17218 measured reflections

5923 independent reflections

5179 reflections with *I* > 2 σ (*I*)

R_{int} = 0.020

θ_{\max} = 27.5°, θ_{\min} = 1.4°

h = -19→17

k = -14→15

l = -34→37

Refinement

Refinement on *F*²

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)]$ = 0.032

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.083$

$S = 1.02$

5923 reflections

271 parameters

0 restraints

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0437P)^2 + 4.4496P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.70 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn	0.247917 (10)	0.577523 (13)	0.418770 (5)	0.04008 (7)
S1	0.22291 (6)	0.39926 (6)	0.37826 (2)	0.05486 (18)
S2	0.28766 (6)	0.56989 (6)	0.31823 (2)	0.0588 (2)
C1	0.21407 (15)	0.50813 (19)	0.48322 (8)	0.0397 (5)
C2	0.26523 (19)	0.4242 (2)	0.50565 (10)	0.0520 (6)
H2	0.3156	0.3979	0.4929	0.062*
C3	0.2432 (2)	0.3786 (3)	0.54658 (11)	0.0616 (7)
H3	0.2786	0.3222	0.5611	0.074*
C4	0.1697 (2)	0.4162 (3)	0.56561 (11)	0.0673 (9)
H4	0.1551	0.3861	0.5933	0.081*
C5	0.1171 (2)	0.4987 (3)	0.54404 (11)	0.0704 (9)
H5	0.0662	0.5235	0.5568	0.085*
C6	0.13936 (19)	0.5452 (2)	0.50331 (10)	0.0542 (6)
H6	0.1039	0.6019	0.4892	0.065*
C7	0.14476 (16)	0.6950 (2)	0.39602 (8)	0.0445 (5)
C8	0.06015 (19)	0.6649 (3)	0.37279 (11)	0.0649 (8)
H8	0.0484	0.5916	0.3654	0.078*
C9	-0.0062 (2)	0.7423 (4)	0.36066 (13)	0.0811 (11)
H9	-0.0623	0.7209	0.3450	0.097*
C10	0.0095 (2)	0.8508 (4)	0.37143 (12)	0.0774 (10)
H10	-0.0357	0.9028	0.3631	0.093*
C11	0.0915 (3)	0.8821 (3)	0.39429 (11)	0.0696 (9)
H11	0.1021	0.9556	0.4019	0.084*
C12	0.1597 (2)	0.8046 (2)	0.40645 (9)	0.0541 (6)
H12	0.2158	0.8270	0.4217	0.065*
C13	0.38064 (15)	0.6470 (2)	0.43524 (8)	0.0419 (5)
C14	0.42367 (19)	0.7084 (3)	0.40417 (10)	0.0585 (7)
H14	0.3981	0.7127	0.3731	0.070*
C15	0.5041 (2)	0.7634 (3)	0.41863 (11)	0.0650 (8)
H15	0.5320	0.8042	0.3973	0.078*
C16	0.54271 (18)	0.7581 (3)	0.46412 (11)	0.0593 (7)
H16	0.5967	0.7954	0.4737	0.071*
C17	0.50182 (19)	0.6978 (3)	0.49562 (11)	0.0599 (7)
H17	0.5279	0.6940	0.5266	0.072*
C18	0.42128 (18)	0.6423 (2)	0.48114 (9)	0.0523 (6)
H18	0.3940	0.6011	0.5027	0.063*
C19	0.24957 (19)	0.4410 (2)	0.32344 (9)	0.0473 (6)

supplementary materials

C20	0.2709 (3)	0.3917 (3)	0.24303 (10)	0.0678 (9)
H20	0.2875	0.4696	0.2433	0.081*
C21	0.1982 (3)	0.3750 (4)	0.20404 (12)	0.0915 (12)
H21A	0.1746	0.3010	0.2054	0.110*
H21B	0.2233	0.3830	0.1748	0.110*
C22	0.1203 (3)	0.4571 (4)	0.20540 (17)	0.1028 (14)
H22A	0.0741	0.4437	0.1796	0.154*
H22B	0.1433	0.5304	0.2034	0.154*
H22C	0.0948	0.4487	0.2341	0.154*
C23	0.3572 (3)	0.3250 (3)	0.23634 (13)	0.0822 (11)
H23A	0.3761	0.3427	0.2067	0.123*
H23B	0.3440	0.2480	0.2374	0.123*
H23C	0.4052	0.3430	0.2608	0.123*
C24	0.2103 (2)	0.2516 (2)	0.29729 (10)	0.0591 (7)
H24A	0.2382	0.2270	0.3277	0.071*
H24B	0.2312	0.2036	0.2741	0.071*
C25	0.1091 (2)	0.2416 (3)	0.29490 (12)	0.0711 (8)
H25A	0.0810	0.2550	0.2632	0.085*
H25B	0.0865	0.2962	0.3150	0.085*
C27	0.0838 (3)	0.1271 (3)	0.31025 (17)	0.0987 (13)
H27A	0.0188	0.1211	0.3082	0.148*
H27B	0.1105	0.1148	0.3419	0.148*
H27C	0.1063	0.0733	0.2903	0.148*
N1	0.24161 (17)	0.36613 (19)	0.28932 (7)	0.0540 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn	0.04146 (10)	0.04203 (11)	0.03675 (10)	-0.00450 (7)	0.00505 (7)	0.00138 (6)
S1	0.0814 (5)	0.0478 (4)	0.0381 (3)	-0.0154 (3)	0.0179 (3)	-0.0017 (3)
S2	0.0903 (5)	0.0459 (4)	0.0421 (4)	-0.0182 (3)	0.0164 (3)	-0.0020 (3)
C1	0.0419 (11)	0.0412 (12)	0.0364 (11)	-0.0039 (10)	0.0066 (9)	-0.0017 (9)
C2	0.0474 (14)	0.0594 (17)	0.0509 (15)	0.0071 (12)	0.0129 (12)	0.0080 (12)
C3	0.0612 (17)	0.0668 (19)	0.0568 (17)	0.0063 (15)	0.0070 (13)	0.0203 (14)
C4	0.080 (2)	0.080 (2)	0.0455 (16)	-0.0051 (17)	0.0205 (15)	0.0131 (14)
C5	0.0710 (19)	0.083 (2)	0.0644 (19)	0.0102 (17)	0.0366 (15)	0.0018 (17)
C6	0.0553 (15)	0.0532 (15)	0.0566 (16)	0.0098 (12)	0.0164 (12)	0.0027 (12)
C7	0.0437 (12)	0.0550 (15)	0.0355 (12)	-0.0011 (11)	0.0074 (9)	0.0085 (10)
C8	0.0526 (15)	0.069 (2)	0.0700 (19)	-0.0120 (14)	-0.0053 (14)	0.0156 (15)
C9	0.0464 (16)	0.109 (3)	0.084 (2)	-0.0040 (18)	-0.0042 (15)	0.035 (2)
C10	0.070 (2)	0.097 (3)	0.068 (2)	0.031 (2)	0.0197 (17)	0.0336 (19)
C11	0.096 (2)	0.0607 (19)	0.0541 (17)	0.0186 (18)	0.0150 (17)	0.0078 (14)
C12	0.0634 (16)	0.0553 (16)	0.0427 (14)	0.0011 (13)	0.0036 (12)	0.0021 (11)
C13	0.0386 (11)	0.0462 (13)	0.0415 (12)	-0.0025 (10)	0.0073 (9)	-0.0015 (10)
C14	0.0583 (16)	0.072 (2)	0.0449 (14)	-0.0156 (14)	0.0058 (12)	0.0049 (13)
C15	0.0580 (17)	0.076 (2)	0.0632 (18)	-0.0208 (15)	0.0179 (14)	0.0008 (15)
C16	0.0394 (13)	0.0664 (19)	0.0721 (19)	-0.0072 (12)	0.0073 (12)	-0.0128 (15)
C17	0.0488 (14)	0.075 (2)	0.0538 (16)	-0.0035 (14)	-0.0026 (12)	-0.0035 (14)

C18	0.0491 (14)	0.0628 (17)	0.0451 (14)	-0.0060 (12)	0.0059 (11)	0.0030 (12)
C19	0.0597 (15)	0.0479 (14)	0.0350 (12)	-0.0078 (11)	0.0089 (11)	-0.0007 (10)
C20	0.095 (2)	0.0675 (19)	0.0435 (15)	-0.0198 (18)	0.0200 (15)	-0.0022 (13)
C21	0.140 (4)	0.084 (3)	0.0484 (18)	-0.018 (3)	0.005 (2)	0.0025 (17)
C22	0.107 (3)	0.090 (3)	0.104 (3)	0.009 (3)	-0.015 (3)	0.023 (3)
C23	0.095 (3)	0.090 (3)	0.070 (2)	0.003 (2)	0.041 (2)	-0.0058 (18)
C24	0.0743 (19)	0.0518 (16)	0.0531 (16)	0.0002 (14)	0.0152 (13)	-0.0078 (13)
C25	0.075 (2)	0.070 (2)	0.069 (2)	-0.0063 (17)	0.0095 (16)	-0.0089 (16)
C27	0.111 (3)	0.066 (2)	0.127 (4)	-0.029 (2)	0.049 (3)	-0.005 (2)
N1	0.0759 (15)	0.0499 (13)	0.0382 (11)	-0.0122 (11)	0.0150 (10)	-0.0048 (9)

Geometric parameters (Å, °)

Sn—C1	2.161 (2)	C14—H14	0.9300
Sn—C7	2.134 (3)	C15—C16	1.366 (4)
Sn—C1	2.161 (2)	C15—H15	0.9300
Sn—C13	2.136 (2)	C16—C17	1.370 (4)
Sn—C1	2.161 (2)	C16—H16	0.9300
Sn—S1	2.4725 (7)	C17—C18	1.388 (4)
S1—C19	1.755 (3)	C17—H17	0.9300
S2—C19	1.682 (3)	C18—H18	0.9300
C1—C6	1.388 (3)	C19—N1	1.337 (3)
C1—C2	1.383 (3)	C20—C21	1.468 (5)
C2—C3	1.383 (4)	C20—N1	1.491 (4)
C2—H2	0.9300	C20—C23	1.548 (5)
C3—C4	1.360 (4)	C20—H20	0.9800
C3—H3	0.9300	C21—C22	1.530 (6)
C4—C5	1.371 (5)	C21—H21A	0.9700
C4—H4	0.9300	C21—H21B	0.9700
C5—C6	1.383 (4)	C22—H22A	0.9600
C5—H5	0.9300	C22—H22B	0.9600
C6—H6	0.9300	C22—H22C	0.9600
C7—C12	1.381 (4)	C23—H23A	0.9600
C7—C8	1.392 (4)	C23—H23B	0.9600
C8—C9	1.374 (5)	C23—H23C	0.9600
C8—H8	0.9300	C24—N1	1.497 (4)
C9—C10	1.371 (6)	C24—C25	1.496 (4)
C9—H9	0.9300	C24—H24A	0.9700
C10—C11	1.360 (5)	C24—H24B	0.9700
C10—H10	0.9300	C25—C27	1.526 (5)
C11—C12	1.394 (4)	C25—H25A	0.9700
C11—H11	0.9300	C25—H25B	0.9700
C12—H12	0.9300	C27—H27A	0.9600
C13—C18	1.385 (3)	C27—H27B	0.9600
C13—C14	1.385 (4)	C27—H27C	0.9600
C14—C15	1.383 (4)		
C7—Sn—C13	113.84 (10)	C16—C17—C18	119.8 (3)
C7—Sn—C1	106.99 (9)	C16—C17—H17	120.1
C13—Sn—C1	105.72 (9)	C18—C17—H17	120.1

supplementary materials

C7—Sn—S1	112.67 (7)	C13—C18—C17	121.3 (3)
C13—Sn—S1	122.09 (7)	C13—C18—H18	119.4
C1—Sn—S1	91.54 (6)	C17—C18—H18	119.4
C19—S1—Sn	97.79 (9)	N1—C19—S2	124.7 (2)
C6—C1—C2	117.5 (2)	N1—C19—S1	117.3 (2)
C6—C1—Sn	121.17 (19)	S2—C19—S1	117.87 (15)
C2—C1—Sn	121.29 (18)	C21—C20—N1	113.0 (3)
C3—C2—C1	121.5 (3)	C21—C20—C23	111.5 (3)
C3—C2—H2	119.2	N1—C20—C23	110.0 (3)
C1—C2—H2	119.2	C21—C20—H20	107.4
C4—C3—C2	119.9 (3)	N1—C20—H20	107.4
C4—C3—H3	120.1	C23—C20—H20	107.4
C2—C3—H3	120.1	C20—C21—C22	111.7 (3)
C3—C4—C5	120.1 (3)	C20—C21—H21A	109.3
C3—C4—H4	120.0	C22—C21—H21A	109.3
C5—C4—H4	120.0	C20—C21—H21B	109.3
C6—C5—C4	120.2 (3)	C22—C21—H21B	109.3
C6—C5—H5	119.9	H21A—C21—H21B	107.9
C4—C5—H5	119.9	C21—C22—H22A	109.5
C5—C6—C1	120.8 (3)	C21—C22—H22B	109.5
C5—C6—H6	119.6	H22A—C22—H22B	109.5
C1—C6—H6	119.6	C21—C22—H22C	109.5
C12—C7—C8	118.0 (3)	H22A—C22—H22C	109.5
C12—C7—Sn	119.54 (19)	H22B—C22—H22C	109.5
C8—C7—Sn	122.4 (2)	C20—C23—H23A	109.5
C9—C8—C7	120.8 (3)	C20—C23—H23B	109.5
C9—C8—H8	119.6	H23A—C23—H23B	109.5
C7—C8—H8	119.6	C20—C23—H23C	109.5
C8—C9—C10	120.6 (3)	H23A—C23—H23C	109.5
C8—C9—H9	119.7	H23B—C23—H23C	109.5
C10—C9—H9	119.7	N1—C24—C25	113.3 (3)
C11—C10—C9	119.7 (3)	N1—C24—H24A	108.9
C11—C10—H10	120.1	C25—C24—H24A	108.9
C9—C10—H10	120.1	N1—C24—H24B	108.9
C10—C11—C12	120.3 (3)	C25—C24—H24B	108.9
C10—C11—H11	119.8	H24A—C24—H24B	107.7
C12—C11—H11	119.8	C24—C25—C27	110.0 (3)
C7—C12—C11	120.6 (3)	C24—C25—H25A	109.7
C7—C12—H12	119.7	C27—C25—H25A	109.7
C11—C12—H12	119.7	C24—C25—H25B	109.7
C18—C13—C14	117.6 (2)	C27—C25—H25B	109.7
C18—C13—Sn	118.10 (18)	H25A—C25—H25B	108.2
C14—C13—Sn	123.77 (18)	C25—C27—H27A	109.5
C15—C14—C13	121.0 (3)	C25—C27—H27B	109.5
C15—C14—H14	119.5	H27A—C27—H27B	109.5
C13—C14—H14	119.5	C25—C27—H27C	109.5
C16—C15—C14	120.4 (3)	H27A—C27—H27C	109.5
C16—C15—H15	119.8	H27B—C27—H27C	109.5
C14—C15—H15	119.8	C19—N1—C20	120.7 (2)

C15—C16—C17	119.9 (3)	C19—N1—C24	121.5 (2)
C15—C16—H16	120.0	C20—N1—C24	117.7 (2)
C17—C16—H16	120.0		
C7—Sn—S1—C19	71.53 (12)	C10—C11—C12—C7	-0.8 (5)
C13—Sn—S1—C19	-69.63 (13)	C7—Sn—C13—C18	114.8 (2)
C1—Sn—S1—C19	-179.32 (11)	C1—Sn—C13—C18	-2.4 (2)
C7—Sn—C1—C6	-2.9 (2)	S1—Sn—C13—C18	-104.5 (2)
C13—Sn—C1—C6	118.8 (2)	C7—Sn—C13—C14	-56.8 (3)
S1—Sn—C1—C6	-117.2 (2)	C1—Sn—C13—C14	-173.9 (2)
C7—Sn—C1—C2	176.2 (2)	S1—Sn—C13—C14	84.0 (2)
C13—Sn—C1—C2	-62.1 (2)	C18—C13—C14—C15	-0.3 (4)
S1—Sn—C1—C2	62.0 (2)	Sn—C13—C14—C15	171.3 (2)
C6—C1—C2—C3	0.0 (4)	C13—C14—C15—C16	0.0 (5)
Sn—C1—C2—C3	-179.2 (2)	C14—C15—C16—C17	0.2 (5)
C1—C2—C3—C4	0.0 (5)	C15—C16—C17—C18	0.0 (5)
C2—C3—C4—C5	0.6 (5)	C14—C13—C18—C17	0.5 (4)
C3—C4—C5—C6	-1.1 (5)	Sn—C13—C18—C17	-171.6 (2)
C4—C5—C6—C1	1.1 (5)	C16—C17—C18—C13	-0.4 (5)
C2—C1—C6—C5	-0.5 (4)	Sn—S1—C19—N1	-178.0 (2)
Sn—C1—C6—C5	178.6 (2)	Sn—S1—C19—S2	4.69 (18)
C13—Sn—C7—C12	-19.9 (2)	N1—C20—C21—C22	66.8 (4)
C1—Sn—C7—C12	96.5 (2)	C23—C20—C21—C22	-168.7 (3)
S1—Sn—C7—C12	-164.36 (18)	N1—C24—C25—C27	-171.8 (3)
C13—Sn—C7—C8	163.4 (2)	S2—C19—N1—C20	2.7 (4)
C1—Sn—C7—C8	-80.2 (2)	S1—C19—N1—C20	-174.4 (2)
S1—Sn—C7—C8	18.9 (2)	S2—C19—N1—C24	177.7 (2)
C12—C7—C8—C9	0.0 (4)	S1—C19—N1—C24	0.6 (4)
Sn—C7—C8—C9	176.8 (3)	C21—C20—N1—C19	-124.4 (3)
C7—C8—C9—C10	-0.3 (5)	C23—C20—N1—C19	110.2 (3)
C8—C9—C10—C11	0.0 (5)	C21—C20—N1—C24	60.4 (4)
C9—C10—C11—C12	0.5 (5)	C23—C20—N1—C24	-65.0 (4)
C8—C7—C12—C11	0.5 (4)	C25—C24—N1—C19	80.9 (3)
Sn—C7—C12—C11	-176.3 (2)	C25—C24—N1—C20	-103.9 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 and Cg2 are the centroids of the C1—C6 and C7—C12 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C14—H14 \cdots S2	0.93	2.75	3.433 (3)	131
C20—H20 \cdots S2	0.98	2.49	3.059 (3)	117
C24—H24a \cdots S1	0.97	2.58	2.938 (3)	102
C25—H25b \cdots S1	0.97	2.84	3.360 (4)	115
C16—H16 \cdots Cg1 ⁱ	0.93	2.78	3.618 (3)	151
C23—H23a \cdots Cg2 ⁱⁱ	0.96	2.91	3.773 (4)	150

Symmetry codes: (i) $x+1/2, y+1/2, z$; (ii) $-x+1/2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

