

## Case Report

# Candidate Genes Associated with Delayed Neuropsychomotor Development and Seizures in a Patient with Ring Chromosome 20

Thiago Corrêa <sup>1</sup>, Amanda Cristina Venâncio,<sup>2</sup> Marcial Francis Galera,<sup>3</sup>  
and Mariluce Riegel <sup>1,4</sup>

<sup>1</sup>Genetics Department, Post-Graduate Program in Genetics and Molecular Biology, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, RS, Brazil

<sup>2</sup>Post-Graduate Program in Health Sciences, Universidade Federal do Mato Grosso (UFMT), Cuiabá, MT, Brazil

<sup>3</sup>Department of Pediatrics, Universidade Federal do Mato Grosso (UFMT), Cuiabá, MT, Brazil

<sup>4</sup>Medical Genetics Service, Hospital de Clínicas, Porto Alegre, RS, Brazil

Correspondence should be addressed to Mariluce Riegel; [mriegel@hcpa.edu.br](mailto:mriegel@hcpa.edu.br)

Received 4 November 2019; Accepted 17 December 2019; Published 21 January 2020

Academic Editor: Muhammad G. Kibriya

Copyright © 2020 Thiago Corrêa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ring chromosome 20 (r20) is characterized by intellectual impairment, behavioral disorders, and refractory epilepsy. We report a patient presenting nonmosaic ring chromosome 20 followed by duplication and deletion in 20q13.33 with seizures, delayed neuropsychomotor development and language, mild hypotonia, low weight gain, and cognitive deficit. Chromosomal microarray analysis (CMA) enabled us to restrict a chromosomal segment and thus integrate clinical and molecular data with systems biology. With this approach, we were able to identify candidate genes that may help to explain the consequences of deletions in 20q13.33. In our analysis, we observed five hubs (ARFGAP1, HELZ2, COL9A3, PTK6, and EEF1A2), seven bottlenecks (CHRNA4, ARFRP1, GID8, COL9A3, PTK6, ZBTB46, and SRMS), and two H-B nodes (PTK6 and COL9A3). The candidate genes may play an important role in the developmental delay and seizures observed in r20 patients. Gene ontology included microtubule-based movement, nucleosome assembly, DNA repair, and cholinergic synaptic transmission. Defects in these bioprocesses are associated with the development of neurological diseases, intellectual disability, neuropathies, and seizures. Therefore, in this study, we can explore molecular cytogenetic data, identify proteins through network analysis of protein-protein interactions, and identify new candidate genes associated with the main clinical findings in patients with 20q13.33 deletions.

## 1. Introduction

Ring chromosomes are rare structural rearrangements in humans, exhibiting an estimated frequency of 1 in 25,000 recognized conceptions [1] and approximately 1 in 30,000–60,000 births [2]. The ring formations are generally *de novo*, being only 1% inheritable [3–5]. Ring chromosome 20 (r20), which was first described in 1972 [6], is a syndrome characterized by refractory epilepsy, intellectual impairment, and behavioral disorders. Facial dysmorphism or other congenital malformations are rarely reported. r20 patients may present normal development until seizure onset, with the cognitive-behavioral decline being observed later, which suggests that the syndrome can be considered an epileptic encephalopathy [7, 8]. Most likely, r20 formation is

the result of intrachromosomal fusions from the direct union of unstable telomeres or the occurrence of two breaks, one in each chromosomal arm, resolved by the junction of the telomere ends of both arms, short and long, forming a circular structure [9, 10]. In the latter case, deletions, duplications, and/or inversions usually occur at the chromosomal ends [11, 12]. The diagnosis of ring chromosome 20 syndrome requires identification of ring formation by conventional cytogenetic techniques with the complement of chromosomal microarray analysis to detect small losses and gains in genetic material.

There are fewer than 20 cases described in the literature of patients carrying subtelomeric deletions in 20q13.33 [13–18]. Common manifestations of these individuals include skeletal and growth abnormalities, behavioral

problems, developmental delay, and seizures [19]. However, there are at least three factors that impair the clinical characterization of affected individuals and the identification of causal genes. First, there are notably few individuals molecularly characterized with high-resolution techniques. Second, there is no pattern regarding the presence of specific clinical manifestations in 20q13.33 deletions. Third, the significance of the ring morphology or chromosomal duplications in clinical abnormalities is unknown. These factors hinder efforts to explain the pathogenesis of 20q13.33 deletions and the probable molecular mechanisms involved in the phenotypic presentation of these individuals.

In this report, we describe an individual carrying a ring chromosome 20 with duplication and deletion in 20q13.33. The integration of cytogenetic, clinical, and protein-protein interactions data enabled us to identify genes that help to explain how the patient's phenotype is affected by the 20q13.33 deletion present on the ring chromosome.

## 2. Case Presentation

The proband is female, aged 2 years and 8 months, and is the only daughter of nonconsanguineous parents. The 35-year-old father and the 29-year-old mother reported gestation with high blood pressure and cesarean section with a gestational age of 41 weeks. The girl was born with a weight of 2860 g, length of 45 cm, and cephalic perimeter of 33 cm. The first convulsion of the child was manifested at 15 days of life as a generalized epileptic crisis, which was repeated 24 hours later. At 21 days, new seizures were characterized as focal seizures with secondary generalization and treatment with phenobarbital. The first neuropediatric assessment occurred at 4 months, showing delayed neuropsychomotor development, mild hypotonia, low weight gain, and cephalic perimeter of 36 cm ( $<-3$  Z scores). The electroencephalogram, cranial resonance, and screening tests for inborn errors of metabolism were unchanged. The child was referred for genetic evaluation. At the age of examination, the patient showed some facial dysmorphic features as enlarged nasal dorsum, bulbous nasal tip, short *columella*, and long nasolabial filter. The treatment with phenobarbital was efficient with control of the epileptic seizures for approximately 1 year, when it presented decompensation of the convulsive conditions, making it necessary to change the anticonvulsant to oxcarbazepine and levetiracetam. The child started walking independently at 26 months, and at 2 years and 6 months of age, the child presented delayed motor development and language and high cognitive deficit.

Karyotyping from the proband was performed on metaphase spreads prepared from peripheral blood samples. The chromosomal analysis was conducted after GTG banding at 550-band resolution, and at least 100 cells were analysed (Figure 1A). The karyotype showed results 46,XX,r(20). The parental karyotype was normal. At least 100 cells from each individual were analyzed. The DNA sample from the child was investigated using chromosomal microarray analysis (CMA) with a 60-mer oligonucleotide-based microarray with a theoretical resolution of 40 kb ( $8 \times 60$  K, Agilent Technologies Inc., Santa Clara, CA, USA).

The images were analyzed using Cytogenomics v 2.0 and 2.7 with the statistical algorithm ADM-2 and a sensitivity threshold of 6.0 (Figure 1B). It is recommended to provide confirmation of CMA results with other methods such as FISH or real-time PCR, especially for the refinement of breakpoints of structural chromosomal abnormalities. The authors pursued to perform it; but unfortunately, there is no sample left of this patient anymore, and the family is not available to obtain a new sample. The protein-protein interaction (PPI) metasearch engine STRING 10.0 (<http://string-db.org/>) was used to create PPI networks based on 40 genes and gene predictions located in the deleted region from our patient (Figure 2A). The list of genes was obtained from CMA analyses and subsequent research into the human assembly of February 2009 (GRCh37/hg19) [20, 21]. The parameters used in STRING were (i) degree of confidence, 0.400, with 1.0 being the highest level of confidence; (ii) 500 proteins in the 1st and 2nd shell; and (iii) all prediction methods enabled, except for textmining and gene fusion. The final PPI network obtained through STRING was analyzed using Cytoscape 3.5 [22]. The MCODE tool was used to identify densely connected regions in the final Cytoscape network. The PPI modules generated by MCODE were further studied by focusing on major biology-associated processes using the Biological Network Gene Ontology (BiNGO) 3.0.3 Cytoscape plugin [23]. The degree of functional enrichment for a given cluster and category was quantitatively assessed (p value) using a hypergeometric distribution. Multiple test correction was also implemented by applying the false discovery rate (FDR) algorithm [24] at a significance level of  $p < 0.05$ . Finally, two major parameters of network centralities (node degree and betweenness) were used to identify hub-bottleneck (H-B) nodes from the PPI network using the Cytoscape plugin CentiScaPe 3.2.1 [25].

## 3. Discussion

The underlying biological mechanism in individuals with ring 20 has not been determined. Hypotheses include (i) gene silencing by the influence of telomere position; (ii) uniparental disomy of chromosome 20; (iii) deleted genes in the chromosomal segment, or (iv) effect of ring instability in cellular functions [26]. In this study, we investigated genotype-phenotype correlations through deleted genes in 20q13.33 using systems biology approaches to explain the associated clinical spectrum in our patient. With this approach, we identified candidate genes that may be involved in the pathophysiology of ring chromosome 20. To measure the importance of the protein-protein interaction network of genes located in the deleted region (Figure 1B), we examined the topological properties of the network using centrality analyses.

The proteins ARFGAP1, HELZ2, and EEF1A2 presented a high node degree in the network. These nodes are considered hubs and have central functions in a biological network [25]. ARFGAP1 serves as a regulator of vesicular trafficking of proteins [27]. HELZ2 is a helicase that acts as a transcriptional coactivator [28]. EEF1A2 promotes the GTP-dependent binding of aminoacyl-tRNA during protein




FIGURE 1: (A) Chromosome banding showing the chromosome 20 pair, with a normal chromosome and the other on the ring (20). (B) CMA profile of chromosome 20 showing the hybridization pattern. The genomic segment with a gain of ~302,774 kb (61,142,577–61,445,350) is shown (vertical blue band). The arrow indicates the shift of the median ratio  $\log^2$  to +1.0. The segment with loss of ~1.4 Mb (61,472,348–62,872,839) is demonstrated (vertical red band). The arrow indicates the shift of the median ratio  $\log^2$  to -1.0.




FIGURE 2: PPI network. A list of 40 genes and gene predictions was obtained from the human assembly of February 2009 (GRCh37/hg19) (Table S12). Interaction data from STRING were used to construct networks using cytoscape software. (A) The primary network is composed of 1,572 nodes (proteins) and 54,925 edges (interactions). (B) The secondary network is composed of 794 nodes and 16,805 edges; yellow nodes are candidate proteins with data available in string (ZGPAT, HELZ2, EEF1A2, TNFRSF6B, ZBTB46, DIDO1, ARFGAP1, RTEL1, SRMS, GMEB2, YTHDF1, PTK6, GID8, BIRC7, COL9A3, COL20A1, KCNQ2, ARFRP1, and CHRNA4). (C) Cluster 2, with  $C_i = 40,863$ , is composed of 510 nodes and 10,388 edges. Nodes with green borders are considered hubs; orange are bottlenecks; and red are hub bottlenecks.

biosynthesis and plays a role in the regulation of actin function and cytoskeletal structure. *EEF1A2* knockout mice showed degeneration of neurons in the spinal cord and brain stem [29], and heterozygous mutations in the gene were associated with intellectual disability, developmental delay, autistic behaviors, and epilepsy [30, 31]. Therefore, the *EEF1A2* protein may be a good candidate to explain some of the symptoms present in individuals with deletion 20q13.33.

*CHRNA4*, *ARFRP1*, *GID8*, and *SRMS* were the nodes identified as bottlenecks in the network. *ARFRP1* plays a role in membrane trafficking between the trans-Golgi network and endosomes. *GID8* is a nuclear retention factor for  $\beta$ -catenin during Wnt signaling, and *SRMS* nonreceptor-type tyrosine kinases are a BRK family of kinases (BFKs) involved in the proliferation or differentiation of keratinocytes [32–34]. *CHRNA4* encodes alpha-4 nicotinic




FIGURE 3: Summary of the bioprocess enrichment identified in cluster 2. The colored horizontal bars show GO terms frequently present.

acetylcholine receptor subunits, and different mutations in the gene cause autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE) [35, 36]. *CHRNA4* is a known gene associated with epilepsy in ring 20 patients [26]. Bottlenecks are related to the control of information between the interactions in the network [25, 37]; therefore, the identification of *CHRNA4* already linked to the syndrome phenotype indicates that the haploinsufficiency of these bottlenecks could play a role in the development of our patient's phenotype.

Hub-bottlenecks (H-B) are nodes with a high degree and betweenness score. Among the nodes classified as H-B are two proteins, PTK6, another member of BFK, interacts directly with SRMS in the network. PTK6 functions as an intracellular signal transducer in epithelial tissues, contributing to the migration, adhesion, and progression of the cell cycle [38, 39]. *COL9A3* is a structural component of hyaline cartilage, and mutations in the gene are associated with multiple epiphyseal dysplasia [40, 41].

Genes associated with the same illness have been observed to interact with each other more frequently than predicted by chance [42]. Therefore, we performed cluster analysis to examine densely connected regions in the final network and observe these novel candidate genes localized in 20q13.33 [43]. We analyzed a total of 11 clusters (data not shown). Interesting relationships were found; for example, the proteins *HELZ2*, *EEF1A2*, *DIDO1*, *YTHDF1*, *PTK6*, *COL9A3*, and *COL20A1* interact with one another in cluster 2 (Figure 2C). Many of these genes are deleted in nonmosaic ring chromosome 20, and the clinical abnormalities identified in these individuals include findings also seen in our patient as seizures, intellectual disability, and developmental delay [8]. Ring chromosome 20 is associated with refractory epilepsy, behavioral problems, and mild-to-severe cognitive impairment. *De novo* microdeletion of 20q13.33 is associated with intellectual disabilities, developmental delay, speech and language delay, seizure, and behavioral problems such as autistic spectrum disorder. However, there is no pattern of abnormalities that would arouse clinical suspicion of a r(20) or *de novo* 20q13.33 microdeletion [19].

Functional enrichment analysis in the clusters revealed that the candidate genes were enriched in several biological processes, including microtubule-based movement, nucleosome assembly, DNA repair, and cholinergic synaptic transmission (Tables S1–S11). Defects in these bioprocesses are associated with the development of neurological diseases, intellectual disability, neuropathies, and seizures [44–48]. In addition, such bioprocesses as cell-matrix adhesion and

integrin-mediated signaling were identified (Figure 3). These pathways are involved in neural stem cell differentiation, neuronal migration, neuroplasticity, maturation, and function of synapses in the peripheral and central nervous system and may have an important contribution to the emergence of intellectual disability and seizures in humans [49–51]. The biological processes involving candidate genes denote the heterogeneity of pathways disrupted by 20q13.33 deletions.

The ring chromosome associated with subtelomeric deletions and duplications can confound and limit genotype-phenotype correlations. In fact, the circular structure of a ring chromosome, as described in this report, can change the ordinary 3D conformation of the chromatin in various ways and thus alter the expression of the genes present in the ring chromosome. The presence of an amplification identified in our ring chromosome analysis hinders efforts to determine its impact on the patient's phenotype. The duplication can be seen as a consequence of the mechanism of ring formation; in this case, an inverted duplication may be stabilized not only through telomere healing and telomere capture but also through circularization in the chromosomal ring [12]. Recently, chromothripsis and chromoanasythesis have been proposed as two independent mechanisms that could explain the combination of deletions and duplications on the same chromosome. Indeed, various molecular approaches, including whole genome sequencing, have shown that the concomitance of amplification, deletion, and ring chromosomes can be the result of a more complex rearrangement with respect to a ring chromosome with only the loss of extremities [52, 53]. In this case, it is expected that the final phenotype of the proband is not only the result of the abnormal dosage of deleted genes but also of the altered expression of duplicate genes present in two copies.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Supplementary Materials

Table 1: list of GO terms identified by BiNGO in Cluster 1.  
 Table 2: list of GO terms identified by BiNGO in Cluster 2.  
 Table 3: list of GO terms identified by BiNGO in Cluster 3.  
 Table 4: list of GO terms identified by BiNGO in Cluster 4.  
 Table 5: list of GO terms identified by BiNGO in Cluster 5.  
 Table 6: list of GO terms identified by BiNGO in Cluster 6.  
 Table 7: list of GO terms identified by BiNGO in Cluster 7.



Table 8: list of GO terms identified by BiNGO in Cluster 8.  
 Table 9: list of GO terms identified by BiNGO in Cluster 9.  
 Table 10: list of GO terms identified by BiNGO in Cluster 10.  
 Table 11: list of GO terms identified by BiNGO in Cluster 11.  
 Table 12: list of genes located in the 0.9 Mb smallest overlap region encompassing band 20q13.33. (*Supplementary Materials*)

## References

- [1] P. A. Jacobs, "Mutation rates of structural chromosome rearrangements in man," *American Journal of Human Genetics*, vol. 33, no. 1, pp. 44–54, 1981.
- [2] G. Kosztolányi, "The genetics and clinical characteristics of constitutional ring chromosomes," *Journal of the Association of Genetic Technologists*, vol. 35, no. 2, pp. 44–48, 2009.
- [3] G. Kosztolányi, "Does 'ring syndrome' exist? An analysis of 207 case reports on patients with a ring autosome," *Human Genetics*, vol. 75, no. 2, pp. 174–179, 1987.
- [4] G. Kosztolányi, K. Méhes, and E. B. Hook, "Inherited ring chromosomes: an analysis of published cases," *Human Genetics*, vol. 87, no. 3, pp. 320–324, 1991.
- [5] J. Knijnenburg, A. van Haeringen, K. B. M. Hansson et al., "Ring chromosome formation as a novel escape mechanism in patients with inverted duplication and terminal deletion," *European Journal of Human Genetics*, vol. 15, no. 5, pp. 548–555, 2007.
- [6] L. Atkins, W. L. Miller, and M. Salam, "A ring-20 chromosome," *Journal of Medical Genetics*, vol. 9, no. 3, pp. 377–380, 1972.
- [7] A. Vignoli, M. P. Canevini, F. Darra et al., "Ring chromosome 20 syndrome: a link between epilepsy onset and neuropsychological impairment in three children," *Epilepsia*, vol. 50, no. 11, pp. 2420–2427, 2009.
- [8] L. K. Conlin, W. Kramer, A. L. Hutchinson et al., "Molecular analysis of ring chromosome 20 syndrome reveals two distinct groups of patients," *Journal of Medical Genetics*, vol. 48, no. 1, pp. 1–9, 2011.
- [9] A. Vazna, M. Havlovicova, and Z. Sedlacek, "Molecular cloning and analysis of breakpoints on ring chromosome 17 in a patient with autism," *Gene*, vol. 407, no. 1–2, pp. 186–192, 2008.
- [10] S. W. Maluf, *Citogenética Humana*, Artmed, Porto Alegre, Brazil, 2011.
- [11] M. J. McGinniss, H. H. Kazazian Jr., G. Stetten et al., "Mechanisms of ring chromosome formation in 11 cases of human ring chromosome 21," *American Journal of Human Genetics*, vol. 50, no. 1, pp. 15–28, 1992.
- [12] E. Rossi, M. Riegel, J. Messa et al., "Duplications in addition to terminal deletions are present in a proportion of ring chromosomes: clues to the mechanisms of formation," *Journal of Medical Genetics*, vol. 45, no. 3, pp. 147–154, 2007.
- [13] M. A. Aldred, S. Aftimos, C. Hall et al., "Constitutional deletion of chromosome 20q in two patients affected with albright hereditary osteodystrophy," *American Journal of Medical Genetics*, vol. 113, no. 2, pp. 167–172, 2002.
- [14] M. Balasubramanian, E. Atack, K. Smith, and M. J. Parker, "A novel de novo 20q13.32–q13.33 deletion in a 2-year-old child with poor growth, feeding difficulties and low bone mass," *Journal of Human Genetics*, vol. 60, no. 6, pp. 313–317, 2015.
- [15] F. Bené, A. Bottani, F. Marcelli et al., "A de novo 1–1.6 Mb subtelomeric deletion of chromosome 20q13.33 in a patient with learning difficulties but without obvious dysmorphic features," *American Journal of Medical Genetics Part A*, vol. 143A, no. 16, pp. 1894–1899, 2007.
- [16] M. Béri-Deixheimer, M.-J. Gregoire, A. Toutain et al., "Genotype-phenotype correlations to aid in the prognosis of individuals with uncommon 20q13.33 subtelomere deletions: a collaborative study on behalf of the 'Association des Cytogénéticiens de Langue Française'," *European Journal of Human Genetics*, vol. 15, no. 4, pp. 446–452, 2007.
- [17] A. E. Roberts, G. F. Cox, V. Kimonis, A. Lamb, and M. Irons, "Clinical presentation of 13 patients with subtelomeric rearrangements and a review of the literature," *American Journal of Medical Genetics*, vol. 128A, no. 4, pp. 352–363, 2004.
- [18] F. Shabtai, E. Ben-Sasson, S. Arieli, and J. Grinblat, "Chromosome 20 long arm deletion in an elderly malformed man," *Journal of Medical Genetics*, vol. 30, no. 2, pp. 171–173, 1993.
- [19] R. N. Traylor, D. L. Bruno, T. Burgess et al., "A genotype-first approach for the molecular and clinical characterization of uncommon de novo microdeletion of 20q13.33," *PLoS One*, vol. 5, no. 8, Article ID e12462, 2010.
- [20] W. J. Kent, C. W. Sugnet, T. S. Furey, and K. M. Roskin, "The human genome browser at UCSC W," *Journal of Medicinal Chemistry*, vol. 19, no. 10, pp. 1228–1231, 1976.
- [21] C. von Mering, L. J. Jensen, B. Snel et al., "STRING: known and predicted protein-protein associations, integrated and transferred across organisms," *Nucleic Acids Research*, vol. 33, pp. D433–437, 2005.
- [22] P. Shannon, A. Markiel, O. Ozier et al., "Cytoscape: a software environment for integrated models of biomolecular interaction networks," *Genome Research*, vol. 13, no. 11, pp. 2498–2504, 2003.
- [23] S. Maere, K. Heymans, and M. Kuiper, "BiNGO: a cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks," *Bioinformatics*, vol. 21, no. 16, pp. 3448–3449, 2005.
- [24] Y. Benjamini and Y. Hochberg, "Controlling the false discovery rate: a practical and powerful approach to multiple testing," *Journal of the Royal Statistical Society: Series B (Methodological)*, vol. 57, no. 1, pp. 289–300, 1995.
- [25] G. Scardoni and C. Laudanna, "Centralities based analysis of complex networks," in *New Frontiers in Graph Theory*, Y. Zhang, Ed., InTech, London, UK, 2002, <http://www.intechopen.com/books/new-frontiers-in-graph-theory/centralities-based-analysis-of-networks>.
- [26] R. D. Daber, L. K. Conlin, L. D. Leonard et al., "Ring chromosome 20," *European Journal of Medical Genetics*, vol. 55, no. 5, pp. 381–387, 2012.
- [27] I. Huber, M. Rotman, E. Pick et al., "[33] Expression, purification, and properties of ADP-ribosylation factor (ARF) GTPase activating protein-1," in *Methods in Enzymology*, vol. 329, pp. 307–316, no. 1995, Elsevier Inc., Amsterdam, Netherlands, 2001.
- [28] T. Tomaru, T. Satoh, S. Yoshino et al., "Isolation and characterization of a transcriptional cofactor and its novel isoform that bind the deoxyribonucleic acid-binding domain of peroxisome proliferator-activated receptor-gamma," *Endocrinology*, vol. 147, no. 1, pp. 377–388, 2006.
- [29] L. A. Griffiths, J. Doig, A. M. D. Churchhouse et al., "Haploinsufficiency for translation elongation factor eEF1A2 in aged mouse muscle and neurons is compatible with normal function," *PLoS One*, vol. 7, no. 7, Article ID e41917, 2012.
- [30] C. Bischoff, S. Kahns, A. Lund et al., "The human elongation factor 1 A-2 gene (EEF1A2): complete sequence and

- characterization of gene structure and promoter activity," *Genomics*, vol. 68, no. 1, pp. 63–70, 2000.
- [31] J. Nakajima, N. Okamoto, J. Tohyama et al., "De novo EEF1A2 mutations in patients with characteristic facial features, intellectual disability, autistic behaviors and epilepsy," *Clinical Genetics*, vol. 87, no. 4, pp. 356–361, 2015.
- [32] Y. Kawachi, H. Nakauchi, and F. Otsuka, "Isolation of a cDNA encoding a tyrosine kinase expressed in murine skin," *Experimental Dermatology*, vol. 6, no. 3, pp. 140–146, 1997.
- [33] Y. Lu, S. Xie, W. Zhang et al., "Twa1/Gid8 is a  $\beta$ -catenin nuclear retention factor in Wnt signaling and colorectal tumorigenesis," *Cell Research*, vol. 27, no. 12, pp. 1422–1440, 2017.
- [34] R. K. Goel, M. Paczkowska, J. Reimand, S. Napper, and K. E. Lukong, "Phosphoproteomics analysis identifies novel candidate substrates of the non-receptor tyrosine kinase, src-related kinase lacking c-terminal regulatory tyrosine and N-terminal myristoylation sites (SRMS)," *Molecular & Cellular Proteomics*, vol. 17, no. 5, pp. 925–947, 2018.
- [35] R. Combi, L. Dalprá, M. L. Tenchini, and L. Ferini-Strambi, "Autosomal dominant nocturnal frontal lobe epilepsy," *Journal of Neurology*, vol. 251, no. 8, pp. 923–934, 2004.
- [36] Y. Chen, L. Wu, Y. Fang et al., "A novel mutation of the nicotinic acetylcholine receptor gene CHRNA4 in sporadic nocturnal frontal lobe epilepsy," *Epilepsy Research*, vol. 83, no. 2-3, pp. 152–156, 2009.
- [37] M. E. J. Newman, "Modularity and community structure in networks," *Proceedings of the National Academy of Sciences*, vol. 103, no. 23, pp. 8577–8582, 2006.
- [38] K. E. Lukong, D. Larocque, A. L. Tyner, and S. Richard, "Tyrosine phosphorylation of Sam68 by breast tumor kinase regulates intranuclear localization and cell cycle progression," *Journal of Biological Chemistry*, vol. 280, no. 46, pp. 38639–38647, 2005.
- [39] W.-S. Shin, H. J. Shim, Y. H. Lee et al., "PTK6 localized at the plasma membrane promotes cell proliferation and migration through phosphorylation of Eps8," *Journal of Cellular Biochemistry*, vol. 118, no. 9, pp. 2887–2895, 2017.
- [40] M. Czarny-Ratajczak, J. Lohiniva, P. Rogala et al., "A mutation in COL9A1 causes multiple epiphyseal dysplasia: further evidence for locus heterogeneity," *The American Journal of Human Genetics*, vol. 69, no. 5, pp. 969–980, 2001.
- [41] C. Jeong, J. Y. Lee, J. Kim et al., "Novel COL9A3 mutation in a family diagnosed with multiple epiphyseal dysplasia: a case report," *BMC Musculoskeletal Disorders*, vol. 15, no. 1, pp. 1–6, 2014.
- [42] A.-L. Barabási, N. Gulbahce, and J. Loscalzo, "Network medicine: a network-based approach to human disease," *Nature Reviews Genetics*, vol. 12, no. 1, pp. 56–68, 2011.
- [43] J. Menche, A. Sharma, M. Kitsak et al., "Uncovering disease-disease relationships through the incomplete interactome," *Science*, vol. 347, no. 6224, Article ID 1257601, 2015.
- [44] A. Friedman, C. J. Behrens, and U. Heinemann, "Cholinergic dysfunction in temporal lobe epilepsy," *Epilepsia*, vol. 48, no. s5, pp. 126–130, 2007.
- [45] P. J. McKinnon, "DNA repair deficiency and neurological disease," *Nature Reviews Neuroscience*, vol. 10, no. 2, pp. 100–112, 2009.
- [46] M. A. M. Franker and C. C. Hoogenraad, "Microtubule-based transport-basic mechanisms, traffic rules and role in neurological pathogenesis," *Journal of Cell Science*, vol. 126, no. 11, pp. 2319–2329, 2013.
- [47] A. J. López and M. A. Wood, "Role of nucleosome remodeling in neurodevelopmental and intellectual disability disorders," *Frontiers in Behavioral Neuroscience*, vol. 9, pp. 1–10, 2015.
- [48] M. Du, J. Li, R. Wang, and Y. Wu, "The influence of potassium concentration on epileptic seizures in a coupled neuronal model in the hippocampus," *Cognitive Neurodynamics*, vol. 10, no. 5, pp. 405–414, 2016.
- [49] D. J. Webb, H. Zhang, D. Majumdar, and A. F. Horwitz, " $\alpha$ 5 integrin signaling regulates the formation of spines and synapses in hippocampal neurons," *Journal of Biological Chemistry*, vol. 282, no. 10, pp. 6929–6935, 2007.
- [50] C. S. Barros, S. J. Franco, and U. Muller, "Extracellular matrix: functions in the nervous system," *Cold Spring Harbor Perspectives in Biology*, vol. 3, no. 1, Article ID a005108, 2011.
- [51] X. Wu and D. S. Reddy, "Integrins as receptor targets for neurological disorders," *Pharmacology & Therapeutics*, vol. 134, no. 1, pp. 68–81, 2012.
- [52] P. Ly and D. W. Cleveland, "Rebuilding chromosomes after catastrophe: emerging mechanisms of chromothripsis," *Trends in Cell Biology*, vol. 27, no. 12, pp. 917–930, 2017.
- [53] L. Nazaryan-Petersen, J. Eisfeldt, M. Pettersson et al., "Replicative and non-replicative mechanisms in the formation of clustered CNVs are indicated by whole genome characterization," *PLoS Genetics*, vol. 14, no. 11, Article ID e1007780, 2018.