



OPEN

# Mitogenome of the extinct Desert 'rat-kangaroo' times the adaptation to aridity in macropodoids

Michael Westerman<sup>1</sup>, Stella Loke<sup>2</sup>, Mun Hua Tan<sup>3</sup> & Benjamin P. Kear<sup>4</sup>


The evolution of Australia's distinctive marsupial fauna has long been linked to the onset of continent-wide aridity. However, how this profound climate change event affected the diversification of extant lineages is still hotly debated. Here, we assemble a DNA sequence dataset of Macropodoidea—the clade comprising kangaroos and their relatives—that incorporates a complete mitogenome for the Desert 'rat-kangaroo', *Caloprymnus campestris*. This enigmatic species went extinct nearly 90 years ago and is known from a handful of museum specimens. *Caloprymnus* is significant because it was the only macropodoid restricted to extreme desert environments, and therefore calibrates the group's specialisation for increasingly arid conditions. Our robustly supported phylogenies nest *Caloprymnus* amongst the bettongs *Aepyprymnus* and *Bettongia*. Dated ancestral range estimations further reveal that the *Caloprymnus-Bettongia* lineage originated in nascent xeric settings during the middle to late Miocene, ~12 million years ago (Ma), but subsequently radiated into fragmenting mesic habitats after the Pliocene to mid-Pleistocene. This timeframe parallels the ancestral divergences of kangaroos in woodlands and forests, but predates their adaptive dispersal into proliferating dry shrublands and grasslands from the late Miocene to mid-Pleistocene, after ~7 Ma. We thus demonstrate that protracted changes in both climate and vegetation likely staged the emergence of modern arid zone macropodoids.

Arid zone marsupials are icons of Australia and have an inferred evolutionary history that extends back over some ~15 Ma<sup>1</sup>. Nevertheless, the precise divergence timings of the major extant clades are ambiguous, as are the possible drivers behind their adaptive radiations<sup>2–13</sup>.

Macropodoids (Macropodiformes: Macropodoidea)—the group encompassing living kangaroos, wallaroos, wallabies, pademelons and tree-kangaroos (Macropodidae), bettongs and potoroos (Potoroidae), the Musky rat-kangaroo (*Hypsiprymnodon moschatus*: Hypsiprymnodontidae), and their stem antecedents<sup>14</sup>—incorporate some of the most distinctive Australian arid zone marsupials, as epitomised by the famous Red kangaroo, *Osphranter rufus*<sup>15</sup>. The well-documented fossil record of this and other 'true kangaroos' (Macropodini) has been used to correlate arid zone macropodoid evolution with the expansion of intracontinental grasslands during the Pliocene and Pleistocene, from ~3–4 Ma<sup>3,9,12</sup>. By contrast, the contemporary diversification of xeric-adapted bettongs is often overlooked, but has considerable significance because it includes the only example of an exclusively desert-inhabiting macropodoid, the Desert 'rat-kangaroo', which is alternatively referred to as the "Oolacunta"<sup>16</sup> or Ngudlukanta<sup>17</sup>, *Caloprymnus campestris* (Fig. 1A).

The first scientific specimens of *C. campestris* were collected from northeastern South Australia (Fig. 1B) in 1842, with three preserved examples subsequently shipped to London for study<sup>18</sup>. These were dubbed 'Bettongia' *campestris* by Gould<sup>19</sup>, although Thomas<sup>20</sup> later recognised 'B.' *campestris* as morphologically distinct from *Bettongia*, and thus established a separate genus, *Caloprymnus*. No further sightings of *C. campestris* were reported after this initial description, and the species was assumed to be extinct for some 90 years until Finlayson<sup>16,21</sup> announced the "Rediscovery of *Caloprymnus campestris*" in 1931–1932, from the remote Kooncheera Dune<sup>17</sup> region in the Sturt Stony Desert of far northeastern South Australia (Fig. 1C). Since then, only a skin recovered

<sup>1</sup>Department of Ecology, Environment and Evolution, La Trobe University, Bundoora, VIC 3086, Australia. <sup>2</sup>Deakin Genomics Centre, School of Life and Environmental Sciences, Deakin University, Burwood, VIC 3125, Australia. <sup>3</sup>Department of Microbiology and Immunology, Bio21 Institute, School of Biosciences, University of Melbourne, Melbourne, VIC 3052, Australia. <sup>4</sup>Museum of Evolution, Uppsala University, 752 36 Uppsala, Sweden. ✉email: m.westerman@latrobe.edu.au; benjamin.kear@em.uu.se



**Figure 1.** (A) Painting of *Caloprymnus campestris* as illustrated by Gould<sup>81</sup> (image in public domain). (B) Estimated historical distribution of *C. campestris* (grey shaded area) and localities from which specimens were collected: (1) Koonchera; (2) Ooroowillanie; (3) Mulka; (4) Killalpaninna (based on data from *Google Maps* and *OZCAM Online Zoological Collections of Australian Museums*: <https://ozcam.org.au/>). (C) Sturt Stony Desert gibber plain habitat of *C. campestris* showing a ‘jump-up’ escarpment and ephemeral drainage channel lined by riparian vegetation in the distance (photograph reproduced with permission from Michael Letnic, University of New South Wales). (D) Preserved skin of *Caloprymnus campestris* (Australian Museum, Sydney [AM] M21674) from Killalpaninna in northeastern South Australia<sup>22</sup> (photograph reproduced with permission from Mark Eldridge, AM).

sometime between 1902 and 1905 (Fig. 1D) has been reidentified<sup>22</sup>, and various unsubstantiated live sightings made<sup>17,23,24</sup>, with the most recent in 2011<sup>24</sup> and 2013<sup>17</sup> prompting unsuccessful surveys for the species in 2018 and 2019<sup>17</sup>. *Caloprymnus campestris* has otherwise been classified as Extinct by the IUCN (<https://www.iucn.org/>) since 1994, with the probable cause being over-predation by feral dogs, cats and foxes<sup>25</sup>.

At latest count, only 25 specimens of *C. campestris* are catalogued in museums worldwide<sup>22</sup>. This dearth of research material has led to uncertainty about potoroid interrelationships<sup>26</sup>, as well as the concomitant chronicle of their arid zone evolution. Here, we therefore analyse the first complete mitochondrial (mt) genome of *C. campestris*, which augments the 12S rRNA (AY245615) and partial cytochrome *b* (AY237246) gene sequences<sup>27</sup> already available from *GenBank* (<https://www.ncbi.nlm.nih.gov/genbank/>). Our novel dataset is used to construct a comprehensive phylogeny of crown potoroid species and subspecies within Macropodoidea. We also apply molecular clock calibrated ancestral range estimations to infer both the timing and context of macropodoid habitat change over the last ~ 25 Ma.

## Materials and methods

**Samples and sequencing.** We obtained non-formalin-fixed liver samples from a male *Caloprymnus campestris* (Museums Victoria, Melbourne, Australia [NMV] C8981) that was collected in 1834 from Mulka cattle station in northeastern South Australia (Fig. 1B). Our DNA extraction, PCR amplifications, sequencing and alignment procedures followed Westerman et al.<sup>27,28</sup>. Whole genome libraries were prepared with the *Nextera* DNA flex library kit (Illumina, CA), incorporating 50 ng of input DNA per sample. Sequencing was performed on the *Illumina* *MiSeq* platform using 2 × 300 bp V3 chemistry to generate 4,445,476 read pairs and



1.55 Gb total sequence data. Raw reads were trimmed for adapters and quality using *Trimmomatic* 0.36<sup>29</sup> (sliding window = 4:15; leading = 3; trailing = 3), and then assembled via genome skimming with *IDBA-UD* 1.1.1<sup>30</sup> (mink = 20; maxk = 300; min\_contig = 500); this yielded an average depth-of-coverage of 76.7x (median = 70x; minimum = 14x; maximum = 311x) and insert length of 109.9 bp. The resulting *C. campestris* mitogenome (A = 34%; C = 24.1%; G = 12.1%; T = 29.8%) was annotated using the *MITOS* webserver<sup>31</sup> with start-stop positions for protein coding genes manually curated using *blastp* homologies extracted from the NCBI non-redundant (*nr*) database.

**Phylogenetic and molecular clock analyses.** Phylogenetic relationships within Macropodoidea were examined using a mitogenome dataset including representatives of all potoroid species, together with *Hypsiprymnodon moschatus* and multiple species-level exemplars for selected macropodid genera (see Supplementary Table S1). The Northern common cuscus, *Phalanger orientalis* (Phalangeridae), and Western pygmy possum, *Cercartetus concinnus* (Burramyidae), were added as non-macropodoid outgroups. To accommodate for recognised gene incongruence<sup>32</sup>, we then compared these results with analyses of nuclear (n), and combined mitogenome/mtDNA/nDNA sequence datasets derived from *GenBank*, which integrated an expanded taxon sample of all potoroid species and subspecies (see Supplementary Tables S1 and S2). The mitogenomes were treated as a single partition, or alternatively sub-partitioned into 12S/16S rRNA stems and loops, pooled 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> protein codon positions, and 3<sup>rd</sup> codon positions with RY coding to allow for heterogeneity and saturation. A General Time Reversible gene partition model, *gamma* distribution and variable site proportions were determined using *jModelTest*<sup>33</sup> (Supplementary Table S3).

Tree building employed Maximum likelihood and Bayesian methods implemented in *RAxML* 7.2.8<sup>34</sup>, *MrBayes* 3.2.7<sup>35</sup> and *BEAST* 2.2.1<sup>36</sup> with node support calculations based on 1000 bootstrap pseudoreplicates (%) and Bayesian Posterior Probabilities (BPP), respectively. Maximum likelihood used a GTR + I +  $\Gamma$  partition model, while non-dated Bayesian MCMC analyses were run for  $6 \times 10^6$  generations with a sample frequency of 1000, eight chains, default temperature of 0.2, and burn-in fixed at  $6 \times 10^4$ . Time-trees were constructed in *BEAST* 2.2.1<sup>36</sup> with relaxed clocks and the minimum–maximum node age constraints listed in the Supplementary Information. Up to 95% of the normal prior distributions were assigned to the interval between minimum and maximum, with 2.5% to each tail. *Gamma* priors (shape = 1; scale = 1) were assigned to the “ucl.d.mean” parameter for each partition. MCMC analyses were run for  $65 \times 10^6$  generations with a burn-in of  $10 \times 10^6$  generations and sampling every  $10 \times 10^3$  generations. ESS values were > 200 for all estimated parameters. *TreeAnnotator* 2.2.1 (<https://www.beast2.org/treeannotator/>) was used to summarise the tree sample with mean node heights.

**Ancestral area analyses.** Distributional areas were optimised onto the time-calibrated *BEAST* consensus tree and analysed using the *R* package *BioGeoBEARS*<sup>37</sup> to compare alternative biogeographical range models, and a Bayesian Binary MCMC (BBM) approach<sup>38,39</sup> to reconstruct ancestral ranges in *RASP* 4<sup>40</sup>. Area codes (Supplementary Table S4) followed standard units<sup>6</sup> but were refined to represent a generalised vegetation map<sup>41</sup>: A = humid forest (rainforest and/or ‘wet’ sclerophyll dominant) prevalent throughout eastern coastal Australia, western Tasmania and New Guinea; B = woodland (‘dry’ sclerophyll dominant) prevalent throughout northern, eastern and southwestern inland Australia and northeastern Tasmania; C = shrubland (*Acacia* and chenopodiaceous shrubland dominant) prevalent throughout central and central-western Australia; and D = grassland-desert (arid grasslands and/or desert dominant) prevalent in central and central-northwestern Australia. The maximum number of ancestral areas was restricted to three because this equalled the maximum number of areas occupied by our terminal taxa at any given node.

*BioGeoBEARS* comparisons proceeded with likelihood ratio testing of ‘Jumping dispersal events (+ J)’, which have been considered inappropriate for dispersal-extinction-cladogenesis (DEC) models<sup>42</sup>. However, the three parameter Bayesian inference of historical biogeography for discrete areas (BAYAREALIKE) + J model ( $P = 0.0006$ ) received overwhelmingly highest support (AICc = 199.6; AICc\_wt = 0.98) for conferring best statistical likelihood on our data (Supplementary Table S5). Finally, we accommodated for connectivity by designating a dispersal multiplier of ‘1’ for adjacent areas (A-B-C)<sup>41</sup> versus non-adjacent areas (A-D)<sup>41</sup>, which were assigned a value of ‘0.5’.

Our BBM analyses utilised 10 MCMC chains with default temperature 0.1, and run over  $5 \times 10^6$  generations with sampling frequency and burn-in fixed at 1000. Model settings included ‘Gamma(+ G)’ for among-site rate variation, and ‘Fixed (JC)’ for state frequencies.

## Results and discussion

The *Caloprymnus campestris* mitogenome (16,866 bp) is ordered with 13 protein-coding genes, two ribosomal (r)RNA genes, 21 transfer (t)RNAs, and a non-coding AT-rich control region, which follows the typical configuration for marsupials<sup>43,44</sup>. The tRNAs are arranged around the origin of the L strand (A-C-W-O<sub>1</sub>-N-Y) and intersected between the NADH2 and COX1 genes. Substitution of the anticodon GCC for tRNA<sup>ASP</sup> (*trnD*) is also consistent with RNA-editing<sup>45</sup>.

Maximum likelihood and Bayesian analyses of our mitogenome dataset produce unanimous resolution of Macropodoidea with Potoroidae as the sister to Macropodidae (Supplementary Figures S1–S6). This pivotal higher-level grouping accords with other crown macropodoid phylogenies<sup>12,46–49</sup>, and warrants a new taxonomic definition<sup>50</sup>, which we coin as Macropodia, new clade, herein (Table 1; Supplementary Information). Bootstrap and BPP support is > 90% for almost all constituent nodes except those uniting: (1) the extinct short-faced kangaroo, *Simosthenurus occidentalis*, with the Banded hare-wallaby, *Lagostrophus fasciatus*, as basally branching macropodids (partitioned/non-partitioned bootstrap = 58/63%; *MrBayes* partitioned/non-partitioned BPP = 0.54/0.56; *BEAST* partitioned/non-partitioned BPP = 1/1); (2) the Quokka, *Setonix brachyurus*, with other


Clade	Definition	Type
Macropodiformes	Most inclusive clade including <i>Balbaroo nalima</i> *, <i>Hypsiprymnodon moschatus</i> , <i>Potorous tridactylus</i> and <i>Macropus giganteus</i> , but excluding <i>Cercartetus concinnus</i> and <i>Phalanger orientalis</i>	Stem
Balbaridae*	Most inclusive clade including <i>Balbaroo nalima</i> *, but excluding <i>Hypsiprymnodon moschatus</i> , <i>Potorous tridactylus</i> and <i>Macropus giganteus</i>	Stem
Macropodoidea	Least inclusive clade including <i>Hypsiprymnodon moschatus</i> , <i>Potorous tridactylus</i> and <i>Macropus giganteus</i>	Crown
Hypsiprymnodontidae	Most inclusive clade including <i>Hypsiprymnodon moschatus</i> and <i>Propleopus oscillans</i> *, but excluding <i>Balbaroo nalima</i> *, <i>Potorous tridactylus</i> and <i>Macropus giganteus</i>	Stem
Hypsiprymnodontinae	Most inclusive clade including <i>Hypsiprymnodon moschatus</i> , but excluding <i>Propleopus oscillans</i> *	Stem
Propleopinae*	Most inclusive clade including <i>Propleopus oscillans</i> *, but excluding <i>Hypsiprymnodon moschatus</i>	Stem
<b>Macropodia, new clade</b>	Least inclusive clade including <i>Potorous tridactylus</i> and <i>Macropus giganteus</i> , but excluding <i>Hypsiprymnodon moschatus</i>	Crown
Potoroidae	Least inclusive clade including <i>Potorous tridactylus</i> and <i>Aepyprymnus rufescens</i> , but excluding <i>Hypsiprymnodon moschatus</i> and <i>Macropus giganteus</i>	Crown
Potorinae	Least inclusive clade including <i>Potorous tridactylus</i> , but excluding <i>Aepyprymnus rufescens</i>	Crown
Bettonginae	Least inclusive clade including <i>Aepyprymnus rufescens</i> , but excluding <i>Potorous tridactylus</i>	Crown
Macropodidae	Most inclusive clade including <i>Simosthenurus occidentalis</i> *, <i>Lagostrophus fasciatus</i> and <i>Macropus giganteus</i> , but excluding <i>Potorous tridactylus</i> and <i>Hypsiprymnodon moschatus</i>	Stem
Sthenurinae*	Most inclusive clade including <i>Simosthenurus occidentalis</i> *, but excluding <i>Lagostrophus fasciatus</i> and <i>Macropus giganteus</i>	Stem
Lagostrophinae	Most inclusive clade including <i>Lagostrophus fasciatus</i> , but excluding <i>Simosthenurus occidentalis</i> * and <i>Macropus giganteus</i>	Stem
Macropodinae	Most inclusive clade including <i>Macropus giganteus</i> , but excluding <i>Simosthenurus occidentalis</i> * and <i>Lagostrophus fasciatus</i>	Stem
Dorcopsini	Least inclusive clade including <i>Dorcopsis hageni</i> , but excluding <i>Dendrolagus lumholtzi</i> and <i>Macropus giganteus</i>	Crown
Dendrolagini	Least inclusive clade including <i>Dendrolagus lumholtzi</i> , but excluding <i>Dorcopsis hageni</i> and <i>Macropus giganteus</i>	Crown
Macropodini	Least inclusive clade including <i>Macropus giganteus</i> , but excluding <i>Dorcopsis hageni</i> and <i>Dendrolagus lumholtzi</i>	Crown

**Table 1.** Phylogenetic definitions for Macropodiformes, including Macropodia, new clade, and other selected constituent subclades. Conceptual explanations and phylogenetic definition registration details are provided in the Supplementary Information. \*Extinct.

macropodines (bootstrap = 48/60%; *MrBayes* BPP = 1/1; *BEAST* BPP = 0.63/0.72); (3) grey kangaroos in the genus *Macropus* with *Osphranter rufus* and brush wallabies representing the genus *Notamacropus* (bootstrap = 80/64%; *MrBayes* BPP = 0.99/1; *BEAST* BPP = 0.99/0.96); and (4) *O. rufus* with *Notamacropus* (bootstrap = 58/55%; *MrBayes* BPP = 0.81/1; *BEAST* BPP = 0.72/0.76). As found by previous studies<sup>5,12,27,46–51</sup>, Potoroinae comprises potoroos within the genus *Potorous* and is distinguished from its sister clade, which we designate Bettonginae<sup>52</sup> to include the Rufous bettong, *Aepyprymnus rufescens*, as the basally branching sister to *C. campestris* and the species of *Bettongia* (Table 1). Alternative monophyly of *C. campestris* with either *A. rufescens*<sup>53,54</sup>, or the species of *Potorous*<sup>27,50</sup> were tested using topological constraints in *PAUP*\* 4.0b10<sup>55</sup> (Supplementary Table S6), but decisively rejected ( $P < 0.0001$ \*\*\*). Taxonomically, therefore, we conclude that the original classification of Gould's Desert 'bettong'<sup>19</sup> as generically consistent with *Bettongia* is feasible, but defer any formal nomenclatural amendment pending a detailed morphological re-evaluation.

Our maximum likelihood, Bayesian and time-tree analyses of the nDNA (Supplementary Figures S7–S9) and combined mitogenome/mtDNA/nDNA datasets (Fig. 2; Supplementary Figures S10–S12) yield broadly compatible topologies, with the basal divergence of potorooids and macropodids, and subsequent split between potoroines and bettongines both occurring from the latest Oligocene to earliest-middle Miocene (Table 2; Supplementary Table S7). Notably, this concurs with divergence times derived using different dating methods and constraints<sup>12,46–50,56</sup>. Furthermore, while our *BioGeoBEARS* and *BBM* ancestral range estimations correlate the latest Eocene (or mid-Eocene using nDNA: Supplementary Table S7) to late Oligocene emergence of crown macropodoids with predominantly humid forest habitats (> 50% probability values from *BAYAREALIKE* + *J* [A] = 65.76%; *BBM* [A] = 61.31%; Supplementary Tables S8 S8 and S9), the initial radiation of potorooids (*BAYAREALIKE* + *J* [B/A] = 45.42/25.55%; *BBM* [B/AB] = 42.62/28.9%), together with the macropodid subclades Sthenurinae (*BAYAREALIKE* + *J* [B] = 82.31%; *BBM* [B] = 66.1%) and Lagostrophinae + Macropodinae (*BAYAREALIKE* + *J* [B] = 70.45%; *BBM* [B/BC] = 41.79/27.29%) are coordinated with earlier Miocene dispersals into woodland dominated mosaics (Fig. 2; Supplementary Tables S8–S11; Supplementary Figures S13 and S14). These potentially included 'mallee-like'<sup>57</sup> sclerophyll communities, which propagated throughout central Australia from the early to middle Miocene<sup>41</sup>.

The globally recognised<sup>58</sup> middle to late Miocene climatic transition from equable to increasingly cool, dry conditions<sup>41,59</sup> coincides with potoroine speciations into mesic environments throughout southern Australia<sup>27,56</sup>. These are tracked by our *BioGeoBEARS* and *BBM* estimates, which infer occupation of primarily woodland and forest habitats after the earliest-late Miocene (Supplementary Tables S7–S9; Supplementary Figures S13 and S14). This is concurrent with the incipient desertification of inland Australia<sup>60</sup>, which may have promoted genetic



**Figure 2.** Time calibrated phylogeny of crown Macropodoidea (filled black diamond) showing divergence of *Caloprymnus campestris* (bold type) within Bettonginae (black open circle), and correlated against a schematic of changing palaeohabitats across the late Oligocene–Holocene interval (modified from Kear et al.<sup>6</sup> and Den Boer et al.<sup>82</sup>). Topology is based on the partitioned mitogenome/mtDNA/nDNA dataset. Bayesian posterior probability (< 1.0) and bootstrap (< 100%) support values (regular type) derived using BEAST 2.2.1<sup>36</sup>/MrBayes 3.2.7<sup>35</sup>/RAxML 7.2.8<sup>34</sup> are indicated at relevant nodes. Branch colours denote major clades: Hypsiprymnodontidae (purple); Macropodia, new clade (burgundy); Potoroidae (pink); Potoroinae (orange); Bettonginae (ochre); Macropodidae (red); Sthenurinae (green); Lagostrophinae (yellow); Macropodinae (light blue); Dorcopsini (grey) Dendrolagini (brown); Macropodini (dark blue). \*Extinct taxa. See Table 2 for node number references (bold type) and the Supplementary Information for other analyses. Graphics produced with Adobe CC2021 by B.P.K.

Node	Divergence	Time estimate
1	Hypsiprymmodontidae v. Macropodia	30.49 (24.75–36.58)
2	Macropodidae v. Potoroidae	21.91 (18.15–25.66)
3	Potorinae v. Bettonginae	18.68 (15.36–22.01)
4	<i>Potorous longipes</i> v. other <i>Potorous</i>	12.71 (10.36–15.23)
5	<i>Potorous platyops</i> * v. other <i>Potorous</i>	9.62 (7.57–11.92)
6	<i>Potorous gilbertii</i> v. other <i>Potorous</i>	8.98 (7.11–11.11)
7	<i>Potorous tridactylus tridactylus</i> v. other <i>Potorous tridactylus</i> subsp.	7.12 (5.46–8.95)
8	<i>Potorous tridactylus apicalis</i> v. <i>Potorous tridactylus trisulcatus</i>	2.2 (1.57–2.97)
9	<i>Aepyprymnus rufescens</i> v. other Bettonginae	14.62 (11.79–17.4)
10	<i>Caloprymnus campestris</i> * v. <i>Bettongia</i> spp.	12.23 (9.77–14.72)
11	<i>Bettongia anhydra</i> * v. other <i>Bettongia</i>	8.8 (6.29–11.73)
12	<i>Bettongia leseuer</i> v. other <i>Bettongia</i>	7.67 (5.86–9.51)
13	<i>Bettongia gaimardi</i> v. other <i>Bettongia</i>	2.46 (1.81–3.17)
14	<i>Bettongia tropica</i> v. <i>Bettongia penicillata</i>	1.76 (1.25–2.33)
15	<i>Simosthenurus occidentalis</i> * v. other Macropodidae	19.21 (15.77–22.62)
16	<i>Lagostrophus fasciatus</i> v. other Macropodidae	18.6 (15.32–21.9)
17	Dorcopsini v. other Macropodinae	13.54 (11.19–15.96)
18	<i>Dorcopsulus vanheurni</i> v. <i>Dorcopsis hageni</i>	7.14 (5.41–9.05)
19	Dendrolagini v. Macropodini	12.6 (10.35–14.78)
20	<i>Thylogale billardieri</i> v. other Dendrolagini	11.51 (9.55–13.68)
21	<i>Petrogale</i> spp. v. <i>Dendrolagus</i> spp.	9.89 (8.01–11.69)
22	<i>Petrogale brachyotis</i> v. <i>Petrogale xanthopus</i>	7.55 (6.04–9.14)
23	<i>Dendrolagus lumholtzi</i> v. <i>Dendrolagus dorianus</i>	7.79 (6.17–9.42)
24	<i>Onychogalea unguifera</i> v. other Macropodini	11.95 (9.81–14.02)
25	<i>Setonix brachyurus</i> v. other Macropodini	11.46 (9.47–13.52)
26	<i>Lagorchestes</i> spp. v. other Macropodini	10.38 (8.59–12.29)
27	<i>Lagorchestes hirsutus</i> v. <i>Lagorchestes conspicillatus</i>	7.53 (5.92–9.12)
28	<i>Wallabia bicolor</i> v. other Macropodini	9.52 (7.86–11.25)
29	<i>Protemnodon anak</i> * v. other Macropodini	8.84 (7.3–10.44)
30	<i>Macropus</i> spp. v. other Macropodini	8.11 (6.68–9.58)
31	<i>Macropus giganteus</i> v. <i>Macropus fuliginosus</i>	3.84 (2.81–4.94)
32	<i>Notamacropus</i> spp. v. <i>Osphranter</i> spp.	7.81 (6.43–9.25)
33	<i>Notamacropus eugenii</i> v. <i>Notamacropus parma</i>	6.57 (5.29–7.9)
34	<i>Osphranter rufus</i> v. other <i>Osphranter</i> spp.	7.33 (6.01–8.7)
35	<i>Osphranter robustus</i> v. <i>Osphranter bernardus</i>	5.22 (4.11–6.37)

**Table 2.** Estimated divergence times (Ma) with confidence intervals for crown macropodoid clades based on the partitioned mitogenome/mtDNA/nDNA dataset. See Fig. 2 for node number references and the Supplementary Information for other dating analyses. \*Extinct.

segregation of the extinct Broad-faced potoroo, *Potorous platyops*, from Gilbert's potoroo, *Potorous gilbertii*, in central-southern<sup>61</sup> and southwestern Australia (BAYAREALIKE + J [B] = 79.64%; BBM [B/AB] = 45.87/44.42%), versus the Long-nosed potoroo, *Potorous tridactylus* (BAYAREALIKE + J [AB] = 77.96%; BBM [AB] = 94.82%), and basally branching Long-footed potoroo, *Potorous longipes*, in southeastern Australia<sup>56</sup>. Additionally, we show that regional subspecies distinctions within *P. tridactylus* were completed by the latest Pliocene to mid-Pleistocene (Table 2; Supplementary Table S7). Curiously, though, Cyt b K2P variation (Supplementary Table S12) implies substantially less genetic difference between the Tasmanian *P. tridactylus apicalis* and northeastern mainland *P. tridactylus tridactylus* (1.93%), in comparison to the southeastern mainland *P. tridactylus trisulcatus* (4.21%). Indeed, these values approximate those contrasting *P. tridactylus tridactylus*/*P. tridactylus trisulcatus* with *P. gilbertii* (2.69/5%), *P. platyops* (4.1/5%), and *P. longipes* (5.84/5.69%), supporting inferences of cryptic taxa<sup>56</sup>, but in our opinion, only up to species-level.

Despite the currently limited DNA sequence coverage for the extinct Finlayson's<sup>62</sup> Desert bettong, *Bettongia anhydra*<sup>63</sup>, we derive unequivocal support (Fig. 2; Supplementary Figures S1–S12) for the monophyly of *Bettongia* spp. (bootstrap = > 90%; BPP = 1), together with close relationships between the woodland-forest dwelling Eastern bettong, *Bettongia gaimardi*, Northern bettong, *Bettongia tropica*, and Brush-tailed bettong, *Bettongia penicillata penicillata* (bootstrap = > 99%; BPP = 1). Only a few hundred Cyt b (or control region) nucleotides are available for the Woylie, *Bettongia penicillata ogilbyi*<sup>64</sup>. Nevertheless, our BioGeoBEARS and BBM estimates suggest a latest middle to probably late Miocene divergence of *B. anhydra* (BAYAREALIKE + J [CD] = 98.54%; BBM [CD] = 55.06%) and the Boodie, *Bettongia leseuer*, (BAYAREALIKE + J [CD] = 98.12%; BBM [BCD] = 81.72%)



in xeromorphic habitats (Table 2; Supplementary Tables S7–S9; Supplementary Figures S13 and S14), followed by Pliocene to as recent as mid-Pleistocene radiations of *B. gaimardi* (BAYAREALIKE + J [CD] = 90.65%; BBM [BCD/BC] = 27.79/23.75%) and *B. tropica* + *B. penicillata* subsp. (BAYAREALIKE + J [CD] = 79.96.12%; BBM [BCD/BC] = 28.32/21.98%) coupled with increasing habitat variegation<sup>41</sup>. We correlate this with vicariant ‘reversions’<sup>5</sup> into eucalypt woodlands and forests<sup>65–67</sup> (Supplementary Tables S10 and S11), which contracted and fragmented with intensifying aridification over the Pliocene–Pleistocene interval<sup>68</sup>.

*Bettongia* is karyotypically conservative, retaining the  $2n = 22$  chromosomal number of most macropodoids<sup>69,70</sup>. Conversely, chromosomal fission in *P. longipes* has produced  $2n = 24$ , while fusions (and inversions) in *P. tridactylus* and *P. gilbertii* manifest unusual reductions to  $2n = 12♀$ ,  $13♂$ <sup>71</sup>. *Aepyprymnus rufescens*, on the other hand, exhibits a unique karyotypic increase to  $2n = 32$ , which is the highest for any marsupial<sup>71</sup>, and presumably reflects its independent evolution since the later-early to early-late Miocene (nDNA favouring a younger later-middle to early-late Miocene range: Table 2; Supplementary Table S7). Although the chromosomal arrangement of *C. campestris* is unknown, our robustly supported (bootstrap = >90%; BPP = 1) earliest-middle to early-late Miocene split from *Bettongia* (Table 2; Supplementary Table S7) suggests a similarly protracted ancestry, yet with genetic differentiation that approaches intrageneric levels within *Bettongia* spp. (Cyt *b* K2P variation being as little as 6.91% compared to *B. penicillata*: Supplementary Table S12). Significantly, our BioGeoBEARS (BAYAREALIKE + J [CD] = 87.73%) and BBM ([CD] = 53.41%) estimates correlate the *C. campestris*–*Bettongia* divergence with a seminal invasion of xeric environments (Supplementary Tables S8–S12; Supplementary Figures S13 and S14), perhaps incorporating arid chenopod shrublands that spread across central Australia from the middle to late Miocene<sup>41,57,60</sup>. The coeval radiation of macropodines is otherwise linked to predominantly woodland and forest settings (Table 2; Supplementary Tables S8–S12; Supplementary Figures S13 and S14). This includes dorcopsins (BAYAREALIKE + J [B] = 54.21%, BBM [AB/B] = 38.93/25.2%) and dendrolagins (BAYAREALIKE + J [B/A] = 49.35/34.51%, BBM [AB/ABC] = 47.93/26.27%) diverging coincident with uplift of the New Guinean landmass<sup>3,72,73</sup>, and macropodins which initially diversified in woodland habitats (BAYAREALIKE + J [B] = 95.79%; BBM [B] = 77.19%), but subsequently expanded into open shrublands and eventually grasslands (e.g., *Osphranter rufus*: BAYAREALIKE + J [B] = 51.33%, BBM [BC/BCD] = 31.62/24.84%) after the late Miocene to as recent as Pliocene to mid-Pleistocene (Table 2; Supplementary Table S7), thereby presaging the modern prevalence of grazing kangaroos<sup>9</sup>.

## Conclusions

Our characterisation of the complete mitogenome for *Caloprymnus campestris* provides an ecological diversification timescale for bettongs and potoroos within the context of crown macropodoid evolution. Most importantly, we show that the unambiguously monophyletic *C. campestris*–*Bettongia* lineage probably originated with the onset of increasingly arid intracontinental climates during the middle to late Miocene<sup>41,57–60,74</sup>, corresponding with the deepest divergences of Australia’s arid zone biota around ~15 Ma<sup>1</sup>. This contrasts with the largely late Miocene to Pleistocene radiation of kangaroos, whose abundance in modern arid zone habitats has been attributed to grazing adaptations and the spread of grasslands during the Pliocene and Pleistocene<sup>3,9,12</sup>. Clearly, therefore, the appearances of Australia’s distinctive arid zone macropodoids were staged over some ~3–6 Ma (based on minimum–maximum confidence interval differences for *C. campestris* versus *Osphranter rufus*: Table 2), and likely occurred in response to a complex interplay of abiotic and biotic drivers involving both climate and vegetation change.

Unfortunately, little is known about the biology of *C. campestris* or other extinct ‘Desert bettongs’, such as *Bettongia anhydra*<sup>63</sup>, and the Nullarbor dwarf bettong<sup>75</sup>, *Bettongia pusilla*<sup>76</sup>. Nonetheless, early eye-witness reports state that *C. campestris* inhabited sparsely vegetated gibber plains<sup>16</sup>. The diet of *C. campestris* is also uncertain<sup>23</sup>, but might have been varied<sup>16,23</sup> similar to the extant arid zone *Bettongia lesueur*<sup>77</sup> and *Bettongia penicillata*<sup>64</sup>, which consume a range of plant matter, fungi and insects<sup>78,79</sup>. *Caloprymnus campestris* was thus probably an important ‘ecosystem engineer’<sup>63</sup> whose tragic loss is compounded by dramatic range reductions and the Near Threatened (*Bettongia gaimardi*, *B. lesueur*, *Potorous tridactylus*), Vulnerable (*Potorous longipes*), Endangered (*Bettongia tropica*), Critically Endangered (*B. penicillata*, *Potorous gilbertii*), or Extinct (*B. anhydra*, *C. campestris*, *Potorous platyops*) IUCN Red listings (<https://www.iucnredlist.org/>) for 10 out of the 11 named non-fossil crown potoroids. The extinction susceptibility of *C. campestris* was presumably exacerbated by its limited distribution (only four recognised collection<sup>22</sup>, and 13 potential sighting localities<sup>17</sup> within a ~350 km radius) and desert specialisation, which when coupled with habitat modification and the introduction of exotic species via European pastoralism<sup>80</sup>, underscores the extreme conservation sensitivity of Australia’s unique arid zone marsupials and the urgent need to document their now dwindling multi-million-year evolutionary histories.

**Ethical approval and informed consent.** No live animal subjects were used for experiments in this study. All extinct animal tissues were obtained and their use approved by the La Trobe University Animal Ethics Committee (AEC). All experiments were performed in accordance with institutional guidelines and regulations.

## Data availability

Raw FASTQ files for the *Caloprymnus campestris* mitogenome assembly have been uploaded onto the Mendeley Data repository (<https://data.mendeley.com/>) under <https://doi.org/10.17632/ft8t7v7gfv.1>. The consensus *C. campestris* mitogenome (MT66337) and other macropodoid DNA sequences are available from GenBank (Supplementary Tables S1 and S2).

Received: 9 November 2021; Accepted: 25 March 2022

Published online: 06 April 2022

## References

- Byrne, M. *et al.* Birth of a biome: insights into the assembly and maintenance of the Australian arid zone biota. *Mol. Ecol.* **17**, 4398–4417 (2008).
- Krajewski, C., Wroe, S. & Westerman, M. Molecular evidence for the pattern and timing of cladogenesis in dasyurid marsupials. *Zool. J. Linn. Soc.* **130**, 375–404 (2000).
- Prideaux, G. J. & Warburton, N. M. An osteology based appraisal of the phylogeny and evolution of kangaroos and wallabies (Macropodidae, Marsupialia). *Zool. J. Linn. Soc.* **159**, 954–987 (2010).
- Westerman, M. *et al.* Phylogenetic relationships of living and recently extinct bandicoots based on nuclear and mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* **62**, 97–108 (2012).
- Mitchell, K. J. *et al.* Molecular phylogeny, biogeography, and habitat preference evolution of marsupials. *Mol. Biol. Evol.* **31**, 2322–2330 (2014).
- Kear, B. P., Aplin, K. P. & Westerman, M. Bandicoot fossils and DNA elucidate lineage longevity amongst xeric-adapted Australasian marsupials. *Sci. Rep.* **6**, 37537 (2016).
- Westerman, M. *et al.* Phylogenetic relationships of dasyuromorphian marsupials revisited. *Zool. J. Linn. Soc.* **176**, 686–701 (2016).
- Kealy, S. & Beck, R. Total evidence phylogeny and evolutionary timescale for Australian faunivorous marsupials (Dasyuromorphia). *BMC Evol. Biol.* **17**, 240 (2017).
- Couzens, A. M. C. & Prideaux, G. J. Rapid Pliocene adaptive radiation of modern kangaroos. *Science* **362**, 72–75 (2018).
- García-Navas, V., Rodríguez-Rey, M. & Westerman, M. Bursts of morphological and lineage diversification in modern dasyurids, a “classic” adaptive radiation. *Biol. J. Linn. Soc.* **123**, 782–795 (2018).
- Travouillon, K. & Phillips, M. J. Total evidence analysis of the phylogenetic relationships of bandicoots and bilbies (Marsupialia: Paramelomorphia): Reassessment of two species and description of a new species. *Zootaxa* **4378**, 224 (2018).
- Celik, M. *et al.* A molecular and morphometric assessment of the systematics of the *Macropus* complex clarifies the tempo and mode of kangaroo evolution. *Zool. J. Linn. Soc.* **186**, 793–812 (2019).
- García-Navas, V., Kear, B. P. & Westerman, M. The geography of speciation in dasyurid marsupials. *J. Biogeogr.* **47**, 2042–2053 (2020).
- Den Boer, W. & Kear, B. P. Is the fossil rat-kangaroo *Palaeopotoropus priscus* the most basally branching stem macropodiform?. *J. Vertebr. Paleontol.* **38**, e1428196 (2018).
- Freedman, C. R., Rothschild, D., Groves, C. & Newman, A. E. M. *Osphranter rufus* (Diprotodontia: Macropodidae). *Mammalogy* **52**, 143–164 (2020).
- Finlayson, H. H. *Caloprymnus campestris*; its occurrence and characters. *Trans. R. Soc. S. Aust.* **56**, 148–167 (1932).
- Vernes, K., Jackson, S. M., Elliott, T. F., Elliott, K. & Carr, S. G. A search for the desert rat-kangaroo or ngudlukanta (*Caloprymnus campestris*) in north-eastern South Australia. *Aust. Mammal.* <https://doi.org/10.1071/AM21022> (2021).
- Finlayson, H. H. Rediscovery of *Caloprymnus campestris* (Marsupialia). *Nature* **129**, 871 (1932).
- Gould, J. On a new species of kangaroo rat. *Proc. Zool. Soc. Lond.* **11**, 81 (1843).
- Thomas, O. *Catalogue of the Marsupialia and Monotremata in the Collection of the British Museum (Natural History)* (British Museum, 1888).
- Finlayson, H. H. Notes on some south and central Australian mammals. *Trans. R. Soc. S. Aust.* **55**, 161–162 (1931).
- Vernes, K., Ingleby, S. & Eldridge, M. D. B. An overlooked, early record of the desert rat-kangaroo (*Caloprymnus campestris*) from Lake Killalpaninna, South Australia. *Aust. Mammal.* **42**, 223–225 (2020).
- Smith, M. J. & Johnson, P. M. In *The Mammals of Australia* 295–296 (New Holland Publishers, 2008).
- Robinson, T. & Forrest, T. A possible sighting of the Desert Rat-kangaroo or *Oolacunta* (*Caloprymnus campestris*) on the Peake Station, South Australia. *S. Aust. Nat.* **86**, 63–75 (2012).
- Carr, S. G. & Robinson, A. C. The present status and distribution of the desert rat-kangaroo *Caloprymnus campestris* (Marsupialia: Potoroidae). *S. Aust. Nat.* **72**, 4–27 (1997).
- Kear, B. P. & Cooke, B. N. A review of macropodoid systematics with the inclusion of a new family. *Mem. Assoc. Australas. Palaeontol.* **25**, 83–101 (2001).
- Westerman, M., Loke, S. & Springer, M. S. Molecular phylogenetic relationships of two extinct potoroid marsupials, *Potoropus platyops* and *Caloprymnus campestris* (Potoroinae: Marsupialia). *Mol. Phylogenet. Evol.* **31**, 476–485 (2004).
- Westerman, M., Loke, S. & Tan, M.-H. Molecular relationships of the red-bellied Dasyure *Phascosorex doriae* a rare marsupial from western New Guinea. *Aust. Mammal.* <https://doi.org/10.1071/AM21011> (2021).
- Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
- Peng, Y., Leung, H. C., Yiu, S.-M. & Chin, F. Y. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**, 1420–1428 (2012).
- Bernt, M. *et al.* MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol. Phylogenet. Evol.* **69**, 313–319 (2013).
- Phillips, M. J., Haouchar, D., Pratt, R. C., Gibb, G. C. & Bunce, M. Inferring kangaroo phylogeny from incongruent nuclear and mitochondrial genes. *PLoS ONE* **8**, e57745 (2013).
- Posada, D. jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* **25**, 1253–1256 (2008).
- Stamatakis, A. RAXML-VI-HPC: maximum likelihood based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688–2690 (2006).
- Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574 (2003).
- Bouckaert, R. *et al.* BEAST 2: A software platform for Bayesian evolutionary analysis. *PLoS Comput. Biol.* **10**, e1003537 (2014).
- Matzke, N. J. Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. *Syst. Biol.* **63**, 951–970 (2014).
- Ali, S. S., Yu, Y., Pfosser, M. & Wetschnig, W. Inferences of biogeographical histories within subfamily Hyacinthoideae using S-DIVA and Bayesian binary MCMC analysis implemented in RASP (Reconstruct Ancestral State in Phylogenies). *Ann. Bot.* **109**, 95–107 (2012).
- Yu, Y., Harris, A. J., Blair, C. & He, X. J. RASP (Reconstruct Ancestral State in Phylogenies): a tool for historical biogeography. *Mol. Phylogenet. Evol.* **87**, 46–49 (2015).
- Yu, Y., Blair, C. & He, X. J. RASP 4: ancestral state reconstruction tool for multiple genes and characters. *Mol. Biol. Evol.* **37**, 604–606 (2020).
- Martin, H. A. Cenozoic climatic change and the development of arid vegetation in Australia. *J. Arid. Environ.* **66**, 533–563 (2006).
- Ree, R. H. & Sanmartín, I. Conceptual and statistical problems with the DEC+J model of founder-event speciation and its comparison with DEC via model selection. *J. Biogeogr.* **5**, 741–749 (2018).
- Nilsson, M., Gullberg, A., Spotorno, A. E., Arnason, U. & Janke, A. Radiation of extant marsupials after the K/T boundary: Evidence from complete mitochondrial genomes. *J. Mol. Evol.* **57**(Suppl), S3–S12 (2003).
- Prada, C. F. & Boore, J. L. Gene annotation errors are common in the mammalian mitochondrial genomes database. *BMC Genomics* **20**, 73 (2019).



45. Janke, A. & Paabo, S. Editing of a tRNA anticodon in marsupial mitochondria changes its codon recognition. *Nucleic Acids Res.* **21**, 1523–1525 (1993).
46. Meredith, R. W., Westerman, M. & Springer, M. S. A phylogeny and timescale for the living genera of kangaroos and kin (Macropodiformes: Marsupialia) based on nuclear DNA sequences. *Aust. J. Zool.* **56**, 395–410 (2008).
47. Llamas, B. *et al.* Late Pleistocene Australian marsupial DNA clarifies the affinities of extinct megafaunal kangaroos and wallabies. *Mol. Biol. Evol.* **32**, 574–584 (2015).
48. Brennan, I. G. Incorporating fossil uncertainty is essential to macroevolutionary inferences: Combined evidence dating of Australian macropods. *bioRxiv*. <https://doi.org/10.1101/772558> (2019).
49. Cascini, M., Mitchell, K. J., Cooper, A. & Phillips, M. Reconstructing the evolution of giant extinct kangaroos: comparing the utility of DNA, morphology and total evidence. *Syst. Biol.* **68**, 520–537 (2019).
50. Beck, R. M. D., Voss, R. S. & Jansa, S. A. Craniodental morphology and phylogeny of marsupials. *PaleoXiv* <https://doi.org/10.31233/osf.io/rph78> (2021).
51. Smith, R. H. *et al.* Germline viral “fossils” guide *in silico* reconstruction of a mid-Cenozoic era marsupial adeno-associated virus. *Sci. Rep.* **6**, 28965 (2016).
52. Bensley, B. A. On the evolution of the Australian Marsupialia: With remarks on the relationships of the marsupials in general. *Trans. Linn. Soc. Lond. Zool.* **9**, 83–217 (1903).
53. Flannery, T. F. In *Kangaroos, Wallabies and Rat-Kangaroos* 1–46 (Surrey Beatty and Sons, 1989).
54. Priddleaux, G. J. *Borongaboodie hatcheri* gen. et sp. Nov. a very large bettong (Marsupialia: Macropodoidea) from the Pleistocene of southwestern Australia. *Rec. West. Aust. Mus* **57**, 317–329 (1999).
55. Swofford, D. *PAUP\** 4.0b10. (Sinauer Associates, Inc., 2002).
56. Frankham, G. J., Handasyde, K. A. & Eldridge, M. D. B. Novel insights into the relationships of the endangered marsupial genus. *Potorous. Mol. Phylogenet. Evol.* **64**, 592–602 (2012).
57. Metzger, C. A. & Retallack, G. J. Paleosol record of Neogene climate change in the Australian outback. *Aust. J. Earth Sci.* **57**, 871–885 (2010).
58. Pound, M. J., Haywood, A. M., Salzmann, U. & Riding, J. B. Global vegetation dynamics and latitudinal temperature gradients during the mid to late miocene (15.97–5.33 Ma). *Earth Sci. Rev.* **112**, 1–22 (2012).
59. McGowran, B. & Li, Q. The Miocene oscillation in southern Australia. *Rec. South Aust. Mus.* **27**, 197–212 (1994).
60. Mao, X. & Retallack, G. Late miocene drying of central Australia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **514**, 292–304 (2019).
61. Seersholm, F. V. *et al.* Ancient DNA from bulk bone reveals past genetic diversity of vertebrate fauna on Kangaroo Island. *Australia. Quat. Sci. Rev.* **262**, 106962 (2021).
62. Finlayson, H. H. Preliminary description of two new forms of *Bettongia* (Marsupialia). *Ann. Mag. Nat. Hist.* **10**, 552–554 (1957).
63. McDowell, M. C. *et al.* Morphological and molecular evidence supports specific recognition of the recently extinct *Bettongia anhydra* (Marsupialia: Macropodidae). *J. Mammal.* **96**, 287–296 (2015).
64. Haouchar, D. *et al.* Ancient DNA reveals complexity in the evolution history and taxonomy of the endangered Australian brush-tailed bettongs (*Bettongia*: Marsupialia: Macropodidae: Potoroinae). *Biodivers. Conserv.* **25**, 2907–2927 (2016).
65. Taylor, R. J. Habitat requirements of the Tasmanian bettong (*Bettongia gaimardi*), a mycophagous marsupial. *Wildl. Res.* **20**, 699–710 (1993).
66. Laurance, W. F. A distributional survey and habitat model for the endangered Northern bettong *Bettongia tropica* in tropical Queensland. *Biol. Conserv.* **82**, 47–60 (1997).
67. Smith, M. *et al.* Conserving the endangered woylie (*Bettongia penicillata ogilbyi*): Establishing a semi-arid population within a fenced safe haven. *Ecol. Manag. Restor.* **21**, 108–114 (2020).
68. Byrne, M. *et al.* Decline of a biome: evolution, contraction, fragmentation, extinction and invasion of the Australian mesic zone biota. *J. Biogeogr.* **38**, 1635–1656 (2011).
69. Sharman, G. B. The mitotic chromosomes of marsupials and their bearing on taxonomy and phylogeny. *Aust. J. Zool.* **9**, 38–60 (1961).
70. Sharman, G. B., Murtagh, C. E., Johnson, P. M. & Weaver, C. M. The chromosomes of a rat-kangaroo attributable to *Bettongia tropica* (Marsupialia: Macropodidae). *Aust. J. Zool.* **28**, 59–63 (1980).
71. Rens, W., O’Brien, P. C. M., Fairclough, H., Graves, J. A. M. & Fergusson-Smith, M. A. Reversal and convergence in marsupial chromosome evolution. *Cytogenet. Genome Res.* **102**, 282–290 (2003).
72. Macqueen, P., Seddon, J. M., Austin, J. J., Hamilton, S. & Goldizen, A. W. Phylogenetics of the pademelons (Macropodidae: *Thylomys*) and historical biogeography of the Australo-Papuan region. *Mol. Phylogenet. Evol.* **57**, 1134–1148 (2010).
73. Eldridge, M. D. B. *et al.* Phylogenetic analysis of the tree-kangaroos (*Dendrolagus*) reveals multiple divergent lineages within New Guinea. *Mol. Phylogenet. Evol.* **127**, 589–599 (2018).
74. Stuu, J.-B.W. *et al.* A 5.3-million-year history of monsoonal precipitation in Northwestern Australia. *Geophys. Res. Lett.* **46**, 6946–6954 (2019).
75. Jackson, S. & Groves, C. *Taxonomy of Australian Mammals* (CSIRO Publishing, 2015).
76. McNamara, J. A. Some smaller macropod fossils from South Australia. *Proc. Linn. Soc. NSW* **117**, 97–106 (1997).
77. Burbidge, A. A., Short, J. & Fuller, P. J. Relict *Bettongia lesueur* warrens in Western Australian deserts. *Aust. Zool.* **31**, 97–103 (2007).
78. Burbidge, A. A. & Short, J. C. In *The Mammals of Australia* 288–290 (New Holland Publishers, 2008).
79. De Torres, P. J. & Start, A. N. In *The Mammals of Australia* 291–292 (New Holland Publishers, 2008).
80. Letnic, M. In *Animals of Arid Australia: Out on Their Own?* 65–75 (Royal Society of NSW, 2007).
81. Gould, J. *The Mammals of Australia* (1845–1863).
82. Den Boer, W., Campione, N. E. & Kear, B. P. Climbing adaptations, locomotory disparity and ecological convergence in ancient stem ‘kangaroos’. *R. Soc. Open Sci.* **6**, 18167 (2019).

## Acknowledgements

We dedicate this paper to the late Joan Dixon (1937–2019) of NMV, who was a distinguished researcher on Australian mammals and provided access to the liver sample (NMV C8981) of *Caloprymnus campestris*. Constructive comments from the Editor, Robin Beck (University of Salford) and two anonymous reviewers improved our manuscript.

## Author contributions

M.W. conceived the study, collected and analysed the data, and wrote the paper. S.L. and M.H.T. analysed the data. B.P.K. analysed the data and wrote the paper.

## Funding

Open access funding provided by Uppsala University. Aspects of this work were supported by a Swedish Research Council Young Researcher grant (2011–3637), and a Swedish Research Council Project grant (2020-3423) to B.P.K.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-09568-0>.

**Correspondence** and requests for materials should be addressed to M.W. or B.P.K.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022