

# Quercetin inhibits TNF- $\alpha$ induced HUVECs apoptosis and inflammation via downregulating NF- $\kappa$ B and AP-1 signaling pathway in vitro

Tielong Chen, PhD<sup>a,\*</sup>, Xudong Zhang, MD<sup>a,b</sup>, Guangli Zhu, MD<sup>a</sup>, Hongfei Liu, MD<sup>a</sup>, Jinru Chen, MD<sup>a</sup>, Yu Wang, MD<sup>a</sup>, Xiaolong He, MD<sup>a</sup>

## Abstract

**Background:** Quercetin, a major flavonol, widely exists in plantage, which has been reported to have an anti-apoptosis and anti-inflammation effects on vascular endothelial cells, but its underlying molecular mechanisms remain unclear.

**Objective:** The aim of this study was to investigate the mechanisms of how quercetin inhibits tumor necrosis factor alpha (TNF- $\alpha$ ) induced human umbilical vein endothelial cells (HUVECs) apoptosis and inflammation.

**Methods and Results:** HUVECs were preconditioned with quercetin for 18 hours, and subsequently treated with TNF- $\alpha$  for 6 hours to induce apoptosis. The expression of intercellular cell adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), E-selectin,  $\beta$ -actin mRNA was then detected by RT-PCR. Flow cytometry was used to estimate the apoptosis rates, and the expression of activator protein 1 (AP-1) and nuclear factor kappa B (NF- $\kappa$ B) was measured by Western blot. TNF- $\alpha$  induced elevated apoptosis rates and upregulation of VCAM-1, ICAM-1, and E-selectin were meaningfully reduced in HUVECs by pretreatment with quercetin. In addition, quercetin also inhibited the activation of AP-1 and NF- $\kappa$ B.

**Conclusion:** Results indicate that quercetin could suppress TNF- $\alpha$  induced apoptosis and inflammation by blocking NF- $\kappa$ B and AP-1 signaling pathway in HUVECs, which might be one of the underlying mechanisms in treatment of coronary heart disease.

**Abbreviations:** AP-1 = activator protein 1, CHD = coronary heart disease, FasL = Fas ligand, HUVECs = human umbilical vein endothelial cells, ICAM-1 = intercellular cell adhesion molecule-1, NF- $\kappa$ B = nuclear factor kappa B, PVDF = polyvinylidene difluoride, TNF- $\alpha$  = tumor necrosis factor alpha, TRAIL = TNF-related apoptosis-inducing ligand, VCAM-1 = vascular cell adhesion molecule 1.

**Keywords:** AP-1, apoptosis, cell adhesion molecules, HUVECs, NF- $\kappa$ B, quercetin

Editor: Wen-Jun Tu.

TC and XZ contributed equally to this study.

This study was supported by the Hangzhou Health Science and Technology Project (2017Z10) and the Hangzhou Science and Technology Bureau Project (20150733Q57, 20160533B63).

The authors have no conflict of interest.

The datasets generated during and/or analyzed during the current study are not publicly available, but are available from the corresponding author on reasonable request.

<sup>a</sup> Department of Cardiology, Hangzhou Hospital of Traditional Chinese Medicine, Hangzhou, Zhejiang, China, <sup>b</sup> Department of Cardiology, Hangzhou Dingqiao's Hospital, Hangzhou, Zhejiang, China.

\* Correspondence: Tielong Chen, Department of Cardiology, Hangzhou Hospital of Traditional Chinese Medicine, No. 453 Stadium Road, Hangzhou 310007, Zhejiang, China (e-mail: ctjppp@foxmail.com).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Chen T, Zhang X, Zhu G, Liu H, Chen J, Wang Y, He X. Quercetin inhibits TNF- $\alpha$  induced HUVECs apoptosis and inflammation via downregulating NF- $\kappa$ B and AP-1 signaling pathway in vitro. *Medicine* 2020;99:38 (e22241).

Received: 2 June 2020 / Received in final form: 10 August 2020 / Accepted: 19 August 2020

<http://dx.doi.org/10.1097/MD.0000000000002241>

## 1. Introduction

Coronary heart disease (CHD), lifetime of which at age 40 years is 1 in 3 for women and 1 in 2 for men in the United States, remains the major cause of death in developed countries.<sup>[1,2]</sup> The underlying mechanism of CHD is initiated by inflammatory processes in the endothelium of the vessel wall.

Inflammation plays a key role in the development of atherosclerosis.<sup>[3,4]</sup> Tumor necrosis factor-alpha (TNF- $\alpha$ ) is one of cytokines involved in systemic inflammation, which could alter leukocyte adhesion molecule expression on cultured endothelial cells such as human umbilical vein endothelial cells (HUVECs).<sup>[5]</sup> Cellular adhesion molecule responding to pathophysiological stimuli can influence the progress of atherosclerosis by mediating the interaction between blood cells (such as leukocytes) and endothelial cells.<sup>[6,7]</sup> Inflammation could be promoted by TNF- $\alpha$  via upregulating expression of adhesion molecules, such as intercellular cell adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), and E-selectin.<sup>[8]</sup>

In addition, TNF- $\alpha$  could activate NF- $\kappa$ B, which could mediate the transcription of a vast array of proteins involved in cell survival and inflammatory response, and anti-apoptotic factors. It has been proven that inhibiting the NF- $\kappa$ B and AP-1 pathway could exert anti-inflammatory effects.<sup>[9,10]</sup>

Flavonoids, the polyphenolic compounds ubiquitous in the plant kingdom inversely correlated with CVD, have the protective effects in atherosclerosis.<sup>[11–13]</sup> One of the most

abundant polyphenolic compounds is quercetin, which has anti-atherosclerotic effects in human and animals,<sup>[14,15]</sup> and has been shown to have the ability to fight against inflammation.<sup>[16,17]</sup>

Studies have shown that quercetin could attenuate cell adhesion molecules expression, such as ICAM-1, VCAM-1, and E-selectin.<sup>[18–20]</sup> It has been proven that the mechanism of leukocyte adhesion in inflammatory lesion might be upregulating the expression of ICAM-1 and VCAM-1 via activation of NF- $\kappa$ B and AP-1.<sup>[21,22]</sup> The protective mechanism against inflammation and atherosclerosis of quercetin might be the inhibitory effect on NF- $\kappa$ B activation and AP-1.<sup>[23,24]</sup> We hypothesize that inhibiting NF- $\kappa$ B and AP-1 activation might be an anti-inflammatory way in the progress of CHD.

The purpose of the present investigation was to study the association between quercetin and adhesion molecules expression. HUVECs were treated with TNF- $\alpha$  to mimic the inflammatory situation. Flow cytometry assay was used to measure apoptosis rates of HUVECs presence or absence of quercetin. PT-PCR and Western blot were used to detect the expression of ICAM-1, VCAM-1, and E-selectin in HUVECs presence or absence of quercetin. We also detected the expression of NF- $\kappa$ B and AP-1 by Annexin-V staining.

## 2. Materials and methods

### 2.1. Materials

HUVECs were obtained from Chinese Academy of Sciences (Shanghai, China). Quercetin was provided by the National Institute for the Control of Pharmaceutical and Biological Products (Beijing, China); our previous study, which was not published, found that quercetin could reverse upregulation of ICAM-1, VCAM-1 and E-selectin expression induced by TNF- $\alpha$ , and the effects were not in a dose-dependent manner, the optimal concentration of quercetin was 30  $\mu$ g/mL. In the present study, quercetin was diluted at the concentration of 30  $\mu$ g/mL. Antibodies of hICAM-1, hVCAM-1, E-selectin, NF- $\kappa$ B, and AP-1 were obtained from R&D systems (Minneapolis, Minnesota, USA). The antibody of  $\beta$ -actin was provided by Santa Cruz Biotechnology (Dallas, Texas, USA). This study was not conducted directly on humans, so ethical approval and informed consent statement were not necessary.

### 2.2. Primers

Primers were designed to assess gene expression of adhesion molecules by RT-PCR analysis. The primer sequences for VCAM-1 were 5' -AGT GGT GGC CTC CTG AAT GG -3' (upstream) and 5' -CTG TGT CTC CTG TCT CCG CT -3' (downstream), yielding a 700-bp fragment. Primer sequences for ICAM were 5' -TAT GGC AAC GAC TCC TTC T -3' (upstream) and 5' -CAT TCA GCG TCA CCT TGG -3' (downstream), yielding a 220-bp fragment. Primer sequences for E-selectin were 5' -CAA GAA GAA GCT TGC CCT ATG -3' (upstream) and 5' -ACT TGA GTC CACT GA AGC CA -3' (downstream), yielding a 128-bp fragment. Sequences for  $\beta$ -actin were 5'-CTG GCA CCA CAC CTT CTA CAA TGA GC -3' (upstream) and 5'-GAG GAT CTT CAT GAG GTA GTC AGT C -3' (downstream), yielding a 325-bp fragment.

### 2.3. HUVECs culture

The HUVECs were supplemented with antibiotics in a humidified 37°C incubator and 10% fetal bovine serum in Dulbecco's

modified Eagle's medium (DMEM; Gibco (California, USA)) with an atmosphere of 5% CO<sub>2</sub> and 95% air. The HUVECs were seeded in 24- or 96-well plates in DMEM, and subsequently randomly divided into 3 groups: (control group: HUVECs were culture with DMEM; TNF- $\alpha$  group: HUVECs were cultured with TNF- $\alpha$  for 6 hours at the dose of 200 ng/mL; quercetin group: HUVECs were cultured with quercetin for 18 hours at the dose of 30  $\mu$ g/mL and then cocultured with TNF- $\alpha$  for 6 hours at the dose of 200 ng/mL). Before treatments, cells were washed with PBS to remove TNF- $\alpha$  or quercetin.

### 2.4. Annexin V assay

We decanted supernatant from a flask of the control group, washed flask twice in warm PBS to remove serum, then detached cells in trypsin without EDTA, washed the flask twice in PBS. Five hundred microliter Annexin VB Binding Buffer was added to suspend the cells, and the cells were subsequently stained with FITC and PIN. We analyzed with flow cytometry to estimate the apoptosis rates. The same progress was done in the TNF- $\alpha$  group and the quercetin group.

### 2.5. RNA extractions and RT-PCR analysis

HUVECs were cultured with TNF- $\alpha$  presence or absence of quercetin. Total RNA from HUVECs was then extracted and analyzed by RT-PCR using a kit. Twenty microliter reaction mixture contained 4- $\mu$ L dNTP Mix, 2- $\mu$ L Primer Mix, 4- $\mu$ L 5xRT buffer, 2- $\mu$ L DTT, 1- $\mu$ L HiFi-MMLV, and Rnase-Free Water. RT was performed at 42°C for 50 minutes and then inactivated at 70°C for 15 minutes. PCR reaction was performed in 50  $\mu$ L containing 2xTaq MasterMix, 10  $\mu$ M of Forward Primer, 10  $\mu$ M of Reverse Primer, Template DNA, and Rnase-Free Water. Basic PCR consists of 3 steps: thermal denaturation of the target DNA, primer annealing of synthetic oligonucleotide primers, and extension of the annealed primers by a DNA polymerase. RT-PCR products were then resolved in 1.7% agarose gels and stained with ethidium bromide. The expression of VCAM-1, ICAM-1, E-selectin,  $\beta$ -actin mRNA was then detected with the electrophoretic analysis and the semi-quantitative analysis.

### 2.6. Western blot analysis

Proteins from HUVECs were separated according to the molecular weight of the proteins using 10% SDS-PAGE gel. The separated gel was transferred to a polyvinylidene difluoride (PVDF) membrane. The membrane was then blocked with 5% non-fat milk powder (blocking agent) in tris buffered saline (TBS). The membrane was then washed three times with TBS containing 0.1% Tween 20 and incubated with a protein-specific antibody at room temperature followed by a secondary antibody. Antibodies to hVCAM-1, hICAM-1, E-selectin, nuclear factor kappa-B (NF- $\kappa$ B), AP-1,  $\beta$ -actin were used as follows. The membranes were rinsed after the secondary antibody incubation, and the bound antibodies were then detected using enhanced chemiluminescence (ECL; Santa Cruz Biotechnology followed by autoradiography. We used the Image pro plus 5 software to semi-quantify protein in every lane.

### 2.7. Flow cytometry

The flow cytometry was used to measure the expression of AP-1 and NF- $\kappa$ B in the 3 groups. We collected cells and suspend them

with the PBS and centrifuge for 5 minutes at the speed of 1500 rpm after added with fixation/permeabilization solutions for 20 minutes, and then decanted the supernatant and added 2 mL Perm/Wash Butter to wash the cells. The NF- $\kappa$ B-PE or AP-1-FITC antibodies were then incubated with the cells for 30 minutes. We finally washed cells with 2 mL Perm/Wash Butter and collected the cells by centrifuge (2000 rpm, 5 minutes), decanted the supernatant, and added 500  $\mu$ L PBS for concussion, and analyzed on FACS can flow cytometer using BD CellQuest Pro software (version 5.1; BD Biosciences, Franklin Lakes, NJ, USA).

## 2.8. Statistical analysis


Data are presented as mean  $\pm$  standard deviation. Multiple comparisons between the groups were performed by 1-way

analysis of variance (ANOVA). *P* values of .05 or less were considered to indicate statistical significance for all subgroup and interaction analyses.

## 3. Results

### 3.1. Quercetin decreases apoptosis rates of endothelial cells

To determine how quercetin reverses TNF- $\alpha$  induced down-regulation of apoptosis rates, we used flow cytometry assay to measure apoptosis rates of HUVECs. Apoptosis rates of HUVECs were increased in TNF- $\alpha$  group compared with control group ( $10.005 \pm 0.903$  vs  $3.602 \pm 0.833\%$ ,  $P < .01$ ), and were decreased with quercetin pretreatment compared with TNF- $\alpha$  group ( $7.154 \pm 0.862$  vs  $10.005 \pm 0.903$ ,  $P < .01$ ) in Figure 1D. In



**Figure 1.** Quercetin decreased TNF- $\alpha$  induced upregulation of apoptosis rates. (A) Apoptosis rates in the control group were measured by flow cytometry assay; (B) Apoptosis rates in the TNF- $\alpha$  group were measured by flow cytometry assay; (C) Apoptosis rates in the quercetin group were measured by flow cytometry assay. (D) Quantitation of apoptosis rates in HUVECs. Results were expressed as mean  $\pm$  standard deviation. \* $P < .01$  vs TNF- $\alpha$  group.

this study, we could see that quercetin had the protective effects on apoptosis of HUVECs induced by TNF- $\alpha$ .

### 3.2. Effects of quercetin on VCAM-1, ICAM-1, E-selectin mRNA expression

Cell adhesion molecules such as VCAM-1, ICAM-1, and E-selectin could mediate the process of leukocytes adhering to endothelial cells, which might be downregulated by quercetin. Figure 2A showed the expression of adhesion molecules detected by RT-PCR. As shown in Figure 2B, the expressions of VCAM-1/ $\beta$ -actin, ICAM-1/ $\beta$ -actin, and E-selectin/ $\beta$ -actin in TNF- $\alpha$  group compared with control group were  $0.982 \pm 0.047$  vs  $0.422 \pm 0.31$  ( $P < .01$ ),  $1.038 \pm 0.024$  vs  $0.314 \pm 0.378$  ( $P < .01$ ), and  $1.09 \pm 0.051$  vs  $0.75 \pm 0.068$  ( $P < .01$ ), respectively. Compared with the TNF- $\alpha$  group, the expressions of VCAM-1/ $\beta$ -actin, ICAM-1/ $\beta$ -actin, and E-selectin/ $\beta$ -actin in the quercetin group were  $0.338 \pm 0.0562$  vs  $0.982 \pm 0.047$  ( $P < .01$ ),  $0.412 \pm 0.061$  vs  $1.038 \pm 0.024$  ( $P < .01$ ), and  $0.812 \pm 0.060$  vs  $1.09 \pm 0.051$  ( $P < .01$ ), respectively. It could be speculated that pretreatment with quercetin could downregulate the upregulation of these adhesion molecules induced by TNF- $\alpha$ .

### 3.3. Effects of quercetin on hVCAM-1, hICAM-1, E-selectin expression

Quercetin could reverse the upregulation of VCAM-1, ICAM-1, and E-selectin induced by TNF- $\alpha$ . Figure 3 showed the expression of hVCAM-1, hICAM-1, and E-selectin detected by Western blot. As shown in Figure 3, the expression of hVCAM-1/ $\beta$ -actin, hICAM-1/ $\beta$ -actin, and E-Selectin/ $\beta$ -actin in the TNF- $\alpha$  group compared with the control group were  $1.122 \pm 0.048$  vs  $0.093 \pm 0.027$  ( $P < .01$ ),  $0.958 \pm 0.031$  vs  $0.059 \pm 0.019$  ( $P < .01$ ), and  $1.120 \pm 0.060$  vs  $0.106 \pm 0.016$  ( $P < .01$ ), respectively. Compared with the TNF- $\alpha$  group, the expression of hVCAM-1/ $\beta$ -actin, hICAM-1/ $\beta$ -actin, and E-selectin/ $\beta$ -actin in the quercetin group

were  $0.702 \pm 0.038$  vs  $1.122 \pm 0.048$  ( $P < .01$ ),  $0.396 \pm 0.032$  vs  $0.958 \pm 0.031$  ( $P < .01$ ), and  $0.743 \pm 0.036$  vs  $1.120 \pm 0.060$  ( $P < .01$ ), respectively. It could be speculated that pretreatment with quercetin could reverse the upregulation of these adhesion molecules induced by TNF- $\alpha$ .


### 3.4. Quercetin inhibited NF- $\kappa$ B and AP-1 pathway activation

NF- $\kappa$ B and AP-1 play important roles in the progress of inflammation. To evaluate the effects, we detected the expression of NF- $\kappa$ B and AP-1 by Annexin-V staining. As shown in Figure 4, NF- $\kappa$ B/ $\beta$ -actin expression was increased in the TNF- $\alpha$  group compared with the control group ( $45.41 \pm 7.037$  vs  $25.03 \pm 3.472$ ,  $P < .01$ ), whereas pretreatment with quercetin decreased the expression of NF- $\kappa$ B/ $\beta$ -actin compared with TNF- $\alpha$  group ( $30.21 \pm 6.30$  vs  $45.41 \pm 7.037$ ,  $P < .01$ ). Figure 5 showed that AP-1/ $\beta$ -actin expression was increased in the TNF- $\alpha$  group compared with the control group ( $49.19 \pm 7.51$  vs  $28.75 \pm 3.68$ ,  $P < .01$ ), and the expression was decreased in the quercetin group compared with the TNF- $\alpha$  group ( $30.16 \pm 4.02$  vs  $49.19 \pm 7.51$ ,  $P < .01$ ). In summary, quercetin might have the ability of reversing upregulation of NF- $\kappa$ B and AP-1 induced by TNF- $\alpha$ , which might be one of the mechanisms of anti-inflammation.


## 4. Discussion

In this study, TNF- $\alpha$  was used to induce cell apoptosis and upregulate expression of adhesion molecules, which contributed to the inflammation of HUVECs. We investigated the underlying mechanism of quercetin in the development of inflammation and the progress of cell apoptosis.

Caspases are responsible for cells into apoptotic bodies. Apoptosis inducers [Fas ligand (FasL), TNF- $\alpha$ ] activate the initiator caspases such as caspase-8 and sequentially initiate the apoptotic cascade.<sup>[25,26]</sup> There are other apoptosis-related




**Figure 2.** Effects of quercetin on the expression of VCAM-1, ICAM-1, E-selectin mRNA. (A) Total RNA from control, TNF- $\alpha$  and quercetin groups were then extracted and analyzed by RT-PCR. (B) Quantitation of VCAM-1, ICAM-1, and E-selectin of control, TNF- $\alpha$ , and quercetin groups, which were detected by RT-PCR. Results were expressed as mean  $\pm$  standard deviation. \* $P < .01$ , <sup>†</sup> $P < .01$ , <sup>‡</sup> $P < 0.01$ , vs TNF- $\alpha$  group.




**Figure 3.** Effects of quercetin on the expression of hVCAM-1, hICAM-1, E-selectin. 25  $\mu$ g HUVECs lysates in control, TNF- $\alpha$  and quercetin groups were loaded onto an SDS-polyacrylamide gel, and hVCAM-1 (A), hICAM-1 (B), and E-selectin (C) were detected by Western blotting. Equal loading of protein was loaded with antibodies against  $\beta$ -actin onto the three groups. Expressions of hVCAM-1, hICAM-1, and E-selectin were detected by Western blotting. Quantitation of hVCAM-1, hICAM-1, and E-selectin expression of HUVECs were detected by Annexin-V. Results were expressed as mean  $\pm$  standard deviation. \* $P$  < .01 vs TNF- $\alpha$  group.

proteins such as an apoptosis-inducing ligand as a TNF-related apoptosis-inducing ligand (TRAIL), and FasL and their receptors. TNF- $\alpha$  has an extra effect on Fas activation, which activates Fas signaling pathway while excited. The results revealed that

apoptosis rate was increased by TNF- $\alpha$ , and was reduced by pretreatment with quercetin. With our present study, quercetin was found to be capable of protecting HUVECs against apoptosis.



**Figure 4.** Quercetin inhibited NF- $\kappa$ B pathway activation. (A–C) Expressions of NF- $\kappa$ B were detected by Annexin-V staining in control group, TNF- $\alpha$  group, and quercetin group, respectively. (D) NF- $\kappa$ B from HUVECs in control, TNF- $\alpha$  and quercetin groups were detected by Western blot. Equal loading of proteins was confirmed by reprobing blots with antibodies against  $\beta$ -actin. (E) Quantitation of the expression of NF- $\kappa$ B was measured by Annexin-V. \* $P$  < .01 vs TNF- $\alpha$  group. Results were expressed as mean  $\pm$  standard deviation.



**Figure 5.** Quercetin inhibited AP-1 pathway activation. (A–C) Expressions of AP-1 were detected by Annexin-V staining in control group, TNF- $\alpha$  group, and quercetin group, respectively. (D) AP-1 from HUVECs in control, TNF- $\alpha$  and quercetin groups were detected by Western blot. Equal loading of proteins was confirmed by reprobing blots with antibodies against  $\beta$ -actin. (E) Quantitation of the expression of AP-1 was measured by Annexin-V. \* $P$  < .01 vs TNF- $\alpha$  group. Results were expressed as mean  $\pm$  standard deviation.

In addition, as a cell signaling protein involved in systemic inflammation, TNF- $\alpha$  could promote expression of adhesion molecules (ICAM-1, VCAM-1, and E-selectin) in HUVECs. Inflammation plays a critical role in the progress of atherosclerosis. The molecules associated with cell adhesion molecules are proteins located on the cell surface helps cells stick to each other and to their surroundings. When stimulated by TNF- $\alpha$ , leukocytes adhere to endothelium via ICAM-1, an endothelial- and leukocyte-associated transmembrane protein, and then transmigrate to the inside tissue.<sup>[27]</sup> E-selectin and VCAM-1 could also be found increased via induction by TNF- $\alpha$ , and helped leukocyte roll on microvascular endothelium.<sup>[28]</sup> Studies showed a different train of thoughts in adhesion mechanisms of atherosclerosis, such as activation of NF- $\kappa$ B and IFN regulatory factor-1, and the downregulation of CCL8 and CXCL10 chemokines.<sup>[29–31]</sup> Expression of ICAM-1, VCAM-1, and E-selectin could be upregulated via modulated by AP-1 and NF- $\kappa$ B.<sup>[21,32]</sup>

NF- $\kappa$ B, a protein complex that controls transcription of DNA, could combine with the target site of  $\kappa$ B and promote transcription. When impulsed by oxygen stress, NF- $\kappa$ B regulates the expression of proinflammatory cytokines, transcription factors, and cell adhesion molecules, which play a key role in the development of microcirculatory disturbance.<sup>[33]</sup> Activation of NF- $\kappa$ B induced by TNF- $\alpha$  could upregulate the expression of

ICAM-1, VCAM-1, and E-selectin, which could integrate with leukocytes, helping leukocytes stick to the vascular wall.

Special attention has been paid to the study on AP-1, another transcription factor, as a key factor in the development of cardiovascular disease. It is a heterodimeric protein composed of proteins belonging to the ATF and JDP, c-Jun, c-Fos families, which could modulate the expression of gene in response to stimuli, such as TNF- $\alpha$ . One of the mechanisms is the MAPK signaling pathway, after being phosphorylated, the c-Fos and c-jun proteins will be activated, following the activation of AP-1.

Quercetin, a major bioflavonoid widely present in fruits and vegetables, has been found to fight against inflammation via suppressing I- $\kappa$ B-phosphorylation, NF- $\kappa$ B translocation, AP-1, and reporter gene transcription.<sup>[24,34]</sup> Studies has shown that quercetin could exert its biological action in other fields via modulating JNK, AP-1, NF- $\kappa$ B signaling pathway.<sup>[35–37]</sup> TNF- $\alpha$  is a key proinflammatory cytokine that acts to provoke inflammation, which induced by TNF- $\alpha$  was reduced with treatment of quercetin.<sup>[38,39]</sup>

Our results showed that the expression of NF- $\kappa$ B and AP-1 were significantly increased by the treatment of TNF- $\alpha$  in the HUVECs compared with the control group. Simultaneously, ICAM-1, VCAM-1, and E-selectin expression were also increased, and both the translation factors and cell adhesion molecules were downregulated after the pretreatment with

quercetin. Hence, the results of our research indicate that quercetin might reverse the upregulation of ICAM-1, VCAM-1, and E-selectin via inhibiting the activation of NF- $\kappa$ B and AP-1. These mechanisms could all relieve the progress of atherosclerosis. The deeper mechanisms of quercetin on atherosclerosis should be further studied.

### Author contributions

Designed the study: Tielong Chen. Performed the study: Xudong Zhang, Guangli Zhu, Hongfei Liu, and Jinru Chen. Analyzed the data: Yu Wang and Xiaolong He. Wrote the paper: Xudong Zhang. Revised the article: Tielong Chen.

### References

- Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart Disease and Stroke Statistics-2017 update: a report from the American Heart Association. *Circulation* 2017;135:e146–603.
- Nichols M, Townsend N, Scarborough P, et al. Cardiovascular disease in Europe 2014: epidemiological update. *Eur Heart J* 2014;35:2929–33.
- Pant S, Deshmukh A, Gurumurthy GS, et al. Inflammation and atherosclerosis: revisited. *J Cardiovasc Pharmacol Ther* 2014;19:170–8.
- Taleb S. Inflammation in atherosclerosis. *Arch Cardiovasc Dis* 2016;109:708–15.
- Zi-Yuan L, Wan-Cheng C, Yong-Hua L, et al. TNF- $\alpha$  enhances vascular cell adhesion molecule-1 expression in human bone marrow mesenchymal stem cells via the NF- $\kappa$ B, ERK and JNK signaling pathways. *Mol Med Rep* 2016;14:643–8.
- Schumski A, Winter C, Döring Y, et al. The ins and outs of myeloid cells in atherosclerosis. *J Innate Immun* 2018;10:479–86.
- Husain K, Hernandez W, Ansari RA, et al. Inflammation, oxidative stress and renin angiotensin system in atherosclerosis. *World J Biol Chem* 2015;6:209–17.
- Zhen-Quan J, Nallasamy P, Liu D, et al. Luteolin protects against vascular inflammation in mice and TNF- $\alpha$ -induced monocyte adhesion to endothelial cells via suppressing I $\kappa$ B/NF- $\kappa$ B signaling pathway. *J Nutr Biochem* 2015;26:293–302.
- Bin F, Sai-Hong D, Jian-Qiu G, et al. Pycnogenol attenuates the release of proinflammatory cytokines and expression of perilipin 2 in lipopolysaccharide-stimulated microglia in part via inhibition of NF- $\kappa$ B and AP-1 activation. *PLoS One* 2015;10:1–20.
- Intayoung P, Limtrakul P, Yodkeeree S. Antiinflammatory activities of crebanine by inhibition of NF- $\kappa$ B and AP-1 activation through suppressing MAPKs and Akt signaling in LPS-induced RAW264 7 macrophages. *Biol Pharm Bull* 2016;39:54–61.
- Terao J, Kawai Y, Murota K. Vegetable flavonoids and cardiovascular disease. *Asia Pac J Clin Nutr* 2008;17:291–3.
- Siasos G, Tousoulis D, Tsigkou V, et al. Flavonoids in atherosclerosis: an overview of their mechanisms of action. *Curr Med Chem* 2013;20:2641–60.
- Sedighi M, Bahmani M, Asgary S, et al. A review of plant-based compounds and medicinal plants effective on atherosclerosis. *J Res Med Sci* 2017;22:30.
- Kleemann R, Verschuren L, Morrison M, et al. Anti-inflammatory, anti-proliferative and anti-atherosclerotic effects of quercetin in human in vitro and in vivo models. *Atherosclerosis* 2011;218:44–52.
- Bondonno NP, Bondonno CP, Hodgson JM, et al. The efficacy of quercetin in cardiovascular health. *Curr Nutr Rep* 2015;4:290–303.
- Ching-Hsia H, Shih-Hung C, Pei-Ming C, et al. Quercetin is a potent anti-atherosclerotic compound by activation of SIRT1 signaling under oxLDL stimulation. *Mol Nutr Food Res* 2015;59:1905–17.
- Leyva-López N, Gutierrez-Grijalva E, Ambríz-Pérez D, et al. Flavonoids as cytokine modulators: a possible therapy for inflammation-related diseases. *Int J Mol Sci* 2016;17:921.
- Pi-Yu C, Ya-Ping H, Wen-Bin H. Inhibitory effect of purple sweet potato leaf extract and its components on cell adhesion and inflammatory response in human aortic endothelial cells. *Cell Adh Migr* 2013;7:237–45.
- Choi YJ, Arzuaga X, Klumper CT, et al. Quercetin blocks caveolae-dependent pro-inflammatory responses induced by co-planar PCBs. *Environ Int* 2010;36:931–4.
- Bin-Wu Y, Ting Y, Xing-Bo S, et al. Quercetin inhibits IL-1 beta-induced ICAM-1 expression in pulmonary epithelial cell line A549 through the MAPK pathways. *Mol Biol Rep* 2009;36:1825–32.
- Si-Min Y, Xu Z, Hai-Li Z, et al. Clematichinenoside inhibits VCAM-1 and ICAM-1 expression in TNF- $\alpha$ -treated endothelial cells via NADPH oxidase-dependent I $\kappa$ B kinase/NF- $\kappa$ B pathway. *Free Radic Biol Med* 2015;78:190–201.
- Youn GS, Kwon D-J, Ju SM, et al. Celastrol ameliorates HIV-1 Tat-induced inflammatory responses via NF- $\kappa$ B and AP-1 inhibition and heme oxygenase-1 induction in astrocytes. *Toxicol Appl Pharmacol* 2014;280:42–52.
- Vicentini FT, He T, Shao Y, et al. Quercetin inhibits UV irradiation-induced inflammatory cytokine production in primary human keratinocytes by suppressing NF- $\kappa$ B pathway. *J Dermatol Sci* 2011;61:162–8.
- Panicker SR, Sreenivas P, Babu MS, et al. Quercetin attenuates monocyte chemoattractant protein-1 gene expression in glucose primed aortic endothelial cells through NF- $\kappa$ B and AP-1. *Pharmacol Res* 2010;62:328–36.
- Gunther C, Martini E, Wittkopf N, et al. Caspase-8 regulates TNF- $\alpha$ -induced epithelial necroptosis and terminal ileitis. *Nature* 2011;477:335–9.
- Beaudouin J, Liesche C, Aschenbrenner S, et al. Caspase-8 cleaves its substrates from the plasma membrane upon CD95-induced apoptosis. *Cell Death Differ* 2013;20:599–610.
- Hai-Bo X, De-Jian W, Cheng P, et al. Rabbit sera containing compound danshen dripping pill attenuate leukocytes adhesion to TNF- $\alpha$ -activated human umbilical vein endothelial cells by suppressing endothelial ICAM-1 and VCAM-1 expression through NF- $\kappa$ B signaling pathway. *J Cardiovasc Pharmacol* 2014;63:323–32.
- Griffin GK, Newton G, Tarrío ML, et al. IL-17 and TNF- $\alpha$  sustain neutrophil recruitment during inflammation through synergistic effects on endothelial activation. *J Immunol* 2012;188:6287–99.
- Kim J, Kim Y, Kim HT, et al. TC1(C8orf4) is a novel endothelial inflammatory regulator enhancing NF- $\kappa$ B activity. *J Immunol* 2009;183:3996–4002.
- Dagia NM, Harii N, Meli AE, et al. Phenyl methimazole inhibits TNF- $\alpha$ -induced VCAM-1 expression in an IFN regulatory factor-1-dependent manner and reduces monocytic cell adhesion to endothelial cells. *J Immunol* 2004;173:2041–9.
- Secchiero P, Corallini F, Di Iasio MG, et al. TRAIL counteracts the proadhesive activity of inflammatory cytokines in endothelial cells by down-modulating CCL8 and CXCL10 chemokine expression and release. *Blood* 2005;105:3413–9.
- Liang Z, Simard MJ, Huot J. Endothelial microRNAs regulating the NF- $\kappa$ B pathway and cell adhesion molecules during inflammation. *FASEB J* 2018;32:4070–84.
- Mauro C, Zazzeroni F, Papa S, et al. The NF- $\kappa$ B transcription factor pathway as a therapeutic target in cancer: methods for detection of NF- $\kappa$ B activity. *Methods Mol Biol* 2009;512:169–207.
- Endale M, Park SC, Kim S, et al. Quercetin disrupts tyrosine-phosphorylated phosphatidylinositol 3-kinase and myeloid differentiation factor-88 association, and inhibits MAPK/AP-1 and IKK/NF- $\kappa$ B-induced inflammatory mediators production in RAW 264.7 cells. *Immunobiology* 2013;218:1452–67.
- Wattel A, Kamel S, Prouillet C, et al. Flavonoid quercetin decreases osteoclastic differentiation induced by RANKL via a mechanism involving NF kappa B and AP-1. *J Cell Biochem* 2004;92:285–95.
- Cheng-Qiu L, Ting W, Chun-Yuan Z, et al. Quercetin attenuates cardiomyocyte apoptosis via inhibition of JNK and p38 mitogen-activated protein kinase signaling pathways. *Gene* 2016;577:275–80.
- Cheng-Wei L, Wen-Chi H, Shing-Chuan S, et al. Quercetin inhibition of tumor invasion via suppressing PKC delta/ERK/AP-1-dependent matrix metalloproteinase-9 activation in breast carcinoma cells. *Carcinogenesis* 2008;29:1807–15.
- Hwang MK, Song NR, Kang NJ, et al. Activation of phosphatidylinositol 3-kinase is required for tumor necrosis factor- $\alpha$ -induced upregulation of matrix metalloproteinase-9: its direct inhibition by quercetin. *Int J Biochem Cell Biol* 2009;41:1592–600.
- Granado-Serrano AB, Martín MA, Bravo L, et al. Quercetin attenuates TNF-induced inflammation in hepatic cells by inhibiting the NF- $\kappa$ B pathway. *Nutr Cancer* 2012;64:588–98.