



# Poplar phyllosphere harbors disparate isoprene-degrading bacteria

Andrew T. Crombie<sup>a,1</sup>, Nasmille L. Larke-Mejia<sup>b</sup>, Helen Emery<sup>b</sup>, Robin Dawson<sup>b</sup>, Jennifer Pratscher<sup>c</sup>, Gordon P. Murphy<sup>d</sup>, Terry J. McGenity<sup>d</sup>, and J. Colin Murrell<sup>b</sup>

<sup>a</sup>School of Biological Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom; <sup>b</sup>School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom; <sup>c</sup>Lyell Centre, Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom; and <sup>d</sup>School of Biological Sciences, University of Essex, Colchester CO4 3SQ, United Kingdom

Edited by Caroline S. Harwood, University of Washington, Seattle, WA, and approved November 1, 2018 (received for review July 24, 2018)

The climate-active gas isoprene (2-methyl-1,3-butadiene) is released to the atmosphere in huge quantities, almost equaling that of methane, yet we know little about the biological cycling of isoprene in the environment. Although bacteria capable of growth on isoprene as the sole source of carbon and energy have previously been isolated from soils and sediments, no microbiological studies have targeted the major source of isoprene and examined the phyllosphere of isoprene-emitting trees for the presence of degraders of this abundant carbon source. Here, we identified isoprene-degrading bacteria in poplar tree-derived microcosms by DNA stable isotope probing. The genomes of isoprene-degrading taxa were reconstructed, putative isoprene metabolic genes were identified, and isoprene-related gene transcription was analyzed by shotgun metagenomics and metatranscriptomics. Gram-positive bacteria of the genus *Rhodococcus* proved to be the dominant isoprene degraders, as previously found in soil. However, a wider diversity of isoprene utilizers was also revealed, notably *Variovorax*, a genus not previously associated with this trait. This finding was confirmed by expression of the isoprene monooxygenase from *Variovorax* in a heterologous host. A *Variovorax* strain that could grow on isoprene as the sole carbon and energy source was isolated. Analysis of its genome confirmed that it contained isoprene metabolic genes with an identical layout and high similarity to those identified by DNA-stable isotope probing and metagenomics. This study provides evidence of a wide diversity of isoprene-degrading bacteria in the isoprene-emitting tree phyllosphere and greatly enhances our understanding of the biodegradation of this important metabolite and climate-active gas.

isoprene | microbiology | plant-microbe interactions | DNA-SIP | phyllosphere

Isoprene (2-methyl-1,3-butadiene) is emitted to the atmosphere at a rate of  $\sim 500 \text{ Tg}\cdot\text{y}^{-1}$ , on a par with methane and approximately one-third of the total volatile organic compound (VOC) emissions (1, 2). The vast majority originates from terrestrial plants ( $400\text{--}600 \text{ Tg}\cdot\text{y}^{-1}$ ), with a small but uncertain flux from marine algae ( $0.1\text{--}12 \text{ Tg}\cdot\text{y}^{-1}$ ) (2–4), bacteria, fungi, and animals (5, 6). A reactive diene, isoprene is rapidly photochemically oxidized (1), with a significant and complex effect on global climate (7). Hydroxyl (OH) and nitrate (NO<sub>3</sub>) radicals and ozone (O<sub>3</sub>) in the atmosphere react with isoprene depending on prevailing conditions (1). In pristine environments, isoprene reacts directly with ozone and hydroxyl radicals, resulting in ozone depletion. However, the high NO<sub>x</sub> levels typical of urban environments result in the formation of tropospheric ozone, with important negative effects on human health and on yields of ozone-sensitive crops (8). Globally, these reactions result in a net radiative forcing of  $0.09 \text{ W}\cdot\text{m}^{-2}$ , with an additional indirect effect since depletion of OH radicals increases the atmospheric lifetime of methane (9). The isoprene oxidation products form secondary organic aerosols and cloud condensation nuclei with implications for planetary albedo, air quality, and climate (10, 11).

Plants produce isoprene in the chloroplast and release it to the atmosphere from the abaxial surface of leaves via the stomata (12). Although isoprene production is not a universal trait among plants, it protects against heat and oxidative stress (13, 14) and has roles in plant/insect signaling and plant energy dynamics (5, 15). High-isoprene-emitting trees worldwide include oil palm (*Elaeis guineensis*), eucalyptus (*Eucalyptus* spp.) and, in temperate regions, English oak (*Quercus robur*), poplar (*Populus* spp.), and willow (*Salix* spp.), with reported emissions of 77 (English oak) and 175 (oil palm)  $\mu\text{g}\cdot\text{g}^{-1}$  (dry leaves)  $\text{h}^{-1}$  (16). Due to its short atmospheric lifetime, isoprene concentrations vary from a few parts per billion by volume (ppbv) (17), to tens of ppbv in high-isoprene-emitting forests (18). Soils acted as a biological sink, both in closed chambers (19, 20) and in continuous systems using isoprene concentrations of 2–200 ppbv (21), suggesting that soil microbes act as a significant sink for isoprene, consuming up to 4% of global emissions.

Several isoprene-consuming bacterial strains have been isolated from soils and marine sediments (6, 22–25), and recently several strains have been characterized biochemically and genetically (26–30). Before this work, all strains characterized by sequence data were members of the Actinobacteria, and all

## Significance

Isoprene is a climate-active gas, produced in huge amounts by trees, yet we know little about its biogeochemical cycle. Bacteria able to grow on isoprene have been isolated from soils and sediments, but the phyllosphere, the principal isoprene source, has remained unexplored. Using targeted cultivation-independent techniques, we show that the phyllosphere of an isoprene-emitting tree contains a diverse and active isoprene-degrading population. We reconstruct the genome of an isoprene-degrading *Variovorax* strain and show that it contains a functional isoprene monooxygenase. This detailed study targets isoprene degraders from the phyllosphere, applies metagenomics to isoprene degradation, and isolates and sequences an isoprene-degrading member of the Proteobacteria.

Author contributions: A.T.C., N.L.L.-M., G.P.M., T.J.M., and J.C.M. designed research; A.T.C., N.L.L.-M., H.E., R.D., and G.P.M. performed research; A.T.C., N.L.L.-M., H.E., J.P., and G.P.M. analyzed data; and A.T.C., T.J.M., and J.C.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

Data deposition: The genome sequences of *Variovorax* sp. WS11 and megaplasmid-cured *Rhodococcus* AD45-ID have been deposited at DNA Data Bank of Japan/European Nucleotide Archive/GenBank under accession nos. PXZZ00000000 and PYHL00000000. Versions described here are PXZZ01000000 and PYHL01000000, respectively. Sequence reads have been deposited at GenBank Sequence Read Archive (accession no. SRP101805).

<sup>1</sup>To whom correspondence should be addressed. Email: a.crombie@uea.ac.uk.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1812668115/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1812668115/-DCSupplemental).

Published online November 29, 2018.

terrestrial examples were rhodococci. They all contain six genes (*isoABCDEF*) encoding a multicomponent isoprene monooxygenase (IsoMO) (Fig. 1) with four additional genes, *isoGHIJ*, just upstream. van Hyleckama Vlieg et al. (26, 27) showed that *isoI* encodes a glutathione S-transferase that conjugates glutathione with the epoxide product of IsoMO, and that *isoH* encodes a dehydrogenase that acts on the product of IsoI. IsoG and IsoJ are involved in the subsequent metabolic pathway but have yet to be characterized. In *Rhodococcus* AD45, the entire cluster *isoGHIJABCDEF* is cotranscribed as an operon (28). Significantly, all previously characterized isoprene gene clusters include the glutathione biosynthesis genes *gshAB*, suggesting a specific role for glutathione in isoprene metabolism. This small thiol is not usually found in Gram-positive bacteria (31), although a role in styrene metabolism has also been suggested recently (32).

The plant microbiome plays an essential role in plant health and development (33), and the phyllosphere, although an unstable environment, constitutes an extensive habitat for microorganisms, mainly bacteria, the diverse community of which typically comprises  $10^6$ – $10^7$  cells per  $\text{cm}^2$  (34). In the intercellular spaces of leaves, near the point of emission from stomata, isoprene concentrations reach 30 ppmv, approximately three orders of magnitude higher than atmospheric concentrations (12, 35–37). Apart from isoprene, plants produce a range of other VOCs. Although leaf-dwelling methylotrophs can reduce plant methanol emissions to the atmosphere (38), the extent to which the plant microbiome moderates release of other VOCs is not known (39, 40). Human intervention is likely to alter global emissions of isoprene, partly because emissions increase with temperature but are inversely related to atmospheric carbon dioxide concentrations, but also due to the development of high-isoprene-emitting agroforestry (41, 42). Therefore, our aim was to better understand the diversity of microbial taxa involved in isoprene consumption and their mechanisms of action. Specifically, this study addresses the following questions: Does the microbiome of a common isoprene-emitting tree harbor isoprene degraders able to take advantage of this abundant carbon source? And are novel genes expressed in response to isoprene?

## Results and Discussion

**Isoprene Degraders Isolated from Leaves and Soil.** Leaf washings from high-isoprene-emitting tree species (oak, poplar, willow) consumed isoprene in microcosms. After 15 d, 65% of the microcosms showed >30% depletion of added isoprene (*SI Appendix, Table S1*), confirming the presence of isoprene-degrading microbes. We obtained three *Rhodococcus* isolates that grew on isoprene as the sole source of carbon and energy—two strains (ACPA1 and ACPA4) from poplar leaves and one (ACS1) from soil beneath trees. All contained isoprene metabolic genes similar to those identified previously (28, 43).


**Stable Isotope Probing Identifies Active Degraders.** To extend the revealed diversity beyond those amenable to cultivation, we used DNA-stable isotope probing (DNA-SIP) in two independent experiments. Since sun-exposed leaves may produce more isoprene than shaded leaves (44), leaves of white poplar (*Populus alba*) taken from locations on the tree exposed to full sun were used. Initially, microbial cells were dislodged from the leaves and

incubated in minimal medium with  $^{13}\text{C}$ -labeled or unlabeled isoprene (headspace concentration  $\sim 500$  ppmv). Added isoprene was consumed rapidly, and cells were harvested after 7 or 8 d (*SI Appendix, Fig. S1*). Extraction, fractionation (based on incorporation of  $^{13}\text{C}$  label), and quantification of DNA showed that labeled DNA from the  $^{13}\text{C}$ -isoprene incubations migrated to “heavy” and “light” regions of the ultracentrifuge tubes depending on incorporation of label, whereas DNA from  $^{12}\text{C}$ -isoprene incubations was restricted (>99%) to light fractions (*SI Appendix, Fig. S2A*). Analysis of 16S rRNA gene amplicons showed that the unenriched bacterial community, highly consistent across six replicates, was composed mainly of Proteobacteria [*Sphingomonas* (mean relative abundance (RA) 36%), *Pseudomonas* (17%), members of the Oxalobacteraceae (8%), *Methylobacterium* (2%) and members of the Comamonadaceae (2%)], Bacteroidetes [*Hymenobacter* (17%) and Sphingobacteriaceae (1%)], and Actinobacteria (4%) (Fig. 2 and *SI Appendix, Fig. S3*), in agreement with previous studies of plant-associated bacteria (45, 46). Following enrichment, the active isoprene-degrading community was dominated by *Rhodococcus* ( $78 \pm 8\%$  mean RA  $\pm$  SD), although several proteobacterial taxa were also labeled, principally members of the Xanthomonadaceae ( $8 \pm 7\%$ ) and Comamonadaceae ( $3 \pm 1.6\%$ ) (Fig. 2 and *SI Appendix, Fig. S3*). So far, all well-described terrestrial isoprene degraders in cultivation are members of the genus *Rhodococcus*, but we have previously noted the presence of sequences related to Proteobacteria in DNA-SIP incubations with labeled isoprene (29), despite the fact that no publicly available proteobacterial genomes contain recognizable isoprene-degradation-related gene sequences. We investigated these nonactinobacterial isoprene degraders in more detail, again using DNA-SIP.


The second DNA-SIP experiment, with isoprene at 150 ppmv, included incubations without isoprene (no substrate) as an additional control. RNA was also extracted from the incubations, although not subjected to ultracentrifugation or separation into labeled and unlabeled fractions. Analysis showed that, as before, labeled and unlabeled DNA were efficiently separated (*SI Appendix, Fig. S2B*). In contrast to the first experiment, nucleic acids were analyzed using a shotgun approach. Community composition was assessed by clade-specific marker genes using MetaPhlan2 (47) (Fig. 2 and *SI Appendix, Fig. S3*). Analysis of the same DNA, from the heavy fraction of the  $^{13}\text{C}$ -isoprene incubations of the first experiment, by both 16S rRNA amplicon and metagenomic sequencing, enabled direct comparison of the microbial community revealed by these two methods. Both confirmed the dominance of *Rhodococcus* sequences (78 and 94% RA by amplicon and shotgun sequencing, respectively), together with a contribution from Comamonadaceae (3 and 1.5%, respectively). However, the Xanthomonadaceae, identified as 8% among 16S rRNA amplicons, were not identified in significant numbers by the shotgun approach (<0.1% RA), possibly indicative of bias in the PCR-based method.

In the second SIP experiment, the time-point-zero community was dominated by the Bacteroidetes *Hymenobacter* and *Pedobacter* (50% RA). *Sphingomonas*, the most abundant genus of the first experiment, was a minor component (3%) in the second. Following isoprene enrichment, the labeled community was again dominated by *Actinobacteria* (mainly *Rhodococcus*, average 74% RA) together with, as before, Proteobacteria, notably Comamonadaceae, principally *Variovorax*, which averaged 16% of the labeled community, albeit with considerable intersample variability (*SI Appendix, Fig. S3*). In contrast, these taxa composed a small fraction (0.5–1%) of the DNA from the pooled light fractions. The community profile of control incubations without added substrate was similar to that of the light fractions, further confirming that the heavy fractions contained the labeled DNA from isoprene consumers (Fig. 2).

The transcriptionally active taxa were characterized by profiling the mRNA transcriptome reads using MetaPhlan2 (*SI Appendix, Fig. S4*). This approach, albeit based on an extremely restricted subset of taxonomically informative marker genes, identified the Bacteroidetes genus *Pedobacter* and the Gammaproteobacterium



**Fig. 1.** The isoprene gene cluster in *Rhodococcus* sp. AD45. Genes present in all previously characterized isoprene degraders are shown in solid colors. In many isolates, *aldH1* is located between *isoJ* and *isoA*. The horizontal line shows genes cotranscribed in *Rhodococcus* AD45 (28).




**Fig. 2.** Community profile of the unenriched (time-point zero), and unlabeled (light) and labeled (heavy) fractions of  $^{13}\text{C}$ -isoprene incubations from two DNA-SIP experiments, analyzed by 16S rRNA gene amplicon (16S) or shotgun metagenomic (MG) sequencing. For the first experiment (left-most four bars), the labeled and unlabeled bacterial communities were characterized by amplicon sequencing, using DNA extracted from unenriched time-point zero cells at the start of the experiment (U-E), DNA from light and heavy fractions of incubations with  $^{13}\text{C}$ -isoprene, and also by shotgun sequencing of the pooled DNA from heavy fractions of  $^{13}\text{C}$ -isoprene incubations. The microbial communities of the second experiment (right-most four bars) were analyzed by shotgun sequencing of the unenriched time-point zero DNA, pooled DNA from samples incubated without substrate (N-S), pooled DNA from the light fractions of  $^{13}\text{C}$ -isoprene incubations, and DNA from the heavy fractions of  $^{13}\text{C}$ -isoprene incubations. Taxa present at  $>10\%$  are shown in boldface type. Taxa identified at higher levels comprise the sum of those not identified more specifically. The 16S and enriched samples show the mean of triplicates. For complete data, including individual replicates and  $^{12}\text{C}$ -isoprene controls, see *SI Appendix*, Fig. S3.

*Pseudomonas* as transcriptionally active, not only in isoprene enrichments, but also in unenriched time-point-zero samples and in incubations without added substrate, suggesting that these taxa were able to scavenge nutrients from endogenous organic matter and/or dead microbial cells. Transcripts of *Rhodococcus* and *Variovorax*, as well as the Bacteroidetes *Chryseobacterium* and *Riemerella* were specifically enriched in isoprene incubations, indicating that these taxa were directly or indirectly stimulated by isoprene (*SI Appendix*, Fig. S4).

**Assembly Identifies *isoA* Sequences.** To identify isoprene-degrading sequences, reads from pooled unenriched time-point-zero samples from each  $^{13}\text{C}$ -isoprene-enriched heavy fraction, from pooled light-fraction DNA, and from no-substrate incubations were coassembled, resulting in a total of 1.84 Gbp of sequence (*SI Appendix*, Table S3). Using, as query, the amino acid sequence of *isoA* (encoding the alpha subunit of IsoMO), we identified 11 sequences with 40–100% inferred amino acid identity to IsoA from *Rhodococcus* AD45 (Fig. 3). These scaffolds were examined in detail, and five contained sufficient sequence data to also identify one or more of the genes, specific for isoprene metabolism in *Rhodococcus* AD45 (28), immediately flanking the monooxygenase (Fig. 4) (36–100% amino acid identity with the *Rhodococcus* AD45 gene products), indicating a likely role in isoprene degradation. Similarly, reads originating from rRNA-depleted RNA were assembled, resulting in 220,637–454,247 transcripts containing 83.0–223.3 Mbp of sequence for each sample (*SI Appendix*, Table S4). Eighteen transcripts containing distinct *isoA*-related sequences were identified (Fig. 3), 8 of which contained other isoprene-related genes in addition to the IsoMO genes *isoABCDEF* (*SI Appendix*, Table S5). Many of these genes and transcripts were closely related to those of the *Rhodococcus* isolates obtained in this study, as well as to other strains identified previously (28, 29) (Figs. 3 and 4). Within-sample quantification of global transcript abundance showed that these *Rhodococcus*-like isoprene gene transcripts were among the most highly expressed, many among the top 0.2 or 1% of community-wide transcripts (Fig. 3 and *SI Appendix*, Table S5). In addition to sequences grouping with those of characterized isolates, three highly expressed transcripts, centered around metagenome scaffold MG\_3829, formed a distinct cluster (97% identity to *isoA* from

*Rhodococcus opacus* PD630), suggesting that there is yet more diversity to discover among these isoprene-degrading Actinobacteria. Apart from these *Rhodococcus*-like *isoA* genes, more divergent sequences were identified, as described below. Significantly, low-level transcription of *isoA* was also observed at time-point zero (unenriched) and in incubations without isoprene (no substrate), suggesting that there is a degree of constitutive expression of these isoprene-degradation genes.

**Genome Reconstruction Through Binning.** To reconstruct individual genomes from the metagenome, scaffolds were assigned to 266 bins based on abundance and nucleotide composition. Bin quality was assessed and refined, and taxonomy was assigned, resulting in 27 genome bins with predicted completeness  $>90\%$  and contamination  $<5\%$ . From the 266 bins, 18 were identified as of interest based on the presence of an *isoA* sequence or predicted RA of  $>2\%$  in the labeled community, and these were examined in greater detail (*SI Appendix*, Tables S6 and S7). In total, nine *isoA*-containing scaffolds were assigned to genome bins, allowing taxonomy to be inferred for most of these sequences (Fig. 3). Several sequences, similar to those of characterized isolates, were assigned to actinobacterial bins (Fig. 3). The most abundant (average 31% RA across three  $^{13}\text{C}$ -isoprene enrichments), predicted to be 99% complete and with 1.1% contamination (*SI Appendix*, Table S6), had 98.7% average nucleotide identity to isolate *Rhodococcus* sp. ACPA4. Several other (less complete) *Rhodococcus* bins were also identified. However, since DNA-SIP revealed a considerable diversity of labeled organisms, we also looked at more dissimilar sequences. Genome bin 197, assigned to *Pseudonocardia*, contained two *isoA* homologs. The first of these, on scaffold MG\_3829, contained a typical isoprene metabolic gene cluster with 75–93% amino acid identity to six isoprene metabolic gene products, including IsoA, from *Rhodococcus* AD45 (Fig. 4). The same genome also contained scaffold MG\_720, with a less similar IsoA homolog (48% amino acid identity with IsoA from *Rhodococcus* AD45). This region of scaffold MG\_720 contained genes with high sequence identity and an identical layout to *Rhodococcus jostii* DSM44719 and *Pseudonocardia dioxanivorans* CB1190 (48) annotated as toluene-4 monooxygenases (*SI Appendix*, Fig. S5). These monooxygenase genes are of low similarity and arranged



**Fig. 3.** The relationship between the *isoA* genes of known isoprene degraders (in boldface type), metagenome scaffold sequences (prefixed MG), and metatranscriptome sequences, together with other representative sequences from the databases. Transcripts are prefixed by "MT" followed by sample identification (Unenriched, time-point zero; No-sub, incubations without isoprene;  $^{12}\text{C}$ - $^{12}\text{C}$ -3, incubations with unlabeled isoprene;  $^{13}\text{C}$ - $^{13}\text{C}$ -3, incubations with labeled isoprene). Scaffolds or transcripts containing isoprene-related genes in addition to *isoABCDEF* are indicated with a double asterisk (\*\*). For each sample, transcripts were ranked by normalized transcript abundance, and highly expressed transcripts are marked with four, three, two, or one red circle, indicating that the *isoA*-containing transcript was among the most abundant 0.2, 1, 10, or 50%, respectively, of all transcripts from that sample (*SI Appendix*, Table S4). Where identical *isoA* sequences were present on different transcripts from the same sample, only the most highly expressed is shown. Partial *isoA* sequences are indicated with the length in parentheses. The taxonomy of genome bins is shown after the scaffold identification. NA, not assigned; NB, not binned. Bootstrap values over 50% (1,000 replications) are shown as solid circles at the nodes. The scale bar indicates nucleotide substitutions per site.


in a different order to *isoABCDEF* from isoprene degraders and are not flanked by any other recognizable isoprene-related genes, suggesting that this region was not responsible for isoprene metabolism by known or predicted pathways. Interestingly, although they could not be identified as directly involved in isoprene degradation, these divergent genes on scaffold MG 720 of genome bin 197 were transcribed at moderately high levels (Fig. 3 and *SI Appendix*, Table S5). Genome bin 095 contained a sequence with homology to a putative toluene monooxygenase from *Myxococcales* bacterium 68–20 (OJY25058.1), with a similar gene layout. Again, although this genome contained no identifiable isoprene-related genes apart from the monooxygenase, the *isoA* homolog was represented in the transcriptome (Fig. 3 and *SI Appendix*, Table S5). These two examples suggest that isoprene is capable of inducing genes not central to its metabolism, that these monooxygenase genes are constitutively expressed at considerable levels, or that they form part of a novel and so-far-undescribed isoprene metabolic pathway.

Genome bin 232, assigned to the betaproteobacterial genus *Variovorax*, contained an *isoA* homolog (scaffold MG 478), which was represented by a moderately expressed transcript and aligned most closely with *xamoA* of the propylene-degrader *Xanthobacter autotrophicus* Py2 (Fig. 3 and *SI Appendix*, Table S5). In contrast to characterized isoprene degraders, however, *X. autotrophicus* Py2 does not use glutathione in its alkene metabolic pathway (49), does not contain any homologs of the isoprene metabolic genes surrounding *isoA-F* at the *xamoA* locus (50), and does not grow on isoprene (30). Scaffold MG 478 comprised 76,719 bp of contiguous DNA, allowing examination of the

genomic context of *isoA* (Fig. 4). In addition to the six genes (*isoABCDEF*) encoding IsoMO, *isoGHII* and *aldh1* were present in an identical layout to those of many isoprene-degrading isolates (Fig. 4), although sequence identity of the gene products with those of *Rhodococcus* AD45 ranged from 42–71%, much lower than those of characterized isolates. Glutathione biosynthesis genes *gshA* and *gshB* were not present in this isoprene cluster but were found on scaffold MG 3916 of this genome bin (85 and 76% amino acid identity to ACS17089.1 and ACS17095.1 from *Variovorax paradoxus* S110, respectively), perhaps reflecting the general use of glutathione in Gram-negative bacteria, as opposed to its often isoprene-specific use in Gram-positive strains.

The co-occurrence of both *isoABCDEF* and *isoGHII* has previously been successful in identifying *bone fide* isoprene degraders, but to verify that this sequence contained the genetic potential for isoprene oxidation, we cloned the putative IsoMO genes into a plasmid vector and induced expression in *Rhodococcus* AD45-ID, a strain of *R. AD45* lacking the megaplasmid containing the *iso* genes, which is incapable of isoprene oxidation. When expressed, the IsoMO from bin 232 indeed oxidized isoprene, in contrast to controls (*SI Appendix*, Fig. S6).

**Targeted Isolation of a *Variovorax* Strain.** By screening the 16S rRNA gene sequences of numerous isolates from isoprene enrichments, we obtained *Variovorax* sp. WS11, isolated from a soil enrichment, which grew on isoprene as the sole source of carbon and energy (*SI Appendix*, Fig. S7). The *Variovorax* sp. WS11 genome contains an isoprene gene cluster with identical layout to the metagenome-derived sequence of bin 232 (Fig. 4). Overall



**Fig. 4.** The isoprene metabolic gene clusters from known isoprene-degrading isolates (in boldface type) together with representative sequences from the assembled metagenome (prefix MG\_ and including bin identification and predicted taxonomy) and metatranscriptomes (prefix MT\_).

average amino acid identity (51) was compared between these genomes and 52 *Variovorax* genome assemblies available in GenBank. The metagenome-derived genome was most similar to *Variovorax* sp. CF079 (GCA\_900101545.1) (82.3%) whereas strain WS11 was most similar to *Variovorax* sp. B2 (GCA\_002891695.1) (88%), and similarity between bin 232 and *V. sp.* WS11 was 79.5%, suggesting that both are novel species. All *Variovorax* genomes in GenBank were searched for isoprene-related gene sequences like those present in the *Variovorax* strains described here, but with negative results, suggesting that much diversity still exists outside the reference sequences.

**Conclusions.** Although isoprene degraders are present in all isoprene-exposed environments tested (6), no studies have identified isoprene degraders residing on the leaves of isoprene-emitting trees at the source of emission. Here, the isoprene degraders retrieved from the poplar phyllosphere were dominated by *Rhodococcus*, but included other Actinobacteria (*Pseudonocardia*) and Proteobacteria (*Variovorax*). Taxonomy alone is insufficient to identify isoprene-degrading bacteria, with extremely closely related strains differing in terms of isoprene-degrading ability, and phylogeny based on 16S rRNA genes is not congruent with *isoA*-based analyses (30). These data suggest lateral transfer of the isoprene metabolic genes and imply that surveys that rely on 16S rRNA gene analysis are not able to identify isoprene degraders.

We used metagenomics to reconstruct the genome of a *Variovorax* strain and conventional methods to obtain related isolate *Variovorax* WS11. Both genomes contained the entire isoprene metabolic gene cluster. Despite being more similar in sequence to alkene monooxygenase from *Xanthobacter autotrophicus* Py2 than IsoMO from known isoprene degraders (Fig. 3), the presence of other genes unique to isoprene metabolism and the expression of the monooxygenase in a heterologous host proved that these are genuine isoprene metabolic gene clusters. *Variovorax* are metabolically versatile bacteria capable of degradation of natural products and xenobiotics, frequently plant-associated and with plant-growth-promoting effects (52, 53), and have been identified as part of the core bacterial microbiome of both *Arabidopsis* and poplar (54, 55).

Interestingly, we also detected a significant level of monooxygenase transcripts, similar or identical to  $^{13}\text{C}$ -labeled DNA scaffolds, which have not, so far, been implicated in isoprene degradation and are possibly indicative of novel isoprene metabolic pathways. This study shows that the leaves of an isoprene-emitting tree provide a habitat for taxonomically disparate isoprene degraders and forms a basis for continued development of molecular tools to detect isoprene degraders. This is a prerequisite for quantification of isoprene-related genes and transcripts and

comparison of the activity of microbes associated with isoprene-emitting and nonemitting environments (including comparisons between high- and low-isoprene-emitting tree species), and hence to establishing the extent to which the tree microbiome is able to take advantage of, and mitigate the release of, this abundant carbon source and climate-active gas.

## Materials and Methods

For full details, see *SI Appendix*.

**Enrichment, Isolation, and Stable Isotope Probing.** Isoprene degraders were enriched from soil or from cells dislodged from leaves by ultrasound, and purified by standard methods. For SIP enrichments, cells were washed from 5 g leaves, resuspended in minimal medium, and supplied with  $^{13}\text{C}$ -labeled and unlabeled isoprene. Isoprene consumption was followed by gas chromatography (GC), and cells were harvested when they had consumed  $\sim 11$  or  $6 \mu\text{mol}\cdot\text{ml}^{-1}$  (isoprene C), for the first or second experiment, respectively. Each treatment was carried out in triplicate.

**Nucleic Acid Extraction and Purification.** DNA and RNA were extracted using standard methods. Total RNA was depleted of rRNA using Ribo-Zero (Illumina). Labeled and unlabeled DNA were separated by density gradient ultracentrifugation and fractionation as described previously (29). Fractions containing labeled (“heavy”) and unlabeled (“light”) DNA were identified based on the data presented in *SI Appendix*, Fig. S2.

**Sequencing of Nucleic Acids.** Communities profiled by 16S rRNA gene amplicons were sequenced using Illumina MiSeq and analyzed using Qiime (56). Metaomic libraries were sequenced using Illumina HiSeq (*SI Appendix*, Tables S3 and S4). Quality-filtered reads were taxonomically profiled using Metaphlan2 v2.5.0 (47), coassembled using IDBA-UD v1.1.1 (57) (*SI Appendix*, Table S3), binned using MaxBin v2.2 (58), and quality-checked and refined using CheckM v1.0.5 (59) and RefineM v0.0.23 (60). Filtered transcript reads were de novo assembled using Trinity v2.3.2 (61) (*SI Appendix*, Table S4). Local Blast databases were searched using tblastn v2.2.28 (62). Reads were mapped to assembled transcripts and quantified using kallisto v0.43.1 (63). Normalized expression levels, as transcripts per million (TPM), of each transcript were ranked for each sample as a percentile.

**Expression of IsoMO.** The IsoMO genes *isoABCDEF* of metagenome bin 232 were PCR-amplified from the pooled heavy fractions of DNA-SIP enrichments and expressed from a vector in a strain of *Rhodococcus* (*R. AD45-ID*) cured of the megaplasmid that contains the isoprene metabolic genes. Isoprene uptake of IsoMO-expressing cell suspensions was quantified by GC.

**Accession Numbers.** The genomes of *Variovorax* sp. WS11 and *Rhodococcus* AD45-ID have been deposited at DNA Database of Japan/European Nucleotide Archive/GenBank under accessions PXZZ00000000 and PYHL00000000. Versions described here are PXZZ01000000 and PYHL01000000, respectively.

Sequence reads have been deposited at GenBank Sequence Read Archive (accession no. SRP101805) (SI Appendix, Tables S2–S4).

**ACKNOWLEDGMENTS.** Plasmid pTipQC1 was a gift from Tomohiro Tamura. This work was funded by the Earth and Life Systems Alliance at the

University of East Anglia; Natural Environment Research Council (NERC) Grants NE/J009725/1 (to J.C.M.) and NE/J009555/1 (to T.J.M.); European Research Council Advanced Grant 694578—IsoMet (to J.C.M.); and NERC Fellowship NE/L010771/1 (to J.P.). We acknowledge receipt of a Colciencias Colombian Government Scholarship (to N.L.L.-M.).

- Atkinson R, Arey J (2003) Gas-phase tropospheric chemistry of biogenic volatile organic compounds: A review. *Atmos Environ* 37(Suppl 2):197–219.
- Guenther AB, et al. (2012) The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions. *Geosci Model Dev* 5:1471–1492.
- Shaw SL, Gantt B, Meshkizde N (2010) Production and emissions of marine isoprene and monoterpenes: A review. *Adv Meteorol* 2010:1–24.
- Dani SKG, et al. (2017) Relationship between isoprene emission and photosynthesis in diatoms, and its implications for global marine isoprene estimates. *Mar Chem* 189:17–24.
- Sanadze GA (2017) Biogenic isoprene emission as expression of dissipativity, a fundamental cell property. *Russ J Plant Physiol* 64:133–140.
- McGenity TJ, Crombie AT, Murrell JC (2018) Microbial cycling of isoprene, the most abundantly produced biological volatile organic compound on Earth. *ISME J* 12:931–941.
- Pacifico F, Harrison SP, Jones CD, Sitch S (2009) Isoprene emissions and climate. *Atmos Environ* 43:6121–6135.
- Ashworth K, Wild O, Hewitt CN (2013) Impacts of biofuel cultivation on mortality and crop yields. *Nat Clim Chang* 3:492–496.
- Folberth GA, Hauglustaine DA, Lathière J, Brocheton F (2006) Interactive chemistry in the Laboratoire de Méteorologie Dynamique general circulation model: Model description and impact analysis of biogenic hydrocarbons on tropospheric chemistry. *Atmos Chem Phys* 6:2273–2319.
- Fiore AM, et al. (2012) Global air quality and climate. *Chem Soc Rev* 41:6663–6683.
- Carlton AG, Wiedinmyer C, Kroll JH (2009) A review of Secondary Organic Aerosol (SOA) formation from isoprene. *Atmos Chem Phys* 9:4987–5005.
- Fall R, Monson RK (1992) Isoprene emission rate and intercellular isoprene concentration as influenced by stomatal distribution and conductance. *Plant Physiol* 100:987–992.
- Sharkey TD, Wiberley AE, Donohue AR (2008) Isoprene emission from plants: Why and how. *Ann Bot* 101:5–18.
- Zeinali N, Altaravneh M, Li D, Al-Nu'airat J, Dlugogorski BZ (2016) New mechanistic insights: Why do plants produce isoprene? *ACS Omega* 1:220–225.
- Loivamäki M, Mumm R, Dicke M, Schnitzler J-P (2008) Isoprene interferes with the attraction of bodyguards by herbaceous plants. *Proc Natl Acad Sci USA* 105:17430–17435.
- Kesselmeier J, Staudt M (1999) Biogenic volatile organic compounds (VOC): An overview on emission, physiology and ecology. *J Atmos Chem* 33:23–88.
- Greenberg JP, et al. (1999) Tethered balloon measurements of biogenic VOCs in the atmospheric boundary layer. *Atmos Environ* 33:855–867.
- Wiedinmyer C, et al. (2005) Ozarks isoprene experiment (OZIE): Measurements and modeling of the “isoprene volcano.” *J Geophys Res Atmos* 110:D18307.
- Cleveland CC, Yavitt JB (1997) Consumption of atmospheric isoprene in soil. *Geophys Res Lett* 24:2379–2382.
- Pegoraro E, et al. (2005) The effect of elevated atmospheric CO<sub>2</sub> and drought on sources and sinks of isoprene in a temperate and tropical rainforest mesocosm. *Glob Change Biol* 11:1234–1246.
- Gray CM, Helmig D, Fierer N (2015) Bacteria and fungi associated with isoprene consumption in soil. *Elem Sci Anth* 3:000053.
- Srivastava N, Singh A, Bhardwaj Y, Dubey SK (2018) Biotechnological potential for degradation of isoprene: A review. *Crit Rev Biotechnol* 38:587–599.
- Cleveland CC, Yavitt JB (1998) Microbial consumption of atmospheric isoprene in a temperate forest soil. *Appl Environ Microbiol* 64:172–177.
- van Ginkel CG, de Jong E, Tilanus JWR, de Bont JAM (1987) Microbial oxidation of isoprene, a biogenic foliage volatile and of 1,3-butadiene, an anthropogenic gas. *FEMS Microbiol Lett* 45:275–279.
- Ewers J, Freier-Schroder D, Knackmuss HJ (1990) Selection of trichloroethene (TCE) degrading bacteria that resist inactivation by TCE. *Arch Microbiol* 154:410–413.
- van Hylckama Vlieg JET, Kingma J, Kruizinga W, Janssen DB (1999) Purification of a glutathione S-transferase and a glutathione conjugate-specific dehydrogenase involved in isoprene metabolism in *Rhodococcus* sp. strain AD45. *J Bacteriol* 181:2094–2101.
- van Hylckama Vlieg JET, Kingma J, van den Wijngaard AJ, Janssen DB (1998) A glutathione S-transferase with activity towards cis-1,2-dichloroepoxyethane is involved in isoprene utilization by *Rhodococcus* sp. strain AD45. *Appl Environ Microbiol* 64:2800–2805.
- Crombie AT, et al. (2015) Regulation of plasmid-encoded isoprene metabolism in *Rhodococcus*, a representative of an important link in the global isoprene cycle. *Environ Microbiol* 17:3314–3329.
- El Khawand M, et al. (2016) Isolation of isoprene degrading bacteria from soils, development of *isoA* gene probes and identification of the active isoprene-degrading soil community using DNA-stable isotope probing. *Environ Microbiol* 18:2743–2753.
- Johnston A, et al. (2017) Identification and characterisation of isoprene-degrading bacteria in an estuarine environment. *Environ Microbiol* 19:3526–3537.
- Johnson T, Newton G, Fahey R, Rawat M (2009) Unusual production of glutathione in *Actinobacteria*. *Arch Microbiol* 191:89–93.
- Heine T, et al. (2018) On the enigma of glutathione dependent styrene degradation in *Gordonia rubripertincta* CWB2. *Appl Environ Microbiol* 84:e00154-18.
- Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P (2013) Structure and functions of the bacterial microbiota of plants. *Annu Rev Plant Biol* 64:807–838.
- Vorholt JA (2012) Microbial life in the phyllosphere. *Nat Rev Microbiol* 10:828–840.
- Brüggemann N, Schnitzler JP (2002) Comparison of isoprene emission, intercellular isoprene concentration and photosynthetic performance in water-limited Oak (*Quercus pubescens* Willd. and *Quercus robur* L.) saplings. *Plant Biol* 4:456–463.
- Sun Z, Hüve K, Vislap V, Niinemets Ü (2013) Elevated [CO<sub>2</sub>] magnifies isoprene emissions under heat and improves thermal resistance in hybrid aspen. *J Exp Bot* 64:5509–5523.
- Singsaas EL, Lerdau M, Winter K, Sharkey TD (1997) Isoprene increases thermotolerance of isoprene-emitting species. *Plant Physiol* 115:1413–1420.
- Abanda-Nkpaw D, Musch M, Tschiersch J, Boettner M, Schwab W (2006) Molecular interaction between *Methylobacterium extorquens* and seedlings: Growth promotion, methanol consumption, and localization of the methanol emission site. *J Exp Bot* 57:4025–4032.
- Junker RR, Tholl D (2013) Volatile organic compound mediated interactions at the plant-microbe interface. *J Chem Ecol* 39:810–825.
- Bringle F, Couëe I (2015) Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. *Front Microbiol* 6:486.
- Rosenstiel TN, Potosnak MJ, Griffin KL, Fall R, Monson RK (2003) Increased CO<sub>2</sub> uncouples growth from isoprene emission in an agriforest ecosystem. *Nature* 421:256–259.
- Rosenkranz M, Pugh TAM, Schnitzler J-P, Arneth A (2015) Effect of land-use change and management on biogenic volatile organic compound emissions: Selecting climate-smart cultivars. *Plant Cell Environ* 38:1896–1912.
- Crombie AT, Emery H, McGenity TJ, Murrell JC (2017) Draft genome sequences of three terrestrial isoprene-degrading *Rhodococcus* strains. *Genome Announc* 5:e01256-17.
- Sharkey TD, Loreto F, Delwiche CF (1991) High carbon dioxide and sun/shade effects on isoprene emission from oak and aspen tree leaves. *Plant Cell Environ* 14:333–338.
- Redford AJ, Bowers RM, Knight R, Linhart Y, Fierer N (2010) The ecology of the phyllosphere: Geographic and phylogenetic variability in the distribution of bacteria on tree leaves. *Environ Microbiol* 12:2885–2893.
- Laforest-Lapointe I, Messier C, Kembel SW (2016) Tree phyllosphere bacterial communities: Exploring the magnitude of intra- and inter-individual variation among host species. *PeerJ* 4:e2367.
- Truong DT, et al. (2015) MetaPhlan2 for enhanced metagenomic taxonomic profiling. *Nat Methods* 12:902–903.
- Sales CM, et al. (2011) Genome sequence of the 1,4-dioxane-degrading *Pseudonocardia dioxanivorans* strain CB1190. *J Bacteriol* 193:4549–4550.
- Allen JR, Clark DD, Krum JG, Ensign SA (1999) A role for coenzyme M (2-mercaptoethanesulfonic acid) in a bacterial pathway of aliphatic epoxide carboxylation. *Proc Natl Acad Sci USA* 96:8432–8437.
- Krum JG, Ensign SA (2001) Evidence that a linear megaplasmid encodes enzymes of aliphatic alkene and epoxide metabolism and coenzyme M (2-mercaptoethanesulfonate) biosynthesis in *Xanthobacter* strain Py2. *J Bacteriol* 183:2172–2177.
- Konstantinidis KT, Tiedje JM (2005) Towards a genome-based taxonomy for prokaryotes. *J Bacteriol* 187:6258–6264.
- Belimov AA, et al. (2009) Rhizosphere bacteria containing 1-aminocyclopropane-1-carboxylate deaminase increase yield of plants grown in drying soil via both local and systemic hormone signalling. *New Phytol* 181:413–423.
- Satola B, Wübbeler JH, Steinbüchel A (2013) Metabolic characteristics of the species *Variovorax paradoxus*. *Appl Microbiol Biotechnol* 97:541–560.
- Bodenhausen N, Horton MW, Bergelson J (2013) Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS One* 8:e56329.
- Beckers B, Op De Beeck M, Weyens N, Boerjan W, Vangronsveld J (2017) Structural variability and niche differentiation in the rhizosphere and endosphere bacterial microbiome of field-grown poplar trees. *Microbiome* 5:25.
- Kuczynski J, et al. (2011) Using QIIME to analyze 16S rRNA gene sequences from microbial communities. *Curr Protoc Microbiol* 27:1E.5.1–1E.5.20.
- Peng Y, Leung HC, Yiu SM, Chin FY (2012) IDBA-UD: A de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–1428.
- Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW (2014) MaxBin: An automated binning method to recover individual genomes from metagenomes using an expectation-maximization algorithm. *Microbiome* 2:1–18.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015) CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055.
- Parks DH, et al. (2017) Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat Microbiol* 2:1533–1542.
- Haas BJ, et al. (2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* 8:1494.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410.
- Bray NL, Pimentel H, Melsted P, Pachter L (2016) Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* 34:525.