

## Case Report

# Oculodentodigital Dysplasia: A Case Report and Major Review of the Eye and Ocular Adnexa Features of 295 Reported Cases

Virang Kumar <sup>1</sup>, Natario L. Couser <sup>2,3,4</sup> and Arti Pandya<sup>5</sup>

<sup>1</sup>Virginia Commonwealth University School of Medicine, Richmond, VA, USA

<sup>2</sup>Department of Ophthalmology, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

<sup>3</sup>Department of Human and Molecular Genetics, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

<sup>4</sup>Department of Pediatrics, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

<sup>5</sup>Department of Pediatrics, Division of Genetics and Metabolism, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Correspondence should be addressed to Virang Kumar; [kumarvk@vcu.edu](mailto:kumarvk@vcu.edu)

Received 22 September 2019; Accepted 23 March 2020; Published 6 April 2020

Academic Editor: Sandra M. Johnson

Copyright © 2020 Virang Kumar et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oculodentodigital dysplasia (ODDD) is a rare genetic disorder associated with a characteristic craniofacial profile with variable dental, limb, eye, and ocular adnexa abnormalities. We performed an extensive literature review to highlight key eye features in patients with ODDD and report a new case of a female patient with a heterozygous missense *GJA1* mutation (c.65G>A, p.G22E) and clinical features consistent with the condition. Our patient presented with multiple congenital anomalies including syndactyly, microphthalmia, microcornea, retrognathia, and a small nose with hypoplastic alae and prominent columella; in addition, an omphalocele defect was present, which has not been reported in previous cases. A systematic review of the published cases to date revealed 91 literature reports of 295 individuals with ODDD. There were 73 different *GJA1* mutations associated with these cases, of which the most common were the following missense mutations: c.605G>A (p.R202H) (11%), c.389T>C (p.I130T) (10%), and c.119C>T (p.A40V) (10%). Mutations most commonly affect the extracellular-1 and cytoplasmic-1 domains of connexin-43 (gene product of *GJA1*), predominately manifesting in microphthalmia and microcornea. The syndrome appears with an approximately equal sex ratio. The most common eye features reported among all mutations were microcornea, microphthalmia, short palpebral fissures, and glaucoma.

## 1. Introduction

Oculodentodigital dysplasia (ODDD, OMIM #164200) is a rare disorder mainly characterized by abnormal craniofacial, dental, ocular, and digital development. The autosomal dominant form has been the most frequently reported inheritance pattern, although a few cases of autosomal recessive inheritance have been described [1–3]. Craniofacial abnormalities may include microcephaly, prominent columella, and underdeveloped nasal alae [2–4]. Dental abnormalities, such as hypoplastic enamel, small teeth, and premature loss of teeth, are often present [2–4]. Digit abnormalities may include syndactyly, camptodactyly, and midphalangeal hypoplasia [2–4]. Ophthalmic manifestations are common, such as microcornea and microphthalmia, and may involve

a wide spectrum of eye and ocular adnexa structures, although previous analyses of prior cases show that full ocular physical exams were not performed on all patients [3, 5].

The gap junction protein alpha 1 (*GJA1*) gene codes for connexin-43, which is a protein that assists in the transmembrane transport of molecules through gap junctions, and mutations in the *GJA1* may cause an alteration of the channel conduction properties [1–3, 6]. We report a case of an 8-month-old female patient with an identified *GJA1* mutation and common clinical features associated with ODDD. This patient had an omphalocele at birth, which has not been reported in previous cases. Her eye features included microphthalmia, microcornea, narrow palpebral fissures, blonde fundus, deep anterior chambers, hyperopia, and epiphora in both eyes secondary to bilateral nasolacrimal



FIGURE 1: Facial photograph of a patient with oculodentodigital dysplasia; note the beaked nose with hypoplastic alae and prominent columella, microphthalmia, microcornea, small palpebral fissures, retrognathia.

duct obstructions. We conducted an extensive literature review to summarize the eye features in patients with ODDD reported to date.

## 2. Case Report

The patient, an 8-month-old female, was born to a noncon-sanguineous couple from a healthy 37-year-old mother of Native American descent and a healthy 30-year-old father of German and Irish descent. Family history is notable for an older sibling with cleft palate, paternal uncle with autism, paternal second cousin with congenital heart defect, and distant paternal great-great uncle with Down syndrome and webbed/fused 4<sup>th</sup> and 5<sup>th</sup> digits of one hand. A normal pregnancy was noted until the second trimester when an omphalocele was detected on ultrasound. A subsequent ultrasound revealed possible syndactyly of the hands. The patient was born at 39 weeks by vaginal delivery with induction. The birth weight was 3.552 kg (75<sup>th</sup> percentile), birth length was 50 cm (68<sup>th</sup> percentile), and birth head circumference was 34.5 cm (70<sup>th</sup> percentile). Apgar scores were 9 at both one minute and five minutes.

Multiple congenital anomalies noted at birth included an omphalocele that measured 4 cm at base and 3.5 cm across with intestines present in the sac, but no liver. The patient had a normocephalic head with sparse wispy hair, a small



FIGURE 2: Complete syndactyly of the 4<sup>th</sup> and 5<sup>th</sup> digits of the right hand.

nose with hypoplastic alae, a prominent columella, small-appearing palpebral fissures, a small cornea, microphthalmia, a wide anterior fontanelle, and retrognathia (Figure 1). Syndactyly of digits 4 and 5 and webbing of digits 3 and 4 of the right (Figure 2) and left hands were present. Cardiac echocardiogram on the day of birth showed the presence of a mild patent ductus arteriosus, mild patent foramen ovale, and a normal aorta. Feeding difficulties were exacerbated by the presence of the omphalocele; surgical correction was performed on day 2 of life.

An ophthalmologic assessment at 4 months of age was notable for deep anterior chambers, bilateral nasolacrimal duct obstruction, microphthalmia, small 8 mm corneas, a blonde fundus, and moderate hyperopia in both eyes.

At her last examination at 8 months of age, the patient continues to have poor feeding with self-limiting volumes but has improved weight gain. The patient is at the 9<sup>th</sup> percentile for weight and 12<sup>th</sup> percentile for length. Cognitive and motor developments are delayed.

Sequencing of the *GJA1* gene (transcript number: NM\_000165.3) from patient genomic DNA revealed a heterozygous missense mutation in the *GJA1* gene: c.65G>A (p.G22E). Deletion/duplication analysis of the *GJA1* gene using the aCGH test was negative.

## 3. Methods

We performed a systematic review of the literature to summarize the ocular findings in individuals with ODDD. A PubMed/Medline search of “oculodentodigital syndrome” led us to find a total of 177 articles. No articles were excluded based on the year published. We reviewed the references to identify other articles that did not appear in our original search. 91 articles describing patients with a description consistent with the clinical syndrome, either with or without molecular confirmation of *GJA1* pathogenic variants, were included. Within these selected articles, we identified 295 cases of ODDD with 73 different *GJA1* mutations, including

TABLE 1: *GJA1* variants without clinical information.

Sources	<i>GJA1</i> variant		Cases
	Nucleotide	Protein	
Paznekas et al. [3]	c.7G>A	p.D3N	1
Paznekas et al. [3]	c.64G>A	p.G22R	1
Paznekas et al. [3]; Richardson et al. [6]	c.79T>C	p.S27P	1
Paznekas et al. [3]	c.163A>G	p.N55D	1
Paznekas et al. [3]	c.174A>C	p.Q58H	1
Paznekas et al. [3]	c.175C>G	p.P59A	1
Paznekas et al. [3]	c.221A>T	p.H74L	1
Paznekas et al. [3]	c.428G>A	p.G143D	1
Paznekas et al. [3]	c.430A>G	p.K144E	1
Paznekas et al. [3]	c.434T>G	p.V145G	1
Paznekas et al. [3]	c.442C>G	p.R148G	1
Paznekas et al. [3]	c.578C>T	p.P193L	1

TABLE 2: Summary of sex distribution.

	Males		Females		Total
Individuals with clinical diagnosis of ODDD (with no molecular confirmation)	14	45%	18	56%	32
Untested individuals with both ODDD phenotype and known relative with molecular confirmation	52	53%	46	47%	98
Individuals with a molecular confirmed <i>GJA1</i> pathogenic variant	72	44%	93	56%	165
Totals	138	47%	157	53%	295

those that exhibited features of ODDD in the absence of molecular confirmation. Such individuals were either clinically diagnosed or were relatives of individuals with molecularly confirmed *GJA1* pathogenic variants. Twelve reported that *GJA1* gene coding alterations were omitted due to insufficient clinical information and data reported and are listed in Table 1 [3, 6].

#### 4. Discussion

Oculodentodigital dysplasia (ODDD) is a rare congenital disorder manifested with developmental anomalies of the eyes, face, dentition, heart, skeletal system, and digits. The syndrome appears to be more common in Caucasian populations with an equal sex ratio [3]. Heterozygous mutation of the *GJA1* gene located at chromosome 6q22.31 has been identified as the most common mutation resulting in ODDD [2, 3]. However, a compound heterozygous individual with missense mutations demonstrated mutations in the *GJA1* gene (p.V41L) and the *GJB2* gene (p.R127H), which encode for connexin-43 and connexin-26, respectively, and has been reported and classified as having overlapping features of Clouston syndrome and ODDD [3, 7].

In addition to the classic phenotypic features of the syndrome, a wide variety of additional physical manifestations

have been observed. Ocular findings of microphthalmia and microcornea have been observed commonly in previous cases [2–4]. Craniofacial anomalies of microcephaly, poor hair growth, hypoplastic nasal alae, and prominent columella have been reported previously [2–4]. Bilateral syndactyly of the 4<sup>th</sup> and 5<sup>th</sup> digits is common [2, 3].

A systematic review of the published cases to date (ranging from 1963 to 2019) revealed 91 literature reports of 295 individuals with ODDD [1–91]. Table 2 [1–91] summarizes the sex distribution across all reviewed reports of ODDD. Patients with ODDD present with an approximately equal sex distribution (47% male and 53% female). Of the 295 individuals reported, 32 were clinically diagnosed with ODDD without molecular confirmation, 98 presented with features of ODDD and had a known relative with molecular confirmation of a *GJA1* pathogenic variant, and 165 individuals had a molecularly confirmed *GJA1* pathogenic variant.

There were 73 different *GJA1* mutations identified from the 165 individuals that had a molecularly confirmed *GJA1* pathogenic variant. Table 3 [1–3, 5–71, 92] summarizes the number of patients with each mutation. Patients with confirmed pathogenic variants and their relatives with no molecular confirmation but with features of ODDD were grouped separately. These two groups comprised 263 of the patients included in this study.

The eye features of all 295 patients are summarized in Table 4 [1–91]. The most common ophthalmic manifestations reported were microcornea ( $n = 111$ ), microphthalmia ( $n = 110$ ), short palpebral fissures ( $n = 56$ ), and glaucoma ( $n = 51$ , 4 closed-angle and 1 open-angle).

Twenty-three patients presented with refractive error, of which isolated myopia was the most frequently noted ( $n = 14$ ), followed by isolated hyperopia ( $n = 6$ ), anisometropia ( $n = 2$ ), and astigmatism ( $n = 1$ ). Forty patients presented with eye movement disorders, with strabismus ( $n = 27$ , 9 esotropic, 1 exotropic) being the most common, followed by nystagmus ( $n = 8$ ), amblyopia ( $n = 3$ ), Duane syndrome ( $n = 2$ ), and Brown syndrome ( $n = 1$ ). Note that 1 patient had both nystagmus and esotropia [71]. Other common findings included epicanthus ( $n = 36$ ), hypotelorism ( $n = 24$ ), hypertelorism ( $n = 22$ ), madarosis ( $n = 19$ ), cataracts ( $n = 17$ ), persistent pupillary membranes ( $n = 13$ ), shallow anterior chambers ( $n = 12$ ), pale/atrophic irides ( $n = 11$ ), telecanthus ( $n = 11$ ), and uveitis ( $n = 10$ ).

A variety of abnormal findings for the retina and optic disc were noted ( $n = 18$ ), with dysplasia of the retina/fundus ( $n = 3$ ) and pale/atrophic optic discs ( $n = 3$ ) being the most common documented findings.

Of the individuals with molecularly confirmed mutations, the most common mutations present were c.605G>A (p.R202H) (11%; with 1 patient also having a c.717G>A synonymous mutation), c.389T>C (p.I130T) (10%), and c.119C>T (p.A40V) (10%). Table 5 [2, 3, 12, 30, 40, 41, 66, 67, 92] summarizes the eye features present in the patients with these mutations.

Less common features of the phenotype observed in our presented case were also reported in other cases as well. These include nasolacrimal duct abnormalities ( $n = 2$ ), pale/atrophic

TABLE 3: Reported *G/AI* mutations and sex distribution in ODDD.

Sources	Multiple mutations?	Nucleotide	Protein	Unspecified	Individuals with a molecular confirmed <i>G/AI</i> pathogenic variant		Untested individuals with both ODDD phenotype and known relative with molecular confirmation		Total individuals with the ODDD phenotype				
					Male	Female	Male	Female	Male	Female	Total		
Cavusoglu et al. 2019	No	c.168_169insT	p.Q57SfsTer6	N/A	1	0	0	0	1	100%	0	0%	1
Aminabadi et al. 2009 & Aminabadi et al. 2010	No	N/A	N/A	Missense mutation exon 2 (unspecified)	1	0	2	1	3	75%	1	25%	4
Dwarakanathan et al. 2015 & Furuta et al. 2012	No	c.75G>T	p.W25C	N/A	1	1	0	0	1	50%	1	50%	2
Quick and Dobersen 2014; National Center for Biotechnology Information 2020	Yes	c.605G>A	p.R202H	N/A	1	0	0	0	1	100%	0	0%	1
Paznekas et al. 2003 & Paznekas et al. 2009	No	c.717G>A	p.R239R	N/A	1	7	4	5	5	29%	12	71%	17
Jamsheer et al. 2010	Yes	c.301C>T	p.R101X	N/A	1	0	0	0	1	100%	0	0%	1
Jamsheer et al. 2010	No	c.6delT	p.G2fsX7	N/A	0	1	0	0	0	0%	1	100%	1
Paznekas et al. 2009; Joss et al. 2008; & Richardson et al. 2006	No	c.301C>T	p.R101X	N/A	0	2	0	0	0	0%	2	100%	2
Paznekas et al. 2009; Richardson et al. 2004; Paznekas et al. 2003; & Gladwin et al. 1997	No	c.97C>T	p.R33X*	N/A	0	0	4	4	4	50%	4	50%	8
Wang et al. 2019	No	c.91A>T	p.I311P	N/A	1	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009 & van Steensel et al. 2005	No	c.780_781delTTG	p.C260fsX306	N/A	1	2	0	0	1	33%	2	67%	3
Paznekas et al. 2009; Paznekas et al. 2003; & Gorlin et al. 1963	No	c.68A>C	p.K23T	N/A	1	0	0	0	1	100%	0	0%	1
Dwarakanathan et al. 2015; Paznekas et al. 2009; & Vreeburg et al. 2007	No	c.689_690delAT	p.Y230fsX236	N/A	0	3	1	0	1	25%	3	75%	4
This study; Gumus 2018; Paznekas et al. 2009; Paznekas et al. 2003; & Traboulsi and Parks 1990	No	c.65G>A	p.G22E	N/A	0	3	0	0	0	0%	3	100%	3
Wiest et al. 2006	No	c.659C>A	p.S220Y	N/A	0	1	0	0	0	0%	1	100%	1
Paznekas et al. 2009; Paznekas et al. 2003; & Norton et al. 1995	No	c.646G>T	p.V216L	N/A	1	0	4	1	5	83%	1	17%	6
Park et al. 2017; Paznekas et al. 2009; & Paznekas et al. 2003	No	c.61G>A	p.G21R	N/A	0	2	0	0	0	0%	2	100%	2
Brice et al. 2013	No	c.617A>G	p.K206R	N/A	1	2	1	1	2	40%	3	60%	5

TABLE 3: Continued.

Sources	Multiple mutations?	G/AI mutation	Nucleotide	Protein	Unspecified	Individuals with a molecular confirmed G/AI pathogenic variant		Untested individuals with both ODDD phenotype and known relative with molecular confirmation		Total individuals with the ODDD phenotype			
						Male	Female	Male	Female	Male	Female	Total	
Paznekas et al. 2009	No	c.602C>T	p.S201F	N/A	0	1	0	0	0	0	1	100%	1
Paznekas et al. 2009 & de la Parra et al. 2007	No	c.5G>T	p.G2V	N/A	1	0	0	0	0	1	100%	0	1
Vitiello et al. 2005 & Vingolo et al. 1994	No	c.581A>C	p.H194P*	N/A	3	5	3	3	6	43%	8	57%	14
Paznekas et al. 2009; Paznekas et al. 2003; & Judisch et al. 1979	No	c.52T>C	p.S18P	N/A	0	0	1	3	1	25%	3	75%	4
Paznekas et al. 2009 & Paznekas et al. 2003	No	c.50A>C	p.Y17S	N/A	3	4	0	0	3	43%	4	57%	7
Paznekas et al. 2009 & Debeer et al. 2005	No	c.504_506delCTT	p.F169del	N/A	0	1	0	0	0	0%	1	100%	1
Wiest et al. 2006 & Thomsen et al. 1998	No	c.461C>A	p.T154N	N/A	0	2	0	1	0	0%	3	100%	3
Paznekas et al. 2009 & van Es et al. 2007	No	c.460A>G	p.T154A*	N/A	0	2	0	0	0	0%	2	100%	2
Paznekas et al. 2009; Richardson et al. 2004; & Paznekas et al. 2003; Gladwin et al. 1997; & Schrandner-Stumpel et al. 1993	No	c.443G>A	p.R148Q	N/A	0	0	2	2	2	50%	2	50%	4
Taşdelen et al. 2018	No	c.442C>T	p.R148Ter	N/A	1	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009; Debeer et al. 2005; & Spaepen et al. 1991	No	c.440Y>C	p.M147T	N/A	0	1	0	0	0	0%	1	100%	1
Paznekas et al. 2009; Richardson et al. 2004; & Brueton et al. 1990	No	c.427G>A	p.G143S	N/A	0	0	8	1	8	89%	1	11%	9
Orosz et al. 2018	No	c.413G>A	p.G138D	N/A	1	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009; Paznekas et al. 2003; & Shapiro et al. 1997	No	c.412G>C	p.G138R	N/A	1	2	2	2	3	43%	4	57%	7
Kogame et al. 2014	No	c.412G>A	p.G138S	N/A	1	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009; Richardson et al. 2004; Paznekas et al. 2003; & Gladwin et al. 1997	No	c.402G>T	p.K134N	N/A	0	0	0	2	0	0%	2	100%	2
Paznekas et al. 2009 & Paznekas et al. 2003	No	c.400A>G	p.K134E	N/A	0	1	0	0	0	0%	1	100%	1
Nishat et al. 2012; Paznekas et al. 2009; Paznekas et al. 2003; & Amador et al. 2008	No	c.389T>C	p.I130T	N/A	7	4	5	1	12	71%	5	29%	17
Paznekas et al. 2009; Musa et al. 2008; Wiest et al. 2006; & Loddenkemper et al. 2002	No	c.338T>C	p.L113P	N/A	2	2	1	0	3	60%	2	40%	5
Paznekas et al. 2009 & Debeer et al. 2005	No	c.330G>C	p.E110D	N/A	2	3	1	2	3	38%	5	63%	8
Paznekas et al. 2009 & Kelly et al. 2006	No	c.32T>C	p.L11P	N/A	0	1	0	0	0	0%	1	100%	1



TABLE 3: Continued.

Sources	Multiple mutations?	G/AI mutation	Nucleotide	Protein	Unspecified	Individuals with a molecular confirmed G/AI pathogenic variant		Untested individuals with both ODDD phenotype and known relative with molecular confirmation		Total individuals with the ODDD phenotype				
						Male	Female	Male	Female	Male	Female	Total		
Gabriel et al. 2011 & Jamsheer et al. 2009	No	c.31C>T	p.L11F	N/A	0	2	0	0	0	0	2	100%	2	
Pornraveetus et al. 2017	No	c.31C>A	p.L11I	N/A	1	0	0	0	0	1	100%	0	0%	1
Jamsheer et al. 2014	No	c.317T>G	p.L106R	N/A	2	0	0	0	0	2	100%	0	0%	2
Paznekas et al. 2009 & Nivelon-Chevallier et al. 1981	No	c.317T>C	p.L106P	N/A	1	0	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009 & Paznekas et al. 2003	No	c.306G>C	p.K102N	N/A	1	2	0	0	0	1	33%	2	67%	3
Paznekas et al. 2009; Paznekas et al. 2003; & Wooldridge et al. 1977	No	c.293A>G	p.Y98C	N/A	1	3	1	1	1	2	33%	4	67%	6
Paznekas et al. 2009	No	c.287T>C	p.V96A	N/A	0	1	0	0	0	0	0%	1	100%	1
Wiest et al. 2006	No	c.287T>A	p.V96E	N/A	0	1	0	0	0	0	0%	1	100%	1
Paznekas et al. 2009 & Kjaer et al. 2004	No	c.286G>A	p.V96M	N/A	2	2	0	0	0	2	50%	2	50%	4
Paznekas et al. 2009 & Honkaniemi et al. 2005	No	c.284A>G	p.H95R	N/A	0	1	0	1	0	0	0%	2	100%	2
Paznekas et al. 2009; Paznekas et al. 2003; & Opijordsmoen and Nyberg-Hansen 1980	No	c.268C>G	p.L90V	N/A	4	0	3	2	7	78%	2	22%	9	
Jamsheer et al. 2014	No	c.257C>A	p.S86Y	N/A	0	1	0	0	0	0	0%	1	100%	1
Pizzuti et al. 2004	No	c.227G>A	p.R76H	N/A	1	0	0	0	0	1	100%	0	0%	1
Izumi et al. 2013	No	c.226C>T	p.R76C	N/A	1	0	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009; Paznekas et al. 2003; & Stanislaw et al. 1998	No	c.226C>A	p.R76S	N/A	0	2	0	2	0	0	0%	4	100%	4
Choi et al. 2018	No	c.221A>C	p.H74P*	N/A	1	0	0	0	0	1	100%	0	0%	1
Paznekas et al. 2009; Richardson et al. 2004; Paznekas et al. 2003; & Gladwin et al. 1997	No	c.206C>A	p.S69Y	N/A	0	0	2	5	2	2	29%	5	71%	7
Paznekas et al. 2009 & Vasconcellos et al. 2005	No	c.176C>A	p.P59H	N/A	4	4	1	0	5	56%	4	44%	9	
Paznekas et al. 2009	No	c.145_147dupCAG	p.Q49dup	N/A	0	1	0	0	0	0	0%	1	100%	1
Paznekas et al. 2009; Paznekas et al. 2003; Weintraub et al. 1975; & Gellis and Feingold 1974	No	c.154_156dupTTT	p.F52dup	N/A	1	0	1	1	1	2	67%	1	33%	3
Hadjichristou et al. 2017 & Paznekas et al. 2009	No	c.146A>C	p.Q49P	N/A	1	1	0	0	0	1	50%	1	50%	2
Izumi et al. 2013	No	c.145C>G	p.Q49E	N/A	0	1	0	0	0	0	0%	1	100%	1
Paznekas et al. 2009 & Paznekas et al. 2003	No	c.145C>A	p.Q49K	N/A	3	2	0	0	3	60%	2	40%	5	

TABLE 3: Continued.

Sources	Multiple mutations?	Nucleotide	Protein	Unspecified	Individuals with a molecular confirmed G/AI pathogenic variant		Untested individuals with both ODDD phenotype and known relative with molecular confirmation		Total individuals with the ODDD phenotype				
					Male	Female	Male	Female	Male	Female	Total		
Amano et al. 2012; Feller et al. 2008; Paznekas et al. 2009; & Iftro et al. 2005	No	c.142G>A	p.E48K	N/A	3	0	0	0	3	100%	0	0%	3
Jamsheer et al. 2014	No	c.139G>C	p.D47H	N/A	0	3	0	0	0	0%	3	100%	3
Turminelli et al. 2016	No	c.125G>C	p.E42Q	N/A	1	0	0	0	1	100%	0	0%	1
Gabriel et al. 2011	No	c.120delGGTT GAGTCAGC	p.V41_ A44del	N/A	0	1	1	2	1	25%	3	75%	4
Paznekas et al. 2009 & Kellermayer et al. 2005	Yes (compound heterozygous with G/B2 mutation)	c.121G>C N/A	p.V41L p.R127H (G/B2 mutation)	N/A	0	1	0	0	0	0%	1	100%	1
Park et al. 2019; Hayashi et al. 2014; Paznekas et al. 2009; Debeer et al. 2005; & Paznekas et al. 2003	No	c.119C>T	p.A40V	N/A	6	4	4	3	10	59%	7	41%	17
Wittlieb-Weber et al. 2015	No	c.175C>T	p.P59S	N/A	1	2	0	0	1	33%	2	67%	3
Attig et al. 2016	No	c.396_398delAAA	p.I132_ K133delinsM	N/A	3	2	0	0	3	60%	2	40%	5
Paznekas et al. 2009	No	c.19T>G	p.L7V	N/A	1	0	0	0	1	100%	0	0%	1
Himi et al. 2009	No	c.13A>T	p.S5C	N/A	0	1	0	0	0	0%	1	100%	1
Pace et al. 2019	No	c.287T>G	p.V96G	N/A	0	1	0	0	0	0%	1	100%	1
Totals	No	c.77T>C	p.L26P	N/A	0	1	0	0	0	0%	1	100%	1
					72	93	52	46	124	47%	139	53%	263

\*Unknown which specific individuals tested.

TABLE 4: Eye and ocular adnexa features reported in ODDD.

Orbit	—	Microphthalmia (110/37%)	Hypotelorism (24/8%)	Hypertelorism (22/7%)	Short axial length (4/1%)				
Anterior chamber		Shallow anterior chamber (12/4%)	Deep anterior chambers (2/<1%)						
Cornea		Microcornea (111/38%)	Thick corneas (4/1%)	Corneal opacities (3/1%)	Corneal farinata (1/<1%)	Band keratopathy (1/<1%)	Corneal keratosis (1/<1%)	Abnormal Descemet's membrane (1/<1%)	Anteriorly deviated Schwalbe's line (1/<1%)
Sclera		Blue sclera (1/<1%)							
Pupil		Persistent pupillary membranes (13/4%)	Eccentric pupils (3/1%)						
Lens		Cataracts (17/6%)	Lens opacities (2/<1%)	White retrolental masses (1/<1%)					
Uvea (iris, ciliary body)		Pale/atrophic irides (11/4%)	Uveitis (10/3%)	General iris abnormalities (7/2%)	Synechiae (4/1%)	Hypoplastic anterior iris stroma (3/1%)	Ciliary body cysts (2/<1%)	Flat iris (1/<1%)	Inferior iris coloboma (1/<1%)
Uvea (choroid)		Thick choroid (2/<1%)	Thin choroid (1/<1%)						
Vitreous		Vitreous degeneration (1/<1%)	Vitreous membrane attachment to optic nerve and lens (1/<1%)	Persistent hyperplastic primary vitreous (1/<1%)					
Retina/fundus		Dysplastic retina/fundus (3/1%)	Pale retina/fundus (2/<1%)	Thread-like retinal vasculature (2/<1%)	Dystrophic retinal epithelium (1/<1%)	Hypoplastic macula (1/<1%)	Absent fundal glow with B-scan ultrasound (1/<1%)		
Optic disc		Pale/atrophic optic disc (3/1%)	Dysplastic optic disc (2/<1%)	Ellipsoid optic disc (1/<1%)	Optociliary vein presence (1/<1%)	Optic disc hypervascularity (1/<1%)			



TABLE 4: Continued.

	Short/narrow palpebral fissures (56/19%)	Epicanthus (36/12%)	Telecanthus (11/4%)	Ptosis (7/2%)	Blepharophimosis (1/<1%)	Entropion (1/<1%)	Ectropion (1/<1%)	Epiblepharon (1/<1%)	Mucosal hypertrophy (1/<1%)
Ocular adnexa	Madarosis (19/6%)	Flared eyebrows (3/1%) (2 medially flared)	Synophrys (1/<1%)						
	Nasolacrimal duct abnormalities (2/<1%)	Hypolacrimation (1/<1%)							
	Myopia (16/5%) (2 anisometropic)	Hyperopia (8/3%) (2 anisometropic)	Astigmatism (1/<1%)						
Refractive errors	Strabismus (27/9%) (9 esotropic, 1 exotropic)	Nystagmus (8/3%)	Amblyopia (3/1%)	Duane syndrome (2/<1%)	Brown syndrome (1/<1%)				
Eye movement disorders	Glaucoma (51/17%) (4 closed-angle, 1 open-angle)	Paracentral scotoma (1/<1%)							
Other	Abnormal ERG (2/<1%)	Delayed visual evoked responses (2/<1%)	Occipital subcortical white matter changes (1/<1%)						

TABLE 5: Common *GJA1* mutations with associated eye features.

Sources	Multiple mutations?	<i>GJA1</i> mutation		Individuals with <i>GJA1</i> mutation (confirmed and affected relatives) Total	Associated eye features
		Nucleotide	Protein		
Quick and Dobersen 2014; National Center for Biotechnology Information 2020	Yes	c.605G>A c.717G>A	p.R202H p.R239R	1	Microphthalmia (1)
Paznekas et al. 2009; Paznekas et al. 2003	No	c.605G>A	p.R202H	17	Microphthalmia (1), microcornea (2)
Nishat et al. 2012; Paznekas et al. 2009; Paznekas et al. 2003; and Amador et al. 2008	No	c.389T>C	p.I130T	17	Microphthalmia (4), hypotelorism (6), cataract (1), pale/atrophic optic disc (1), and short palpebral fissures (4)
Park et al. 2019; Hayashi et al. 2014; Paznekas et al. 2009; Debeer et al. 2005; and Paznekas et al. 2003	No	c.119C>T	p.A40V	17	Microphthalmia (9), hypertelorism (3), hypotelorism (4), short axial length (4), cataract (1), microcornea (8), thick cornea (4), macular hypoplasia (1), shallow anterior chamber (4), myopia (4), strabismus (6) (1 esotropic), glaucoma (6), and epicanthus (3)

retina/fundus ( $n = 2$ ), and deep anterior chambers ( $n = 2$ ). Additionally, including this study, the three patients with the p.G22E mutation have the following findings: microphthalmia ( $n = 3$ ), cataracts ( $n = 1$ ), microcornea ( $n = 2$ ), blonde fundus ( $n = 1$ ), persistent pupillary membrane ( $n = 1$ ), deep anterior chamber ( $n = 1$ ), hyperopia ( $n = 1$ ), strabismus ( $n = 2$ , 1 esotropic), amblyopia ( $n = 1$ ), glaucoma ( $n = 1$ ), short palpebral fissures ( $n = 1$ ), nasolacrimal duct abnormalities ( $n = 1$ ), and epicanthus ( $n = 1$ ) [2, 3, 21, 22].

Some unique genotype-phenotype correlations were noted upon further analysis. Three patients presented with eccentric pupils, but only 2 of these patients were reported with an associated mutation. Both mutations (p.Q49dup and p.Q49P) seem to affect the same amino acid in connexin-43 [3, 61, 72]. Additionally, uveitis was reported in 10 patients, 9 of which were associated with similar mutations. Eight of these patients were within the same study and had the p.H194P mutation, another patient had no molecular confirmation of a *GJA1* mutation, and the other patient was reported with a missense mutation on exon 2 [4, 9, 10, 27, 28]. However, since the majority of these patients were reported within the same study, the apparent genotype-phenotype correlation of p.H194P and uveitis might be due to underreporting of uveitis from other sources with different pathogenic variants or may be due to other factors of the family not identified within the study.

Further analysis of the genotype-phenotype correlation was conducted by pairing the phenotypic manifestations of each mutation with the corresponding defects in the connexin-43 domains. The domains were defined by the amino acid ranges provided on UniProt (P17302-CXA1\_HUMAN) [93]. Table 6 [1–3, 5–71, 92, 93] provides a summary of the phenotypes associated with mutations from each domain.

The domains most commonly affected by *GJA1* mutations are the extracellular-1 loop and the cytoplasmic-1 loop of connexin-43, accounting for 19 and 20 mutations, respectively. Disruptions in the extracellular-1 loop presented primarily as microphthalmia ( $n = 32$ ) and microcornea ( $n = 30$ ). A similar pattern can be seen in the cytoplasmic-1 loop, as the most common presentations were microphthalmia ( $n = 20$ ) and microcornea ( $n = 18$ ). Other clinical findings, however, may be able to distinguish mutations resulting from these domains. The next most common findings associated with mutations in the extracellular-1 loop were glaucoma ( $n = 15$ ) and hypertelorism ( $n = 11$ ), as opposed to short palpebral fissures ( $n = 14$ ) and hypotelorism ( $n = 14$ ) for the cytoplasmic-1 loop.

Mutations affecting the cytoplasmic N-terminus and the transmembrane-1 domain shared similar features to the ones in the extracellular-1 and cytoplasmic-1 domains, as microphthalmia and microcornea were the most common clinical findings. However, the mutations in the cytoplasmic N-terminus and transmembrane-1 domain presented with microcornea ( $n = 17$  and  $n = 21$ , respectively) more frequently than microphthalmia ( $n = 5$  and  $n = 14$ , respectively). The opposite pattern is true for the extracellular-1 and cytoplasmic-1 domains.

The mutations in the extracellular-2 loop demonstrate a different phenotypic pattern, as microphthalmia ( $n = 14$ ) occurs the most frequently, while microcornea is less frequent ( $n = 4$ ). Mutations in the transmembrane-2 domain also display a unique pattern, with hypertelorism ( $n = 5$ ) being the most frequent clinical finding. Other domains listed in Table 6 also demonstrate some unique clinical patterns, but this may be due to variability from the small number of samples. The patterns mentioned previously, however, still provide insight into the role of different connexin-43

TABLE 6: Mutant connexin-43 domains and associated phenotype.

<i>GJA1</i> mutation	Protein domain (amino acid range) (obtained from UniProt-P17302)	Associated phenotype (no. of individuals)
p.G2fsX7 (with p.R101X) p.G2V p.L11P p.L11F p.L11I p.L7V p.S5C	Cytoplasmic N-terminus (1-13)	Microcornea (7), microphthalmia (5), epicanthus (4), strabismus (3) (1 esotropic), short palpebral fissures (2), telecanthus (2), amblyopia (1), dysplastic fundus (1), opticiliary vein (1), dysplastic optic disc (1), pale/atrophic optic disc (1), persistent pupillary membrane (1), myopia (3), hyperopia (1) (anisometropic), glaucoma (1), ptosis (1), entropion (1), madarosis (1), hypertelorism (1), and cataract (1)
p.W25C p.R33X p.I31M p.K23T p.G22E p.G21R p.S18P p.Y17S p.L26P	Transmembrane-1 (14-36)	Microcornea (21), microphthalmia (14), short palpebral fissures (11), persistent pupillary membrane (6), madarosis (6), epicanthus (6), glaucoma (5), anterior iris stroma hypoplasia (3), hypertelorism (2), cataract (2), iris abnormalities (2), blonde fundus (1), iridoschisis (1), deep anterior chamber (1), hyperopia (2), strabismus (7) (3 esotropic), amblyopia (1), nystagmus (1), ptosis (1), epiblepharon (1), nasolacrimal duct obstruction (1), and flared eyebrows (1) (medially flared)
p.Q57SfsTer6 p.R76H p.R76C p.R76S p.H74P p.S69Y p.P59H p.Q49dup p.F52dup p.Q49P p.Q49E p.Q49K p.E48K p.D47H p.E42Q p.V41_A44del p.V41L (with p.R127H (GJB2 mutation)) p.A40V p.P59S	Extracellular-1 (37-76)	Microphthalmia (32), microcornea (30), glaucoma (15) (2 closed-angle, 1 open-angle), hypertelorism (11), epicanthus (10), strabismus (9) (3 esotropic), short palpebral fissures (9), iris atrophy (peripupillary) (8), cataract (6), shallow anterior chamber (6), hypotelorism (5), short axial length (4), myopia (4), corneal farinata (4), telecanthus (3), iris abnormalities (2), eccentric pupils (2), persistent pupillary membrane (2), dysplastic fundus (1), dysplastic optic (1), macular hypoplasia (1), synechiae (1), ciliary body cysts (1), deep anterior chamber (1), hyperopia (1), ptosis (1), blepharophimosis (1), madarosis (1), nasolacrimal duct abnormalities (1), and low-voltage ERG (1)
p.Y98C p.V96A p.V96E p.V96M p.H95R p.L90V p.S86Y p.V96G	Transmembrane-2 (77-99)	Hypertelorism (5), microcornea (2), microphthalmia (3), glaucoma (3), strabismus (2) (1 esotropic), short palpebral fissures (2), eyelid mucosal hypertrophy (1), telecanthus (1), epicanthus (1), optic disc atrophy (1), hyperopia (1), myopia (1), strabismus (1), paracentral scotoma (1), madarosis (1), and delayed visual evoked potentials (1)

TABLE 6: Continued.

<i>GJA1</i> mutation	Protein domain (amino acid range) (obtained from UniProt-P17302)	Associated phenotype (no. of individuals)
p.R101X (with p.G2fsX7) p.R101X p.T154N p.T154A p.R148Q p.R148Ter p.M147T p.G143S p.G138D p.G138R p.G138S p.K134N p.K134E p.I130T p.L113P p.E110D p.L106R p.L106P p.K102N p.I132_K133delinsM	Cytoplasmic-1 (100-154)	Microphthalmia (20), microcornea (18), short palpebral fissures (14), hypotelorism (14), glaucoma (9), myopia (7), epicanthus (5), cataract (3), strabismus (3), shallow anterior chamber (3), hypertelorism (2), opaque lens (1), optic disc hypervascularity (1), pale/atrophic optic disc (1), pale irides (1), iris abnormalities (2), astigmatism (1), Duane syndrome (1), ptosis (1), occipital subcortical white matter changes (1), and delayed visual evoked responses (1)
p.F169del	Transmembrane-3 (155-177)	Short palpebral fissures (1)
p.R202H (with p.R239R) p.R202H p.K206R p.S201F p.H194P	Extracellular-2 (178-208)	Microphthalmia (18), uveitis (8), glaucoma (8), microcornea (4), opaque cornea (2), thick choroid (2), cataract (1), shallow anterior chamber (1), nystagmus (2), and ptosis (1)
p.S220Y p.V216L	Transmembrane-4 (209-231)	Microphthalmia (1), glaucoma (1), microcornea (1), and persistent pupillary membrane (1)
p.Y230fsX236	Transmembrane-4 & cytoplasmic C-terminus (209-382)	Hypertelorism (2), hypotelorism (1), and flared eyebrows (2) (1 medially flared)
p.R239R (with p.R202H) p.I311P p.C260fsX306	Cytoplasmic C-terminus (232-382)	Short palpebral fissures (3), epicanthus (2), hypotelorism (2), microcornea (2), pale irides (2), myopia (2), hyperopia (1) (1 anisometric), corneal opacity (1), microphthalmia (1), retinal dysplasia (1), choroid thinning (1), glaucoma (1), madarosis (1), and loss of flash ERG (1)
Missense mutation exon 2 (unspecified)	Unknown	Microphthalmia (1), cataract (1), microcornea (1), uveitis (1), glaucoma (1), epicanthus (1), telecanthus (1), short palpebral fissures (1), and ptosis (1)

domains in providing phenotypic variability among patients with ODDD.

In conclusion, this report provides a comprehensive review of the eye and ocular adnexa abnormalities that are currently known to be associated with the ODDD phenotype. Limitations of this report include the possibility of an incomplete ophthalmologic evaluation and/or lack of reporting of eye features in all of the evaluated case reports or misdiagnosis in the individuals with the ODDD phenotype without molecular confirmation. As such, it is possible that the

reported common eye features within this summary may be over or underrepresented. Ophthalmic manifestations are commonly associated within the phenotype, and a wide spectrum of eye and ocular adnexa structures may be affected. The rarity of this condition provides further incentive to further investigate the phenotype.

### Consent

Consent has been obtained.

## Conflicts of Interest

Virang Kumar and Arti Pandya declare that they have no conflicts of interest. Natario L. Couser, MD, MS, is a principal investigator at the Virginia Commonwealth University site of Retrophin, Inc., and book editor in Elsevier.

## Supplementary Materials

Supplementary Material 1: all GJA1 mutations with associated eye and ocular adnexa features. This dataset groups patients with ODDD by GJA1 mutation and reports the associated eye and ocular adnexa features. (*Supplementary Materials*)

## References

- [1] A. Dwarakanathan, M. Bhat, S. GN, and S. Shetty, "Missense and deletion mutations in GJA1 causing oculodentodigital dysplasia in two Indian families," *Clinical Dysmorphology*, vol. 24, no. 4, pp. 159–162, 2015.
- [2] W. A. Paznekas, S. A. Boyadjiev, R. E. Shapiro et al., "Connexin 43 (GJA1) mutations cause the pleiotropic phenotype of oculodentodigital dysplasia," *American Journal of Human Genetics*, vol. 72, no. 2, pp. 408–418, 2003.
- [3] W. A. Paznekas, B. Karczeski, S. Vermeer et al., "GJA1 mutations, variants, and connexin 43 dysfunction as it relates to the oculodentodigital dysplasia phenotype," *Human Mutation*, vol. 30, no. 5, pp. 724–733, 2009.
- [4] G. Kayalvizhi, B. Subramanian, and G. Suganya, "Clinical manifestations of oculodentodigital dysplasia," *Journal of the Indian Society of Pedodontics and Preventive Dentistry*, vol. 32, no. 4, pp. 350–352, 2014.
- [5] D. R. de la Parra and J. C. Zenteno, "A new GJA1 (connexin 43) mutation causing oculodentodigital dysplasia associated to uncommon features," *Ophthalmic Genetics*, vol. 28, no. 4, pp. 198–202, 2007.
- [6] R. Richardson, D. Donnai, F. Meire, and M. J. Dixon, "Expression of Gja1 correlates with the phenotype observed in oculodentodigital syndrome/type III syndactyly," *Journal of Medical Genetics*, vol. 41, no. 1, pp. 60–67, 2004.
- [7] R. Kellermayer, M. Keller, P. Ratajczak et al., "Bigenic connexin mutations in a patient with hidrotic ectodermal dysplasia," *European Journal of Dermatology*, vol. 15, no. 2, pp. 75–79, 2005.
- [8] D. Cavusoglu, N. O. Dundar, P. Arican, B. Ozyilmaz, and P. Gencpinar, "A hypomyelinating leukodystrophy with calcification: oculodentodigital dysplasia," *Acta Neurologica Belgica*, 2019.
- [9] N. A. Aminabadi, A. T. Ganji, A. Vafaei, M. Pourkazemi, and S. G. Oskouei, "Oculodentodigital dysplasia: disease spectrum in an eight-year-old boy, his parents and a sibling," *The Journal of Clinical Pediatric Dentistry*, vol. 33, no. 4, pp. 337–341, 2009.
- [10] N. A. Aminabadi, M. Pourkazemi, S. G. Oskouei, and Z. Jamali, "Dental management of oculodentodigital dysplasia: a case report," *Journal of Oral Science*, vol. 52, no. 2, pp. 337–342, 2010.
- [11] N. Furuta, M. Ikeda, K. Hirayanagi, Y. Fujita, M. Amanuma, and K. Okamoto, "A novel GJA1 mutation in oculodentodigital dysplasia with progressive spastic paraplegia and sensory deficits," *Internal Medicine*, vol. 51, no. 1, pp. 93–98, 2012.
- [12] J. S. Quick and M. Dobersen, "Cardiac arrhythmia and death of teenager linked to rare genetic disorder diagnosed at autopsy," *The American Journal of Forensic Medicine and Pathology*, vol. 35, no. 2, pp. 103–105, 2014.
- [13] A. Jamsheer, M. Badura-Stronka, A. Sowińska, S. Debicki, K. Kiryluk, and A. Latos-Bieleńska, "A severe progressive oculodentodigital dysplasia due to compound heterozygous GJA1 mutation," *Clinical Genetics*, vol. 78, no. 1, pp. 94–97, 2010.
- [14] S. K. Joss, S. Ghazawy, S. Tomkins, M. Ahmed, J. Bradbury, and E. Sheridan, "Variable expression of neurological phenotype in autosomal recessive oculodentodigital dysplasia of two sibs and review of the literature," *European Journal of Pediatrics*, vol. 167, no. 3, pp. 341–345, 2008.
- [15] R. J. Richardson, S. Joss, S. Tomkin, M. Ahmed, E. Sheridan, and M. J. Dixon, "A nonsense mutation in the first transmembrane domain of connexin 43 underlies autosomal recessive oculodentodigital syndrome," *Journal of Medical Genetics*, vol. 43, no. 7, article e37, 2006.
- [16] A. Gladwin, D. Donnai, K. Metcalfe et al., "Localization of a gene for oculodentodigital syndrome to human chromosome 6q22-q24," *Human Molecular Genetics*, vol. 6, no. 1, pp. 123–127, 1997.
- [17] Z. Wang, L. Sun, P. Wang et al., "Novel ocular findings in oculodentodigital dysplasia (ODDD): a case report and literature review," *Ophthalmic Genetics*, vol. 40, no. 1, pp. 54–59, 2019.
- [18] M. A. M. van Steensel, L. Spruijt, I. van der Burgt et al., "A 2-bp deletion in the GJA1 gene is associated with oculodentodigital dysplasia with palmoplantar keratoderma," *American Journal of Medical Genetics Part A*, vol. 132a, no. 2, pp. 171–174, 2005.
- [19] R. J. Gorlin, L. H. Meskin, and J. W. S. Geme, "Oculodentodigital Dysplasia," *The Journal of Pediatrics*, vol. 63, no. 1, pp. 69–75, 1963.
- [20] M. Vreeburg, E. A. de Zwart-Storm, M. I. Schouten et al., "Skin changes in oculo-dento-digital dysplasia are correlated with C-terminal truncations of connexin 43," *American Journal of Medical Genetics. Part A*, vol. 143, no. 4, pp. 360–363, 2007.
- [21] E. Gumus, "A rare symptom of a very rare disease: a case report of a oculodentodigital dysplasia with lymphedema," *Clinical Dysmorphology*, vol. 27, no. 3, pp. 91–93, 2018.
- [22] E. I. Traboulsi and M. M. Parks, "Glaucoma in oculo-dento-osseous dysplasia," *American Journal of Ophthalmology*, vol. 109, no. 3, pp. 310–313, 1990.
- [23] T. Wiest, O. Herrmann, F. Stögbauer et al., "Clinical and genetic variability of oculodentodigital dysplasia," *Clinical Genetics*, vol. 70, no. 1, pp. 71–72, 2006.
- [24] K. K. Norton, J. C. Carey, and D. H. Gutmann, "Oculodentodigital dysplasia with cerebral white matter abnormalities in a two-generation family," *American Journal of Medical Genetics*, vol. 57, no. 3, pp. 458–461, 1995.
- [25] K. W. Park, H. S. Ryu, J. Kim, and S. J. Chung, "Oculodentodigital dysplasia presenting as spastic paraparesis: the first genetically confirmed Korean case and a literature review," *Journal of Movement Disorders*, vol. 10, no. 3, pp. 149–153, 2017.
- [26] G. Brice, P. Ostergaard, S. Jeffery, K. Gordon, P. S. Mortimer, and S. Mansour, "A novel mutation in GJA1 causing oculodentodigital syndrome and primary lymphoedema in a three generation family," *Clinical Genetics*, vol. 84, no. 4, pp. 378–381, 2013.



- [27] C. Vitiello, P. D'Adamo, F. Gentile, E. M. Vingolo, P. Gasparini, and S. Banfi, "A novel GJA1 mutation causes oculodentodigital dysplasia without syndactyly," *American Journal of Medical Genetics Part A*, vol. 133a, no. 1, pp. 58–60, 2005.
- [28] E. M. Vingolo, K. Steindl, R. Forte et al., "Autosomal dominant simple microphthalmos," *Journal of Medical Genetics*, vol. 31, no. 9, pp. 721–725, 1994.
- [29] G. F. Judisch, A. Martin-Casals, J. W. Hanson, and W. H. Olin, "Oculodentodigital dysplasia. Four new reports and a literature review," *Archives of Ophthalmology*, vol. 97, no. 5, pp. 878–884, 1979.
- [30] P. Debeer, H. van Esch, C. Huysmans et al., "Novel GJA1 mutations in patients with oculo-dento-digital dysplasia (ODDD)," *European Journal of Medical Genetics*, vol. 48, no. 4, pp. 377–387, 2005.
- [31] M. Thomsen, U. Schneider, M. Weber, and F. U. Niethard, "The different appearance of the oculodentodigital dysplasia syndrome," *Journal of Pediatric Orthopaedics. Part B*, vol. 7, no. 1, pp. 23–26, 1998.
- [32] R. J. J. van Es, D. Wittebol-Post, and F. A. Beemer, "Oculodentodigital dysplasia with mandibular retrognathism and absence of syndactyly: a case report with a novel mutation in the connexin 43 gene," *International Journal of Oral and Maxillofacial Surgery*, vol. 36, no. 9, pp. 858–860, 2007.
- [33] C. T. Schrandt-Stumpel, J. B. de Groot-Wijnands, C. de Die-Smulders, and J. P. Fryns, "Type III syndactyly and oculodentodigital dysplasia: a clinical spectrum," *Genetic Counseling*, vol. 4, no. 4, pp. 271–276, 1993.
- [34] E. Tasdelen, C. D. Durmaz, and H. G. Karabulut, "Autosomal recessive oculodentodigital dysplasia: a case report and review of the literature," *Cytogenetic and Genome Research*, vol. 154, no. 4, pp. 181–186, 2018.
- [35] A. Spaepen, C. Schrandt-Stumpel, J. P. Fryns, C. de Die-Smulders, M. Borghgraef, and H. van den Berghe, "Hallermann-Streiff syndrome: clinical and psychological findings in children. Nosologic overlap with oculodentodigital dysplasia?," *American Journal of Medical Genetics*, vol. 41, no. 4, pp. 517–520, 1991.
- [36] L. A. Brueton, S. M. Huson, B. Farren, and R. M. Winter, "Oculodentodigital dysplasia and type III syndactyly: separate genetic entities or disease spectrum?," *Journal of Medical Genetics*, vol. 27, no. 3, pp. 169–175, 1990.
- [37] O. Orosz, M. Fodor, I. Balogh, and G. Losonczy, "Relative anterior microphthalmos in oculodentodigital dysplasia," *Indian Journal of Ophthalmology*, vol. 66, no. 2, pp. 334–336, 2018.
- [38] R. E. Shapiro, J. W. Griffin, and O. C. Stine, "Evidence for genetic anticipation in the oculodentodigital syndrome," *American Journal of Medical Genetics*, vol. 71, no. 1, pp. 36–41, 1997.
- [39] T. Kogame, T. Dainichi, Y. Shimomura, M. Tanioka, K. Kabashima, and Y. Miyachi, "Palmoplantar keratosis in oculodentodigital dysplasia with a GJA1 point mutation out of the C-terminal region of connexin 43," *The Journal of Dermatology*, vol. 41, no. 12, pp. 1095–1097, 2014.
- [40] S. Nishat, Q. Mansoor, A. Javaid, and M. Ismail, "Oculodentodigital syndrome with syndactyly type III in a Pakistani consanguineous family," *Journal of Dermatological Case Reports*, vol. 6, no. 2, pp. 43–48, 2012.
- [41] C. Amador, A. M. Mathews, M. del Carmen Montoya, M. E. Laughridge, D. B. Everman, and K. R. Holden, "Expanding the neurologic phenotype of oculodentodigital dysplasia in a 4-generation Hispanic family," *Journal of Child Neurology*, vol. 23, no. 8, pp. 901–905, 2008.
- [42] F. U. Musa, P. Ratajczak, J. Sahu et al., "Ocular manifestations in oculodentodigital dysplasia resulting from a heterozygous missense mutation (L113P) in GJA1 (connexin 43)," *Eye (London, England)*, vol. 23, no. 3, pp. 549–555, 2009.
- [43] T. Loddenkemper, K. Grote, S. Evers, M. Oelerich, and F. Stögbauer, "Neurological manifestations of the oculodentodigital dysplasia syndrome," *Journal of Neurology*, vol. 249, no. 5, pp. 584–595, 2002.
- [44] S. C. Kelly, P. Ratajczak, M. Keller, S. M. Purcell, T. Griffin, and G. Richard, "A novel GJA 1 mutation in oculo-dento-digital dysplasia with curly hair and hyperkeratosis," *European Journal of Dermatology*, vol. 16, no. 3, pp. 241–245, 2006.
- [45] L. A. Gabriel, R. Sachdeva, A. Marcotty, E. J. Rockwood, and E. I. Traboulsi, "Oculodentodigital dysplasia: new ocular findings and a novel connexin 43 mutation," *Archives of Ophthalmology*, vol. 129, no. 6, pp. 781–784, 2011.
- [46] A. Jamsheer, M. Wisniewska, A. Szpak et al., "A novel GJA1 missense mutation in a Polish child with oculodentodigital dysplasia," *Journal of Applied Genetics*, vol. 50, no. 3, pp. 297–299, 2009.
- [47] T. Porntaveetus, C. Srichomthong, A. Ohazama, K. Suphapeetiporn, and V. Shotelersuk, "A novel GJA1 mutation in oculodentodigital dysplasia with extensive loss of enamel," *Oral Diseases*, vol. 23, no. 6, pp. 795–800, 2017.
- [48] A. Jamsheer, A. Sowińska-Seidler, M. Socha, A. Stembalska, C. Kiraly-Borri, and A. Latos-Bieleńska, "Three novel GJA1 missense substitutions resulting in oculo-dento-digital dysplasia (ODDD) - further extension of the mutational spectrum," *Gene*, vol. 539, no. 1, pp. 157–161, 2014.
- [49] A. Nivelon-Chevallier, D. Audry, F. Audry, and R. Dumas, "Oculo-dental-digital dysplasia: report of a case with spastic paraplegia," *Journal de Génétique Humaine*, vol. 29, no. 2, pp. 171–179, 1981.
- [50] W. E. Wooldridge, D. D. Anthony, E. R. Olson, G. P. Bates, and T. J. Sammon, "Oculodentodigital dysplasia," *Missouri Medicine*, vol. 74, no. 8, pp. 379–80, 383, 1977, 383.
- [51] K. W. Kjaer, L. Hansen, H. Eiberg, P. Leicht, J. M. Opitz, and N. Tommerup, "Novel connexin 43 (GJA1) mutation causes oculo-dento-digital dysplasia with curly hair," *American Journal of Medical Genetics*, vol. 127a, no. 2, pp. 152–157, 2004.
- [52] J. Honkaniemi, J. P. Kalkkila, P. Koivisto, V. Kähärä, T. Latvala, and K. Simola, "Letter to the editor: novel GJA1 mutation in oculodentodigital dysplasia," *American Journal of Medical Genetics. Part A*, vol. 139, no. 1, pp. 48–49, 2005.
- [53] S. Opjordsmoen and R. Nyberg-Hansen, "Hereditary spastic paraplegia with neurogenic bladder disturbances and syndactyly," *Acta Neurologica Scandinavica*, vol. 61, no. 1, pp. 35–41, 1980.
- [54] A. Pizzuti, E. Flex, R. Mingarelli, C. Salpietro, L. Zelante, and B. Dallapiccola, "A homozygous GJA1 gene mutation causes a Hallermann-Streiff/ODDD spectrum phenotype," *Human Mutation*, vol. 23, no. 3, p. 286, 2004.
- [55] K. Izumi, A. M. Lippa, A. Wilkens, H. A. Feret, D. McDonald-McGinn, and E. H. Zackai, "Congenital heart defects in oculodentodigital dysplasia: report of two cases," *American Journal of Medical Genetics Part A*, vol. 161a, no. 12, pp. 3150–3154, 2013.
- [56] C. L. Stanislaw, C. Narvaez, R. G. Rogers, and C. S. Woodard, "Oculodentodigital dysplasia with cerebral white matter



- abnormalities: an additional case," *Proceedings of the Greenwood Genetic Centre*, vol. 17, no. 1, pp. 20–24, 1998.
- [57] J. Choi, A. Yang, A. Song et al., "Oculodentodigital dysplasia with a novel mutation in GJA1 diagnosed by targeted gene panel sequencing: a case report and literature review," *Annals of Clinical and Laboratory Science*, vol. 48, no. 6, pp. 776–781, 2018.
- [58] J. P. Vasconcellos, M. B. Melo, R. B. Schimiti, N. C. Bressanim, F. F. Costa, and V. P. Costa, "A novel mutation in the GJA1 gene in a family with oculodentodigital dysplasia," *Archives of Ophthalmology*, vol. 123, no. 10, pp. 1422–1426, 2005.
- [59] D. M. Weintraub, J. L. Baum, and H. M. Pashayan, "A family with oculodentodigital dysplasia," *The Cleft Palate Journal*, vol. 12, pp. 323–329, 1975.
- [60] S. S. Gellis and M. Feingold, "Oculodentodigital dysplasia. Picture of the month," *American Journal of Diseases of Children*, vol. 128, no. 1, pp. 81–82, 1974.
- [61] C. Hadjichristou, V. Christophidou-Anastasiadou, A. Bakopoulou et al., "Oculo-dento-digital dysplasia (ODDD) due to a GJA1 mutation: report of a case with emphasis on dental manifestations," *The International Journal of Prosthodontics*, vol. 30, no. 3, pp. 280–285, 2017.
- [62] K. Amano, M. Ishiguchi, T. Aikawa et al., "Cleft lip in oculodentodigital dysplasia suggests novel roles for connexin43," *Journal of Dental Research*, vol. 91, 7\_suppl, pp. S38–S44, 2012.
- [63] L. Feller, N. H. Wood, M. D. Sluiter et al., "Report of a black South African child with oculodentodigital dysplasia and a novel GJA1 gene mutation," *American Journal of Medical Genetics Part A*, vol. 146a, no. 10, pp. 1350–1353, 2008.
- [64] A. Iтро, A. Marra, V. Urciuolo, P. Difalco, and A. Amodio, "Oculodentodigital dysplasia. A case report," *Minerva Stomatologica*, vol. 54, no. 7-8, pp. 453–459, 2005.
- [65] G. Tumminelli, I. di Donato, V. Guida, A. Rufa, A. de Luca, and A. Federico, "Oculodentodigital dysplasia with massive brain calcification and a new mutation of GJA1 gene," *Journal of Alzheimer's Disease*, vol. 49, no. 1, pp. 27–30, 2016.
- [66] D. Y. Park, S. Y. Cho, D. K. Jin, and C. Kee, "Clinical characteristics of autosomal dominant GJA1 missense mutation linked to oculodentodigital dysplasia in a Korean family," *Journal of Glaucoma*, vol. 28, no. 4, pp. 357–362, 2019.
- [67] R. Hayashi, T. Bito, M. Taniguchi-Ikeda, M. Farooq, M. Ito, and Y. Shimomura, "Japanese case of oculodentodigital dysplasia caused by a mutation in the GJA1 gene," *The Journal of Dermatology*, vol. 41, no. 12, pp. 1109–1110, 2014.
- [68] C. A. Wittlieb-Weber, K. M. Haude, C. T. Fong, and J. M. Vinocur, "A novel GJA1 mutation causing familial oculodentodigital dysplasia with dilated cardiomyopathy and arrhythmia," *HeartRhythm Case Reports*, vol. 2, no. 1, pp. 32–35, 2016.
- [69] A. Attig, M. Trabelsi, S. Hizem et al., "Oculo-dento-digital dysplasia in a Tunisian family with a novel GJA1 mutation," *Genetic Counseling*, vol. 27, no. 3, pp. 433–439, 2016.
- [70] M. Himi, T. Fujimaki, T. Yokoyama, K. Fujiki, T. Takizawa, and A. Murakami, "A case of oculodentodigital dysplasia syndrome with novel GJA1 gene mutation," *Japanese Journal of Ophthalmology*, vol. 53, no. 5, pp. 541–545, 2009.
- [71] N. P. Pace, V. Benoit, D. Agius et al., "Two novel GJA1 variants in oculodentodigital dysplasia," *Molecular Genetics & Genomic Medicine*, vol. 7, no. 9, article e882, 2019.
- [72] U. C. Parashari, S. Khanduri, S. Bhadury, and F. A. Qayyum, "Radiographic diagnosis of a rare case of oculo-dento-digital dysplasia," *South African Journal of Radiology*, vol. 15, no. 4, pp. 134, 2011.
- [73] P. Beighton, H. Hamersma, and M. Raad, "Oculodento-osseous dysplasia: heterogeneity or variable expression?," *Clinical Genetics*, vol. 16, no. 3, pp. 169–177, 1979.
- [74] D. C. Doshi, P. K. Limdi, N. V. Parekh, and N. R. Gohil, "Oculodentodigital dysplasia," *Indian Journal of Ophthalmology*, vol. 64, no. 3, pp. 227–230, 2016.
- [75] M. Frasson, N. Calixto, S. Cronemberger, R. A. L. Pessoa de Aguiar, L. L. Leão, and M. J. Burle de Aguiar, "Oculodentodigital dysplasia: study of ophthalmological and clinical manifestations in three boys with probably autosomal recessive inheritance," *Ophthalmic Genetics*, vol. 25, no. 3, pp. 227–236, 2004.
- [76] F. D. Gillespie, "A hereditary syndrome: dysplasia oculodentodigitalis," *Archives of Ophthalmology*, vol. 71, no. 2, pp. 187–192, 1964.
- [77] D. H. Gutmann, E. H. Zackai, D. McDonald-McGinn, K. H. Fischbeck, and J. Kamholz, "Oculodentodigital dysplasia syndrome associated with abnormal cerebral white matter," *American Journal of Medical Genetics*, vol. 41, no. 1, pp. 18–20, 1991.
- [78] G. J. Kurlander, N. W. Lavy, and J. A. Campbell, "Roentgen differentiation of the oculodentodigital syndrome and the Hallermann-Streif syndrome in infancy," *Radiology*, vol. 86, no. 1, pp. 77–86, 1966.
- [79] D. S. Levine, "Delayed gastric emptying and chronic diarrhea in a patient with oculodentodigital dysplasia syndrome," *Journal of Pediatric Gastroenterology and Nutrition*, vol. 5, no. 2, pp. 329–333, 1986.
- [80] M. Martínez-García, A. Bustamante-Aragónés, I. Lorda, and M. J. Trujillo-Tiebas, "Displasia oculodentodigital: consejo genético, opciones reproductivas y estudio molecular de un caso clínico referido para diagnóstico preimplantacional," *Medicina Clínica*, vol. 138, no. 13, pp. 592–593, 2012.
- [81] J. K. Mills, L. Wheeler, and S. N. Oishi, "A case of familial syndactyly associated with eye and dental abnormalities," *Jaapa*, vol. 28, no. 12, pp. 40–43, 2015.
- [82] S. Mosaed, B. H. Jacobsen, and K. Y. Lin, "Case report: imaging and treatment of ophthalmic manifestations in oculodentodigital dysplasia," *BMC Ophthalmology*, vol. 16, no. 1, 2016.
- [83] F. Owlia, M. H. Akhavan Karbassi, R. Hakimian, and M. S. Alemrajabi, "A highlighted case for emphasizing on clinical diagnosis for rare syndrome in third world," *Iranian Journal of Child Neurology*, vol. 11, no. 4, pp. 77–80, 2017.
- [84] P. Scheutzel, "Oculodentodigital syndrome: report of a case," *Dento Maxillo Facial Radiology*, vol. 20, no. 3, pp. 175–178, 1991.
- [85] J. A. Schneider, G. G. Shaw, and D. van Reken, "Congenital heart disease in oculodentodigital dysplasia," *Virginia Medical*, vol. 104, no. 4, pp. 262–263, 1977.
- [86] M. G. Schuller, M. L. Barnett, K. Strassburger, D. L. Friedman, and E. M. Sonnenberg, "Oculodentodigital dysplasia," *Oral Surgery, Oral Medicine, and Oral Pathology*, vol. 61, no. 4, pp. 418–421, 1986.
- [87] N. L. Sharma, R. C. Sharma, A. Goyal, B. K. Goyal, and K. R. Lakhanpal, "Oculodentodigital dysplasia with cutaneous keratotic papules," *Indian Journal of Dermatology, Venereology and Leprology*, vol. 48, no. 5, pp. 271–273, 1982.
- [88] H. S. Sugar, "Oculodentodigital dysplasia syndrome with angle-closure glaucoma," *American Journal of Ophthalmology*, vol. 86, no. 1, pp. 36–38, 1978.

- [89] P. Tejada, Y. W. Eduardo, E. Gutiérrez, A. Barceló, and J. Sánchez, “Glaucoma hereditario asociado a displasia oculo-dentodigital,” *Archivos de la Sociedad Española de Oftalmología*, vol. 86, no. 9, pp. 292–294, 2011.
- [90] C. J. Thoden, S. Ryoppy, and P. Kuitunen, “Oculodentodigital dysplasia syndrome. Report of four cases,” *Acta Paediatrica Scandinavica*, vol. 66, no. 5, pp. 635–638, 1977.
- [91] E. I. Traboulsi, B. M. Faris, V. M. D. Kaloustian, J. M. Opitz, and J. F. Reynolds, “Persistent hyperplastic primary vitreous and recessive oculo-dento- osseous dysplasia,” *American Journal of Medical Genetics*, vol. 24, no. 1, pp. 95–100, 1986.
- [92] National Center for Biotechnology Information, “ClinVar,” March 2020, <https://www.ncbi.nlm.nih.gov/clinvar/variation/VCV000137482.1>.
- [93] T. U. Consortium, “UniProt: a worldwide hub of protein knowledge,” *Nucleic Acids Research*, vol. 47, no. D1, pp. D506–D515, 2018.