Published in final edited form as: *J Diabetes Clin Res.*; 1(2): 53–58. doi:10.33696/diabetes.1.009.

Cut-off Value of Random Blood Glucose among Asian Indians for Preliminary Screening of Persons with Prediabetes and Undetected Type 2 Diabetes Defined by the Glycosylated Haemoglobin Criteria

Priscilla Susairaj¹, Chamukuttan Snehalatha¹, Arun Raghavan¹, Arun Nanditha¹, Ramachandran Vinitha¹, Krishnamoorthy Satheesh¹, Desmond G Johnston², Nicholas J Wareham³, Ambady Ramachandran^{1,*}

¹India Diabetes Research Foundation and Dr. A. Ramachandran's Diabetes Hospitals, Chennai, India

²Faculty of Medicine, Imperial College, London, UK

³Medical Research Council Epidemiology Unit, Institute of Metabolic Science, University of Cambridge, UK

Abstract

Aim—The increased morbidity and mortality due to type 2 diabetes can be partly due to its delayed diagnosis. In developing countries, the cost and unavailability of conventional screening methods can be a setback. Use of random blood glucose (RBG) may be beneficial in testing large numbers at a low cost and in a short time in identifying persons at risk of developing diabetes. In this analysis, we aim to derive the values of RBG corresponding to the cut-off values of glycosylated hemoglobin (HbA1c) used to define prediabetes and diabetes.

Methods—Based on their risk profile of developing diabetes, a total of 2835 individuals were screened for a large diabetes prevention study. They were subjected to HbA1c testing to diagnose prediabetes and diabetes. Random capillary blood glucose was also performed. Correlation of RBG with HbA1c was computed using multiple linear regression equation. The optimal cut-off value for RBG corresponding to HbA1c value of 5.7% (39 mmol/mol), and 6.5% (48 mmol/mol) were computed using the receiver operating curve (ROC). Diagnostic accuracy was assessed from the area under the curve (AUC) and by using the Youden's index.

Author Contribution Statement

Conflicts of Interest

All authors have no conflict of interest to declare.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited https://creativecommons.org/licenses/by/4.0/.

Correspondence should be addressed to Prof. A. Ramachandran; dr.ramachandran@ardiabetes.org.

CS, AN, DGJ, NJW, AR-Conceived and designed the experiments, PS, KS, NJW, AR - Performed the experiments, CS, PS, KS, -Analyzed the data, CS, AN, RV, PS, ArunR, DGJ, NJW, AR - Contributed to the writing of the manuscript, PS, CS, AN, AR - Wrote the first draft of the manuscript. PS, KS- Enrolled patients. All the authors agree with the manuscript's results and conclusions.

Results—RBG showed significant correlation with HbA1c (r=0.40, p<0.0001). Using the ROC analysis, a RBG cut-off value of 140.5 mg/dl (7.8 mmol/L) corresponding to an HbA1c value of 6.5% (48mmol/mol) was derived. A cut-off value could not be derived for HbA1c of 5.7% (39 mmol/mol) since the specificity and sensitivity for identifying prediabetes were low.

Conclusion—Use of a capillary RBG value was found to be a simple procedure. The derived RBG cut-off value will aid in identifying people with undiagnosed diabetes. This preliminary screening will reduce the number to undergo more cumbersome and invasive diagnostic testing.

Keywords

Random blood glucose; Screening for diabetes; Prediabetes; Glycosylated hemoglobin

Introduction

Type 2 diabetes (T2D) remains undiagnosed for many years in large number of persons living in developing countries [1]. The cost of diagnostic tests and the invasive procedures involved in conventional screening methods remain a major setback to timely testing. According to the recent estimates by the International Diabetes Federation (IDF), there are more than 42.2 million people with undiagnosed diabetes only in India [2], which accounts for more than half the number living with diabetes. Recommendations have been given by the World Health Organization (WHO) [3] and American Diabetes Association (ADA) [4] on screening strategies for high risk groups. Previously reported studies in the Western [5,6] and Chinese Populations [7] have evaluated cut-off points using fasting plasma glucose (FPG), HbA1c and random blood glucose (RBG) to define both prediabetes and undiagnosed diabetes. A large community-based screening program in India has studied the correlation of random capillary blood glucose with oral glucose tolerance test (OGTT) values to define cut-points for identifying diabetes and prediabetes [8]. Screening using random capillary blood glucose offers great benefits for testing large numbers, at low cost and in a short time. However, definitive cut-off points for RBG corresponding to the respective glycated hemoglobin (HbA1c) values for prediabetes and diabetes specific for Asian Indians are not clearly defined. In this analysis, we aim to derive these cut-off values of RBG corresponding to HbA1c used to define prediabetes (5.7%, 39 mmol/mol) and diabetes (6.5%, 48 mmol/mol) [9].

Materials and Methods

This is a post – hoc analysis of the primary prevention trial done in South India (Clinical Trials.Gov No. NCT01570946, Clinical Trials Registry – India Number: CTRI/ 2014/07/004799). The study design, recruitment strategy and the outcome of screening were described previously [10,11]. Briefly, the study cohort; men and women aged 35 to 55 years were prescreened for risk factors of diabetes using non-invasive diabetes risk assessment and then screened using HbA1c testing. Persons at a higher risk of conversion to diabetes, having HbA1c values between 6.0% (42 mmol/mol) and <6.5% (48 mmol/mol) were enrolled in the primary prevention study. They were randomized to receive either standard care with lifestyle advice at baseline or to the intervention arm to receive frequent text messages on their mobile phones on healthy lifestyle practices, in addition to the above. The study was

approved by the Ethics Committee of India Diabetes Research Foundation on 28th February 2012.

Recruitment of study participants was conducted between April 2012 and September 2013. From the 15 study sites a total of 6030 persons were identified to undergo initial prescreening using a non-invasive questionnaire method. At the prescreening, 2835 individuals were identified as having 3 risk factors for diabetes and were selected for the next stage of screening using HbA1c testing. The HbA1c test was performed using a pointof-care device (Bio-Rad in2itTM (I) System). This instrument has been standardized to the recommendations of the Diabetes Control and Complications Trial and certified by the National Glycohemoglobin Standardization Program. The HbA1c values obtained by the point-of-care analyzer and immunoturbidimetric assay used in our laboratory (Tina Quant II assay, Roche Diagnostics, Germany) yielded a significant correlation (r=0.887, p<0.0001).

Concurrently, capillary random plasma glucose was also measured using a glucometer (Accu-Check Performa, Roche Diagnostics, Mannheim Germany), calibrated to infer venous plasma glucose values [12]. Glucose estimations done with the glucometer and in the central laboratory using hexokinase method yielded a correlation coefficient of 0.89, p<0.0001. The testing devices were calibrated each day at the worksite and quality control was performed from time to time. Prescreening and screening tests were done on the same day 30 to 45 minutes post breakfast. RBG values were available for 2835 individuals.

Statistical analysis

Correlation of RBG with HbA1c was computed using multiple linear regression equation corrected for age and gender. The optimal cut-off values for RBG corresponding to HbA1c values of 5.7% (39 mmol/mol), (cut-off for prediabetes) and 6.5% (48 mmol/mol), (cut-off for diagnosing diabetes) were computed using the receiver operating characteristic curve (ROC). The sensitivity and specificity of RBG for detecting diabetes and prediabetes based on HbA1c were calculated by the area under the curve (AUC). Diagnostic accuracy was calculated using the Youden's index.

Results

Prescreening was conducted for a total of 6030 persons, to verify eligibility for the prevention trial. Among them 2835, (men: 2226 women: 609; mean age 45.9 ± 5.6 years) with 3 risk factors for diabetes were eligible for the HbA1c and RBG testing. Table 1 shows the characteristics of the study cohort and the presence of risk variables. Of the risk factors, obesity (81.7%), sedentary lifestyle (63.2%) and abdominal obesity in men (60.6%) were predominantly noted.

RBG showed significant correlation with HbA1c (r=0.40, p<0.0001). The regression equation was: y(HbA1c)=0.013*RBG+4.012. Using the ROC analysis, the cut-off value for RBG corresponding to an HbA1c of 5.7% (39 mmol/mol) (n=1121) was 113.5 mg/dl (6.3 mmol/L) (sensitivity 61%, specificity 61%, p<0.0001). The AUC was $0.662 \pm SE 0.10 (95\% CI 0.642-0.682)$, (Figure 1).

Cut-off value for RBG was 113.5 mg/dl (6.3 mmol/L) (sensitivity 61%, specificity 61%, p<0.0001). AUC = 0.662 ± SE 0.10 (95% CI 0.642-0.682). ROC: Receiver Operating Characteristic Curve; RBG: Random Blood Glucose; AUC: Area Under the Curve.

Figure 2 shows that the cut-off value for RBG corresponding to HbA1c of 6.5% (48 mmol/mol) (n=283) was 140.5 mg/dl (7.8 mmol/L) (sensitivity 69%, specificity 83%, p<0.0000). The AUC was $0.823 \pm SE 0.16$ (95% CI 0.792-0.854). Based on the derived sensitivity and specificity, the Youden's index wasabove 50%.

Cut-off value for RBG was 140.5 mg/dl (7.8 mmol/L) (sensitivity 69%, specificity 83%, p<0.0000). AUC = $0.823 \pm SE \ 0.16$ (95% CI 0.792-0.854). ROC: Receiver Operating Characteristic Curve; RBG: Random Blood Glucose; AUC: Area Under the Curve.

Discussion

In this analysis, we intended to study the utility of RBG as a preliminary screening tool to identify persons at risk of developing diabetes and requiring further confirmatory diagnostic testing. The use of HbA1c test definitely has several advantages over that of blood glucose; but the higher cost of the test may limit its utility in large epidemiological studies. It may also be impractical in places where reliable test methods of HbA1c are not available. Therefore, the feasibility of using RBG in place of HbA1c or even a timed glucose measurement such as FPG or 2h PG values may be more advantageous and practical. In order to overcome this limitation, we assessed the cut-off values for RBG corresponding to the HbA1c values of 5.7% (39 mmol/mol) and 6.5% (48 mmol/mol) using ROC procedures. RBG values of 113.5 mg/dl (6.3 mmol/L) and 140.5 mg/dl (7.8 mmol/L) respectively were derived for prediabetes and diabetes defined by the above HbA1c values. However, it was noted that a RBG cut-off value of 113.5 mg/dl (6.3 mmol/L) did not show an acceptable level of sensitivity and specificity, and hence cannot be used to define prediabetes.

An earlier study conducted in the US, tested the performance of four screening methods for diabetes and dysglycemia based on risk score, capillary blood glucose measurement and a combination of both [6]. The study reported that a casual capillary blood glucose value of 140 mg/dl had a sensitivity of 65% and specificity of 96% for diagnosis of diabetes based on fasting and 2-hour glucose criterion. Similarly, a casual blood glucose value of 120mg/dl had a sensitivity of 62% and specificity of 90% for fasting plasma glucose of 110mg/dl.

Based on a comparison of RBG and corresponding oral glucose values, in a large number of randomly chosen individuals without a history of diabetes, it was suggested for Asian Indians with a RBG of >110 mg/dl (6.1 mmol/L) at screening can be recommended for definitive testing [8]. This research team, also from the same city had reported that RBG cutpoint of 140 mg/dl (7.8 mmol/L) corresponded to the 2h PG 200mg/dl (11.1 mmol/L) used in diagnosis of diabetes. Values of 119 mg/dl (6.6 mmol/L) corresponded to the diagnostic value of 2h PG of 140 mg/dl (7.8 mmol/L) and 113 mg/dl (6.3 mmol/L) corresponded to impaired fasting glucose (ADA) value of 100mg/dl (5.6 mmol/L).

Our cut-off value for diabetes 140.5 mg/dl (7.8 mmol/L) does agree with the reported observations from the various studies discussed here. However, the derived cut-off value of

113.5 mg/dl (6.3 mmol/L) did not have acceptable sensitivity and specificity to identify prediabetes.

Conclusion

The use of RBG testing with convenience of sampling at any time during the day could substantially reduce time and cost of screening both at a community level and in a clinical setting. However, the limitation lies in the accuracy of the glucose monitors used and other factors such as postprandial time and age that may influence the performance of the RBG test as reported by Rolka et al. [6]. It is also noted that the relation between blood glucose and the HbA1c value be influenced by several factors such as race and level of hemoglobin.

We report than an RBG value of 140 mg/dl (7.8 mmol/L) could be used to identify people who should be subjected to confirmatory OGTT to diagnose diabetes. RBG measurement may be applied if HbA1c or timed glucose testing are not practical for large scale screening of undiagnosed diabetes.

Acknowledgments

We thank the epidemiology team of the India Diabetes Research Foundation in coordinating for data collection and Mrs. Vijaya Balaji for the statistical support rendered in this study.

Source of Financial Support

This trial (Ref. No. 58/1/6/ICMR-MRC/2009-NCD-II) was funded by the Indian Council of Medical Research– Medical Research Council Joint Initiative Chronic Non-Communicable Diseases Research Funding.

References

- Ramachandran A, Ma RC, Snehalatha C. Diabetes in Asia. Lancet. 2010; 375:408–8. [PubMed: 19875164]
- 2. International Diabetes Federation. IDF Diabetes Atlas. 8th edn. Brussels, Belgium: International Diabetes Federation; 2017. http://diabetesatlas.org/en/ [Last Accessed 14th August 2019]
- 3. Report of a World Health Organization Consultation. Use of glycated haemoglobin (HbA1c) in the diagnosis of diabetes mellitus. Diabetes Res Clin Pract. 2011; 93:299–9.
- American Diabetes Association. Standards of Medical Care in Diabetes—2018. Diabetes Care. 2018; 41(Supplement 1):S13–S27. [PubMed: 29222373]
- Ziemer DC, Kolm P, Foster JK, Weintraub WS, Vaccarino V, Rhee MK, et al. Random plasma glucose in serendipitous screening for glucose intolerance: screening for impaired glucose tolerance study 2. J Gen Intern Med. 2008; 23:528–5. [PubMed: 18335280]
- Rolka DB, Narayan KM, Thompson TJ, Goldman D, Lindenmayer J, Alich K, et al. Performance of recommended screening tests for undiagnosed diabetes and dysglycemia. Diabetes Care. 2001; 24:1899–3. [PubMed: 11679454]
- Zhou X, Pang Z, Gao W, Wang S, Zhang L, Ning F, et al. Performance of an A1C and fasting capillary blood glucose test for screening newly diagnosed diabetes and pre-diabetes defined by an oral glucose tolerance test in Qingdao, China. Diabetes care. 2010; 33:545–50. [PubMed: 20007941]
- Somannavar S, Datta M, Ganesan A, Deepa M, Mohan V. Random Capillary Blood Glucose Cut Points for Diabetes and Pre-Diabetes Derived From Community-Based Opportunistic Screening in India. Diabetes Care. 2009; 32:641–3. [PubMed: 19073758]
- 9. International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. Diabetes Care. 2009; 32:1327–4. [PubMed: 19502545]

- Priscilla S, Nanditha A, Simon M, Satheesh K, Kumar S, Shetty AS, et al. A pragmatic and scalable strategy using mobile technology to promote sustained lifestyle changes to prevent type 2 diabetes in India—Outcome of screening. Diabetes Res Clin Pract. 2015; 110:335–40. [PubMed: 26547503]
- Thomson H, Oliver N, Godsland IF, Darzi A, Srivanichakorn W, Majeed A, et al. Protocol for a clinical trial of text messaging in addition to standard care versus standard care alone in prevention of type 2 diabetes through lifestyle modification in India and the UK. BMC Endocr Disord. 2018; 18:63. [PubMed: 30200935]
- Freckmann G, Schmid C, Baumstark A, Pleus S, Link M, Haug C. System accuracy evaluation of 43 blood glucose monitoring systems for self-monitoring of blood glucose according to DIN EN ISO 15197. J Diabetes Sci Technol. 2012; 6:1060–5. [PubMed: 23063032]



Figure 1. ROC showing predictive performance of RBG corresponding to an HbA1c value of 5.7% (39mmol/mol).

Susairaj et al.



Figure 2. ROC showing predictive performance of RBG corresponding to an HbA1c value of 6.5% (48 mmol/mol).

Table 1

Baseline characteristics of study cohort.

	Total n=2835 Men: 2226 (78.5%) Women: 609 (21.5%)
Characteristics (mean ± SD)	
Age in years	45.9 ± 5.6
Weight (kg)	75.4 ± 11
Body mass index (BMI) (kg/m ²)	27.7 ± 3.4
Waist Circumference (cm)	95.2 ± 7.9
Blood Pressure (mmHg)	
Systolic	129.3 ± 17.2
Diastolic	82.4 ± 10.7
HbA1c (%, mmol/mol)	$5.9 \pm 0.89, 41$
Random Blood Glucose (mg/dl, mmol/L)	126.9 ± 42.8, 7.04
Presence of Risk Factors n (%)	
Positive Family History	1508 (53.2)
Overweight BMI 23.0-24.5 kg/m ²	419 (14.8)
Obese BMI 25.0 kg/m ²	2317 (81.7)
Abdominal obesity in men (90cm)	1718 (60.6)
Abdominal obesity in women (80cm)	577 (20.4)
History of Prediabetes	196 (6.9)
Hypertension – newly diagnosed (140/90mmHg)	663 (23.4)
Hypertension – Known	621 (21.9)
Sedentary Lifestyle	1793 (63.2)
Data presented are mean ± SD, n (%)	•