




Complete Circular Genome Sequence of a *mecB*- and *mecD*-Containing Strain of *Micrococcus canis*

Sybille Schwendener,^a  Vincent Perreten^a

^aInstitute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Bern, Switzerland

ABSTRACT The complete genome sequence of *Micrococcus canis* strain 19/EPI0118, isolated from a veterinary clinic environment in Switzerland, was determined using hybrid assembly of Oxford Nanopore and Illumina reads. 19/EPI0118 harbored the methicillin resistance genes *mecB* and *mecD* on a staphylococcal cassette chromosome *mec* element and a *Micrococcus* chromosomal resistance island, respectively.

Micrococcus canis is a Gram-positive bacterium which has been found on the skin of healthy dogs and in infection sites (1, 2). Some *M. canis* strains carry the methicillin resistance gene *mecB* on staphylococcal cassette chromosome *mec* (SCC*mec*) elements or on a plasmid (3, 4). The *mecD* gene has so far only been associated with chromosomal resistance islands McRI_{*mecD*} in *Micrococcus caseolyticus* and an SCC*mec* element in *Micrococcus bohemicus* (5–7). Recently, a methicillin-resistant *M. canis* strain (19/EPI0118) isolated from the environment of a companion animal clinic in Switzerland was found to contain both the *mecB* and *mecD* genes (8). The strain was also resistant to fusidic acid (8). We report here the complete genome sequence of 19/EPI0118 and the genetic elements containing *mecB*, *mecD*, and *fusC*.

Genomic DNA of 19/EPI0118 (cryopreserved in our collection) was extracted from culture grown overnight at 37°C on Trypticase soy agar containing 5% sheep blood, using the MasterPure complete DNA and RNA purification kit (Lucigen, Middleton, WI). DNA sequencing was performed using a NEBNext Ultra II directional DNA library with TruSeq adapters on an Illumina NovaSeq 6000 system (2 × 150-bp paired-end reads) at Eurofins Genomics (Germany) and in-house with a MinION system (Oxford Nanopore Technologies [ONT]; R9.4.1 SpotON flow cell, MinION MK1b device). The ONT library was prepared using the 1D ligation sequencing kit (SQK-LSK109) and the native barcoding expansion kit (EXP-NBD104) (Oxford Nanopore Technologies, UK). The ONT reads were base called and demultiplexed using Guppy software v4.4.1 and end trimmed and size filtered using Cutadapt v2.5. The Illumina reads were used without filtering after confirmation of high-quality and adaptor-free sequences by FastQC v0.11.7 analysis. The genome sequence of 19/EPI0118 was *de novo* assembled and circularized using Unicycler v0.4.8 run with default parameters, using paired-end Illumina reads (8,580,460 reads, 1,287,069,000 total bases, 543× coverage) and ONT reads larger than 10 kb (22,513 reads; mean, 20,422 bp; 460,021,332 total bases; 194× coverage) (9). All calculations were performed on the high-performance computing (HPC) cluster UBELIX at the University of Bern (<http://www.id.unibe.ch/hpc>). The genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline service (10).

The complete genome of 19/EPI0118 contained a circular chromosome of 2,359,135 bp and four small circular plasmids (3,561, 2,426, 1,888, and 1,436 bp) with a GC content of 36.5%. A total of 2,362 protein-coding sequences and 78 RNAs, including 58 tRNAs and 5 16S-23S rRNA clusters, were predicted. The *mecB* gene was found on a pseudo (Ψ) SCC*mec* (ΨSCC*mecB*_{19/EPI0118}) integrated at the 3' end of the chromosomal

Citation Schwendener S, Perreten V. 2021. Complete circular genome sequence of a *mecB*- and *mecD*-containing strain of *Micrococcus canis*. Microbiol Resour Annuoc 10:e00408-21. <https://doi.org/10.1128/MRA.00408-21>.

Editor Frank J. Stewart, Montana State University

Copyright © 2021 Schwendener and Perreten. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Vincent Perreten, vincent.perreten@vetsuisse.unibe.ch.

Received 22 April 2021

Accepted 18 May 2021

Published 3 June 2021




FIG 1 The genetic structure of Ψ SCC $_{mecB_{19/EPI0118}}$ and $McRI_{mecD-1}$ elements in *M. canis* 19/EPI0118. (A) (Ψ)SCC $_{mecB}$ elements in *M. canis*. Chromosomal regions from the 23S rRNA (pseudouridine[1915]-N[3])-methyltransferase gene (*rlmH*) to the UvrABC system protein A gene (*uvrA*) were compared for *M. canis* strains KM45013^T (GenBank accession number [CP021059](#)), 19/EPI0118 ([CP072837](#)), and Epi0076A ([CP047363](#)). The *rlmH* gene (gray) contains at the 3' end the integration site sequence (ISS) for SCC. Further copies of ISS that subdivide composite (Ψ)SCC $_{mecB}$ elements are indicated. The ISS have the following sequences for 19/EPI0118: ISS1, 5'-GAAAGTTATCATAAATGA; ISS2, 5'-GAAAGTTATCATAAGTGA; ISS3, 5'-TGGGTATATCAAAAATAA; they flank the Ψ SCC $_{mecB_{19/EPI0118}}$ ([CP072837](#), positions 32,310 to 45,199) and Ψ SCC ([CP072837](#), positions 45,200 to 69,684). (B) Comparison of $McRI_{mecD-1}$ of *M. canis* 19/EPI0118 ([CP072837](#), positions 259,551 to 277,684) with $McRI_{mecD-1}$ of *M. caseolyticus* IMD0819 ([CP021058](#)). The chromosomal region downstream of the tRNA pseudouridine(38-40) synthase gene (*truA*) was compared. The attachment site (*att*) for $McRI_{mecD}$ at the 3' end of the 30S ribosomal protein S9 gene (*rpsI*) (gray) and copies of it that delimitate chromosomal islands are indicated. The core *att* sequence is 5'-GAACGTAA(A/G)AA(A/G)CCAGGTCTTAAAGGCGCTCGTCGTTACACAGTCTCAAACGTTAAT (positions containing an A or G base are given in parenthesis). Genes are represented by arrows and are color-coded as follows: red, *mec* operon genes and fusidic acid resistance gene; green, *ccr* genes; light green, integrase (*int*) and associated genes of $McRI_{mecD}$; yellow, other recombinase genes; blue, putative hemolysin genes (*hlgBC*). Figures were generated using Easyfig software (11). Gray connections indicate regions with between 72% and 100% nucleotide sequence identity.

rlmH gene, followed by a Ψ SCC (Fig. 1A). Cassette chromosome recombinase (*ccr*) genes were not present, and the *mecB* operon lacked the β -lactamase *bla_{Zm}* gene. The fusidic acid resistance gene *fusC* was found at the 3' end of Ψ SCC $_{mecB_{19/EPI0118}}$ (Fig. 1A). The *mecD* gene was part of an $McRI_{mecD-1}$ integrated at the 3' end of the *rpsI* gene (Fig. 1B). This element was, except for three mismatches, identical to the $McRI_{mecD-1}$ of *M. caseolyticus* IMD0819 (5).

These findings showed that a *Macrocooccus* strain can carry two methicillin resistance genes in its chromosome and suggests exchange of $McRI_{mecD}$ between *M. canis* and *M. caseolyticus*.

Data availability. The complete genome sequence of *M. canis* strain 19/EPI0118 has been deposited in GenBank under accession numbers [CP072837](#), [CP072838](#), [CP072839](#), [CP072840](#), and [CP072841](#). The associated BioProject and BioSample accession numbers are [PRJNA720016](#) and [SAMN18630837](#), respectively. The raw reads were deposited in the SRA database with accession numbers [SRR14161194](#) (Illumina) and [SRR14161193](#) (ONT).

ACKNOWLEDGMENTS

We are grateful to Alexandra Collaud for technical assistance.

This project was financed by internal funds of the Institute of Veterinary Bacteriology of the University of Bern.

REFERENCES

1. Gobeli Brawand S, Cotting K, Gómez-Sanz E, Collaud A, Thomann A, Brodard I, Rodriguez-Campos S, Strauss C, Perreten V. 2017. *Macrocococcus canis* sp. nov., a skin bacterium associated with infections in dogs. *Int J Syst Evol Microbiol* 67:621–626. <https://doi.org/10.1099/ijsem.0.001673>.
2. Cotting K, Strauss C, Rodriguez-Campos S, Rostaer A, Fischer NM, Roosje PJ, Favrot C, Perreten V. 2017. *Macrocococcus canis* and *M. caseolyticus* in dogs: occurrence, genetic diversity and antibiotic resistance. *Vet Dermatol* 28:559–e133. <https://doi.org/10.1111/vde.12474>.
3. Gómez-Sanz E, Schwendener S, Thomann A, Gobeli Brawand S, Perreten V. 2015. First staphylococcal cassette chromosome *mec* containing a *mecB*-carrying gene complex independent of transposon Tn6045 in a *Macrocococcus canis* isolate from a canine infection. *Antimicrob Agents Chemother* 59:4577–4583. <https://doi.org/10.1128/AAC.05064-14>.
4. Chanchaithong P, Perreten V, Schwendener S. 2019. *Macrocococcus canis* contains recombinogenic methicillin resistance elements and the *mecB* plasmid found in *Staphylococcus aureus*. *J Antimicrob Chemother* 74:2531–2536. <https://doi.org/10.1093/jac/dkz260>.
5. Schwendener S, Cotting K, Perreten V. 2017. Novel methicillin resistance gene *mecD* in clinical *Macrocococcus caseolyticus* strains from bovine and canine sources. *Sci Rep* 7:43797. <https://doi.org/10.1038/srep43797>.
6. Schwendener S, Nigg A, Collaud A, Overesch G, Kittl S, Phumthanakorn N, Perreten V. 2019. Typing of *mecD* islands in genetically diverse methicillin-resistant *Macrocococcus caseolyticus* strains from cattle. *Appl Environ Microbiol* 85:e01496-19. <https://doi.org/10.1128/AEM.01496-19>.
7. Schwendener S, Keller JE, Overesch G, Perreten V. 2021. Novel SCCmec element containing the methicillin resistance gene *mecD* in *Macrocococcus bohemicus*. *J Glob Antimicrob Resist* 24:360–362. <https://doi.org/10.1016/j.jgar.2021.02.001>.
8. Schmidt JS, Kuster SP, Nigg A, Dazio V, Brilhante M, Rohrbach H, Bernasconi OJ, Budel T, Campos-Madueno EI, Gobeli Brawand S, Schuller S, Endimiani A, Perreten V, Willi B. 2020. Poor infection prevention and control standards are associated with environmental contamination with carbapenemase-producing *Enterobacterales* and other multidrug-resistant bacteria in Swiss companion animal clinics. *Antimicrob Resist Infect Control* 9:93. <https://doi.org/10.1186/s13756-020-00742-5>.
9. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
10. Tatusova T, DiCuccio M, Badretdin A, Chetvermin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
11. Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer. *Bioinformatics* 27:1009–1010. <https://doi.org/10.1093/bioinformatics/btr039>.