

A novel knee prosthesis model of implant-related osteomyelitis in rats

Niels H Sørensen¹, Nina Vendel Jensen², Birgit Meinecke Nürnberg³, Asger Lundorff Jensen⁴, Janne Koch⁵, Steen Seier Poulsen⁶, Gerald Pier⁷, and Helle Krogh Johansen⁸

¹Hand Section, Department of Planned Surgery Z, Gentofte University Hospital; ²Department of Anaesthesiology, Intensive Care and Operations, Gentofte University Hospital; ³Department of Pathology, Roskilde University Hospital; ⁴Faculty of Health and Medical Science, University of Copenhagen; ⁵Faculty of Health and Medical Science, University of Copenhagen; ⁶Biomedical Department, Panum Institute, Copenhagen University, Denmark; ⁷Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; ⁸Department of Clinical Microbiology, Rigshospitalet, Copenhagen University, Denmark.

Correspondence: nsn@dadlnet.dk

Submitted 12-03-24. Accepted 12-09-07

Background and purpose There have been numerous reports of animal models of osteomyelitis. Very few of these have been prosthesis models that imitate human conditions. We have developed a new rat model of implant-related osteomyelitis that mimics human osteomyelitis, to investigate the pathology of infection after orthopedic implant surgery.

Methods 2 wild-type strains of *Staphylococcus aureus*, MN8 and UAMS-1, and their corresponding mutants that are unable to produce poly-N-acetyl glucosamine (PNAG) (*ica::tet*) were injected into the medullary canals of the femur and tibia at 3 different doses: 10^2 , 10^3 , and $> 10^4$ CFU/rat. We measured clinical signs, inflammatory markers, radiographic signs, histopathology, and bacteriology in the infected animals.

Results An inoculum of at least 10^4 cfu of either wild-type bacterial strain resulted in histological, bacteriological, and radiographic signs of osteomyelitis with loosening of the prosthesis. An inoculum of 10^3 CFU gave signs of osteomyelitis but the prosthesis remained in situ. Bacterial inocula of 10^2 cfu gave no signs of osteolysis.

Interpretation We have established a new knee prosthesis model that is suitable for reliable induction of experimental implant-associated osteomyelitis with the prosthesis in situ, using a small inoculum of *S. aureus*. At a dose of 10^3 CFU/rat, bacteria unable to produce PNAG (*ica::tet*) had only minor defects in their virulence.

Infections associated with in-dwelling orthopedic devices can be difficult to cure without removing the device, and they are quite expensive to manage (Darouiche 2004). Usual infection-control measures, laminar air flow, and use of systemic antimicrobial prophylaxis have not completely eliminated orthopedic implant-related infections (Darouiche 2003). *Staphylo-*

coccus aureus is the predominant bacterium associated with infected metal implants (Eron 1985). There is an urgent need to explore other approaches to enhance diagnosis, prevention and treatment of infection, and osteomyelitis—for which appropriate animal models are extremely useful (Monzón et al. 2002). The aim of the present study was to establish an implant model of acute osteomyelitis associated with metallic implants without any promoters of infection other than the implant itself.

Materials and methods

S. aureus strains and bacterial challenge (Table 1)

We used the MN8 strain of *S. aureus*, originally obtained from a patient with toxic shock syndrome-1 (TSST-1), and the genetically related UAMS-1 strain of *S. aureus*, a primary isolate from a patient admitted to the Arkansas Children's Hospital with an osteoarticular and soft-tissue infection and septic shock (Lucke et al. 2003, Cassat et al. 2006). UAMS-1 is a clinical isolate from osteomyelitis and is therefore clearly relevant to the model. Strain MN8 is closely related to UAMS-1 genetically: both belong to the EMRSA-16 group of *S. aureus*, which are characterized by an identical mutation in the alpha-toxin gene that results in loss of production of this factor. *ica*-deficient strains were used to evaluate the role of PNAG antigen, which is synthesized by proteins encoded at the *ica* locus (Cramton et al. 1999), to determine whether it is a dominant feature of implant-related infections caused by *S. aureus*. To date, 100% of over 100 clinical isolates from all sources tested have been found to express poly-N-acetyl glucosamine (PNAG), which contributes to this organism's ability to make biofilms. Biofilms are thought to be a major component of the pathogenesis of implant-related infections.

Table 1. Inoculum, *ica:tet* status, and strain

Group	n	Inoculum (CFU)	<i>ica</i> status	Strain
Control	5	0		÷
G1	3	10 ²	+	MN8
G2	9	10 ³	+	MN8
G3	4	10 ³	–	MN8
G4	11	≥ 10 ⁴	+	MN8
G5	3	10 ²	+	UAMS-1
G6	6	10 ³	+	UAMS-1
G7	4	10 ³	–	UAMS-1
G8	9	≥ 10 ⁴	+	UAMS-1

Thus, PNAG may be a virulence factor in implant infections and is a known target for immunotherapy and vaccination.

41 rats were infected with 2 wild-type strains of *S. aureus* MN8 or UAMS-1, both of which are *ica*-positive, using inoculating doses of bacteria of 10⁵, 10⁴, 10³, or 10² CFU/rat (MN8: n = 23; UAMS-1: n = 18). There were at least 3 rats in each challenge group.

Viable counts were done on the different bacterial strains to determine the exact dose inoculated. Both strains were also tested using mutants with deleted *ica* genes (*ica::tet*), which are referred to as *ica*-negative strains, at an infecting dose of 10³ CFU/rat with four rats in each group (Cramton et al. 1999, Beenken et al. 2004, Kropec et al. 2005). 5 rats were followed without being infected (the control group). The animals were followed for 2–6 weeks and then killed.

Anesthesia

All rats were sedated with a subcutaneous injection of hypnorm/dormicum, 0.3 mL/100 g body weight, given preoperatively and reinjected every 15 min (0.15 mL/100 g). After the operation, a femoralis block of the operated extremity was placed below the inguinal ligament using 1% lidocain/0.5% bupivacain in 1 mL. The rats were killed with an intracardiac injection of 2 mL pentobarbital (200 mg/mL).

Surgery

We used 54 Sprague-Dawley male rats that were 7–9 weeks-old (Taconic Europe) and weighing about 300 g. The experiments were approved by the Animal Ethics Committee of Denmark (reg. no. 2005/561-1049 from September 19, 2005).

The skin over the left knee was sterilized twice with alcohol. The fur was removed with a hair razor. The knee was opened with a parapatellar medial incision and the tendon with the patella was dislocated laterally. The articulating cartilage was osteotomized with bone scissors from the distal femur, with the proximal tibia including the menisci and cruciate ligaments protecting the collateral ligaments. A 2-mm-wide and a 10-mm-deep hole was bored with a hand drill into the femur and tibia to fit the joint components. The joint capsule and skin were closed with Ethibond 4-0 and Vicryl 5-0 after placement of the knee prosthesis (press-fit model) without bone cement.



Figure 1. Non-constrained knee prosthesis.

Prosthesis

A rat-sized, non-constrained knee prosthesis was used (Figure 1) The femoral component was made from a metal alloy and the tibial component was milled from high-density polyethylene stock.

Infection process

The infection was initiated by direct injection of 10 µL of the appropriate *S. aureus* bacterial suspension into the medullary canals of the femur and tibia. The volume of the suspension fitted into the marrow hole. Afterwards the condylar prosthesis was inserted.

Radiographic evaluation

AP and lateral radiographs were taken on days 0 (OP), 7, 14, 21, and 42. To assess development and progression of bone infection, a modified scoring system was used (Schmidmaier et al. 2006). The following 6 parameters were scored: (I) periosteal reaction, (II) osteolysis, (III) soft-tissue swelling, (IV) deformity, (V) general impression or destruction, (VI) loosening of the prosthesis, and (VII) sequestrum formation. The score for each of the first 5 parameters used a scale of 0 to 4: 0 (absent), 1 (mild), 2 (moderate), 3 (severe). Parameter VI “loosening of the prosthesis” was judged as 0 (none), 1 (1 component), or 2 (both components), and parameter VII was judged as 0 (absent) or 1 (present). The maximum score that could be achieved was 18 (Figures 2 and 3).

Microbiological evaluation

The prosthesis components were explanted and rolled over non-selective media (5% Danish blood agar and chocolate agar plates (State Serum Institute, Copenhagen, Denmark)). Isolated bacteria were identified as previously described (Højby and Frederiksen 2000). The same procedure was used for bone from the femurs and tibias and for the synovialis, which were rolled over Danish blood agar and chocolate agar in 3 directions and incubated at 37°C. Bacterial growth was judged as no growth, growth in the first streak (+), growth in the first 2 streaks (++), or growth in all 3 streaks (+++).




Figure 2. Prosthesis in situ without osteolysis or bone destruction. Control group




Figure 3. Severe inflammation with loosening of the prosthesis and bone destruction. *S. aureus* strain MN8, 10^4 CFU (score = 13).




Figure 4. Severe inflammation with intramedullary abscess in the femur (score = 8).

Bone and soft-tissue histology (Figure 4)

After removal of the prosthesis, the remnants of the tibia, femur, and synovialis were fixed in 4% buffered paraformaldehyde and decalcified in 10% formic acid/EDTA for 7 days. Samples were embedded in paraffin and transverse sections of 5 μ m, including the implantation site, were cut on a microtome. The sections were stained with hematoxylin and eosin. For the first 30 rats, histological analysis of tissues from the blood, liver, and spleen was also carried out.

Semiquantitative scoring of all specimens was performed blind by pathologists who were not aware of the treatment groups (SSP and BMN).

For histological scoring of severity of inflammation, transverse sections of the tibia and femur (with the prosthesis removed) and tissue from the synovialis were investigated. Each of the 3 tissues (femur, tibia, and synovialis) was given a score ranging from 0 to 4, depending on the severity of inflammation. 0 meant no signs of inflammation, 1 was slight focal accumulation of inflammatory cells (neutrophils), 2 was a moderate but consistent inflammation in the transverse sections and/or moderate inflammation of the entire circumference around the cavity after prosthesis removal, 3 was the start of formation of an abscess in the cavity, and 4 was abscess formation and destruction of bone material with the synovialis completely infiltrated by neutrophils. The scores from the 3 separate tissues were added, giving a maximum score of 12.

Biochemical analysis

α -1 acid glycoprotein (AGP)—an acute-phase protein—was measured on days 0 (preoperatively), 7, 14, 21, and 42 (normal range: 0–130) (Matsumoto et al. 2007). Hemoglobin and leucocytes were also measured. There were no differences in the groups before and after the operation, irrespective of the bacterial strain and the infectious dose used.

Clinical evaluation

Body weight and temperature were determined, and well-being of the animal was estimated using food intake as a parameter. Macroscopic judgement of the knee was scored as described previously (Petty et al. 1988). Briefly, normal = 0, moderate swelling and edema = 0.5, swelling and abscess = 1.0, and abscess and illness of the animal = 1.5. If abscesses and/or fistulae were present, pus was aspirated and cultured on blood agar plates, and any bacteria were identified (Høiby 2000).

Statistics

All results are expressed as means. They were analyzed using the Kruskal-Wallis test for one-way analysis of variance. Statistical calculations were performed using SPSS for Windows software version 12.2. All p-values less than 0.05 were considered to be statistically significant.

Results

Radiography

No osteolysis or signs of inflammation were seen in any of the radiographs in the control group, including loosening of the prosthesis (Figure 1). Radiographic changes in groups G1 and G5 (rats injected with 10^2 bacteria) indicated slight inflammation with little osteolysis and no loosening of the prosthesis. The modified scores (Schmidmaier et al. 2006) after 1 and 2 weeks were 2.9 and 3.6, respectively. Radiographs from most rats in groups G2 and G6 (injected with 10^3 bacteria, *ica*+) showed slight to moderate inflammation with osteolysis, a periosteal reaction, and soft-tissue swelling but the prosthesis remained in situ. The scores for G2 were 1.4 and 5.6 after 1 and 2 weeks, respectively, and for G6 they were 0 and 5.3 after 1 and 2 weeks, respectively. The rats infected with 10^3 CFU of the 2 different *ica*+ *S. aureus* strains MN8 and UAMS-1

Table 2. Radiographic scores

Week	Group	Inoculum (CFU)	Strain	Mean score
1	Control	0	–	0.5
2	Control	0	–	2.6
1	G1	10 ²	MN8	2.3
2	G1	10 ²	MN8	3.2
1	G2	10 ³	MN8	1.4
2	G2	10 ³	MN8	5.6
1	G3	<i>ica</i> - 10 ³	MN8	6.0
2	G3	<i>ica</i> - 10 ³	MN8	7.4
1	G4	≥ 10 ⁴	MN8	5.0
2	G4	≥ 10 ⁴	MN8	7.1
1	G5	10 ²	UAMS	3.2
2	G5	10 ²	UAMS	4.0
1	G6	10 ³	UAMS	0
2	G6	10 ³	UAMS	5.3
1	G7	<i>ica</i> - 10 ³	UAMS	6.0
2	G7	<i>ica</i> - 10 ³	UAMS	5.5
1	G8	≥ 10 ⁴	UAMS	7.2
2	G8	≥ 10 ⁴	UAMS	7.8

(groups G3 and G7) had the same level of virulence as judged from the same scores of 7.4 and 5.5 after 2 weeks, respectively. The radiographic images of most rats in groups G4 and G8 (injected with > 10⁴ *ica*+ *S. aureus* MN8 or UAMS-1, respectively) showed moderate to severe inflammation and most of the prostheses had loosened or were no longer in place (n = 12). For G4, mean scores after 1 and 2 weeks were 5.0 and 7.1, respectively, and for G8 mean scores of 7.2 and 7.8 were measured after 1 and 2 weeks (Table 2).

Microbiology

No bacteria were cultured from the control group; thus, the prostheses were sterile. In G1 and G5 inoculated (with 10² CFU *S. aureus*), only 2 of 6 cultures obtained from the knees yielded a few colonies of *S. aureus* and these were seen only with the UAMS-1 strain. In G2 (wild-type strain MN8, 10³ CFU) 8 of 12 implant cultures had a moderate number of colonies (++) and in G6 (wild-type strain UAMS-1, 10³ CFU), all 6 cultures were positive (++) in G3 (*ica*- *S. aureus* MN8, 10³ CFU), 3 of 4 cultures were positive at the ++ level. In G7 (*ica*- *S. aureus* UAMS-1, 10³ CFU), all 4 cultures yielded colonies at the ++ level. In G4 and G8 (wild-type MN8 and wild-type UAMS-1, > 10⁴ CFU), all cultures were positive for *S. aureus* at the +++ level. No bacteria could be cultured from spleen, liver, or blood.

Histopathology

The histopathology was consistent with the radiographic findings. No signs of inflammation were found in the control group. No histological signs of inflammation or bacteria were found in the liver, spleen, or blood in any of the groups. In the 2 groups that were given an inoculum of 10² (G1 and G5), all histological sections showed minor bone inflammation around the prosthesis with little bone destruction. In G2 and G6 (wild-

Table 3. Histopathology scores

Group	n	Inoculum (CFU)	Mean score (0–12)	Strain
Control	5		0.0	
G1	3	10 ²	2.7	MN8
G2	6	10 ³	5.4	MN8
G3	4	10 ³ <i>ica</i> -neg	6.5	MN8
G4	11	≥ 10 ⁴	> 9	MN8
G5	3	10 ⁴	5.7	UAMS-1
G6	6	10 ³	5.3	UAMS-1
G7	4	10 ³ <i>ica</i> -neg	7.2	UAMS-1
G8	8	10 ⁴	> 9	UAMS-1

type MN8 and wild-type UAMS-1 strains), the histopathology sections showed typical signs of bone inflammation (osteomyelitis) such as destruction of cortical and cancellous bone (osteolysis) and new bone formation. In G3 and G7 (the *S. aureus ica*-groups) the same findings were obtained but with less osteolysis. In G4 and G8 (wild-type MN8 and wild-type UAMS-1, > 10⁴ CFU), there were massive signs of bone infection with destruction and abscess formation. 4 rats were killed or died preoperatively because of aspiration (Table 3). The 2 groups infected with more than 10⁴ CFU were treated as 1 group for further analysis.

Biochemistry and blood leukocyte counts

In the control group, the preoperative AGP levels were < 200 µg/mL. In G2 (wild-type MN8, 10³ CFU), the AGP increased from a mean of 170 µg/mL to 542 µg/mL after 2 weeks, and then decreased to a mean of 424 after 4 weeks. In G6 (wild-type UAMS-1, 10³ CFU), the AGP increased from a mean of 168 to 218 after 2 weeks. The rats cleared the infection after 2 weeks. Hemoglobin and leucocyte levels were not statistically significantly different from those in the control group, and the biochemistry parameters after 2 weeks and the leukocyte counts and hemoglobin levels were similar. Only the AGP parameter was relevant for measurement of inflammation.

Clinical results

There was no statistically significant difference in mean weight loss between the control group and the G1 to G3 (CFU ≤ 10³ and G5 to G7 rats. However, the weight loss in G4 and G8 rats given the highest dose (> 10⁴ CFU) was higher than in the control group. There was no soft-tissue swelling around the knee in the groups given 10² CFU (G1 and G5). Body temperature was similar in the different groups.

Discussion

Various experimental animal models of osteomyelitis have been developed since 1884 when Rodet and Lexer (Rodet 1885, Lexer 1894) demonstrated that osteomyelitis similar to

that seen in humans could be created experimentally by injection of staphylococci to produce bone abscesses. Mader (1985) and Norden (1988), together with several other authors, demonstrated that osteomyelitis could be induced in an implant model by injection of bacteria into the tibia and femur cavities using either rats or rabbits (Mader 1985, Rissing 1990, An et al. 2006). Osteomyelitis studies related to orthopedic implant and prosthetic joint infection have been conducted using various animals such as rats (Zak et al. 1982, Rissing et al. 1985), rabbits (Nelson et al. 1990), dogs (Fokushima et al. 2005), and guinea pigs (Ofluoglu et al. 2007). The in vivo animal models of osteomyelitis have certain advantages, which have provided us with a clearer understanding of the osteomyelitis process and treatments. The limitations of the models were mostly due to differences in the immune systems of the animals, the non-physiological way the bacteria were inoculated into the animals, the short observation periods, and the high bacterial doses used (Mader 1985). Rats are the second most widely used animal for these studies, which is why we also used them. In addition, it is simple to evaluate the histological and microbiological bone processes in rats. Development of the surgical procedure and use of loop-glasses and a specially designed small non-constrained knee prosthesis allows evaluation of the osteomyelitis process in spite of the small size of the animal.

In most osteomyelitis models, lesions are induced by sclerosing agents, arachidonic acid, and foreign bodies containing bacteria to facilitate bone infection. Nelson et al. (1990) and Fukushima et al. (2005) did not use sclerosing agents and foreign bodies in their rat osteomyelitis models. They infected the animals with a combination of *Pseudomonas aeruginosa* and *S. aureus*. Fukushima et al. (2005) examined the relationship between the inoculating dose of bacteria and the progression of osteomyelitis, and found that development of significant histological and radiographic signs of osteomyelitis required inocula of at least 6×10^3 CFU. Other authors have used foreign-body implants such as stainless steel pins and K-wires implanted into the bone marrow together with the bacteria to study the osteomyelitis process (Hudetz et al. 2008, Li et al. 2008).

In our rat model the foreign body was implanted as a knee prosthesis, to mimic the human condition in which the osteomyelitis process is initiated in the bone marrow around the prosthesis. Osteomyelitis was seen after 1 week; it was localized to the primary infected area without spreading to the liver, spleen, or blood. We found that inoculation of at least 10^3 CFU of 2 wild-type *S. aureus* isolates resulted in significant histological, bacteriological, and radiographic signs of osteomyelitis with loosening of the prostheses. Smeltzer et al. (1997) found that development of radiographic and histologic signs of osteomyelitis required inocula of at least 10^4 CFU. They used the same strain as we used (UAMS-1). We also found that the prosthesis remained in situ with an inoculum of 10^3 CFU, but the prosthesis was expelled after inoculation

with $> 10^4$ CFU (for either *S. aureus* strain). For both MN8 and UAMS-1, 10^3 CFU was the optimal inoculum for study of the osteomyelitis process around the prostheses.

The *ica* gene is a major operon for expression of a potential virulence factor, a polysaccharide involved in staphylococcal biofilm formation (PNAG). However, the role of biofilms in staphylococcal bone infection is still unclear (Knobben et al. 2007), and they may not require the PNAG polysaccharide for formation. Both of the *ica*-strains gave rise to osteomyelitis that was no different to that achieved with the wild-type strains at the same size of inoculum (Table 2). Hudetz et al. (2008) found that the presence of *ica* genes had a strong effect on biofilm formation in vitro and a weak effect in vivo.

Overall, it appears that *S. aureus* causes persistent infection in a knee prosthesis implanted into rats, requiring a small number of bacteria. The tissue pathology and infection were independent of the presence of the *ica* genes and PNAG-dependent biofilm formation. We have also shown that after 2 weeks, the rats can clear the inflammation with doses between 10^3 and 10^5 , indicating that it is not possible to study the course of inflammation for longer time. The immune system of the rat is quite different from the human immune system, which may explain their ability to clear the infection after injection of such relatively high doses. Even so, our study shows that a dose of 10^3 CFU of *S. aureus* would be a suitable experimental condition with which to study active and passive immunization against *S. aureus* in the same knee prosthesis model.

NHS: planning, surgery on the animals (with prostheses), and writing. NVJ: surgery on the animals (with prostheses), organization of data in tables, and planning of surgery. BMN: conducted the pathology and histological scoring analysis. ALJ: designed the analysis and testing to generate the biochemical data. JK: helped with veterinary problems, took care of the animals, and obtained clinical data on the animals. SSP: participated in generating the pathology and histological scoring data, along with the histopathological micrographs. GP: provided the bacterial strains, helped with immunological problems, edited the manuscript. HKJ: generated all of the microbiological data and scored inflammation.

We thank Prof. Mark Smeltzer for sending the UAMS-1 strains of *S. aureus*, Ulla Johansen for technical assistance, and Prof. Lars Dahlin for valuable comments on the manuscript.

No competing interests declared.

An Y H, Kang Q K, Arciola C R. Animal models of osteomyelitis. Review. Int J Artificial Organs. 2006; 29 (4): 407-20.

Beenken K E, Dunman P M, McAleese F, et al. Global gene expression in *Staphylococcus aureus* biofilms. J Bacteriol 2004; 186 (14): 4665-84.

Cassat J, Dunman P M, Smeltzer M S, et al. Transcriptional profiling of *Staphylococcus aureus* clinical isolate and its isogenic agr and sar-A mutants reveals global differences in comparison to the laboratory strain RN6390. Microbiology 2006; 152: 3075-90.

- Cramton S E, Ulrich M, Götz F, et al. Anaerobic Conditions Induce Expression of Polysaccharide Intercellular Adhesin in *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Infect Immun* 2001; 69 (6): 4079-85.
- Darouiche R O. Antimicrobial approaches for preventing infections associated with surgical implants. *Clin Infect Dis* 2003; 36: 1284-9.
- Darouiche R O. Treatment of infections associated with surgical implants. *N Engl J Med* 2004; 350: 1422-9.
- Eron L J. Prevention of infection following orthopaedic surgery. *Antibiot Chemother* 1985; 33: 140-64.
- Fokushima N, Yokoyama K, Sasahara T, et al. Establishment of rat model of acute staphylococcal osteomyelitis: relationship between inoculation dose and development of osteomyelitis. *Arch Orthop Trauma Surg incl Arthroscopy and Sports Med* 2005; 125 (3): 169-76.
- Hudetz D, Hudetz S U, Harris L G, et al. Week effect of metal type and ica genes on staphylococcal infection of titanium and stainless steel implants. *Clin Microbiol Infect* 2008; 14: 1135-45.
- Høiby N, Frederiksen B. Microbiology of cystic fibrosis. In: *Cystic Fibrosis*, 2nd edition (eds Hodson M E, Geddes D M). London: Arnold: 2000:83-107.
- Knobben B A, van der Mai H C, van Horn J R, et al. Transfer of bacteria between biomaterials surfaces in the operating room – an experimental study. *J Biomed Mater Res A* 2007; 80: 790-9.
- Kropec A, Maira-Litran T, Jefferson K K, et al. Poly-N-acetylglucosamine production in *Staphylococcus aureus* is essential for virulence in murine models of systemic infection. *Infect Immun* 2005; 73 (10): 6868-76.
- Lexer E. Zur experimentellen erzeugung osteomyelitischer herde. *Arch Klin Chir* 1894; 48: 181-200.
- Li D, Gromow K, Søballe K, et al. quantitative mouse model of implant-associated osteomyelitis and the kinetics of microbial growth, osteolysis and humoral immunity. *J Orthop Research* 2008; 26: 96-105.
- Lucke M, Schmidmaier G, Wildemann B, et al. A new model of implant-related Osteomyelitis in rats. *J Biomed Mater Res B Appl Biomater* 2003; 67 (1): 593- 602.
- Mader J T. Animal models of osteomyelitis. *Am J Med* 1985; 78: 213-7.
- Matsumoto K, Nishi K, Kadowaki D, et al. Alpha1-acid glycoprotein suppresses Rat acute inflammatory paw oedema through the inhibition of neutrophils activation and prostaglandin E2 generation. *Biol Pharm Bull* 2007; 30 (7): 1226-30.
- Monzón M, Garcia-Alvarez F, Laclériga A, Amorena B. Evaluation of four experimental osteomyelitis infection models by using precolonized implants and bacterial suspensions. *Acta Orthop Scand* 2002; 73 (1): 11-9.
- Nelson D R, Buxton T B, Luu Q N, Rissing J P. The promotional effect of bonewax on experimental *Staphylococcus aureus* osteomyelitis. *J Thorac Cardiovasc Surg* 1990; 99: 977-80.
- Norden C W. Lessons learned from animal model of osteomyelitis. *Rev Infect Dis* 1988; 10: 103-10.
- Ofluoglu E A, Zileli M, Aydin D, et al. Implant-related infection model of rat spine. *Arch Orthop Trauma Surg* 2007; 127: 391-6.
- Petty W, Spanner S, Shuster J. *J Bone Joint Surg (Am)* 1988; 70: 536-9.
- Rissing J P. Animal models of osteomyelitis; knowledge, hypothesis, and speculation. *Infect. Dis Clin North Am* 1990; 4: 377-90.
- Rissing J P, Buxton T B, Weinstein R S, et al. Model of experimental chronic osteomyelitis in rats. *Infect. Immun* 1985; 47: 581-6.
- Rodet A. Physiologie pathologique – etude expérimentale sur l'ostéomyélite infectieuse. *C R Acad Sci* 1885; 99: 569-71.
- Schmidmaier G, Luke M, Wildermann B, et al. Prophylaxes and treatment of implant-related infections by antibiotic-coated implants; a review *Int J Care Injury*; 2006; 37: 105-12.
- Smeltzer M S, Thomas J R, Hickmon S G, et al. Characterization of rabbit model of staphylococcal osteomyelitis. *J Orthop Res* 1997; 15: 414-21.
- Zak O, Zak F, Rich R, et al. Experimental staphylococcal osteomyelitis in rats; therapy with rifampin and cloxacillin alone or in combination. In: Perty P, Grassi GG(eds) *Current chemotherapy and immunotherapy*. Am Soc for Microbiology, Washington DC 1982: 973-4.