

The Impact of Anesthetic Management Under Bispectral Index Monitoring on the Early Recovery Quality of Elderly Patients Undergoing Laparoscopic Surgery: A Blinded Randomized Controlled Trial

Min Shi , Yufang Long*, Zhihu Zhou, Liqiao Huang, Danping Wu, Xu Zhang 

Department of Anesthesiology, The Second Affiliated Hospital of Guilin Medical University, Guilin, 541199, People's Republic of China

*These authors contributed equally to this work

Correspondence: Xu Zhang, Department of Anesthesiology, The Second Affiliated Hospital of Guilin Medical University, No. 212 Renmin Road, Guilin, 541199, People's Republic of China, Tel +86 189 3477 9973, Email zhangxu@glmc.edu.cn

Purpose: To comprehensively evaluate the impact of bispectral index (BIS) monitoring-guided anesthesia depth on the early recovery quality in elderly patients undergoing laparoscopic surgery.

Methods: Ninety patients aged ≥ 60 years scheduled for laparoscopic surgery under general anesthesia were randomized into three groups: Group C (empirically guided anesthesia), Group B1 (BIS-guided, target range 40–60), and Group B2 (BIS-guided, target range 50–60). Blinded researchers evaluated recovery quality (QoR-15), pain (VAS), and sleep (RCSQ) preoperatively and on postoperative days 1, 2, 3, and 7 (POD_{1,2,3,7}). Postoperative delirium was assessed with CAM (POD₁–POD₃), and cognitive function (MMSE) was measured preoperatively, POD₃, and POD₇. Intraoperative data included vital signs, BIS values, anesthetic dosage, emergence/extubation times, PACU stay, and adverse events within three days post-surgery. Time to first ambulation and hospital stay were also recorded.

Results: Compared with Group C, Group B1 and B2 had lower propofol consumption, shorter emergence/extubation times, and higher BIS values (T₂–T₅ and overall mean) ($P < 0.05$). QoR-15 scores improved on POD₂ in Group B1 and on both POD₁ and POD₂ in Group B2 ($P < 0.05$). The RCSQ scores increased on POD₁ and POD₃ in Group B1 ($P < 0.05$) and on POD₁, POD₂ and POD₃ in Group B2 ($P < 0.05$). In addition, Group B2 had a shorter PACU stay and time to first postoperative ambulation ($P < 0.05$). No differences were found in the incidence of postoperative delirium, POCD, or MMSE scores among the three groups. Compared to Group B1, Group B2 exhibited shorter emergence and extubation times, elevated BIS values at T₃ and T₅, a higher mean BIS value throughout surgery, and enhanced QoR-15 scores on POD₁ and POD₂ ($P < 0.05$).

Conclusion: BIS monitoring-guided anesthesia management can enhance early recovery from laparoscopic surgery in elderly patients with BIS values within a safe range, which may be particularly advantageous for this demographic during laparoscopic procedures.

Keywords: bispectral index, BIS, elderly patients, quality of postoperative recovery, cognitive function, sleep quality

Introduction

With the development of surgical treatment techniques, more elderly patients receive operations and the postoperative survival time has been prolonged, thus the quality of postoperative recovery has received increasing attention.^{1,2} Because of the decline in organ function and comorbidities, elderly patients have a poor ability to tolerate surgery and anesthesia, which can lead to various postoperative complications, thus anesthesia management appears to have certain particularities and complexities.³ Large deviations in intraoperative anesthetic management and preoperative physiological

compensatory function will contribute to a significant increase in the incidence of serious postoperative complications as well as mortality.⁴ With the purpose of improving perioperative management and quality of early postoperative recovery among elderly patients; the primary task for anesthesiologists is to optimize anesthesia management to shorten the perioperative non-physiological process and create conditions for the rapid recovery of postoperative organ and system function under the premise of maintaining the preoperative physiological status and compensatory function of elderly patients. One solution is to optimize the anesthesia plan in terms of providing short-acting sedation and analgesic drugs to control the depth of anesthesia and provide an effective anti-stress response, ensuring that elderly patients are fully awake when extubating, facilitating postoperative recovery of physiological state and early out-of-bed mobilization, and avoiding mechanical ventilation support.

Depth control is an important part of general anesthesia management and is closely related to patient safety and postoperative prognosis. Especially for elderly patients, the depth of anesthesia is difficult to control due to their increased sensitivity to general anesthesia drugs and decreased rate of drug clearance.⁵ Bispectral index (BIS) monitoring is considered as an accurate, sensitive, real-time and convenient indicator for judging the depth of sedation under anesthesia because of its association with the depth of sedation and with the state of consciousness of patients.^{6,7} Intraoperative monitoring of BIS level is helpful to predict the prognosis of patients and reduce the complications related to depth of sedation,⁸ thus perioperative use of BIS to monitor the depth of anesthesia may shed light on promoting the quality of postoperative recovery in elderly surgical patients.

Enhanced recovery after surgery (ERAS) provides a new platform and mode for patients undergoing surgical treatment, and a large number of clinical practices based on the ERAS concept have demonstrated that individualized and accurate anesthesia management is essential to improve the quality of postoperative recovery in elderly patients.^{9,10} Recent studies have shown that the use of BIS on the depth of anesthesia monitoring can reduce the dosage of intraoperative sedative drugs, shorten emergence time and extubation time, and decrease adverse stress reactions.¹¹ However, most previous clinical studies only evaluated the quality of early postoperative recovery in terms of emergence time, pain, or other adverse reactions, which have been insufficient to comprehensively reflect the postoperative recovery quality of patients. The QoR-15 scale is a patient-centered, highly reliable and valid questionnaire with good compliance that broadly assesses the quality of postoperative recovery of patients from five dimensions, which are emotional state, physical comfort, psychological support, physical independence, and pain.¹² Currently, no known research has focused on using QoR-15 to assess the effect of BIS monitoring on the quality of postoperative recovery in elderly patients. Moreover, previous studies on elderly patients have focused on a wider range of BIS values (40–60), it remains poorly understood whether slightly higher BIS values (50–60) are beneficial.

Postoperative cognitive function recovery is essential for evaluating the quality of recovery in elderly patients. Postoperative delirium (POD) and postoperative cognitive dysfunction (POCD) are common postoperative complications in elderly surgical patients. Early assessment and active intervention are effective measures to prevent and manage POD and POCD.^{13–15} However, the effect of intraoperative depth of anesthesia monitoring through BIS grade on postoperative cognitive function remains controversial. Furthermore, there is a close relationship between sleep quality and cognitive function, and postoperative sleep disturbance is a significant factor affecting postoperative recovery in elderly patients,¹⁶ which deserves further attention.

In summary, by applying the QoR-15, Confusion Assessment Method (CAM), Mini-Mental State Examination (MMSE), and Richards-Campbell Sleep Questionnaire (RCSQ), this study aimed to comprehensively investigate the effect of depth of anesthesia monitoring through the BIS on early recovery quality among elderly patients after laparoscopic surgery to provide a reference for improving the quality of anesthesia and early postoperative recovery among elderly patients and create good conditions for improving postoperative prognosis and outcome.

Materials and Methods

Study Subject

The institutional review board of the Second Affiliated Hospital of Guilin Medical University approved this prospective, randomized, double-blind study, which was registered in the Chinese Clinical Trial Registry (ChiCTR2200055416).

Ninety patients aged ≥ 60 years who received general anesthesia for laparoscopic surgery were recruited between January and June 2022. All the participants and their families provided written informed consent after a detailed understanding of the study objectives, procedures, and potential risks. This study adhered to the Declaration of Helsinki, ensuring ethical standards in medical research involving human subjects, encompassing informed consent, participant safety, and scientific validity.

The inclusion criteria were as follows: (1) age ≥ 60 years, (2) scheduled for elective laparoscopic surgery, (3) expected anesthesia time of >2 hours, (4) grade I to III according to the American Society of Anesthesiologists (ASA) physical status classification, and (5) signed informed consent. Patients were excluded if they (1) had a history of severe neurological or psychiatric disorders, (2) had alcohol and drug abuse, (3) had severe organ dysfunction, (4) had preoperative sleep disorders, (5) received other test drugs or participated in other clinical trials 3 months before surgery, and (6) had severe hearing and visual impairment that could not cooperate with the examination. The withdrawal criteria were as follows: (1) serious intraoperative or postoperative adverse reactions that affected the quality of surgery or were life-threatening, (2) operation time > 6 h, (3) intraoperative blood loss > 1000 mL, and (4) voluntary withdrawal from the clinical trial.

Grouping

Ninety patients were randomly assigned using a simple randomization method into three groups: Group C (control group with empirically guided anesthesia), Group B1 (BIS-guided anesthesia with a target range of 40–60), and Group B2 (BIS-guided anesthesia with a target range of 50–60). BIS values were monitored in all patients using a bispectral index monitor (Covidien IIC, Mansfield, MA, USA). The patients in Group C received an empirically guided depth of anesthesia. The BIS values were blinded to the attending anesthesiologist throughout the anesthesia procedure, with the instrument screen either backward to the anesthesiologist or covered with a sticker. Patients in Groups B1 and B2 received anesthesia management guided by BIS monitoring, with target BIS values ranging from 40 to 60 and 50 to 60, respectively. The group assignments were blinded to the patients, PACU staff, follow-up staff, and the data recorders. Only the anesthesiologist in charge was blinded, and the participants were managed as assigned.

Anesthetic Procedures and Postoperative Management

All patients underwent routine fasting and liquid fasting before surgery, and were administered preoperative medication according to the actual condition. After admission to the operating room, venous access was established and noninvasive blood pressure, electrocardiogram, and pulse oxygen saturation were monitored. Before induction of anesthesia, all patients were monitored using the BIS, and data were recorded every minute. After the BIS values were stable, anesthesia was induced by intravenous injection of midazolam 0.04–0.1 mg/kg, sufentanil 0.3–0.6 μ g/kg, cisatracurium 0.1–0.2 mg/kg, and etomidate 0.15–0.2 mg/kg. Mechanical ventilation was performed after successful tracheal intubation with a tidal volume of 8–10 mL/kg and respiratory rate of 10–14 breaths/min. Pressure of end-tidal CO₂ (PetCO₂) was monitored, and the respiratory rate was adjusted as necessary to maintain it within the range of 35–45 mmHg. Anesthesia maintenance was achieved through total intravenous anesthesia, combining propofol (4–12 mg/kg/h) and remifentanyl (0.05–0.2 μ g/kg/min), supplemented with intermittent doses of cisatracurium (0.03–0.05 mg/kg) for muscle relaxation.

Intraoperative management was performed according to groups following routine clinical anesthesia. Based on the principles of patient safety and stable vital signs, noninvasive blood pressure changes were maintained within 20% of those before induction. In addition to common causes, anesthesia should be appropriately deepened or reduced if abnormal vital signs are related to the depth of anesthesia.

After all anesthetics were withdrawn at the end of surgery, patient-controlled intravenous analgesia (PCIA) was administered. The patients were then admitted to the PACU and the endotracheal tube was removed after adequate recovery of consciousness and spontaneous breathing.

Observation Indicators

Preoperative general indicators included age, sex, height, weight, body mass index (BMI), ASA of Anesthesiologists Classification, and educational level.

Surgery- and anesthesia-related information included operation time, anesthesia time, dosage of anesthetic drugs, blood loss, urine volume, infusion volume, time from the end of surgery to awakening and extubation, and duration of PACU stay. Mean arterial pressure (MAP), heart rate (HR), and BIS values were recorded before anesthesia induction (T_0), 5 min after intubation (T_1), at the beginning of surgery (T_2), 1 h after initiation of surgery (T_3), 2 h after initiation of surgery (T_4), and at the end of surgery (T_5). The BIS data were exported from 5 minutes after the completion of intubation to the end of surgery, and the mean BIS values during this period were calculated. Intraoperative events included bradycardia ($HR < 50$ bpm), tachycardia ($HR > 100$ bpm), hypertension ($MAP > 105$ mmHg or MAP increase $\geq 20\%$ of baseline value), hypotension ($MAP < 60$ mmHg or MAP decrease $\geq 20\%$ of baseline value), and cardiovascular medication remedies.

Postoperative quality of recovery, intensity of pain, and sleep quality were evaluated using the QoR-15, Visual Analog Score (VAS), and RCSQ at Preop, POD₁, POD₂, POD₃, and POD₇. CAM was used to assess delirium within three days after surgery. The MMSE was conducted to observe the occurrence of POCD preoperatively and on, POD₃ and POD₇. Adverse reactions, as well as surgery- and anesthesia-related complications, were followed up and recorded within 3 days after surgery. The time to first postoperative ambulation and postoperative hospital stay were recorded.

Statistical Analysis

This randomized controlled study used the QoR-15 score on POD1 as the primary outcome, hypothesizing superiority in BIS-guided groups (B1 and B2) over Group C. Pilot data ($n=10$ per group) estimated scores of 100.6 ± 7.85 (C), 107.7 ± 9.36 (B1), and 109.5 ± 7.72 (B2), consistent with ranges reported by Liu et al.¹⁷ The minimum clinically important difference for QoR-15 was 8.0.¹⁸ Using PASS 15 software (NCSS LLC, Kaysville, Utah, USA) with a mean difference of 9 (B2 vs C), pooled SD of 8.5, $\alpha=0.05$, and 90% power (two-tailed), we calculated 27 patients per group. Adjusting for a 10% dropout rate, we enrolled 30 patients per group, totaling 90 patients.

Data normality was assessed using the Kolmogorov–Smirnov test. Continuous variables are expressed as mean \pm standard deviation or median and interquartile range (IQR) for normally and non-normally distributed variables, respectively. Categorical variables are presented as n (%). Numerical differences between the two groups were assessed using the chi-square test or Fisher's exact test for categorical variables and analysis of variance or the Kruskal–Wallis test for continuous variables. Statistical significance (P value) was assessed using a 2-tailed test in all instances, and the threshold for significance was set at $P < 0.05$. All statistical analyses were performed using SPSS (version 25.0 (SPSS Inc., Chicago, IL, USA)) and GraphPad Prism software (version 8.0; GraphPad Software, San Diego, California, USA).

Results

Preoperative General Information

A total of 132 patients were screened, and 90 eligible patients were divided into Groups C ($n=30$), B1 ($n=30$), and B2 ($n=30$). Based on the inclusion and exclusion criteria, 83 patients were included and analyzed in this study. Figure 1 shows a detailed flowchart.

As shown in Table 1, there were no significant differences among the three groups in terms of sex, age, height, weight, BMI, ASA classification, or education level at baseline ($P > 0.05$).

Intraoperative Information

Surgery and Anesthesia

No significant differences were found among the three groups in terms of specialty distribution, surgical duration, anesthetic duration, intraoperative sufentanil, remifentanil, cisatracurium dosage, infusion volume, blood loss, urine volume, or incidence of intraoperative hemodynamic adverse events ($P > 0.05$). Compared with Group C, the dosage of propofol was significantly reduced in Group B1 and Group B2 ($P < 0.05$). In addition, emergence and extubation times




Figure 1 Flowchart of patient recruitment, randomization and analysis.

were significantly shortened in Group B1 and Group B2 ($P<0.05$), and PACU stay was significantly shorter in Group B2 ($P<0.05$). In addition, compared with Group B1, Group B2 had statistically shorter emergence time and extubation time ($P<0.05$). The details are presented in Table 2 and Figure 2.

Table 1 Patients Characteristics

	C (n=28)	B1 (n=28)	B2 (n=27)	χ^2/F	P
Gender (male/female)	11/17	13/15	11/16	0.326	0.849
Age (yr)	67.82±4.73	68.00±5.56	68.52±5.42	0.130	0.878
Height (cm)	162.07±7.78	161.64±6.08	160.00±7.92	0.613	0.544
Weight (kg)	64.76±9.97	61.33±8.93	63.63±10.31	0.905	0.409
BMI (kg/m ²)	24.55±2.64	23.42±2.77	24.79±3.26	1.763	0.178
ASA class I/II/III	6/18/4	5/19/4	4/18/5	0.630	0.960
Education Level (yr)	6 (1.75)	6 (1)	6 (2)	0.188	0.910

Table 2 Information of Operation and Anesthesia

	C (n=28)	B1 (n=28)	B2 (n=27)	$\chi^2/F/H$	P
Distribution of specialties					
Gynaecology	12(42.9%)	11(39.3%)	10(37.0%)	1.552	0.956
Urology	5(17.9%)	7(25%)	8(29.6%)		
Hepatology	5(17.9%)	6(21.4%)	5(18.5%)		
Gastroenterology	6(21.4%)	4(14.35)	4(14.8%)		
Surgical duration (min)	164(49.75)	158.5(60.5)	161(61)	0.042	0.979
Anesthetic duration (min)	190(51.75)	194.5(63)	196(71)	0.399	0.819
Infusion volume (mL)	2217.9±448.1	2314.3±535.6	2351.9±443.0	0.579	0.563
Urine volume (mL)	450(150)	412.5(187.5)	500(150)	0.329	0.849
Bleeding volume (mL)	60(50)	55(70)	80(50)	3.443	0.179
Sufentanil usage (μg)	45(10)	40(10)	40(15)	2.909	0.233
Remifentanil usage (μg)	1565(436)	1497.5(642)	1363(294)	2.628	0.269
Propofol usage (mg)	1034.9±153.1	914.4±212.1*	863.5±171.1*	6.568	0.002
Cis atracurium usage (mg)	19.21±2.71	18.54±3.61	19.07±4.10	0.290	0.749
Emergence time (min)	38(11)	34.5(7.5)*	23(15)*#	24.436	0.000
Extubation time (min)	49.6±9.2	41.4±6.5*	35.7±10.8*#	16.879	0.000
PACU stay duration (min)	80.5(15.5)	75(10)	67(18) *	15.455	0.000
Cardiovascular events	3(10.7%)	2(7.1%)	2(7.4%)	0.286	1.000
Hypertension	0(0)	0(0)	0(0)	/	/
Hypotension	2(7.1%)	1(3.57%)	1(3.7%)	0.498	1.000
Tachycardia	0(0)	0(0)	1(3.7%)	2.099	0.325
Bradycardia	1(3.57%)	1(3.57%)	0(0)	0.988	1.000

Notes: * $P<0.05$, compared to Group C; # $P<0.05$, compared to Group B1.

BIS Value

Compared with Group C, Group B1 and Group B2 had notably higher BIS values at the intraoperative T₂, T₃, T₄, and T₅ time points and higher average BIS values over the course of surgery ($P<0.05$). In addition, compared with Group B1,




Figure 2 (A) Propofol usage. (B) Emergence time. (C) Tracheal extubation time. (D) PACU stay duration.

Notes: * $P<0.05$, compared with Group C; # $P<0.05$, compared with Group B1.

Table 3 BIS Value at Different Time Points and Overall Average Value During Operation

	C (n=28)	B1 (n=28)	B2 (n=27)	H	P
T ₀	97(3)	95(5.75)	96(4)	2.033	0.362
T ₁	42.5(6)	42(4.75)	43(5)	5.315	0.070
T ₂	45(5.75)	47.5(8.5)*	52(7)*	22.298	0.000
T ₃	41.5(9.5)	48.5(7.75)*	54(5)*#	42.770	0.000
T ₄	42.5(6.5)	48(6.75)*	53(5)*	51.399	0.000
T ₅	45(6.75)	50(10.5)*	56(4)*#	36.380	0.000
Overall average	42(4.75)	47(5.5)*	53(4)*#	54.178	0.000

Notes: * $P<0.05$, compared to Group C; # $P<0.05$, compared to Group B1.

Group B2 had significantly higher BIS values at the intraoperative T₃ and T₅ time points ($P<0.05$). Table 3 and Figure 3 provide details.

MAP and HR


No significant difference in MAP was found among the three groups at any time point ($P>0.05$). Compared to T₀, the MAP of Group C decreased significantly at all time points from T₁ to T₅ ($P<0.05$). In Group B1, the MAP decreased from T₁ to T₄ time points ($P<0.05$), whereas the MAP in Group B2 decreased at T₁, T₂ and T₃ ($P<0.05$). Further details are provided in Table 4.

There was no significant difference in the HR among the three groups at any time point ($P>0.05$). Compared with T₀, the HR from T₁ to T₅ time points in Group C and Group B1 decreased significantly ($P<0.05$). In Group B2, the HR decreased considerably from T₁ to T₄ time points considerably decreased ($P<0.05$). For more details, please refer to Table 4.

Postoperative Recovery Information

As shown in Table 5, there were no significant differences among the three groups in the VAS scores and postoperative analgesic remedies (defined as VAS >5) ($P>0.05$).

No significant difference in QoR-15 score was found among the three groups in Preop ($P>0.05$). Compared with Preop, QoR-15 scores significantly decreased on POD₁, POD₂, POD₃, and POD₇ in Group C ($P<0.05$) while on POD₁, POD₂, and POD₃ in Group B1 and Group B2 ($P<0.05$). Compared with Group C, Group B1 had significantly higher

**Figure 3** BIS values at different time points.

Notes: * $P<0.05$, compared with Group C; # $P<0.05$, compared with Group B1.

Table 4 MAP and HR of T₀-T₅

	C (n=28)	B1 (n=28)	B2 (n=27)	F	P
MAP					
T ₀	93.14±8.58	90.25±8.99	91.19±7.91	0.842	0.435
T ₁	79.68±9.48 ^Δ	80.46±8.53 ^Δ	81.11±8.81 ^Δ	0.177	0.838
T ₂	80.29±9.12 ^Δ	82.36±8.84 ^Δ	83.07±9.84 ^Δ	0.675	0.512
T ₃	80.75±9.19 ^Δ	79.39±7.87 ^Δ	82.26±9.44 ^Δ	0.721	0.489
T ₄	80.79±9.09 ^Δ	80.25±9.38 ^Δ	84.33±9.80	1.515	0.226
T ₅	83.54±8.13 ^Δ	84.07±8.70	85.59±8.58	0.434	0.649
HR					
T ₀	78.75±7.06	80.29±6.20	79.11±9.72	0.298	0.743
T ₁	65.89±5.04 ^Δ	67.25±5.65 ^Δ	68.74±5.36 ^Δ	1.945	0.150
T ₂	70.32±5.45 ^Δ	72.82±5.95 ^Δ	73.30±6.27 ^Δ	2.039	0.137
T ₃	69.29±6.00 ^Δ	69.46±5.78 ^Δ	71.86±6.07 ^Δ	1.636	0.201
T ₄	71.07±7.90 ^Δ	71.29±6.84 ^Δ	70.85±5.78 ^Δ	0.027	0.973
T ₅	72.82±6.57 ^Δ	73.04±6.61 ^Δ	74.59±7.86	0.517	0.598

Note: ^ΔP<0.05, compared to T₀.

Table 5 VAS Score and Remedial Analgesia

	C (n=28)	B1 (n=28)	B2 (n=27)	χ ² /H	P
Preop	0(0)	0(0)	0(0)	0.003	0.988
POD ₁	3.5(1)	4(1)	4(1)	0.116	0.944
POD ₂	3(1)	3(1)	3(1)	0.463	0.793
POD ₃	2(1)	2(0.75)	2(0)	1.889	0.389
POD ₇	1.5(1)	1(1)	1(1)	2.613	0.271
Remedial analgesia	3(10.7%)	4(14.3%)	3(11.1%)	0.202	0.904

QoR-15 scores on POD₂ ($P<0.05$), and Group B2 had significantly higher QoR-15 scores on both POD₁ and POD₂ ($P<0.05$). Moreover, compared with Group B1, Group B2 had significantly higher QoR-15 scores on POD₁ and POD₂ ($P<0.05$). For more information, please refer to [Tables 6](#) and [Figure 4](#).

The RCSQ scores on Preop of the three groups were not significantly different ($P>0.05$). Compared with Preop, the RCSQ scores significantly decreased on POD₁, POD₂ and POD₃ in all three groups ($P<0.05$). Compared to Group C, Group B1 had remarkably higher RCSQ scores on POD₁ and POD₃ ($P<0.05$), and Group B2 had noticeably higher RCSQ scores on both POD₁, POD₂, and POD₃ ($P<0.05$). In addition, no difference in the RCSQ score was found on POD₁, POD₂, POD₃, and POD₇ between Group B1 and Group B2 ($P>0.05$). Further details are provided in [Tables 7](#) and [Figure 5](#), respectively.

Table 6 Perioperative Qor-15 Scores

	C (n=28)	B1 (n=28)	B2 (n=27)	F	P
PREOP	134.43±7.26	130.89±8.79	132.78±5.32	1.652	0.198
POD ₁	104.54±9.85 ^Δ	107.82±6.79 ^Δ	114.15±8.57* ^{#Δ}	9.065	0.000
POD ₂	107.50±8.65 ^Δ	113.82±6.94* ^Δ	120.37±8.54* ^{#Δ}	17.463	0.000
POD ₃	118.61±8.76 ^Δ	121.04±8.26 ^Δ	122.85±8.26 ^Δ	1.757	0.179
POD ₇	124.43±7.02 ^Δ	126.57±9.00	128.44±5.42	2.076	0.132

Notes: * $P<0.05$, compared with Group C; # $P<0.05$, compared with Group B1; ^Δ $P<0.05$, compared with Preop.




Figure 4 Perioperative Qor-15 scores.

Notes: * $P<0.05$, compared with Group C; # $P<0.05$, compared with Group B1; ^ $P<0.05$, compared with Preop.

The incidence of POD and POCD at any time point, their total incidence (POD or POCD: C, 17.9%; B1, 14.3%; B2, 11.1%), and MMSE scores exhibited no significant differences across the three groups ($P>0.05$), with no notable trends in incidence rates. Similarly, postoperative complication rates were lower in Groups B1 (3.6%) and B2 (7.4%) compared to Group C (10.7%), though not significantly ($P>0.05$). No respiratory complications occurred in any group, and the incidence of circulatory system complications also showed no significant differences across groups ($P>0.05$), with postoperative hospital stay remaining similar ($P>0.05$). However, the time to first ambulation in Group B2 was significantly shorter than in Group C ($P<0.05$), suggesting enhanced early mobility, whereas Group B1 showed no significant difference from Group C ($P>0.05$). Additional details are presented in [Tables 8 and 9](#).

Discussion

This study indicated that the depth of anesthesia monitoring under BIS could promote early recovery after laparoscopic surgery in elderly patients, which has significant implications for the selection of intraoperative anesthetic monitoring as well as the improvement of anesthetic management among elderly patients.

In clinical practice, anesthesiologists generally judge the depth of anesthesia through a series of vital sign changes such as blood pressure, heart rate, pupil size, muscle tone and nerve reflexes.¹⁹ However, such vital signs sometimes do not accurately reflect the depth of anesthesia level because of the differences in stimulation intensity, surgical types and individuals, also, there will use adequate analgesics, muscle relaxants, vasoactive drugs, anticholinergics and other drugs during anesthesia, thus it becomes difficult and unreliable to identify and judge the depth of anesthesia through simple clinical observation. Too light anesthesia can lead to a significant stress response, circulatory system hyperexcitability, supply and demand imbalance of oxygen in various vital organs, endocrine and metabolic disorders, and even intraoperative awareness and more serious adverse consequences, such as post-traumatic stress disorder, causing great

Table 7 Perioperative RCSQ Scores

	C (n=28)	B1 (n=28)	B2 (n=27)	F	P
PREOP	75.46±5.19	74.82±5.11	77.00±3.99	1.488	0.232
POD ₁	60.43±4.32 ^Δ	63.32±5.46* ^Δ	65.00±6.30* ^Δ	5.046	0.009
POD ₂	65.07±5.93 ^Δ	67.25±5.57 ^Δ	69.15±6.18* ^Δ	3.297	0.042
POD ₃	66.86±4.78 ^Δ	70.39±6.15* ^Δ	71.89±5.26* ^Δ	6.264	0.003
POD ₇	74.75±5.63	75.86±5.51	74.63±6.03	0.388	0.680

Notes: * $P<0.05$, compared with Group C; ^Δ $P<0.05$, compared with Preop.




Figure 5 Perioperative RCSQ scores.

Notes: * $P < 0.05$, compared with Group C; # $P < 0.05$, compared with Group B1; $\Delta P < 0.05$, compared with Preop.

physical and mental damage to patients. On the other hand, anesthesia that is too deep can trigger adverse events including insufficient stress response, severe respiratory and circulatory function inhibition, delayed awakening, and postoperative neurocognitive dysfunction, and in severe cases, irreversible damage or death.²⁰ Therefore, the evaluation of depth of anesthesia is of great significance for the safety and postoperative recovery of patients, especially for elderly patients with vulnerable physiological status, and accurate monitoring of the depth of anesthesia has special value.²¹

Table 8 POD Incidence, POCD Incidence, and MMSE Scores

	C (n=28)	B1 (n=28)	B2 (n=27)	χ^2/F	P
Incidence of POD					
POD ₁	3(10.7%)	3(10.7%)	2(7.4%)	0.229	0.892
POD ₂	2(7.1%)	2(7.1%)	1(3.7%)	0.381	0.827
POD ₃	1(3.6%)	0(0)	0(0)	1.988	0.370
Total cases	5(17.9%)	4(14.3%)	3(11.1%)	0.507	0.776
Incidence of POCD					
POD ₃	4(14.3%)	4(14.3%)	3(11.1%)	0.160	0.923
POD ₇	3(10.7%)	2(7.1%)	1(3.7%)	1.008	0.604
Total cases	5(17.9%)	4(14.3%)	3(11.1%)	0.507	0.776
MMSE scores					
PREOP	24.68±2.16	24.71±1.96	24.93±2.15	0.112	0.894
POD ₃	23.14±2.49	23.32±2.83	23.41±2.14	0.080	0.923
POD ₇	24.04±2.19	24.93±2.61	24.33±2.02	1.104	0.336

Table 9 Complications, First Postoperative Ambulation, and Postoperative Hospital Stay

	C (n=28)	B1 (n=28)	B2 (n=27)	$\chi^2/F/H$	P
Postoperative complications	3(10.7%)	1(3.6%)	2(7.4%)	1.067	0.690
Nausea and vomiting	2(7.1%)	1(3.6%)	1(3.7%)	0.498	1.000
Respiratory complications	0(0)	0(0)	0(0)	/	/
Circulatory complications	1(3.6%)	0(0)	1(3.7%)	1.044	0.770
First postoperative ambulation (h)	24(14.5)	20(6)	18(10)*	8.380	0.015
Postoperative hospital stay (d)	7.57±1.83	7.25±1.90	7.96±1.99	0.962	0.386

Note: * $P < 0.05$, compared with Group C.

According to previous studies on BIS monitoring among elderly patients, the first concern related to the depth of anesthesia was its influence on the quality of anesthesia recovery. Anesthesia recovery is most strongly linked to anesthetic methods, selection of anesthetic drugs, and management. Anesthetic management is particularly crucial under the established anesthetic and drug regimens. Lewis et al²² found that using BIS monitoring to guide the depth of anesthesia could reduce intraoperative awareness and dosage of intraoperative sedative drugs, shorten anesthesia emergence and extubation times, and facilitate early postoperative recovery. Yu et al²³ applied intraoperative BIS monitoring to maintain the depth of anesthesia at a specific level in elderly patients, and reported that this method could better inhibit adverse stress responses, reduce anesthesia-related hypotension, and maintain hemodynamic stability. Monitoring the depth of anesthesia to improve the management of intraoperative hypotension in elderly patients might be a convenient intervention with the potential to reduce the incidence of intraoperative anesthesia-related hypotension in this vulnerable group.²⁴ Similar to previous findings, our results showed that compared with Group C, dosage of intravenous anesthetic propofol, emergence time and extubation time were all less in Groups B1 and B2, suggesting that depth of anesthesia under BIS monitoring could reduce intraoperative anesthetic drug consumption and improve the quality of recovery. In addition, no significant difference was observed in the incidence of intraoperative hemodynamic adverse events among the three groups, whereas the time required for emergence and extubation was shorter and PACU stay was shorter in Group B2 than in Group B1, indicating that slightly higher BIS values were more favorable for anesthesia recovery in elderly patients undergoing laparoscopic surgery under the premise of anesthesia safety.

The recovery of postoperative cognitive function is important for evaluating the quality of recovery in elderly patients, especially POD and POCD, which have a high incidence in elderly surgical patients. Age is the most significant factor of postoperative cognitive function recovery, while surgery and general anesthesia are generally considered as common risk factors for POCD, but the specific mechanism remains unknown.^{25,26} Clinical studies have suggested that the depth of anesthesia has a certain correlation with the occurrence and recovery of POCD,²⁷ while BIS monitoring of depth of anesthesia can promote early postoperative cognitive function recovery and reduce the occurrence of acute POD.²⁸ However, Wildes et al²⁹ disagreed with the above results through a randomized controlled study of 1232 elderly patients aged 60 years and above undergoing major surgery, from which they found no superior effect of electroencephalography-guided anesthesia on POD reduction compared with conventional anesthesia. Some investigators therefore believed that current evidence could not support its routine use to prevent POD in elderly patients or any other patients.³⁰ A meta-analysis also reported no significant correlation between depth of anesthesia and POCD.³¹ Studies on the effect of different depth of anesthesia on cognitive function also remain controversial, including the view of targeted light anesthesia could reduce the incidence of POD and the risk of POCD,³² and avoiding too deep anesthesia through intraoperative monitoring might help improve postoperative cognitive function to benefit patients at high risk of POCD.³³ Quan et al³⁴ performed deep anesthesia with BIS values at low levels (30~45) and decreased the occurrence of early POCD after abdominal surgery in elderly patients. The diverse conclusions might stem from the heterogeneity of studies, but the development of deep anesthesia could independently predict POD and should be vigilant for explosive suppression during surgery.^{35,36} Current evidence suggests that early risk identification, as well as careful adjustment of the depth of anesthesia and treatment of pain guided by processed electroencephalographic monitoring, were the most effective strategies for reducing the risk of POD.³⁷

In our study, there were no significant differences in POD, POCD, and MMSE scores among the three groups, which may be because unlike the more invasive surgical types, minimally invasive laparoscopic surgery had no significant effect on postoperative cognitive function in elderly patients. Second, the relatively small sample size limited the statistical power and may not have been sufficient to detect subtle changes. Third, patients included in this study generally had a lower education level and could not complete complex scales such as the Montreal Cognitive Assessment (MoCA); therefore, the MMSE was finally selected. MMSE is the most commonly used cognitive function assessment scale in previous studies, with the disadvantages of being relatively simple, less sensitive, and having learning and capping effects when repeatedly evaluated. Future studies are needed to synthesize more scales and diagnostic tools for systematic cognitive function assessment.

Sleep quality has an indispensable relationship with cognitive function, and sleep problems and disorders may lead to impaired cognition.³⁸ Epidemiological studies have shown that about 50% of the elderly have sleep problems, but it is

often not fully appreciated in elderly people in clinical practice.³⁹ The general anesthesia can change postoperative sleep architecture, especially in elderly patients with a high incidence of postoperative sleep disorders; general anesthesia can cause deleterious effects such as increased risk of delirium and cardiovascular events, and poor recovery.¹⁶ It may take a week or more to recover after surgery due to decreased subjective sleep quality and efficiency and increased daytime fatigue.⁴⁰ Song et al⁴¹ proved that compared with subarachnoid anesthesia, general anesthesia is associated with greater impairment of melatonin rhythm and sleep patterns, as well as a higher incidence of POD in elderly patients. This might be due to the effect of propofol on sleep rhythm, and the use of propofol at the recommended sedation level in critically among patients with assisted ventilation mode would inhibit rapid eye movement sleep and further deteriorate their sleep quality, but very few reports focused on the effect of propofol on the postoperative sleep quality of elderly patients during routine surgical anesthesia application.⁴² As one of the instruments used to evaluate sleep quality, the RCSQ scale contains five items, with higher scores representing better sleep quality.⁴³ Compared with the more commonly used Pittsburgh Sleep Quality Index, the RCSQ is more concise and easier to understand and capture, and its items are relevant to sleep quality and have shown good reliability and validity, providing a comprehensive sleep assessment over a short length of hospital stay.⁴⁴ In this study, RCSQ scores in all three groups returned to the preoperative baseline on POD₇. Group B1 had higher RCSQ scores on POD₁ and POD₃ than Group C, whereas Group B2 had higher RCSQ scores from POD₁ to POD₃ than Group C. It could be concluded that the postoperative sleep quality was better than that of Group C to some extent when monitoring the depth of anesthesia through BIS. Maintaining an intraoperative BIS value of 50–60 could improve postoperative sleep quality more effectively, which is beneficial for postoperative recovery.

Anesthesia recovery and cognitive function recovery are part of the postoperative recovery quality, whereas in the evaluation of perioperative recovery quality, the subjective feelings of patients are often easily ignored. The effectiveness and safety of current anesthesia also place greater emphasis on patient-centered outcome measures.⁴⁵ QoR-15 is a widely used and validated patient-reported outcome measure for quality of recovery after surgery developed by Stark et al from QoR-40,¹² which has equivalent psychometric properties but is more feasible to use and meets the requirements of validity, reliability, interpretability, acceptability, and feasibility.^{18,46} Items of QoR-15 include five dimensions, which are emotional state, physical comfort, psychological support, physical independence, and pain, and the anesthesiologist is able to assess the patient's recovery from multiple dimensions reported by the patient after surgery and further evaluate the quality of anesthesia.⁴⁷ On the basis of Consensus-based Standards for the selection of health Measurement Instruments (COSMIN), a recent systematic review suggested the use of QoR-15 as a standard outcome measure for quality of recovery in surgical and anesthetic clinical trials since the primary aim of surgical and anesthetic interventions was to improve patient outcomes.⁴⁸ Whereas in previous studies on the influence of depth of anesthesia on postoperative recovery, the results were generally physician-centered evaluations that focused only on some morbidity parameters, and the effect remained unknown if taking the patient's physical and psychological status into account. Our results indicate that monitoring and regulation of the depth of anesthesia could impact patients' perception and evaluation of recovery quality. Compared to Group C, Group B1 had significantly higher QoR-15 scores on POD₂, and Group B2 had significantly higher QoR-15 scores on both POD₁ and POD₂. In addition, compared with Group B1, Group B2 had significantly higher QoR-15 scores on POD₁ and POD₂. It could be inferred that lighter anesthesia resulted in a better evaluation of early postoperative recovery quality in elderly patients undergoing laparoscopic surgery.

Pain is an essential factor that affects the postoperative status. Significant postoperative pain can reduce patients' postoperative comfort and satisfaction, and also lead to complications such as sympathetic hyperexcitability, increased stress response, cognitive impairment, sleep disorders, postoperative activity limitation, increased risk of thromboembolism, and prolonged hospital stay, all of which slow patients' postoperative recovery.¹⁷ In our study, the same postoperative analgesia method was used in the three groups, and no significant differences were found in VAS score and analgesic remedy at each time point. However, Group B2 had a significantly shorter time to first postoperative ambulation than Group C, indicating that the QoR-15 score could better reflect the overall level of recovery of the patients. The mean QoR-15 scores did not return to the preoperative baseline on POD₇ in all of the three groups, which might be because after minimally invasive laparoscopic surgery, the body could not adapt and effectively compensate for the negative effects within a short period of time, such as pain, disruption of dietary patterns, energy metabolism and endocrine dysfunction, sleep disorders, and reduced physical activity. Some scholars believed that this might be a long-

term and slow repair process,³⁵ at the same time, whether baseline data could truly reflect the preoperative status of patients also required further verification.⁴⁸

Through this trial, we also found that compared with Groups B1 and B2, more anesthetic drugs were used in Group C, and the BIS values at each time point and the mean BIS values during surgery in Group C were lower, indicating that the depth of anesthesia guided by experience might generally be deeper and that deep anesthesia lasts for a longer period than expected in elderly patients, which could be a potential risk for elderly patients. Therefore, it is necessary to monitor and adjust the depth of anesthesia within a certain range during the perioperative period to avoid long-term deep anesthesia in elderly patients with poor health. Through this trial, we also found that compared with Groups B1 and B2, more anesthetic drugs were used in Group C, and the BIS values at each time point and the mean BIS values during surgery in Group C were lower, indicating that the depth of anesthesia guided by experience might generally be deeper and that deep anesthesia lasts for a longer period than expected in elderly patients, which could be a potential risk for elderly patients. Therefore, it is necessary to monitor and adjust the depth of anesthesia within a certain range during the perioperative period to avoid long-term deep anesthesia in elderly patients with poor health. Our results demonstrate that BIS-guided anesthesia significantly enhanced immediate postoperative recovery, notably earlier ambulation in Group B2 compared to Group C [18(10) vs 24(14.5) hours; $P=0.015$], but did not significantly reduce other complications (eg, total incidence: B1 3.6%, B2 7.4%, C 10.7%; $P=0.690$; nausea/vomiting lower in B1 and B2). This is consistent with Chiang et al¹¹ and Gruenewald et al¹⁹ who found BIS-guided techniques primarily improve short-term outcomes like mobility without broadly affecting complication rates unless optimized for high-risk patients. This suggests that anesthesia depth's primary benefit lies in immediate recovery, supporting its targeted application in clinical practice.

In light of these results, an intriguing question arises regarding the potential role of total intravenous anesthesia (TIVA) in this context. TIVA, which allows for precise titration of anesthetic agents such as propofol, may offer an alternative or complementary approach to controlling anesthesia depth more effectively than inhaled agents alone. Studies such as those by Li et al⁴⁹ suggest that TIVA, when paired with depth-of-anesthesia monitoring, could provide finer control over sedation levels, potentially enhancing outcomes like mobility while minimizing oversedation-related risks. This precision is particularly advantageous in procedures requiring tight control, such as in elderly or hemodynamically unstable patients.⁵⁰ Whether TIVA could outperform BIS-guided volatile anesthesia in our study population—particularly in terms of refining immediate postoperative recovery—remains an open question worthy of further investigation. To fully elucidate these possibilities, future research should compare BIS-guided volatile anesthesia with TIVA in similar cohorts, focusing not only on mobility but also on other short-term recovery parameters, such as cognitive function and patient-reported outcomes. Additionally, while our findings suggest that anesthesia depth management has a limited scope in reducing overall complications, its role in specific patient subgroups (eg, the elderly or those with comorbidities) warrants further exploration. In conclusion, our study highlights the nuanced impact of anesthesia depth on postoperative recovery and opens the door to considering alternative strategies, such as TIVA, to optimize patient outcomes in the immediate postoperative phase.

This study has some limitations. First, all included participants were elderly patients with lower educational levels who underwent elective laparoscopic surgery, thus, our results may not be generalizable to more educated, healthier, and younger patients and other surgical modalities. Second, we only performed follow-up within 7 days after surgery because of limited resources, while recovery after anesthesia and operation might take weeks or even years, and the impairment and recovery of cognitive function are also slow processes. In addition, these assessments are not realistic for emergency surgery because of the need to obtain baseline data. Finally, the ability of TIVA to maintain a stable pharmacokinetic profile might theoretically improve depth control, but its practical superiority would depend on factors such as patient-specific responses, surgical type, and resource availability, which were not directly assessed in our study. Nevertheless, our results still indicate that using the BIS to monitor the depth of anesthesia and maintain a higher BIS value within a safe range in laparoscopic surgery in elderly patients could help improve the quality of anesthesia recovery and promote sleep quality recovery and early autonomous activity in a short period of time after surgery, which might be one of the important means of enhancing the quality of early postoperative recovery in elderly patients. These results remain of great value for the evaluation and prediction of long-term health status of elderly patients after surgery.

Conclusion

In summary, this study demonstrated the guiding significance of BIS in monitoring the depth of anesthesia after laparoscopic surgery among elderly patients in terms of reducing the amount of general anesthesia, shortening emergence time and extubation time, and decreasing the length of PACU stay. Furthermore, it improved the postoperative sleep quality to a certain extent, thereby promoting early postoperative recovery from the patient's perspective. Experience-guided anesthesia in elderly patients has the potential to result in deep anesthesia or even prolonged deep anesthesia, whereas lighter anesthesia may be more beneficial in elderly patients undergoing laparoscopic surgery. These findings have important implications for the selection of intraoperative monitoring tools and optimization of anesthetic management.

Abbreviations

QoR, quality of recovery; VAS, visual analog scale; CAM, confusion assessment method; MMSE, Mini-Mental State Examination; RCSQ, Richards-Campbell sleep questionnaire; BIS, bispectral index; POD, postoperative delirium; POCD, postoperative cognitive dysfunction; PACU, post-anesthesia care unit; MAP, mean arterial pressure; HR, heart rate.

Acknowledgments

This work was supported by the Project of Improving the Basic Scientific Research Ability of Young and Middle-aged Teachers in Colleges and Universities in Guangxi (No. 2020KY12023) and the Guangxi Medical and Health Key Discipline Cultivation Project. Special thanks to Huailong Yang and Xinyi Liu, who participated in this study and made contributions to the final completion of this manuscript.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

There are no conflicts of interest reported by the authors. An unauthorized version of the Chinese MMSE was used by the study team without permission, however this has now been rectified with PAR. The MMSE is a copyrighted instrument and may not be used or reproduced in whole or in part, in any form or language, or by any means without written permission of PAR (www.parinc.com).

References

1. Liu Y, Xiao W, Z ML, et al. Geriatric anesthesia-related morbidity and mortality in China: Current status and trend. *Chin Med J*. 2017;130(22):2738–2749. doi:10.4103/0366-6999.218006
2. Chan SP, Ip KY, Irwin MG. Peri-operative optimisation of elderly and frail patients: a narrative review. *Anaesthesia*. 2019;74 Suppl 1:80–89. doi:10.1111/anae.14512
3. Pandin P, Estruc I, Van Hecke D, et al. Brain aging and anesthesia. *J Cardiothorac Vasc Anesth*. 2019;33 Suppl 1:S58–S66. doi:10.1053/j.jvca.2019.03.042
4. Ludbrook GL, Goldsman AG. Coordinated perioperative care—a high value proposition? *Br J Anaesth*. 2017;118(1):3–5. doi:10.1093/bja/aew331
5. Akhtar S. Pharmacological considerations in the elderly. *Curr Opin Anaesthesiol*. 2018;31(1):11–18. doi:10.1097/ACO.0000000000000544
6. M SM, Beese U, H HE, et al. Bispectral index values and propofol concentrations at loss and return of consciousness in patients with frontal brain tumours and control patients. *Br J Anaesth*. 2014;112(1):110–117. doi:10.1093/bja/aet342
7. J AA, Kamata K, Jäntti V, et al. Comparison of bispectral index and entropy values with electroencephalogram during surgical anaesthesia with sevoflurane. *Br J Anaesth*. 2015;115(2):258–266. doi:10.1093/bja/aev206
8. Sessler DI, Sigl JC, Kelley SD, et al. Hospital stay and mortality are increased in patients having a “triple low” of low blood pressure, low bispectral index, and low minimum alveolar concentration of volatile anesthesia. *Anesthesiology*. 2012;116(6):1195–1203. doi:10.1097/ALN.0b013e31825683dc
9. Griffiths R, Beech F, Brown A, et al. Peri-operative care of the elderly 2014: association of anaesthetists of Great Britain and Ireland. *Anaesthesia*. 2014;69 Suppl 1:81–98. doi:10.1111/anae.12524

10. Rasmussen LS, Jørgensen CC, Kehlet H. Enhanced recovery programmes for the elderly. *Eur J Anaesthesiol.* 2016;33(6):391–392. doi:10.1097/EJA.0000000000000452
11. Chiang MH, Wu SC, Hsu SW, et al. Bispectral index and non-bispectral index anesthetic protocols on postoperative recovery outcomes. *Minerva Anesthesiol.* 2018;84(2):216–228. doi:10.23736/S0375-9393.17.12033-X
12. Stark PA, Myles PS, Burke JA. Development and psychometric evaluation of a postoperative quality of recovery score: the QoR-15. *Anesthesiology.* 2013;118(6):1332–1340. doi:10.1097/ALN.0b013e318289b84b
13. Jin Z, Hu J, Ma D. Postoperative delirium: perioperative assessment, risk reduction, and management. *Br J Anaesth.* 2020;125(4):492–504. doi:10.1016/j.bja.2020.06.063
14. Duning T, Ilting-Reuke K, Beckhuis M, et al. Postoperative delirium - treatment and prevention. *Curr Opin Anaesthesiol.* 2021;34(1):27–32. doi:10.1097/ACO.0000000000000939
15. Cottrell JE, Hartung J. Anesthesia and cognitive outcome in elderly patients: a narrative viewpoint. *J Neurosurg Anesthesiol.* 2020;32(1):9–17. doi:10.1097/ANA.0000000000000640
16. Luo M, Song B, Zhu J. Sleep disturbances after general anesthesia: current perspectives. *Front Neurol.* 2020;11:629. doi:10.3389/fneur.2020.00629
17. Liu Q, Lin JY, Zhang YF, et al. Effects of epidural combined with general anesthesia versus general anesthesia on quality of recovery of elderly patients undergoing laparoscopic radical resection of colorectal cancer: a prospective randomized trial. *J Clin Anesth.* 2020;62:109742. doi:10.1016/j.jclinane.2020.109742
18. Myles PS, Myles DB, Gallagher W, et al. Minimal clinically important difference for three quality of recovery scales. *Anesthesiology.* 2016;125(1):39–45. doi:10.1097/ALN.0000000000001158
19. Gruenewald M, Harju J, Preckel B, et al. Comparison of adequacy of anaesthesia monitoring with standard clinical practice monitoring during routine general anaesthesia: an international, multicentre, single-blinded randomised controlled trial. *Eur J Anaesthesiol.* 2021;38(1):73–81. doi:10.1097/EJA.0000000000001357
20. Wang J, Zhang L, Huang Q, et al. Monitoring the end-tidal concentration of sevoflurane for preventing awareness during anesthesia (MEETS-PANDA): a prospective clinical trial. *Int J Surg.* 2017;41:44–49. doi:10.1016/j.ijssu.2017.03.015
21. Eagleman SL, Drover CM, Li X, MacIver MB, Drover DR. Offline comparison of processed electroencephalogram monitors for anaesthetic-induced electroencephalogram changes in older adults. *Br J Anaesth.* 2021;126(5):975–984. doi:10.1016/j.bja.2020.12.042
22. Lewis SR, Pritchard MW, Fawcett LJ, et al. Bispectral index for improving intraoperative awareness and early postoperative recovery in adults. *Cochrane Database Syst Rev.* 2019;9(9):CD003843. doi:10.1002/14651858.CD003843.pub4
23. Yu H, Zhang L, Ma Y, et al. Early postoperative recovery in operating room after desflurane anesthesia combined with Bispectral index (BIS) monitoring and warming in lengthy abdominal surgery: a randomized controlled study. *BMC Anesthesiol.* 2018;18(1):110. doi:10.1186/s12871-018-0577-6
24. Wickham A, Highton D, Martin D, et al. Care of elderly patients: a prospective audit of the prevalence of hypotension and the use of BIS intraoperatively in 25 hospitals in London. *Perioper Med.* 2016;5:12. doi:10.1186/s13741-016-0036-1
25. Zhang Y, Bao HG, Lv YL, et al. Risk factors for early postoperative cognitive dysfunction after colorectal surgery. *BMC Anesthesiol.* 2019;19(1):6. doi:10.1186/s12871-018-0676-4
26. Rengel KF, Pandharipande PP, Hughes CG. Special considerations for the aging brain and perioperative neurocognitive dysfunction. *Anesthesiol Clin.* 2019;37(3):521–536. doi:10.1016/j.anclin.2019.04.010
27. Sumner M, Deng C, Evered L, et al. Processed electroencephalography-guided general anaesthesia to reduce postoperative delirium: a systematic review and meta-analysis. *Br J Anaesth.* 2023;130(2):e243–e253. doi:10.1016/j.bja.2022.01.006
28. Bao L, Liu T, Zhang Z, et al. The prediction of postoperative delirium with the preoperative bispectral index in older aged patients: a cohort study. *Aging Clin Exp Res.* 2023;35(7):1531–1539. doi:10.1007/s40520-023-02408-9
29. Wildes TS, Mickle AM, Ben Abdallah A, et al. Effect of electroencephalography-guided anesthetic administration on postoperative delirium among older adults undergoing major surgery: the ENGAGES randomized clinical trial. *JAMA.* 2019;321(5):473–483. doi:10.1001/jama.2018.22005
30. Evered LA, Goldstein PA. Reducing perioperative neurocognitive disorders (PND) through depth of anesthesia monitoring: a critical review. *Int J Gen Med.* 2021;14:153–162. doi:10.2147/IJGM.S242230
31. Lu X, Jin X, Yang S, et al. The correlation of the depth of anesthesia and postoperative cognitive impairment: a meta-analysis based on randomized controlled trials. *J Clin Anesth.* 2018;45:55–59. doi:10.1016/j.jclinane.2017.12.002
32. Li Y, Zhang B. Effects of anesthesia depth on postoperative cognitive function and inflammation: a systematic review and meta-analysis. *Minerva Anesthesiol.* 2020;86(9):965–973. doi:10.23736/S0375-9393.20.14251-2
33. Cao YH, Chi P, Zhao YX, et al. Effect of bispectral index-guided anesthesia on consumption of anesthetics and early postoperative cognitive dysfunction after liver transplantation: an observational study. *Medicine.* 2017;96(35):e7966. doi:10.1097/MD.00000000000007966
34. Quan C, Chen J, Luo Y, et al. BIS-guided deep anesthesia decreases short-term postoperative cognitive dysfunction and peripheral inflammation in elderly patients undergoing abdominal surgery. *Brain Behav.* 2019;9(4):e01238. doi:10.1002/brb3.1238
35. M RF, Franck M, Lendner J, et al. Monitoring depth of anaesthesia in a randomized trial decreases the rate of postoperative delirium but not postoperative cognitive dysfunction. *Br J Anaesth.* 2013;110 Suppl 1:i98–105. doi:10.1093/bja/aet055
36. Hesse S, Kreuzer M, Hight D, et al. Association of electroencephalogram trajectories during emergence from anaesthesia with delirium in the postanaesthesia care unit: an early sign of postoperative complications. *Br J Anaesth.* 2019;122(5):622–634. doi:10.1016/j.bja.2018.09.016
37. Swarbrick CJ, Partridge JSL. Evidence-based strategies to reduce the incidence of postoperative delirium: a narrative review. *Anaesthesia.* 2022;77 Suppl 1:92–101. doi:10.1111/anae.15607
38. Dzierzewski JM, Dautovich N, Ravvits S. Sleep and cognition in older adults. *Sleep Med Clin.* 2018;13(1):93–106. doi:10.1016/j.jsmc.2017.09.009
39. Rodriguez JC, Dzierzewski JM, Alessi CA. Sleep problems in the elderly. *Med Clin North Am.* 2015;99(2):431–439. doi:10.1016/j.mcna.2014.11.013
40. van Zuylen ML, Meewisse AJG, Ten Hoop W, et al. Effects of surgery and general anaesthesia on sleep-wake timing: CLOCKS observational study. *Anaesthesia.* 2022;77(1):73–81. doi:10.1111/anae.15564
41. Song Y, Liu Y, Yuan Y, et al. Effects of general versus subarachnoid anaesthesia on circadian melatonin rhythm and postoperative delirium in elderly patients undergoing hip fracture surgery: a prospective cohort clinical trial. *EbioMedicine.* 2021;70:103490. doi:10.1016/j.ebiom.2021.103490

42. Kondili E, Alexopoulou C, Xirouchaki N, et al. Effects of propofol on sleep quality in mechanically ventilated critically ill patients: a physiological study. *Intensive Care Med.* **2012**;38(10):1640–1646. doi:10.1007/s00134-012-2623-z
43. Tonna JE, Dalton A, Presson AP, et al. The effect of a quality improvement intervention on sleep and delirium in critically ill patients in a surgical ICU. *Chest.* **2021**;160(3):899–908. doi:10.1016/j.chest.2021.03.030
44. Hoey LM, Fulbrook P, Douglas JA. Sleep assessment of hospitalised patients: a literature review. *Int J Nurs Stud.* **2014**;51(9):1281–1288. doi:10.1016/j.ijnurstu.2014.02.001
45. Léger M, Campfort M, Cayla C, et al. Postoperative quality of recovery measurements as endpoints in comparative anaesthesia studies: a systematic review. *Br J Anaesth.* **2021**;126(6):e210–e212. doi:10.1016/j.bja.2021.03.008
46. Bu XS, Zhang J, Zuo YX. Validation of the Chinese version of the quality of recovery-15 score and its comparison with the post-operative quality recovery scale. *Patient.* **2016**;9(3):251–259. doi:10.1007/s40271-015-0148-6
47. Myles PS, Boney O, Botti M, et al. Systematic review and consensus definitions for the standardised endpoints in perioperative medicine (StEP) initiative: patient comfort. *Br J Anaesth.* **2018**;120(4):705–711. doi:10.1016/j.bja.2017.12.037
48. Kleif J, Waage J, Christensen KB, et al. Systematic review of the QoR-15 score, a patient-reported outcome measure measuring quality of recovery after surgery and anaesthesia. *Br J Anaesth.* **2018**;120(1):28–36. doi:10.1016/j.bja.2017.11.013
49. Li Z, Cai J, Li J, Xu X, Zheng L. Comparative evaluation of the bispectral index (BIS) and BISpro during propofol anaesthesia. *J Int Med Res.* **2021**;49(4):3000605211001705. doi:10.1177/03000605211001705
50. Biggs D, Boncompagni G, Pedemonte JC, Fuentes C, Cortinez LI. The effect of age on electroencephalogram measures of anesthesia hypnosis: a comparison of BIS, Alpha Power, Lempel-Ziv complexity and permutation entropy during propofol induction. *Front Aging Neurosci.* **2022**;14:910886. doi:10.3389/fnagi.2022.910886

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress
Taylor & Francis Group