

POSTER PRESENTATION

Open Access

P075: Emergence of carbapenem-resistant enterobacteriaceae in surgical and intensive care units of a hospital with low usage of carbapenem in Kano, North West Nigeria

I Yusuf^{1*}, AH Arzai², MI Getso³, A Sherif⁴, M Haruna⁵

From 2nd International Conference on Prevention and Infection Control (ICPIC 2013) Geneva, Switzerland. 25-28 June 2013

Introduction

Carbapenem Resistant Enterobacteriaceae (CRE) have emerged in surgical (SW) and intensive care units (ICUs) of tertiary health care centers in North West Nigeria despite their low or no usage in the hospitals.

Objectives

1. To determine the antibacterial susceptibility of pathogens from patients with severe bacterial infections that require rapid and aggressive antimicrobial treatments to imipenem (IMP) and meropenem (MEM) 2. Screen the pathogens for carbapenemase and metallo β lactamase production 3. Tests their susceptibility to colistin and tigecycline. 4. To evaluate the views of by different health care practitioners on causes, effects and control of CRE.

Methods

Isolates from patients admitted for atleast 7 days were screened for susceptibility to IMP and MEM using the CLSI 2012 break points. Carbapenemase and Metallo-β-lactamase (MBL) production were detected phenotypically by Modified Hodge Test (MHT) and Combine Disc Test (CDT) respectively.

Results

Resistance to IMP and MEM was 10.5% and 12.5% respectively. Carbapenemase production was 34.5% (highest so far in Kano). About 14.4% of the carbapenemase producers produce MBL. All the isolates were susceptible to colistin and tigecycline, except 2 *E. coli* and

1 *K. pneumoniae* which are resistant and also none carbapenemase producers. Of 486 medical practitioners studied in the region, only 1.2% have previous knowledge that MBL causes resistance. About 24.5%, 5.9% and 3.9% have knowledge of ESBL, AmpC and carbapenemase as a cause of rising resistance in the state. In addition, only 38.2% doctors have ever prescribe carbapenems in their professional practices and 12.8% of the doctors only prescribe in a ratio of 1:20 due to its high cost.

Conclusion

Carbapenem resistance is in increase in hospitals with a low or no usage of carbapenem and the resistances are mediated by carbapenemase especially MBL. Awareness of this problem is low among the medical practitioners that have direct contact with the patients and can contribute to its wide spread in the community.

Disclosure of interest

None declared.

Author details

¹Microbiology, Bayero University Kano, Nigeria. ²Microbiology, Bayero University, Kano, Nigeria. ³Department of Medical Microbiology, Faculty of Medicine, Nigeria. ⁴Department of Medical Microbiology, Bayero University Kano, Kano, Nigeria. ⁵Department of Biology, Kano state University of Science and Technology, Wudil, Nigeria.

Published: 20 June 2013

Reference

1. CLSI: Performance Standards for Antimicrobial Susceptibility Testing. CLSI, Wayne, PA; 2012, M100-S22.

¹Microbiology, Bayero University Kano, Nigeria
Full list of author information is available at the end of the article

doi:10.1186/2047-2994-2-S1-P75

Cite this article as: Yusuf *et al.*: P075: Emergence of carbapenem-resistant enterobacteriaceae in surgical and intensive care units of a hospital with low usage of carbapenem in Kano, North West Nigeria. *Antimicrobial Resistance and Infection Control* 2013 **2**(Suppl 1):P75.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

