

RESEARCH

Open Access

Toxoplasma infection in pregnant women: a current status in Songklanagarind hospital, southern Thailand

Hemah Andiappan¹, Veeranoot Nissapatorn^{1*}, Nongyao Sawangjaroen², Waenurama Chemoh², Yee Ling Lau¹, Thulasi Kumar¹, Subashini Onichandran¹, Chitkasaem Suwanrath³ and Verapol Chandeying^{3,4}

Abstract

Background: Toxoplasmosis, being one of the TORCH's infections in pregnant women, is caused by *Toxoplasma gondii*, an obligate intracellular protozoan parasite. This parasitic infection in pregnancy congenitally causes severe outcomes to their fetus and newborn. This study aimed to determine the seroprevalence and stages of *Toxoplasma* infection in pregnant women and its associated risks exposures.

Methods: The study was conducted within the pregnant women attending the antenatal clinic (ANC) at Songklanagarind hospital, Hat Yai, Songkhla province, Thailand. The sera of a total of 760 consecutive pregnant women were screened using standard commercial ELISA kits for detection of anti-*Toxoplasma* IgG and IgM antibodies. IgG avidity in the seropositive for both anti-*Toxoplasma* IgG and IgM antibodies were also assessed. The pregnant women's socio-demographic, obstetrics and risk factors associated with *Toxoplasma* seropositivity data were analyzed using univariate and multivariate analyses.

Results: From the total 760 pregnant women, 190 (25%, 95% CI = 22.05-28.20) were positive for anti-*Toxoplasma* antibodies. Of these, 167 (22.0%, 95% CI = 19.0-25.0) were positive for only anti-*Toxoplasma* IgG antibody and 23 (3.0%, 95% CI = 2.0-4.0) were positive for both anti-*Toxoplasma* IgG and IgM antibodies. All these samples were high avidity, indicated the infection occurred prior to four to five months. By applying statistical univariate analysis, age group, occupation and sources of drinking water showed a significant association with *Toxoplasma* seropositivity ($p < 0.05$). Multivariate logistic regression analysis further indicated that the significant factors associated with *Toxoplasma* seropositivity are age ≥ 26 (OR = 1.65, 95% CI = 1.11-2.44), working as laborer (OR = 1.57, 95% CI = 1.13-2.18) and drinking unclean (piped/tap/rain) water (OR = 1.75, 95% CI = 1.08-2.84).

Conclusion: The pregnant women in the active age group, working as laborers and exposure to unclean drinking water from various sources were at higher risk of *Toxoplasma* infection. Therefore, health education and the awareness of risk exposures regarding this parasitic disease are required to minimize the effects of this parasitic infection in pregnant women as well as in the general population.

Keywords: Toxoplasmosis, *Toxoplasma gondii*, Seroprevalence, Risk factors, Pregnant women, Thailand

* Correspondence: nissapat@gmail.com OR veeranoot@um.edu.my

¹Department of Parasitology, Faculty of Medicine, University of Malaya, Kuala Lumpur 50603, Malaysia

Full list of author information is available at the end of the article

Background

Toxoplasma gondii, an obligate intracellular protozoan parasite [1], is capable of causing severe and life threatening conditions in pregnant women and immunocompromised individuals. The sources of this parasitic infection are by the ingestion of raw and/or undercooked meat containing parasite cysts in the animal tissues, by consuming oocysts infected water and/or food, or having contact with cat fecal contaminated soil [2]. *Toxoplasma* infection in pregnant women, poses great concern to the fetus. Severe impairment occurs internally, i.e. hydrocephalus, intracerebral calcification, retinochoroiditis and mental retardation, however, clinical presentation in the newborn is asymptomatic at birth in general [3].

Detection of anti-*Toxoplasma* antibodies in pregnant women is the most widely used approach in diagnosis of this parasitic infection [4]. The detected antibodies, which indicate recent or past infection in the pregnant women, are important to confirm whether the fetus is at risk [5]. The presence of anti-*Toxoplasma* IgG antibodies represents past infection meanwhile the detection of anti-*Toxoplasma* IgM antibodies indicates recent infection [6]. However, the specified IgM antibodies remain for several months or years after initial infection [7]. This limitation causes a problem in diagnosis of whether the maternal infection occurs prior to or after conception. Misinterpretation of IgM positive results in conventional single-serum assay may lead to misdirection in treatment and termination of pregnancy [8]. The assessment of IgG avidity for *Toxoplasma* infection in pregnant women has been introduced in recent studies to assist in discrimination between past and recently acquired infection. The result of this avidity test is most helpful in determining the infection of *Toxoplasma* in pregnant woman, especially for those who are in their first trimester [9].

Therefore, this study was conducted to determine the current seroprevalence of *Toxoplasma* infection among pregnant women attending the ANC at Songklanagarind hospital, southern Thailand, to investigate the association between plausible risk factors of *Toxoplasma* infection with the seropositive pregnant women and to validate the stages of *Toxoplasma* infection in these pregnant women using avidity measurement.

Methods

Study site and population

A prospective cross-sectional study was carried out at the ANC of Songklanagarind hospital, Hat Yai, Songkhla province, Thailand from December 2012 to August 2013. This public hospital attached to Prince of Songkla University, with its capacity of 850 in-patient beds, is located in the south of Thailand and was built to

facilitate the teaching, research, and training for medical personnel in various disciplines, and for the provision of healthcare to the general public, particularly among Southern Thais. The study included 760 eligible pregnant women who gave informed consent before this study. The questionnaire [10] was designed to detect socio-demographic and biologically plausible risk factors associated with *Toxoplasma* infection, and clinical history and presenting signs and symptoms relating to toxoplasmosis (if any). This study was conducted with the approval from the ethical committee of the Faculty of Medicine, Prince of Songkla University, Thailand (Ethics number: EC 52-268-12-1-3).

Serum collection

Approximately 5 mL of venous blood samples were drawn and their sera were collected and kept at -20°C until further testing.

Screening for anti-*Toxoplasma* IgG and IgM antibodies

Anti-*Toxoplasma* IgG and IgM antibodies were screened by using a standard ELISA commercial kit (IgG-Trinity Biotech and IgM-Trinity Biotech, New York) in accordance with the manufacturer's instructions. A positive sample for the anti-*Toxoplasma* IgG and IgM antibodies was also tested for its avidity using a standard ELISA commercial kit (IgG-NovaLis Dietzenbach, Germany); high avidity ($>40\%$) indicated a past infection (of at least 4–5 months) and a low avidity ($<40\%$) indicated a recently acquired infection (within 4–5 months).

Statistical analysis

Data obtained from both the questionnaire and laboratory tests were entered, edited, and analyzed using the statistical software SPSS version 17.0 (SPSS, Inc., Chicago, IL). The data with quantitative variables were expressed as the mean (\pm SD) and range, whereas qualitative variables were estimated and presented as frequencies and percentages. Univariate analyses and the 2 test were used to investigate the association between *Toxoplasma* seropositivity as a dependent variable and possible demographic and risk factors as independent variables; $p < 0.05$ was regarded as being statistically significant.

Results

A total of 760 pregnant women were recruited. Their age range was from 14 to 47 years with a mean of 29.5 ± 6.34 years. The majority of these women was between 26 and 35 years (408, 53.7%), had tertiary education (369, 48.6%) and worked as laborers (317, 41.7%). Most of these pregnant women were in their first trimester of pregnancy (621, 81.7%), had no history of receiving antibiotic treatment (664, 87.4%), have one or no child (637, 83.8%) and

had no experience of miscarriage (592, 77.9%) as shown in Table 1.

Overall, the seroprevalence of *Toxoplasma* infection in our study was 190 (25.0%, 95% CI = 22.05-28.20) of which 167 (22.0%, 95% CI = 19.0-25.0) were positive for only anti-*Toxoplasma* IgG antibodies and 23 (3.0%, 95% CI = 2.0-4.0) were positive for both anti-*Toxoplasma* IgG and IgM antibodies. Serum samples positive for only anti-*Toxoplasma* IgM antibodies (19, 2.5%) were re-screened by collecting second serum samples after intervals of four weeks apart. These serum samples were later reported as false positive since there was no seroconversion found.

The 23 samples that were seropositive for both anti-*Toxoplasma* IgG and IgM antibodies were subsequently assessed by IgG avidity test. All these samples showed high avidity, indicating they may have acquired the infection four to five months earlier. It was noted that 20 of these samples were in their first trimester and the remaining three were in their second trimester of pregnancy.

For univariate analysis, this study verified that age group, occupation and source of drinking water had significant associations with seropositive pregnant women ($p < 0.05$), as shown in Table 1. After the multivariate logistic regression analysis was performed, it was confirmed that age ≥ 26 (OR = 1.65, 95% CI = 1.11-2.44), working as a laborer (OR = 1.57, 95% CI = 1.13-2.18) and drinking unclean (piped/tap/rain) water (OR = 1.75, 95% CI = 1.08-2.84) were identified as significant risk factors for *Toxoplasma* acquisition, as shown in Table 2.

Discussion

Toxoplasma infection in pregnant women shows variation in seroprevalence globally. From our findings, the overall seroprevalence of chronic *Toxoplasma* infection was 25% in the recruited Thai pregnant women. This finding fell within the range of 2.6% - 28% of seroprevalence that have been reported in Thailand [10-18]. However, our prevalence rate is still higher compared to previous studies conducted recently in our Asian counterparts such as China, Japan and Taiwan being 3.98%, 10.3% and 11.8%, respectively [19-21]. Globally, the seroprevalence of *Toxoplasma* infection remains high in many countries across the continents, e.g., 84.7% in Congo [22], 83.6% in Ethiopia [23], 45% in India [24], 59% in Brazil [25] and 30.9% in Tanzania [26]. The seroprevalence may vary in a global view, but the risk of this parasitic infection in human populations, especially in pregnant women, still holds a great interest. Our findings imply that we need to take a holistic approach to educate our pregnant women on toxoplasmosis in order to reduce the infection rate and the overall disease burden in our society.

In this study, 2.5% of pregnant women were false positive for anti-*Toxoplasma* IgM antibodies after re-screening their second serum sample collected four weeks later to test for seroconversion. This phenomenon could be due to host natural IgM antibodies reacting with *Toxoplasma* antigens without infection [1,8]. Meanwhile, the pregnant women who were positive for both anti-*Toxoplasma* IgG and IgM antibodies have high IgG avidity, indication of past infection. The specified IgM antibodies may remain for several months or years after initial infection [7]. The majority of these pregnant women were in their first trimester of pregnancy. This is of great concern as the effect of *Toxoplasma* infection is more severe to the fetus during the first trimester of pregnancy [27]. In the incidence of positive for both anti-*Toxoplasma* IgG and IgM antibodies, possibilities of acute infection or false IgM positive results were predicted. IgG avidity measurement assists as a confirmatory tool in the determination of infection stages in the suspected pregnant women with the availability of a single serum sample [28]. A high avidity of IgG antibodies indicates that there is no risk for congenital toxoplasmosis in the fetus, especially for pregnant women in their first trimester, regardless of the IgM antibodies results [5].

Our epidemiological data showed that age ≥ 26 years, working as a laborer and drinking unclean water were significant factors associated with *Toxoplasma* infection. This could be generally explained that the seroprevalence of *Toxoplasma* infection increases by age [29] and also specifically by their low-socioeconomic and poor hygiene practice which can play as an important role in the transmission of the parasites [30-32].

Pregnant women with more than one child have been shown a significant association with *Toxoplasma* infection [33] and this could be due to lack of cleanliness among their children [34]. However, no significant association was found between obstetric histories and *Toxoplasma* seropositivity in our study. Most of these pregnant women were not aware of toxoplasmosis and this could lead to high *Toxoplasma* seropositivity, as they do not know how to protect themselves from this parasitic infection [35,36].

Concerning other plausible risk factors, some of our pregnant women had a history of close contacts with cats (37%) but had no significant association with *Toxoplasma* infection. This finding is, however, contrary to a few of the previous studies reported, where having close contact with felines was shown as one of the vital factors in transmission of *Toxoplasma* infection [37,38]. Studies have also shown that *Toxoplasma* infection can be transmitted by other animals to their owners [39-41]. *Toxoplasma* cysts infected animals' tissues for human consumption is a mode of disease transmission [42-44]. Most of our pregnant women had consumed undercooked meat but there

Table 1 Socio-demographic, obstetric profiles and risk factors associated with *Toxoplasma* seropositivity in pregnant women by univariate analysis

Variable	No. of pregnant women		P-value
	Total (n, %) N = 760	<i>Toxoplasma</i> seropositive (n, %) N = 190	
Demographic profile			
Age			
Range 14–47 years			
Mean 29.5 ± 6.34 years			
Age group			0.042
14–25	214 (28.2)	40 (18.7)	
26–35	408 (53.7)	112 (27.5)	
36–47	138 (18.2)	38 (27.5)	
Education			0.311
Primary	48 (6.3)	16 (33.3)	
Secondary	343 (45.1)	80 (23.3)	
Tertiary	369 (48.6)	94 (25.5)	
Occupation			0.002
Labourer	317 (41.7)	95 (29.9)	
Non-Labourer	275 (36.2)	69 (25.0)	
Unemployed	168 (22.1)	26 (15.5)	
Obstetric history			
Antibiotic usage			
No	664 (87.4)	167 (25.2)	0.801
Yes	96 (12.6)	23 (23.9)	
Trimester			
First	621 (81.7)	162 (26.1)	0.255
Second	129 (17.0)	27 (20.9)	
Third	10 (1.3)	1 (10.0)	
No. children			0.460
≤1	637 (83.8)	156 (24.5)	
≥2	123 (16.2)	34 (27.6)	
History of abortion			
No	592 (77.9)	148 (25.0)	1.000
Yes	168 (22.1)	42 (25.0)	
Awareness of toxoplasmosis			
No	708 (93.2)	180 (25.4)	0.320
Yes	52 (6.8)	10 (19.2)	
Risk factors of toxoplasmosis			
Close contact with cats			
No	482 (63.4)	114 (23.7)	0.258
Yes	278 (36.6)	76 (27.3)	
Consumption of undercooked meat			
No	275 (36.2)	64 (23.3)	0.408
Yes	485 (63.8)	126 (25.9)	

Table 1 Socio-demographic, obstetric profiles and risk factors associated with *Toxoplasma* seropositivity in pregnant women by univariate analysis (Continued)

Receiving blood transfusion			0.414
No	758 (99.7)	190 (25.1)	
Yes	2 (0.26)	0 (0)	
Drinking water from various source			0.013
Boiled	75 (9.8)	24 (32.0)	
Mineral/Filtered	597 (78.6)	135 (22.6)	
Piped/tap/rain	88 (11.6)	31 (35.2)	
Drinking milk			0.693
No	97 (12.8)	27 (27.8)	
Boiled	8 (1.1)	3 (37.5)	
Pasteurized	649 (85.4)	158 (24.3)	
Non-Pasteurized	6 (0.8)	2 (33.3)	
Having contact with soil			0.395
No	316 (41.6)	74 (23.4)	
Yes	444 (58.4)	116 (26.1)	
Close contact with other animals			0.967
No	369 (48.6)	92 (24.9)	
Yes	391 (51.4)	98 (25.0)	

is no significant association with high *Toxoplasma* infection. This could be due to absence of contamination by *Toxoplasma* cysts in the consumed meat and needs to be further studied. Our finding is consistent with a previous study reported in this region [11], showing that drinking unclean water (pipe/tap/rain) had significant association with *Toxoplasma* seropositive pregnant women, indicating the water may be contaminated with *Toxoplasma* oocysts. However, high seropositive (32%) were also found in pregnant women who consumed boiled water but were not statistically significant. Based on the above results, this could be a good indicator of other confounding factors contributing to *Toxoplasma* infection and requires further studies.

Conclusion

Our findings showed high *Toxoplasma* infection rates in this group of pregnant women and showed significantly

higher risk with age group, low socioeconomic status and drinking unclean water. The following guidelines should be implemented to eliminate *Toxoplasma* infection and eventually eradicate its disease burden; Firstly, a routine screening for toxoplasmosis among women in the reproductive age group and pregnant women, especially for those in their early pregnancy is strongly encouraged for monitoring and preventive purposes. Secondly, health education on toxoplasmosis (brochures were attached as Additional file 1) and its risk exposures is required to increase the awareness about this disease and to minimize the effects of this *Toxoplasma* infection in the general population and pregnant women in particular. Lastly, serological diagnosis through the detection of anti-*Toxoplasma* antibodies and IgG avidity measurement in pregnant women assist the dating of the infection and as well as determining the decision for course of treatment especially those in their early pregnancies.

Table 2 Multivariate logistic regression analysis for demographic profiles and risk factors associated with *Toxoplasma* seropositive pregnant women

Variable	Adjusted odds ratio	95% CI		P value
Age ≥26	1.65	1.11	2.44	0.012
Working as labourer	1.57	1.13	2.18	0.007
Drinking Piped/tap/rain water	1.75	1.08	2.84	0.018

Adjusted variables include age group, occupation and source of drinking water in this statistical analysis.

Additional file

Additional file 1: Health care education on Toxoplasmosis in pregnant women. Brochures in English Language. Brochures in Thai language.

Competing interest

All authors declare having no conflicts of interest.

Authors' contributions

All authors were involved in the design of the study. VN was the principal investigator. VN and NS drafted the protocol, consent forms, and paper with input from the other authors. HA, TK, SO and WC were responsible for sample collection, serological analysis and data analysis. HA and WC were responsible for statistical analysis and interpretation. LYL, NS, CS, VC and VN were involved in data analysis, statistical analysis and interpretation and results discussion and inputs. All authors contributed to the writing of the paper and approved the final version.

Acknowledgments

We thank all the participating pregnant women for their consistent support and cooperation throughout this study. We also thank the doctors, nurses, and staffs of ANC at Songklanagarind Hospital, Hat Yai, Songkhla province, Thailand for their assistance and contribution in this study.

Funding

This study was supported by UM High Impact Research Grant UM-MOHE UM.C/625/1/HIR/MOHE/MED/18 from the Ministry of Higher Education Malaysia, and University of Malaya Research Grant (UMRG 488/12HTM and UMRG 544/14HTM).

Author details

¹Department of Parasitology, Faculty of Medicine, University of Malaya, Kuala Lumpur 50603, Malaysia. ²Department of Microbiology, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand. ³Department of Obstetrics and Gynecology, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand. ⁴Faculty of Medicine, University of Phayao, Phayao 56000, Thailand.

Received: 28 March 2014 Accepted: 19 May 2014

Published: 22 May 2014

References

1. Sensini A: *Toxoplasma gondii* infection in pregnancy: opportunities and pitfalls of serological diagnosis. *Clin Microbiol Infect* 2006, **12**:504–512.
2. Cook AJ, Gilbert RE, Buffolano W, Zufferey J, Petersen E, Jennum PA, Foulon W, Semprini AE, Dunn DT: Sources of toxoplasmosis infection in pregnant women: European multicentre case-control study. *BMJ* 2000, **321**:142–147.
3. Singh S: Mother-to-child transmission and diagnosis of *Toxoplasma gondii* infection during pregnancy. *Indian J Med Microbiol* 2003, **21**:69–76.
4. Giannoulis C, Zournatzi B, Giomisi A, Diza E, Tzafettas I: Toxoplasmosis during pregnancy: case report and review of the literature. *Hippokratia* 2008, **12**:139–143.
5. Montoya JG, Remington JS: Management of *Toxoplasma gondii* infection during pregnancy. *Clin Infect Dis* 2008, **47**:554–566.
6. Lopes FM, Gonçalves DD, Mitsuka-Breganó R, Freire RL, Navarro IT: *Toxoplasma gondii* infection in pregnancy. *Braz J Infect Dis* 2007, **11**:496–506.
7. Leite M, Siciliano S, Rocha LS, Justa MT, César KR, Granato CF: Correlation between specific IgM levels and percentage IgG-class antibody avidity to *Toxoplasma gondii*. *Rev Inst Med Trop Sao Paulo* 2008, **50**:237–242.
8. Liesenfeld O, Press C, Montoya JG, Gill R, Isaac-Renton JL, Hedman K, Remington JS: False-positive results in immunoglobulin M (IgM) *Toxoplasma* antibody tests and importance of confirmatory testing: the Platelia Toxo IgM test. *J Clin Microbiol* 1997, **35**:174–178.
9. Remington JS, Thulliez P, Montoya JG: Recent developments for diagnosis of toxoplasmosis. *J Clin Microbiol* 2004, **42**:941–945.
10. Nissapatorn V, Suwanrath C, Sawangjaroen N, Ling LY, Chandeying V: Toxoplasmosis-serological evidence and associated risk factors among pregnant women in southern Thailand. *Am J Trop Med Hyg* 2011, **85**:243–247.
11. Sakae C, Natphopsuk S, Settheetham-Ishida W, Ishida T: Low prevalence of *Toxoplasma gondii* infection among women in northeastern Thailand. *J Parasitol* 2013, **99**:172–173.
12. Sukthana Y: Difference of *Toxoplasma gondii* antibodies between Thai and Austrian pregnant women. *Southeast Asian J Trop Med Public Health* 1999, **30**:38–41.
13. Chintana T: Pattern of antibodies in toxoplasmosis of pregnant women and their children in Thailand. *Southeast Asian J Trop Med Public Health* 1991, **22**(Suppl):107–110.
14. Puthavathana P, Pimolpan K, Louisirirotchanaikul S, Wasi C, Thongcharoen P: Sero-epidemiology of TORCH agents among pregnant Thais. *Asian Pac J Allergy Immunol* 1983, **1**:11–14.
15. Wanachiwanawin D, Sutthent R, Chokephaibulkit K, Mahakittikun V, Ongrotchanakun J, Monkong N: *Toxoplasma gondii* antibodies in HIV and non-HIV infected Thai pregnant women. *Asian Pac J Allergy Immunol* 2001, **19**:291–293.
16. Maleewong W, Lulitanond V, Pipitgool V, Auwijitaroon Y, Kutsarejariya S, Morakote N: Prevalence of *Toxoplasma* antibodies in blood donors and pregnant women in Khon Kaen Province. *J Med Assoc Thai* 1989, **72**:256–259.
17. Chintana T, Sukthana Y, Bunyakai B, Lekka A: *Toxoplasma gondii* antibody in pregnant women with and without HIV infection. *Southeast Asian J Trop Med Public Health* 1998, **29**:383–386.
18. Sukthana Y, Chintana T, Supatanapong W, Siripan C, Lekka A, Cheabchalrad R: Predictive value of latex agglutination test in serological screening for *Toxoplasma gondii*. *Southeast Asian J Trop Med Public Health* 2001, **32**:314–318.
19. Hua HY, Tang F, Liu YX, You L, Dong MH, Chen YE, Chen H, Hu YH, Ding GS, Sun BC, Guo JH, Gao Q: Survey of *Toxoplasma gondii* infection among pregnant women in Jiangsu Province, China. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* 2013, **25**:56–58.
20. Sakikawa M, Noda S, Hanaoka M, Nakayama H, Hojo S, Kakinoki S, Nakata M, Yasuda T, Ikenoue T, Kojima T: Anti-*Toxoplasma* antibody prevalence, primary infection rate, and risk factors in a study of toxoplasmosis in 4,466 pregnant women in Japan. *Clin Vaccine Immunol* 2012, **19**:365–367.
21. Chou CS, Lin LY, Chen KM, Lai SC: Flowcytomix analysis for *Toxoplasma gondii* infection in pregnant women in central Taiwan. *J Obstet Gynaecol* 2011, **31**:375–379.
22. Doudou Y, Renaud P, Coralie L, Jacqueline F, Hypolite S, Hypolite M, Patrick M, Andria Ida L, Van Sprundel M, Marleen B, Van Geertruyden JP, Pascal L: Toxoplasmosis among pregnant women: high seroprevalence and risk factors in Kinshasa, Democratic Republic of Congo. *Asian Pac J Trop Biomed* 2014, **4**:69–74.
23. Zemene E, Yewhalaw D, Abera S, Belay T, Samuel A, Zeynudin A: Seroprevalence of *Toxoplasma gondii* and associated risk factors among pregnant women in Jimma town, Southwestern Ethiopia. *BMC Infect Dis* 2012, **12**:337.
24. Singh S, Pandit AJ: Incidence and prevalence of toxoplasmosis in Indian pregnant women: a prospective study. *Am J Reprod Immunol* 2004, **52**:276–283.
25. Ferezin RI, Bertolini DA, Demarchi IG: Prevalence of positive serology for HIV, hepatitis B, toxoplasmosis and rubella in pregnant women from the northwestern region of the state of Paraná. *Rev Bras Ginecol Obstet* 2013, **35**:66–70.
26. Mwambe B, Mshana SE, Kidenya BR, Massinde AN, Maziqo HD, Micheal D, Majinge C, Groß U: Sero-prevalence and factors associated with *Toxoplasma gondii* infection among pregnant women attending antenatal care in Mwanza, Tanzania. *Parasit Vectors* 2013, **6**:222.
27. Kieffer F, Wallon M: Congenital toxoplasmosis. *Handb Clin Neurol* 2013, **112**:1099–1101.
28. Ashburn D, Joss AW, Pennington TH, Ho-Yen DO: Do IgA, IgE and IgG avidity tests have any value in the diagnosis of *Toxoplasma* infection in pregnancy? *J Clin Pathol* 1998, **51**:312–315.
29. Nowakowska D, Wujcicka W, Sobala W, Spiewak E, Gaj Z, Wilczynski J: Age associated prevalence of *Toxoplasma gondii* in 8281 pregnant women in Poland between 2004 and 2012. *Epidemiol Infect* 2014, **142**:656–661.
30. Alvarado-Esquivel C, Torres-Castorena A, Liesenfeld O, Estrada-Martinez S, Urbina-Álvarez JD: High seroprevalence of *Toxoplasma gondii* infection in a

- subset of Mexican patients with work accidents and low socioeconomic status. *Parasit Vectors* 2012, **5**:13.
31. Dattoli VC, Veiga RV, Cunha SS, Pontes-de-Carvalho L, Barreto ML, Alcantara-Neves NM: Oocyst ingestion as an important transmission route of *Toxoplasma gondii* in Brazilian urban children. *J Parasitol* 2011, **97**:1080–1084.
 32. Francisco Fde M, De Souza SL, Gennari SM, Pinheiro SR, Muradian V, Soares RM: Seroprevalence of toxoplasmosis in a low-income community in the Sao Paulo municipality, SP, Brazil. *Rev Inst Med Trop Sao Paulo* 2006, **48**:167–170.
 33. Jenum PA, Kapperud G, Stray-Pedersen B, Melby KK, Eskild A, Eng J: Prevalence of *Toxoplasma gondii* specific immunoglobulin G antibodies among pregnant women in Norway. *Epidemiol Infect* 1998, **120**:87–92.
 34. Mohan B, Dubey ML, Malla N, Kumar R: Seroepidemiological study of toxoplasmosis in different sections of population of Union Territory of Chandigarh. *J Commun Dis* 2002, **34**:15–22.
 35. Ferguson W, Mayne PD, Cafferkey M, Butler K: Lack of awareness of risk factors for primary toxoplasmosis in pregnancy. *Ir J Med Sci* 2011, **180**:807–811.
 36. Pappas G, Roussos N, Falagas ME: Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis. *Int J Parasitol* 2009, **39**:1385–1394.
 37. Gebremedhin EZ, Abede AH, Tessema TS, Tullu KD, Medhin G, Vitale M, Di Marco V, Cox E, Dorny P: Seroepidemiology of *Toxoplasma gondii* infection in women of child-bearing age in central Ethiopia. *BMC Infect Dis* 2013, **13**:101.
 38. Ngui R, Lim YA, Amir NF, Nissapatorn V, Mahmud R: Seroprevalence and sources of toxoplasmosis among Orang Asli (indigenous) communities in Peninsular Malaysia. *Am J Trop Med Hyg* 2011, **85**:660–666.
 39. Yang N, Mu M, Li H, Hu J, Gao W, Yang S, He J: Seroprevalence of *Toxoplasma gondii* infection in pet dogs in Shenyang, northeastern China. *J Parasitol* 2013, **99**:176–177.
 40. Hemsworth S, Pizer B: Pet ownership in immunocompromised children—a review of the literature and survey of existing guidelines. *Eur J Oncol Nurs* 2006, **10**:117–127.
 41. Schurer JM, Hill JE, Fernando C, Jenkins EJ: Sentinel surveillance for zoonotic parasites in companion animals in indigenous communities of Saskatchewan. *Am J Trop Med Hyg* 2012, **87**:495–498.
 42. Hill DE, Dubey JP: *Toxoplasma gondii* prevalence in farm animals in the United States. *Int J Parasitol* 2013, **42**:107–113.
 43. Kang SW, Doan HT, Noh JH, Choe SE, Yoo MS, Kim YH, Reddy KE, Nguyen TT, Van Quyen D, Nguyen LT, Kweon CH, Jung SC: Seroprevalence of *Toxoplasma gondii* and *Trichinella spiralis* infections in wild boars (*Sus scrofa*). *Parasitol Int* 2013, **62**:583–585.
 44. Wang L, Cheng HW, Huang KQ, Xu YH, Li YN, Du J, Yu L, Luo QL, Wei W, Jiang L, Shen JL: *Toxoplasma gondii* prevalence in food animals and rodents in different regions of China: isolation genotyping and mouse pathogenicity. *Parasit Vectors* 2013, **6**:273.

doi:10.1186/1756-3305-7-239

Cite this article as: Andiappan et al.: *Toxoplasma* infection in pregnant women: a current status in Songklanagarind hospital, southern Thailand. *Parasites & Vectors* 2014 **7**:239.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

