

Invited Mini Review

Synapses in neurodegenerative diseases

Jae Ryul Bae¹ & Sung Hyun Kim^{2,*}¹Department of Biomedical Science, Graduate School, Kyung Hee University, ²Department of Physiology, School of Medicine, Kyung Hee University, Seoul 02447, Korea

Synapse is the basic structural and functional component for neural communication in the brain. The presynaptic terminal is the structural and functionally essential area that initiates communication and maintains the continuous functional neural information flow. It contains synaptic vesicles (SV) filled with neurotransmitters, an active zone for release, and numerous proteins for SV fusion and retrieval. The structural and functional synaptic plasticity is a representative characteristic; however, it is highly vulnerable to various pathological conditions. In fact, synaptic alteration is thought to be central to neural disease processes. In particular, the alteration of the structural and functional phenotype of the presynaptic terminal is a highly significant evidence for neural diseases. In this review, we specifically describe structural and functional alteration of nerve terminals in several neurodegenerative diseases, including Alzheimer's disease (AD), Parkinson's disease (PD), Amyotrophic lateral sclerosis (ALS), and Huntington's disease (HD). [BMB Reports 2017; 50(5): 237-246]

INTRODUCTION

Synapse is a fundamental unit for brain function. Flow of neural information between neurons is ignited from presynaptic terminals by the release of a small chemical ingredient called neurotransmitter. This is stored in the small endosomal compartment (synaptic vesicle (SV)), and it's released by the fusion of SV, facilitated by a series of neuronal-activity triggered action of molecular players in the release area (active zone) at the presynaptic terminals.

The presynaptic terminal is composed of several structural and functional components. Cytomatrix proteins (Bassoon and Piccolo) and cytoskeletal protein (actin) provide the structural framework. The active zone (AZ) is the critical site for neurotransmitter release. Many essential molecular machineries

are localized in the AZ, such as SNARE components for fusion, voltage-gated Ca²⁺ channels, cell adhesion molecules, etc. Each nerve terminal possesses around 100-200 synaptic vesicles. The synaptic vesicle (SV), a tiny endosomal compartment (~40 nm diameter), contains the neurotransmitter which associates directly and/or indirectly with more than a hundred proteins for proper functioning. For appropriate physiological functions, a number of proteins exist at the nerve terminals. Physiologically, the regulation and maintenance of neurotransmitter release remain critical questions. Several distinct SV pools distributed in the presynaptic terminal and SV exocytosis is tightly regulated by Ca²⁺ and its molecular players. Subsequently, SV retrieval continuously maintains the synaptic communication via several endocytic pathways. However, the morphological and physiological intact is easily altered in various neurological diseases. From synaptic vesicle and synaptic protein depletion to neurotransmission and Ca²⁺ dynamics impairment, a number of alterations in the structure and function of nerve terminals are exhibited in neurological diseases. Furthermore, these presynaptic dysfunctions are thought to be the very early symptoms of neuronal disorders.

This review specifically describes the structural and functional presynaptic alterations in neurodegenerative diseases. Alzheimer's disease (AD) is a high impact neurodegenerative disease. Several pathogenic factors have been identified, such as amyloid beta (A β) plaque, neurofibrillary tangle, and ApoE4. However, the exact pathological etiology requires to be further explored. It is important to understand synaptic alterations by these factors at the initial stage, before the eventual occurrence of neuronal cell death. Parkinson's disease (PD) is the second most common neurodegenerative disease, exhibiting degeneration of dopaminergic neurons in the substantia nigra pars compacta. Consequently, this results in dopamine depletion in the brain, leading to several neurological symptoms, tremors, bradykinesia, and rigidity. A number of sporadic and familial factors have been discovered. Some of the evidences have reported that these factors are deeply implicated with presynaptic function, although it is still much less known how PD is initiated. Other neurodegenerative diseases, such as Huntington disease (HD) and amyotrophic lateral sclerosis (ALS), are also involved in synaptic dysfunction. We describe in depth normal and pathological phenotype of these factors at presynaptic terminals.

*Corresponding author. Tel: +82-2-961-0286; Fax: +82-2-964-2195; E-mail: sunghyunkim@khu.ac.kr

<https://doi.org/10.5483/BMBRep.2017.50.5.038>

Received 7 March 2017

Keywords: Alzheimer's disease, Neurotransmission, Parkinson's disease, Presynaptic terminals, Synaptic vesicle

NERVE TERMINALS IN ALZHEIMER'S DISEASE

Alzheimer's disease (AD), the most common type of dementia, is a rapidly emergent and prominent neurodegenerative disease. The patient progressively loses their memory with a decline in cognition, eventually reaching mortality due to death of the brain cells. Several causative genetic factors have been revealed. Oligomerization of the amyloid beta (A β) plaque from amyloid precursor protein (APP) by BACE and γ -secretase, is a well-known factor for AD. Mutation or modification of the Tau protein aggregates to form neurofibrillary tangle (NFT) or paired helical filaments (PHF), called Tauopathy, which is also a known cause of AD. A critical genetic factor for late-onset AD is apolipoprotein E, particularly ϵ 4 isoforms (ApoE4). Although these genetic factors are identified and characterized, a number of complicities are still emerging, and remain elusive. Here, we describe the genetic factors involved in the function and dysfunction at the presynaptic terminals (Table 1).

Amyloid precursor protein (APP) and Amyloid beta (A β)

Amyloid precursor protein (APP) is an essential source for amyloid beta 40 or 42 (A β 40 or 42) which are known major pathogenic factors in AD. APP normally participates in presynaptic function, although primary function of APP is still not explored. APP is enriched in nerve terminals with Rab5 positive large vesicular organelle (1) or a small set of synaptic vesicles (2), and is involved in structure and function of nerve terminals. APP modulates the initial nerve terminal formation. Cultured neurons from APP knock-out brain revealed up-regulation of synaptophysin, a presynaptic marker. Consistently, immunohistochemistry from a slice of APP KO brain showed a high intensity of synaptophysin, indicating that APP is a negative regulator of synaptic formation. Secondly, APP is also involved in physiological modulation of synaptic functions.

APP KO neurons significantly increase the readily releasable pool (RRP) of synaptic vesicles (3). According to computational analysis of APP, it is likely to serve as a hub protein in the presynaptic active zone (PAZ), and is a context regulator in the hippocampal active zone network (4).

Amyloid beta (A β) is a fragment peptide from APP, cleaved by BACE and γ -secretase. Oligomeric aggregation of A β peptide is a critical pathogenic factor in AD. Several reports exhibit the A β tangle effects in nerve terminal phenotype. Treatment of A β oligomer in neurons results in significantly decreasing the presynaptic protein expression, but not post-synapse (5), indicating that A β affects the initial structural formation of presynaptic terminals. Physiologically soluble A β binds to APP, inducing the APP-APP homodimer. Consequently, there is a boost of Ca²⁺ influx, eventually resulting in an increase in the release probability (6, 7), indicating that A β is a positive regulator of neurotransmission at nerve terminal. However, in pathological conditions, increased A β can perturb the release probability by altering spike probability of neurons (8). Internalized A β is localized at the nerve terminal, subsequently disrupting the synaptic vesicle protein VAMP2 function for vesicle fusion (9). In addition, it induces the depletion of presynaptic mitochondria and decreases its motility, thereby decreasing the size of synaptic vesicle pool.

Another important feature of the A β at synapses is that the synaptic activity for neurotransmission and release of A β is tightly correlated, and a nerve terminal is a major place for A β release. The brain interstitial fluid (ISF) reveals that synaptic activity influences the A β levels. The more the synaptic activity, higher is the A β level in ISF. This result correlates with APP endocytosis. The cleavage of APP to produce A β occurs in the endosomes or a small fraction of SV, and not on the surface of the plasma membrane (2). Synaptic vesicle exocytosis is thus required for more endocytosis of APP.

Table 1. Summary of presynaptic phenotype by AD genetic factors

Alzheimer's disease		
Factor	Phenotype at presynaptic terminal	Ref
Amyloid Precursor Protein (APP)	• Negative regulator of synapse formation	3
	• Negative regulator for readily releasable pool of synaptic vesicle	3
	• Molecular Hub in presynaptic active zone (PAZ)	4
Amyloid beta (A β)	• Downregulation of presynaptic protein expression	5
	• Increase release probability (soluble A β - normal condition)	7
	• Disruption of vesicle fusion ability by inhibiting VAMP2 function (pathologic A β)	9
BACE1 γ -secretase/Presenilin	• Negative regulator for excitatory synaptic transmission (homeostatic synaptic plasticity)	15
	• Presynaptic short-term plasticity, synaptic facilitation	17
	• Homeostatic synaptic scaling of excitatory synapses	18
Tau	• Synaptic stability (presynaptic proteins, synaptic vesicle)	19, 20
	• Downregulation of amount glutamate	27
ApoE4	• Modulation of spontaneous vesicle release	28

Hence, the production and release of A β are modulated by activity-dependent synaptic transmission and endocytosis at the nerve terminals (10, 11).

Beta-secretase (BACE)

β -site amyloid precursor protein-cleaving enzyme 1 (BACE1) is a key enzyme that produces Amyloid beta under pathological conditions. Localized at the synaptic vesicles, BACE1 is important for synaptic functions. Numerous potential substrates have been identified, which contain several synaptic proteins, in addition to APP (12). Furthermore, BACE1 was biochemically detected in the synaptic vesicle enriched fraction, indicating that synaptic vesicle is the likely location for APP processing (13, 14). BACE1 KO mice revealed that basal excitatory synaptic transmission was augmented. It is likely that the pathway downstream of BACE1 at the synapse was decreased, which could be due to the scaling of homeostatic synaptic plasticity (15). Synaptic adhesion protein Neuroligin1 and voltage-gated sodium channel were also known substrates for BACE1; however, it is as yet unknown how BACE1 regulates these substrates.

γ -secretase and presenilin

γ -secretase, along with BACE1, is essential for the production of a 42 peptide of amyloid beta. Several functions of γ -secretase have been reported at synapses. Localization studies revealed that γ -secretase is present in the synaptic endosomal fraction of rat brain, which is highly overlapped with the localization of BACE1 protein (16). In neurons with conditional knockout of presenilin, one of the subunits of the γ -secretase complex, presynaptic short-term plasticity and synaptic facilitation were severely altered, mainly mediated by presynaptic functions; these impairments result from intracellular Ca²⁺ release at the presynaptic terminals (17). In addition, hippocampal neurons derived from presenilin KO mice failed to the homeostatic scaling of excitatory synapses (18). Collectively, presenilin regulates neurotransmission at the nerve terminals.

Tau

Tau was originally discovered as a microtubule-associated protein. The neurofibrillary tangle (NFT) or paired helical filament (PHF) one of the major hallmarks of AD, is formed by Tau protein aggregation. However, studies report the functioning of tau at synapses. Tau is involved in axonal transport and synaptic protein stability for regulation of microtubule stability (19). In addition, it also provides the structural support to form and maintain synapses (20). Truncated tau contains specific phosphor-pattern, and can be localized both at the pre- and post-synaptic compartments. Particularly, at the presynaptic terminal, it impairs the stability of microtubules, resulting in a reduction of synaptic vesicles (21).

In pathological conditions, the Tau protein strongly influences synaptic dysfunction. The brain of the rTg4510

mouse, a human mutant P301L tau overexpressed mouse model, revealed age-dependent synaptic loss at both the pre- and post-synaptic regions, subsequently resulting in synaptic dysfunction. Tauopathy exhibits a strong impairment of synaptic transmission, and in combination with APP models, the synaptic impairment was aggravated, suggesting that the two pathological protein (Tau and APP) act in concert with synaptic function and dysregulation (22, 23).

Apolipoprotein (APOE)

ApoE is a lipoprotein mainly involved in the transport of lipoprotein, cholesterol, and lipid-related materials. It is well known that ApoE is strongly related to the pathology of AD, and is also associated with another AD factor, such as Amyloid-beta. Particularly, the apolipoprotein E4 (ApoE4) allele is the major causative allele of ApoE, having a functional role in nerve terminals. Hippocampal neurons with ApoE4 allele expression have a high sensitivity to environmental factors that lower the levels of presynaptic proteins such as synaptophysin (24, 25), although the synaptic area in the dentate gyrus increases (26). In addition, ApoE4 targets the replacement mice, showed down-regulation of glutaminase which converts the glutamine to glutamate, and up-regulation of the vesicular glutamate transporter. Consequently, neurons replaced with ApoE4 release decreased levels of glutamate at nerve terminals (27). Interestingly, this effect on presynaptic terminals appear to be restricted only in ApoE4 alleles, and not in other E2 and E3 alleles, thereby suggesting that structural and functional regulation is specifically influenced by the ApoE4 allele. Recently, it has been discovered that several ApoE receptors (e.g. Apoer2 and Vldlr) are expressed at the nerve terminal membranes. Reelin, a ligand for ApoE receptor, signaled a transient increase of intracellular Ca²⁺, resulting in elevation of spontaneous vesicle release by VAMP7 mediated fusion (28).

Also, ApoE4 and amyloid beta were closely associated in AD pathogenesis. In a patient with AD, the ApoE4 was colocalized with oligomeric A β , and enhanced the synaptic localization of oligomeric A β . These findings suggest that ApoE4 is a stimulator for oligomeric A β toxicity for synapses (29). The proteomic response in nerve terminals is more susceptible than in the cell body, suggesting that ApoE has a nerve terminal region-specific functional effect.

NERVE TERMINALS IN PARKINSON'S DISEASE

Parkinson's disease (PD) is the second most common neurodegenerative disorder. It is a movement disorder characterized by bradykinesia, postural instability, and rigidity, following the progressive loss of dopaminergic neurons in the midbrain. Pathogenesis of PD is classified into sporadic and familial cases, which develop due to environmental and genetic factors. About two dozen genetic factors of PD have been identified so far. Few genetic factors, including α -synuclein,

Table 2. Summary of presynaptic phenotype by PD genetic factors

Parkinson's disease		
Factor	Phenotype at presynaptic terminal	Ref
α -synuclein	<ul style="list-style-type: none"> • Impairment of dopamine release in SNpc • Impairment of synaptic vesicle endocytosis and reclustering • Reduction of synaptic vesicle recycling pool 	38 39,40 41
LRRK2	<ul style="list-style-type: none"> • Impairment of release and decreased DA uptake in SNpc • Impairment of synaptic endocytosis in presynaptic terminals 	48 45
Parkin	<ul style="list-style-type: none"> • Reduction of dopamine release • Impairment of synaptic plasticity in striatal cells 	55 53
PINK1	<ul style="list-style-type: none"> • Impairment of synaptic plasticity and release of dopaminergic neuron 	62
DJ-1	<ul style="list-style-type: none"> • Defect of LTD through inhibitory effects of D2 receptor 	68
Synaptojanin1	<ul style="list-style-type: none"> • Slowed endocytosis rate for small stimulation by defect of phosphatase activity 	74,75
Endophilin	<ul style="list-style-type: none"> • Regulation of Parkin expression 	76

LRRK2 (Leucine-rich repeat kinase 2), Parkin, PINK-1 (PTEN Induced Putative Kinase 1) and DJ-1, have been thoroughly researched in the pathogenesis of PD. Accumulating evidence shows that genetic factors of PD are associated with alteration of the synaptic functions (30, 31) (Table 2).

α -synuclein

α -synuclein is a small protein containing 140 amino acids, and contributes to early-onset PD (32). Generally, α -synuclein localizes at the presynaptic terminal. It is associated with synaptic vesicles, controlled synaptic vesicle trafficking, and SNARE complex formations at the nerve terminal (33-35). In pathological conditions, α -synuclein is implicated in the alteration of synaptic functions. Human α -synuclein overexpressing animal models show protein aggregations at the nerve terminals (36, 37), and overexpression of human α -synuclein by viral vector injection into substantia nigra in animal models leads to impaired dopamine release (38). Furthermore, neurotransmission inhibition might be related to the impairment of synaptic vesicle endocytosis (39) or synaptic vesicle reclustering after synaptic vesicle endocytosis (40). In addition, the overexpressing pathogenic mutants of α -synuclein (A30P and A53T) in primary midbrain neurons, led to abnormal neurite growth and reduced the recycling pool of synaptic vesicles (41). This evidence suggests that α -synuclein aggregation alters synaptic formation and functions.

LRRK2

LRRK2 is a large multidomain protein which includes kinase, GTPase, and protein-protein interaction domains. It is one of the prominent familial PD factors, particularly the gain-of-function mutant of LRRK2 (G2019S) is strongly associated with familial PD as well as sporadic PD (42, 43). Several studies report that LRRK2 is implicated in the structural and functional regulation of synapses through kinase-dependent mechanisms. It regulates the presynaptic and postsynaptic morphology by

the phosphorylation-dependent interaction of Futsch and 4E-BP in fly models (44). LRRK2 participates in synaptic vesicle endocytosis by phosphorylating endophilin (45), which is related with delayed endocytosis of synaptic vesicles, and subsequently affecting the neurotransmission impairment (46). LRRK2 also phosphorylates the NSF (N-ethylmaleimide-Sensitive Factor) D2 domain (Threonine 645), which plays a key role in SNARE complex disassembly after synaptic vesicle exocytosis. NSF phosphorylation by LRRK2 exhibits an elevated rate of SNARE disassembly (47). BAC transgenic animals for LRRK2 G2019S mutation, characterized by elevated kinase activity, show impairment of striatal dopamine release and a decrease of dopamine uptake, without dopaminergic neuron loss in the substantia nigra pars compacta (SNpc) (48). Furthermore, a neuron with LRRK2 G2019S expression shows elevated release probability with increased synaptic density (49), and altered glutamatergic synaptic plasticity (50).

Parkin

Parkin is an E3 ubiquitin ligase, and has an important role in cellular homeostasis due to its regulation of mitophagy and protein degradation. However, the loss-of-function mutation of Parkin is associated with juvenile-onset PD (51, 52). Parkin has been implicated in the modulation of synaptic functions. Parkin KO mice show decrease of evoked dopamine release in the striatum, and the striatal medium spiny neuron exhibit impairments of synaptic plasticity, which are long-term depression and long-term potentiation (53). Parkin also negatively regulates the number and strength of excitatory synapse (54), and neurotransmission is impaired by reduced AMPA receptor endocytosis due to loss of function of Parkin (55). Several studies report that functional loss of Parkin impaired degradation of synaptic proteins, including α -synuclein, synphilin-1, and CDCrel-1, thereby contribute to protein aggregation (56-58).

PINK1

Inherited nonsense and missense mutation of PINK1 (PTEN-induced putative kinase1) is a known early-onset familial PD factor (59). PINK1 has a N-terminal mitochondrial targeting motif and a conserved kinase domain (60), and is closely related to mitochondrial function and mitochondrial quality control (61). Pathologic mutation of PINK1 results in abnormal morphology of mitochondria. In addition, there is impairment of dopamine release, which presumably relates to synaptic mitochondrial dysfunction by pathogenic PINK1 (62). Loss of PINK1 also impairs the normal development of dopaminergic neuron, consequently leading to locomotor dysfunction (63). PINK1-deficient mice show normal number of dopaminergic neurons, but the release of dopamine is significantly decreased, suggesting that PINK1 has a role in synaptic transmission (64).

DJ-1

Generally, DJ-1 acts as a sensor for cellular redox homeostasis (65). However, functional mutation of DJ-1 is a causative familial factor for autosomal recessive early-onset PD (66). Localization studies reveal that DJ-1 is localized in the synaptic membrane. The binding affinity for synaptic membrane reduces with pathogenic DJ-1 compared to WT DJ-1 (67), indicating that it is likely involved in synaptic functions. In fact, DJ-1 depleted mice show signs of LTD (long-term depression), through the inhibitory effects of the D2 receptor by loss of DJ-1 (68).

Synaptojanin-1

Synaptojanin-1, a known phosphoinositide phosphatase, has a role in endocytosis process. It interacts with several endocytic proteins such as dynamin, Dap160/intersectin, and BAR proteins including endophilin and amphiphysin (69, 70), suggesting a key role in synaptic vesicle recycling processing, particularly clathrin-coated pit uncoating (71). Recently, the Sac1 domain mutation of synaptojanin-1 (p.Arg258Gln) has been reported in a family with early-onset progressive Parkinsonism (72, 73). Although synaptojanin-1 mutation mediated pathogenesis of PD has been less explored, the pathogenic phenotype is exhibited in mutations of synaptojanin-1 associated with PD, as well as early onset refractory seizures and neurological decline (74, 75), suggesting that the loss-of-function of synaptojanin-1 may contribute to the pathogenesis of PD and other neurological diseases by impaired synaptic vesicle recycling.

Endophilin

Endophilin is a key factor in synaptic vesicle recycling. Recently, however, some papers report that it is related to PD genetic factors, including LRRK2, parkin, and synaptojanin-1 (45, 76, 77). Endo-A, a fly ortholog of endophilin, is a substrate for LRRK2. BAR domain (Serine75) in Endo-A is phosphorylated, and recruitment of Endo-A to the endocytic

complex during endocytosis gets modulated. Consequently, hyper-phosphorylation of BAR domain of Endo-A in LRRK2 G2019S mutant shows impairment of synaptic endocytosis in presynaptic terminals (45). In addition, endophilin phosphorylation by LRRK2 increases the recruitment of atg3 to the membrane area of presynaptic terminals, resulting in macroautophagy induction, affecting the membrane curvature induction for autophagy (78). Interestingly, endophilin mutant mice exhibited strongly increased parkin expression, suggesting that endophilin genetically interacts with parkin (76).

NERVE TERMINALS IN OTHER NEURODEGENERATIVE DISEASES

ALS

Amyotrophic lateral sclerosis (ALS) is a motor neuron disorder characterized by progressive loss of motor neurons in the cortex, brainstem and spinal cord. The loss of motor neuron leads to muscle atrophy and weakness, thereby eventually resulting in death. Superoxide dismutase-1 (SOD-1), one of the most prominent ALS genetic factors, is an antioxidant enzyme involved in the conversion of free superoxide radicals to oxygen and hydrogen peroxide. Both, a dominant and a recessive mutation of SOD-1, have been identified in ALS patients (79-81). It has been reported that the mutations of SOD-1 were implicated in synaptic dysfunctions. Both wild type of SOD-1 and pathogenic SOD-1 were localized at the pre and post-synapse. The G93A SOD-1 mutant, one of the pathogenic SOD-1 mutants, shows mislocalization in pre-synaptic terminals as well as post-synapse, thereby impairing axonal transport and contributing neuronal cell death (82, 83). SOD1 mutant mice also show length-dependent axonopathy with synaptic degeneration (84), and decreased synaptophysin-positive presynaptic bouton in the remaining motor neurons (85). Mutation of TDP-43, a DNA/RNA-binding protein which modulates RNA splicing and micro RNA biogenesis (86, 87), has been identified in familial ALS. Transgenic animals of the mutant with human TDP-43 exhibit reduced levels of

Table 3. Summary of presynaptic phenotype by ALS and HD genetic factors

ALS and Huntington's disease		
Factor	Phenotype at presynaptic terminal	Ref
ALS SOD-1	• Axonal transport	83
	• Synaptic degeneration	84,85
TDP-43	• Expression regulation of presynaptic protein	88
	• Attenuation of synaptic transmission	89
FUS	• Active zone formation, synaptic transmission	93
HD HTT	• Synaptic transmission, release probability	94
	• Synaptic vesicle dynamics	95,96




Fig. 1. Summarization of diagram for alteration of presynaptic terminals in various neurodegenerative diseases.

synaptophysin (a presynaptic protein) in the brain, as well as cognitive and motor deficits in behavioral tests (88); also, the synaptic transmission was attenuated (89). FUS (Fused-in-Sarcoma) is also one of the DNA/RNA-binding proteins having a similar structure and functions as TDP-43 (90). The mutation in nuclear localization signal (NLS) of FUS leads to increased cytoplasmic FUS position, which induces the aggregation of FUS mutants, as a pathogenesis of ALS (91, 92). FUS mutations were also linked to synaptic dysfunctions. Overexpression FUS mutant disrupts formation of presynaptic active zones, consequently reducing the synaptic transmission with decreased quantal size (93) (Table 3).

Huntington's disease

Huntington's disease (HD) is an inherited autosomal dominant neurodegenerative disorder. It is mainly caused by mutation of the huntingtin (htt) protein, which has an abnormally high copy of polyglutamine (polyQ) repeat at the N-terminus. General symptoms of HD are motor dysfunction and cognitive deficits, which correlate with the neurodegeneration of specific regions such as the striatum and cerebral cortex. Some of the presynaptic alterations in HD have been reported in various genetic models. An HD model system expressing 128 polyQ expansion in *Drosophila*, revealed that it had significantly higher neurotransmitter release and release probability (94). Presynaptic specific protein alterations are also reported. For example, rabphilin 3A expression level decreased (95); however, levels of SCAMP5, one of the synaptic vesicle proteins was increased (96), suggesting that these alterations of presynaptic protein levels result in an impairment of synaptic vesicle fusion or endocytosis process (Table 3).

CONCLUSION

We here reviewed the structural and functional alterations of presynaptic terminals by genetic factors in several neurodegenerative diseases (Fig. 1). In AD, APP, an original source for A β peptide, is a molecular hub in PAZ. It negatively regulates the nerve terminal formation and readily releasable synaptic vesicle pool. Pathological A β (aggregate A β) strongly inhibits the synaptic vesicle fusion machinery; however, soluble A β increases the release probability. BACE1 and presenilin are also important regulators for presynaptic physiology. In addition, other genetic factors for AD (Tau and ApoE4) are also involved in synaptic stability and synaptic release. In PD, numerous studies for the genetic factors has revealed the implication in presynaptic functions. α -synuclein expression controlled the release probability and recycling pool size, and LRRK2 modulates dopamine release and synaptic vesicle endocytosis by phosphorylating several endocytic proteins (e.g. endophilin). Interestingly, recently accumulating reports show that endocytic proteins (e.g. synaptojanin1, endophilin) are strongly related in PD, indicating that the synaptic vesicle endocytosis process might be an important pathway related in the pathogenesis of PD.

A number of the genetic factors for neurodegenerative diseases are closely related with synaptic function and its alteration. However, most studies just display the phenotype of synaptic dysfunctions, without detailed mechanisms of how the genetic factors result in these dysfunctions. By far, most studies for the pathogenesis of neurodegenerative diseases tend to focus on mechanisms of how neuronal cell death or neurodegeneration occur. Most of the neurodegenerative diseases are generally thought to be chronic diseases.

Ultimately, neurons are likely to die after experiencing a number of abnormal processes during neurodegeneration. Synapses possess high variability and plasticity, and are also highly vulnerable to pathological conditions. It is likely to reveal abnormal phenotype or alteration of the synaptic function at the early onset of neurodegeneration, suggesting that an in depth investigation for synaptic dysfunction may provide a new approach to the understanding of the early pathogenesis of neurodegenerative diseases.

ACKNOWLEDGEMENTS

This work is supported by Basic Science Research Program and Brain Science Research Program through the Nation Research Foundation of Korea (NRF) funded by the Ministry of science, ICT & Future Planning (NRF-2013R1A1A1063174, NRF-2017R1A2B4007019), and a grant from Kyung Hee University in 2016 (KHU-20160702).

CONFLICTS OF INTEREST

The authors have no conflicting financial interests.

REFERENCES

1. Ikin AF, Annaert WG, Takei K et al (1996) Alzheimer amyloid protein precursor is localized in nerve terminal preparations to Rab5-containing vesicular organelles distinct from those implicated in the synaptic vesicle pathway. *J Biol Chem* 271, 31783-31786
2. Groemer TW, Thiel CS, Holt M et al (2011) Amyloid precursor protein is trafficked and secreted via synaptic vesicles. *PLoS One* 6, e18754
3. Priller C, Bauer T, Mitteregger G, Krebs B, Kretschmar HA and Herms J (2006) Synapse formation and function is modulated by the amyloid precursor protein. *J Neurosci* 26, 7212-7221
4. Lassek M, Weingarten J, Wegner M et al (2016) APP Is a Context-Sensitive Regulator of the Hippocampal Presynaptic Active Zone. *PLoS Comput Biol* 12, e1004832
5. Jang BG, In S, Choi B and Kim MJ (2014) Beta-amyloid oligomers induce early loss of presynaptic proteins in primary neurons by caspase-dependent and proteasome-dependent mechanisms. *Neuroreport* 25, 1281-1288
6. Abramov E, Dolev I, Fogel H, Ciccotosto GD, Ruff E and Slutsky I (2009) Amyloid-beta as a positive endogenous regulator of release probability at hippocampal synapses. *Nat Neurosci* 12, 1567-1576
7. Fogel H, Frere S, Segev O et al (2014) APP homodimers transduce an amyloid-beta-mediated increase in release probability at excitatory synapses. *Cell Rep* 7, 1560-1576
8. Romani A, Marchetti C, Bianchi D et al (2013) Computational modeling of the effects of amyloid-beta on release probability at hippocampal synapses. *Front Comput Neurosci* 7, 1
9. Russell CL, Semerdjieva S, Empson RM, Austen BM, Beesley PW and Alifragis P (2012) Amyloid-beta acts as a regulator of neurotransmitter release disrupting the interaction between synaptophysin and VAMP2. *PLoS One* 7, e43201
10. Cirrito JR, Yamada KA, Finn MB et al (2005) Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron* 48, 913-922
11. Cirrito JR, Kang JE, Lee J et al (2008) Endocytosis is required for synaptic activity-dependent release of amyloid-beta in vivo. *Neuron* 58, 42-51
12. Munro KM, Nash A, Pignoni M, Lichtenthaler SF and Gunnarsen JM (2016) Functions of the Alzheimer's Disease Protease BACE1 at the Synapse in the Central Nervous System. *J Mol Neurosci* 60, 305-315
13. Lundgren JL, Ahmed S, Schedin-Weiss S et al (2015) ADAM10 and BACE1 are localized to synaptic vesicles. *J Neurochem* 135, 606-615
14. Del Prete D, Lombino F, Liu X and D'Adamio L (2014) APP is cleaved by Bace1 in pre-synaptic vesicles and establishes a pre-synaptic interactome, via its intracellular domain, with molecular complexes that regulate pre-synaptic vesicles functions. *PLoS One* 9, e108576
15. Petrus E and Lee HK (2014) BACE1 is necessary for experience-dependent homeostatic synaptic plasticity in visual cortex. *Neural Plast* 2014, 128631
16. Frykman S, Hur JY, Franberg J et al (2010) Synaptic and endosomal localization of active gamma-secretase in rat brain. *PLoS One* 5, e8948
17. Zhang C, Wu B, Beglopoulos V et al (2009) Presenilins are essential for regulating neurotransmitter release. *Nature* 460, 632-636
18. Pratt KG, Zimmerman EC, Cook DG and Sullivan JM (2011) Presenilin 1 regulates homeostatic synaptic scaling through Akt signaling. *Nat Neurosci* 14, 1112-1114
19. Spires-Jones TL and Hyman BT (2014) The intersection of amyloid beta and tau at synapses in Alzheimer's disease. *Neuron* 82, 756-771
20. Voelzmann A, Okenve-Ramos P, Qu Y et al (2016) Tau and spectraplakins promote synapse formation and maintenance through Jun kinase and neuronal trafficking. *Elife* 5
21. Jadhav S, Katina S, Kovac A, Kazmerova Z, Novak M and Zilka N (2015) Truncated tau deregulates synaptic markers in rat model for human tauopathy. *Front Cell Neurosci* 9, 24
22. Kopeikina KJ, Polydoro M, Tai HC et al (2013) Synaptic alterations in the rTg4510 mouse model of tauopathy. *J Comp Neurol* 521, 1334-1353
23. Kopeikina KJ, Wegmann S, Pitstick R et al (2013) Tau causes synapse loss without disrupting calcium homeostasis in the rTg4510 model of tauopathy. *PLoS One* 8, e80834
24. Levi O, Jongen-Reelo AL, Feldon J, Roses AD and Michaelson DM (2003) ApoE4 impairs hippocampal plasticity isoform-specifically and blocks the environmental stimulation of synaptogenesis and memory. *Neurobiol Dis* 13, 273-282
25. Zhu Y, Nwabuisi-Heath E, Dumanis SB et al (2012) APOE genotype alters glial activation and loss of synaptic markers in mice. *Glia* 60, 559-569
26. Cambon K, Davies HA and Stewart MG (2000) Synaptic

- loss is accompanied by an increase in synaptic area in the dentate gyrus of aged human apolipoprotein E4 transgenic mice. *Neuroscience* 97, 685-692
27. Dumanis SB, DiBattista AM, Miessau M, Moussa CE and Rebeck GW (2013) APOE genotype affects the presynaptic compartment of glutamatergic nerve terminals. *J Neurochem* 124, 4-14
 28. Bal M, Leitz J, Reese AL et al (2013) Reelin mobilizes a VAMP7-dependent synaptic vesicle pool and selectively augments spontaneous neurotransmission. *Neuron* 80, 934-946
 29. Koffie RM, Hashimoto T, Tai HC et al (2012) Apolipoprotein E4 effects in Alzheimer's disease are mediated by synaptotoxic oligomeric amyloid-beta. *Brain* 135, 2155-2168
 30. Picconi B, Piccoli G and Calabresi P (2012) Synaptic dysfunction in Parkinson's disease. *Adv Exp Med Biol* 970, 553-572
 31. Belluzzi E, Greggio E and Piccoli G (2012) Presynaptic dysfunction in Parkinson's disease: a focus on LRRK2. *Biochem Soc Trans* 40, 1111-1116
 32. Stefanis L (2012) alpha-Synuclein in Parkinson's disease. *Cold Spring Harb Perspect Med* 2, a009399
 33. Goedert M (2001) Alpha-synuclein and neurodegenerative diseases. *Nat Rev Neurosci* 2, 492-501
 34. Norris EH, Giasson BI and Lee VM (2004) Alpha-synuclein: normal function and role in neurodegenerative diseases. *Curr Top Dev Biol* 60, 17-54
 35. Rizo J and Sudhof TC (2012) The membrane fusion enigma: SNAREs, Sec1/Munc18 proteins, and their accomplices—guilty as charged? *Annu Rev Cell Dev Biol* 28, 279-308
 36. Tanji K, Mori F, Mimura J et al (2010) Proteinase K-resistant alpha-synuclein is deposited in presynapses in human Lewy body disease and A53T alpha-synuclein transgenic mice. *Acta Neuropathol* 120, 145-154
 37. Spinelli KJ, Taylor JK, Osterberg VR et al (2014) Presynaptic alpha-synuclein aggregation in a mouse model of Parkinson's disease. *J Neurosci* 34, 2037-2050
 38. Lundblad M, Decressac M, Mattsson B and Bjorklund A (2012) Impaired neurotransmission caused by overexpression of alpha-synuclein in nigral dopamine neurons. *Proc Natl Acad Sci U S A* 109, 3213-3219
 39. Xu J, Wu XS, Sheng J et al (2016) alpha-Synuclein Mutation Inhibits Endocytosis at Mammalian Central Nerve Terminals. *J Neurosci* 36, 4408-4414
 40. Nemani VM, Lu W, Berge V et al (2010) Increased expression of alpha-synuclein reduces neurotransmitter release by inhibiting synaptic vesicle reclustering after endocytosis. *Neuron* 65, 66-79
 41. Scott D and Roy S (2012) alpha-Synuclein inhibits intersynaptic vesicle mobility and maintains recycling-pool homeostasis. *J Neurosci* 32, 10129-10135
 42. Mills RD, Mulhern TD, Liu F, Culvenor JG and Cheng HC (2014) Prediction of the repeat domain structures and impact of parkinsonism-associated variations on structure and function of all functional domains of leucine-rich repeat kinase 2 (LRRK2). *Hum Mutat* 35, 395-412
 43. Martin I, Kim JW, Dawson VL and Dawson TM (2014) LRRK2 pathobiology in Parkinson's disease. *J Neurochem* 131, 554-565
 44. Lee S, Liu HP, Lin WY, Guo H and Lu B (2010) LRRK2 kinase regulates synaptic morphology through distinct substrates at the presynaptic and postsynaptic compartments of the *Drosophila* neuromuscular junction. *J Neurosci* 30, 16959-16969
 45. Matta S, Van Kolen K, da Cunha R et al (2012) LRRK2 controls an EndoA phosphorylation cycle in synaptic endocytosis. *Neuron* 75, 1008-1021
 46. Arranz AM, Delbroek L, Van Kolen K et al (2015) LRRK2 functions in synaptic vesicle endocytosis through a kinase-dependent mechanism. *J Cell Sci* 128, 541-552
 47. Belluzzi E, Gonnelli A, Cimaru MD et al (2016) LRRK2 phosphorylates pre-synaptic N-ethylmaleimide sensitive fusion (NSF) protein enhancing its ATPase activity and SNARE complex disassembling rate. *Mol Neurodegener* 11, 1
 48. Li X, Patel JC, Wang J et al (2010) Enhanced striatal dopamine transmission and motor performance with LRRK2 overexpression in mice is eliminated by familial Parkinson's disease mutation G2019S. *J Neurosci* 30, 1788-1797
 49. Beccano-Kelly DA, Kuhlmann N, Tatarnikov I et al (2014) Synaptic function is modulated by LRRK2 and glutamate release is increased in cortical neurons of G2019S LRRK2 knock-in mice. *Front Cell Neurosci* 8, 301
 50. Beccano-Kelly DA, Volta M, Munsie LN et al (2015) LRRK2 overexpression alters glutamatergic presynaptic plasticity, striatal dopamine tone, postsynaptic signal transduction, motor activity and memory. *Hum Mol Genet* 24, 1336-1349
 51. Leroy E, Anastasopoulos D, Konitsiotis S, Lavedan C and Polymeropoulos MH (1998) Deletions in the Parkin gene and genetic heterogeneity in a Greek family with early onset Parkinson's disease. *Hum Genet* 103, 424-427
 52. Lucking CB, Abbas N, Durr A et al (1998) Homozygous deletions in parkin gene in European and North African families with autosomal recessive juvenile parkinsonism. The European Consortium on Genetic Susceptibility in Parkinson's Disease and the French Parkinson's Disease Genetics Study Group. *Lancet* 352, 1355-1356
 53. Kitada T, Pisani A, Karouani M et al (2009) Impaired dopamine release and synaptic plasticity in the striatum of parkin^{-/-} mice. *J Neurochem* 110, 613-621
 54. Helton TD, Otsuka T, Lee MC, Mu Y and Ehlers MD (2008) Pruning and loss of excitatory synapses by the parkin ubiquitin ligase. *Proc Natl Acad Sci U S A* 105, 19492-19497
 55. Cortese GP, Zhu M, Williams D, Heath S and Waites CL (2016) Parkin Deficiency Reduces Hippocampal Glutamatergic Neurotransmission by Impairing AMPA Receptor Endocytosis. *J Neurosci* 36, 12243-12258
 56. Khandelwal PJ, Dumanis SB, Feng LR et al (2010) Parkinson-related parkin reduces alpha-Synuclein phosphorylation in a gene transfer model. *Mol Neurodegener* 5, 47
 57. Zhang Y, Gao J, Chung KK, Huang H, Dawson VL and Dawson TM (2000) Parkin functions as an E2-dependent ubiquitin- protein ligase and promotes the degradation of the synaptic vesicle-associated protein, CDCrel-1. *Proc*

- Natl Acad Sci U S A 97, 13354-13359
58. Chung KK, Zhang Y, Lim KL et al (2001) Parkin ubiquitinates the alpha-synuclein-interacting protein, synphilin-1: implications for Lewy-body formation in Parkinson disease. *Nat Med* 7, 1144-1150
 59. Valente EM, Abou-Sleiman PM, Caputo V et al (2004) Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 304, 1158-1160
 60. Beilina A, Van Der Brug M, Ahmad R et al (2005) Mutations in PTEN-induced putative kinase 1 associated with recessive parkinsonism have differential effects on protein stability. *Proc Natl Acad Sci U S A* 102, 5703-5708
 61. Plun-Favreau H, Klupsch K, Moiso N et al (2007) The mitochondrial protease HtrA2 is regulated by Parkinson's disease-associated kinase PINK1. *Nat Cell Biol* 9, 1243-1252
 62. Morais VA, Verstreken P, Roethig A et al (2009) Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Mol Med* 1, 99-111
 63. Xi Y, Ryan J, Noble S, Yu M, Yilbas AE and Ekker M (2010) Impaired dopaminergic neuron development and locomotor function in zebrafish with loss of pink1 function. *Eur J Neurosci* 31, 623-633
 64. Kitada T, Pisani A, Porter DR et al (2007) Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci U S A* 104, 11441-11446
 65. Junn E, Jang WH, Zhao X, Jeong BS and Mouradian MM (2009) Mitochondrial localization of DJ-1 leads to enhanced neuroprotection. *J Neurosci Res* 87, 123-129
 66. Bonifati V, Rizzu P, van Baren MJ et al (2003) Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299, 256-259
 67. Usami Y, Hatano T, Imai S et al (2011) DJ-1 associates with synaptic membranes. *Neurobiol Dis* 43, 651-662
 68. Goldberg MS, Pisani A, Haburcak M et al (2005) Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-linked gene DJ-1. *Neuron* 45, 489-496
 69. Slepnev VI and De Camilli P (2000) Accessory factors in clathrin-dependent synaptic vesicle endocytosis. *Nat Rev Neurosci* 1, 161-172
 70. McPherson PS, Garcia EP, Slepnev VI et al (1996) A presynaptic inositol-5-phosphatase. *Nature* 379, 353-357
 71. Mani M, Lee SY, Lucast L et al (2007) The dual phosphatase activity of synaptojanin1 is required for both efficient synaptic vesicle endocytosis and reavailability at nerve terminals. *Neuron* 56, 1004-1018
 72. Krebs CE, Karkheiran S, Powell JC et al (2013) The Sac1 domain of SYNJ1 identified mutated in a family with early-onset progressive Parkinsonism with generalized seizures. *Hum Mutat* 34, 1200-1207
 73. Quadri M, Fang M, Picillo M et al (2013) Mutation in the SYNJ1 gene associated with autosomal recessive, early-onset Parkinsonism. *Hum Mutat* 34, 1208-1215
 74. Hardies K, Cai Y, Jardel C et al (2016) Loss of SYNJ1 dual phosphatase activity leads to early onset refractory seizures and progressive neurological decline. *Brain* 139, 2420-2430
 75. Cao M, Wu Y, Ashrafi G et al (2017) Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs Clathrin Uncoating at Synapses and Triggers Dystrophic Changes in Dopaminergic Axons. *Neuron* 93, 882-896 e885
 76. Cao M, Milosevic I, Giovedi S and De Camilli P (2014) Upregulation of Parkin in endophilin mutant mice. *J Neurosci* 34, 16544-16549
 77. Schuske KR, Richmond JE, Matthies DS et al (2003) Endophilin is required for synaptic vesicle endocytosis by localizing synaptojanin. *Neuron* 40, 749-762
 78. Soukup SF, Kuenen S, Vanhauwaert R et al (2016) A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals. *Neuron* 92, 829-844
 79. Felbecker A, Camu W, Valdmanis PN et al (2010) Four familial ALS pedigrees discordant for two SOD1 mutations: are all SOD1 mutations pathogenic? *J Neurol Neurosurg Psychiatry* 81, 572-577
 80. Robberecht W, Aguirre T, Van den Bosch L, Tilkin P, Cassiman JJ and Matthijs G (1996) D90A heterozygosity in the SOD1 gene is associated with familial and apparently sporadic amyotrophic lateral sclerosis. *Neurology* 47, 1336-1339
 81. Andersen PM (2006) Amyotrophic lateral sclerosis associated with mutations in the CuZn superoxide dismutase gene. *Curr Neurol Neurosci Rep* 6, 37-46
 82. Lee DY, Jeon GS, Shim YM, Seong SY, Lee KW and Sung JJ (2015) Modulation of SOD1 Subcellular Localization by Transfection with Wild- or Mutant-type SOD1 in Primary Neuron and Astrocyte Cultures from ALS Mice. *Exp Neurobiol* 24, 226-234
 83. Bae JR and Kim SH (2016) Impairment of SOD1-G93A motility is linked to mitochondrial movement in axons of hippocampal neurons. *Arch Pharm Res* 39, 1144-1150
 84. Tallon C, Russell KA, Sakhalkar S, Andrapallayal N and Farah MH (2016) Length-dependent axo-terminal degeneration at the neuromuscular synapses of type II muscle in SOD1 mice. *Neuroscience* 312, 179-189
 85. Zang DW, Lopes EC and Cheema SS (2005) Loss of synaptophysin-positive boutons on lumbar motor neurons innervating the medial gastrocnemius muscle of the SOD1G93A G1H transgenic mouse model of ALS. *J Neurosci Res* 79, 694-699
 86. Gregory RI, Yan KP, Amuthan G et al (2004) The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432, 235-240
 87. Xu ZS (2012) Does a loss of TDP-43 function cause neurodegeneration? *Mol Neurodegener* 7, 27
 88. Medina DX, Orr ME and Oddo S (2014) Accumulation of C-terminal fragments of transactive response DNA-binding protein 43 leads to synaptic loss and cognitive deficits in human TDP-43 transgenic mice. *Neurobiol Aging* 35, 79-87
 89. Handley EE, Pitman KA, Dawkins E et al (2016) Synapse Dysfunction of Layer V Pyramidal Neurons Precedes Neurodegeneration in a Mouse Model of TDP-43 Proteinopathies. *Cereb Cortex* 1-18
 90. Nolan M, Talbot K and Anson O (2016) Pathogenesis of FUS-associated ALS and FTD: insights from rodent

- models. *Acta Neuropathol Commun* 4, 99
91. Da Cruz S and Cleveland DW (2011) Understanding the role of TDP-43 and FUS/TLS in ALS and beyond. *Curr Opin Neurobiol* 21, 904-919
 92. Lagier-Tourenne C and Cleveland DW (2009) Rethinking ALS: the FUS about TDP-43. *Cell* 136, 1001-1004
 93. Machamer JB, Collins SE and Lloyd TE (2014) The ALS gene FUS regulates synaptic transmission at the *Drosophila* neuromuscular junction. *Hum Mol Genet* 23, 3810-3822
 94. Romero E, Cha GH, Verstreken P et al (2008) Suppression of neurodegeneration and increased neurotransmission caused by expanded full-length huntingtin accumulating in the cytoplasm. *Neuron* 57, 27-40
 95. Deak F, Shin OH, Tang J et al (2006) Rabphilin regulates SNARE-dependent re-priming of synaptic vesicles for fusion. *EMBO J* 25, 2856-2866
 96. Parker JA, Metzler M, Georgiou J et al (2007) Huntingtin-interacting protein 1 influences worm and mouse presynaptic function and protects *Caenorhabditis elegans* neurons against mutant polyglutamine toxicity. *J Neurosci* 27, 11056-11064