




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Dual priming oligonucleotide (DPO)-based real-time RT-PCR assay for accurate differentiation of four major viruses causing porcine viral diarrhea

Shuo Jia^{a,1}, Baohua Feng^{a,1}, Zhuo Wang^a, Yingying Ma^a, Xuwen Gao^a, Yanping Jiang^a, Wen Cui^a, Xinyuan Qiao^a, Lijie Tang^{a,b}, Yijing Li^{a,b}, Li Wang^{a,b,*}, Yigang Xu^{a,b,**}

^a Heilongjiang Key Laboratory for Animal Disease Control and Pharmaceutical Development, College of Veterinary Medicine, Northeast Agricultural University, Harbin, PR China

^b Northeastern Science Inspection Station, China Ministry of Agriculture Key Laboratory of Animal Pathogen Biology, Harbin, PR China

ARTICLE INFO

Keywords:

Porcine viral diarrhea
Dual priming oligonucleotide (DPO)
Real-time RT-PCR assay
Accurate differentiation

ABSTRACT

Currently in China, porcine epidemic diarrhea virus (PEDV), transmissible gastroenteritis virus (TGEV), porcine rotavirus (PoRV), and porcine deltacoronavirus (PDCoV) are the major causes of porcine viral diarrhea, and mixed infections in clinics are common, resulting in significant economic losses in pig industry. Here, a dual priming oligonucleotide (DPO)-based multiplex real-time SYBR Green RT-PCR assay were developed for accurately differentiating PEDV, TGEV, PoRV, and PDCoV in clinical specimens targeting the N gene of TGEV, PEDV, and PDCoV, and the VP7 gene of PoRV. Results showed that the DPO primer allowed a wider annealing temperature range (40–65 °C) and had a higher priming specificity compared to conventional primer, in which more than 3 nucleotides in the 3'- or 5'-segment of DPO primer mismatched with DNA template, PCR amplification efficiency would decrease substantially or extension would not proceed. DPO-based multiplex real-time RT-PCR method had analytical detection limit of 8.63×10^2 copies/ μ L, 1.92×10^2 copies/ μ L, 1.74×10^2 copies/ μ L, and 1.76×10^2 copies/ μ L for PEDV, TGEV, PoRV, and PDCoV in clinical specimens, respectively. A total of 672 clinical specimens of piglets with diarrheal symptoms were collected in Northeastern China from 2017 to 2018 followed by analysis using the assay, and epidemiological investigation results showed that PEDV, TGEV, PoRV, and PDCoV prevalence was 19.05%, 5.21%, 4.32%, and 3.87%, respectively. The assay developed in this study showed higher detection accuracy than conventional RT-PCR method, suggesting a useful tool for the accurate differentiation of the four major viruses causing porcine viral diarrhea in practice.

1. Introduction

Currently, porcine viral diarrhea is a major cause of high morbidity and high mortality in piglets and an important threat to the development of pig industry worldwide, resulting in significant economic losses [1]. Porcine epidemic diarrhea virus (PEDV), transmissible gastroenteritis virus (TGEV), and porcine rotavirus (PoRV) are the most common viral agents causing piglet diarrhea [2]. Moreover, porcine deltacoronavirus (PDCoV), since first reported in 2009, has become an important cause of newborn piglet diarrhea [3–9]. Sometimes, mixed infections in clinics are frequent. TGEV is a common cause of epidemic and endemic viral enteritis in neonates and senile pigs [10,11]. Like

TGEV, PEDV can cause severe enteritis in newborn piglets, but has higher mortality rate that may reach 100% during epidemics [12,13]. PDCoV, as a newly emerging porcine intestinal coronavirus, can also cause enteritis and severe diarrhea in piglets [9], leading to numerous economic losses for farmers in China. Its infection appears to be clinically milder with lower neonatal mortality rate of 30–40% compared to typical PEDV infection. PoRV is also a common cause of viral diarrhea diseases among newborn piglets [14,15]. Infection with any of these viruses develops into similar clinical symptoms, including severe diarrhea and dehydration, and in some cases, mixed infections of these pathogens are common; therefore, it is difficult to distinguish them clinically [4,16–18]. Thus, it is of great significance to develop a useful

* Corresponding author. Heilongjiang Key Laboratory for Animal Disease Control and Pharmaceutical Development, College of Veterinary Medicine, Northeast Agricultural University, Harbin, PR China.

** Corresponding author. Heilongjiang Key Laboratory for Animal Disease Control and Pharmaceutical Development, College of Veterinary Medicine, Northeast Agricultural University, Harbin, PR China.

E-mail addresses: wanglicau@163.com (L. Wang), yigangxu@neau.edu.cn (Y. Xu).

¹ Shuo Jia and Baohua Feng contributed equally to this work.

<https://doi.org/10.1016/j.mcp.2019.101435>

Received 1 July 2019; Received in revised form 28 July 2019; Accepted 12 August 2019

Available online 12 August 2019

0890-8508/© 2019 Elsevier Ltd. All rights reserved.

Table 1
Primers used in this study.

Viruses	Primer ID	Sequences (5'–3')	Reference	Product size
PEDV	PEDV-N-F ^a	ATGGCTTCTGTCAGCTTTCA	GenBank DQ355221.1	1326 bp
	PEDV-N-R ^a	TTAATTTCTGTATCGAAGATCTCG		
	PEDV-DPO-F ^b	GGTATTGGAGAAAATCCTGACAGGTTTT ^d GCAACAGCA	251 bp	
	PEDV-DPO-R ^b	GACGCATCAACACCTTTTTCGTTTTCCGCATC		
	PEDV-C-F ^c	GGTATTGGAGAAAATCCTGACAGGCATAAGCAACAGCA		251 bp
PEDV-C-R ^c	GACGCATCAACACCTTTTTCGACAAAATCCGCATC			
TGEV	TGEV-N-F ^a	ATGGCCAACCGGGACAA	GenBank KU729220.1	1149 bp
	TGEV-N-R ^a	TTAGTTCGTTACCTCATCAATTATC		
	TGEV-DPO-F ^b	CTGTTCTTGC CGCACTTAAAA ^d GGTGTGTTGAC	180 bp	
	TGEV-DPO-R ^b	TAGCTCCATAAAATCTTGTACATC ^d TTACCTGCAG		
	TGEV-C-F ^c	CTGTTCTTGC CGCACTTAAAAAGTTAGGTGTGTTGAC		180 bp
TGEV-C-R ^c	TAGCTCCATAAAATCTTGTACATCACCTTTACCTGCA			
PoRV	PoRV-VP7-F ^a	ATGTATGGTATTGAATATAACACAG	GenBank JN388691.1	981 bp
	PoRV-VP7-R ^a	CTAGACTCGGTAATAAAAAGGCAG		
	PoRV-DPO-F ^b	TGTTTTTAACAAAAGGATTTTT ^d ACAGGGTCA	318 bp	
	PoRV-DPO-R ^b	CTGTAGTCGAACATCCTA ^d TTAGAGTCTGTC		
	PoRV-C-F ^c	TGTTTTTAACAAAAGGATGGCCAACAGGGTCA		318 bp
PoRV-C-R ^c	CTGTAGTCGAACATCCTATCCCGAGAGTCTGC			
PDCoV	PDCoV-N-F ^a	ATGGCCGACCAAGTAGTCCCTA	GenBank MH715491.1	1029 bp
	PDCoV-N-R ^a	CTACGCTGCTGATTCCTGCTTT		
	PDCoV-DPO-F ^b	AGGCTACTCATCCTCAGTTTCGTGGTTTT ^d GAGTTCGGC	400 bp	
	PDCoV-DPO-R ^b	GGGCCACTTACACGCTCCTGAGGTC ^d TTAGCGTTG		
	PDCoV-C-F ^c	AGGCTACTCATCCTCAGTTTCGTGGCAATGGAGTTCGGC		400 bp
PDCoV-C-R ^c	GGGCCACTTACACGCTCCTGAGGTTCTCTAGCGTTG			

^a Primers were used to amplify the full-length of the target gene for positive standard control.

^b DPO primers were used to establish detection method.

^c Conventional primers were used as control for comparison with DPO primers.

^d "T" means deoxyinosine.

method for rapid and accurate differentiation of these viruses causing viral diarrhea in piglets.

Although virus isolation, immunofluorescence assay, and enzyme-linked immunosorbent assays are standard diagnostic methods for viruses [19], these techniques are time-consuming, relatively low in specificity and sensitivity, and therefore are not suitable for early diagnosis or epidemiological investigation of enterovirus-associated diseases. Currently, polymerase chain reaction (PCR)-based methods have been proven convenient and highly sensitive for detecting porcine diarrhea-associated viruses [20,21]. Of them real-time SYBR Green PCR method that combines conventional PCR and fluorescent signal detection with advantages of high sensitivity, good repeatability, and intuitive results is widely used for epidemiological investigation and pathogen identification [22–24]. Moreover, the closed-tube detection of SYBR Green-based real-time PCR assay can effectively prevent aerosol pollution and reduce the incidence of false positives. However, the specificity of SYBR Green fluorescent dye is not so good that the melting curve of PCR product has to be used to confirm the amplification specificity. Therefore, primer design is the key to determining the detection specificity of SYBR Green-based real-time PCR method. However, the fundamental solution for eliminating false positive amplification still remains a challenge, which requires repeated optimization of primer parameters to achieve a high specificity, including primer specificity, primer length, melting temperature, GC content, less secondary structure, and PCR annealing temperature, in particular the development of a multiplex PCR assay.

Dual priming oligonucleotide (DPO) system, with a long 5'-segment, a short 3'-segment, and polydeoxyinosine (poly I) linker bridging 5'- and 3'-segments, is an approach to designing PCR primer first reported by Chun et al. [25], which can effectively eliminate non-specific priming without disrupting amplification of target sequences, showing higher specificity than conventional primers [25–28]. Moreover, compared with conventional primer, the design process of DPO system is simpler and does not require repeated optimization of primer parameters. The position of 3'-segment is firstly determined at a site containing 6–12 bases with 40–80% GC content, then five deoxyinosines are designated

for the poly I linker, and then the sequence upstream of 3'-segment is automatically extended to 18–25 bases until $T_m > 65^\circ\text{C}$ to generate the 5'-segment [25]. Notably, DPO system allows a wide annealing temperature to effectively amplify target genes, indicating a promising tool for developing multiplex PCR assay. Several publications reported the successful development of DPO-based multiplex PCR assay for simultaneous detection of multiple viral pathogens [26,27,29,30].

In this study, combining the strong specificity of DPO primers and high sensitivity of SYBR Green fluorescent dye, a DPO system-based multiplex real-time RT-PCR assay for the accurate differentiation of PEDV, TGEV, PoRV, and PDCoV was developed, which would provide technical support for identifying these viral pathogens causing diarrhea in piglet, and for improvement in epidemiological investigation.

2. Materials and methods

2.1. Viruses, primers, and clinical specimens

Viral strains including PEDV, TGEV, PoRV, PDCoV, bovine viral diarrhea virus (BVDV), bovine parvovirus (BPV), infectious bursal disease virus (IBDV), infectious hematopoietic necrosis virus (IHNV), bovine rotavirus (BRV), feline infectious peritonitis virus (FIPV), and bovine respiratory syncytial virus (BRSV) were used for specificity analysis in this study. A total of 672 clinical specimens of newborn piglets with diarrhea symptoms were collected from pig farms located in Northeastern China (Heilongjiang, Jilin, and Liaoning provinces) from 2017 to 2018, and were subjected to detection by the assay developed in this study. N gene of PEDV, TGEV, and PDCoV, and the VP7 gene of PoRV were used as target genes to design the primers using Oligo 6.0 software, and the details of primers are shown in Table 1.

2.2. Preparation of plasmid standards

Total RNA of PEDV, TGEV, PoRV, and PDCoV was extracted respectively from cell cultures using the TRIzol® Plus RNA Purification Kit (Invitrogen, USA) [31], followed by reverse transcription using the




Fig. 1. Comparison of priming specificity between DPO primer and conventional primer. A: Point mutations targeting gene sequences of the 5'- or 3'-segments of DPO primer (3'-segment with three/five sites mutation namely N3/N5; 5'-segment with three/five sites mutation namely SN3/SN5), followed by PCR amplification using conventional PCR assay and SYBR Green-based real-time PCR assay with conventional primer (CP) and DPO primer, respectively. B: PCR products were analyzed by agarose gel electrophoresis. (Left) M: DNA Marker DL2000; 1: N3+CP; 2: N3+DPO; 3: SN3+CP; 4: SN3+DPO; 5: N gene + CP; 6: N gene + DPO; 7: Negative control. (Right) M: DNA Marker DL2000; 1: N5+CP; 2: N5+DPO; 3: SN5+CP; 4: SN5+DPO; 5: N gene + CP; 6: N gene + DPO; 7: Negative control. C: SYBR Green fluorescence detection results. Detection result of three nucleotide mismatches (a/c: amplification plot; b/d: melt curve). 1: N3/SN3+conventional primer; 2: N3/SN3+DPO. Detection result of five nucleotide mismatches (e/g: amplification plot; f/h: melt curve). 1: N5/SN5+CP; 2: N5/SN5+DPO.

Superscript Reverse Transcriptase Reagent Kit (Takara, Japan) according to the manufacturer's instructions, generating complementary DNA (cDNA). Next, the full-length of target genes were amplified with primers (as shown in Table 1) using the corresponding cDNA as template. PCR system in a total volume of 25 μ L was as follows: 10 \times exTaq Buffer 2.5 μ L, exTaq DNA polymerase 0.5 μ L, dNTP mixture (10 mM each) 3 μ L, cDNA 5 μ L, primers (TGEV-N-F/R, PEDV-N-F/R, PoRV-VP7-F/R, or PDCoV-N-F/R) (10 μ M) 1 μ L, and ddH₂O 12 μ L. PCR amplification was carried out using a SensoQuest LabCycler (SensoQuest, Germany) with the conditions that 95 $^{\circ}$ C for 5 min, followed by 35 cycles at 95 $^{\circ}$ C for 30 s, 55 $^{\circ}$ C for 30 s, 72 $^{\circ}$ C for 45 s, and a final extension at 72 $^{\circ}$ C for 10 min. Then, PCR product was cloned into plasmid pMD-19T [32], giving rise to recombinant plasmids pMD-

TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N, respectively.

2.3. Specificity and annealing temperature sensitivity analysis of the DPO primer

We used the N gene of TGEV as a model to evaluate the specificity of DPO primer. As shown in Fig. 1-A, the gene sequences targeting the 5'- or 3'-segments of DPO primer were subjected to point mutation at three sites (3'-segment with three site mutations, namely N3; 5'-segment with three site mutations, namely SN3) or five sites (3'-segment with five site mutations, namely N5; 5'-segment with five site mutations namely SN5). Also, the gene sequences (PCR amplification region) with point




Fig. 2. Comparison of effective annealing temperature between conventional primer and DPO primer (A) and specific PCR product of target gene of each virus (B). M: DNA Marker; numbers 1–8: Annealing temperature at 40 °C, 43.2 °C, 48.2 °C, 51 °C, 54 °C, 56.8 °C, 61.8 °C, and 65 °C, respectively; NC: negative control.

mutations were synthesized, followed by detection using conventional PCR and SYBR Green-based real-time PCR with the conventional primer TGEV-C-F/R and DPO primer TGEV-DPO-F/R, respectively. For annealing temperature sensitivity analysis of DPO primer, a multiplex PCR assay was carried out using the mix of plasmids pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N as template with the primers TGEV-DPO-F/R, PEDV-DPO-F/R, PoRV-DPO-F/R, and PDCoV-DPO-F/R at annealing temperature from 40 °C to 65 °C to observe the simultaneous amplification effect of each target gene. In parallel, conventional PCR primers were used as control.

2.4. Development of DPO-based real-time RT-PCR assay

First, DPO-based singleplex real-time RT-PCR assay for detecting PEDV, TGEV, PoRV, and PDCoV was developed, respectively. PCR system in a total volume of 20 μ L was as follows: 10 μ L of SYBR Green I Universal Master Mix, 0.5 μ L of DPO primer pair (10 μ M), 5 μ L/1 μ L of cDNA/plasmid DNA template, and DEPC H₂O added to make up the 20 μ L volume. PCR conditions were 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s, and 60 °C for 1 min performed in an Applied Biosystems 7500 (ABI, USA). Second, the standard curve for viral quantification was constructed using 10-fold serially diluted plasmids pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N with primary concentration of 1.52×10^8 copies/ μ L, 1.64×10^8 copies/ μ L, 2.27×10^8 copies/ μ L, and 1.58×10^8 copies/ μ L, respectively. Subsequently, DPO-based multiplex (duplex, triplex, and quadruplex) RT-PCR assays were developed for detecting different combinations of PEDV, TGEV, PoRV, and PDCoV, following optimization of the reaction system based on single real-time RT-PCR assay. And, the empty plasmid pMD-19T without target genes was used as an inhibition control for monitoring PCR inhibition in this real-time PCR assay.

2.5. Sensitivity, specificity and repeatability analysis of DPO-based real-time RT-PCR assay

To evaluate the limit of detection of the DPO-based real-time RT-PCR assay for PEDV, TGEV, PoRV, and PDCoV in clinical specimens, known PEDV-, TGEV-, PoRV-, or PDCoV-positive feces sample was subjected to 10-fold serial dilution with PBS buffer followed by the extraction of total RNA from the diluted sample. The DPO-based real-time RT-PCR assay was performed to detect the presence of viral nucleic acids, and the standard curve equation for each virus was used to calculate copy number of viral nucleic acids in each diluted feces sample. For the evaluation of detection specificity, PEDV, TGEV, PoRV, PDCoV, BVDV, BRV, BPV, IBDV, BRSV, IHNV, and FIPV were subjected to detection using DPO-based real-time RT-PCR assay developed in this study, and the plasmid standard pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N as the positive control, respectively. To assess the repeatability of detection results, plasmids pMD-TGEV-N (1.52×10^8 copies/ μ L), pMD-PEDV-N (1.64×10^8 copies/ μ L), pMD-PoRV-VP7 (2.27×10^8 copies/ μ L), and pMD-PDCoV-N (1.58×10^8

copies/ μ L) were 10-fold serially diluted, and 10^1 , 10^3 , and 10^5 fold diluted plasmids were used to evaluate the intra-assay repeatability and the inter-assay repeatability.

2.6. Application of DPO-based real-time RT-PCR assay

To evaluate the practicability of DPO-based real-time RT-PCR assay developed in this study, 672 piglet diarrhea samples were collected in Heilongjiang, Jilin, and Liaoning provinces located in Northeastern China from 2017 to 2018, followed by the extraction of total RNA using the TRIzol method as described above, and the assay was carried out. The results of analysis were determined in accordance with the following rules: (i) the cycle threshold (Ct) value of samples detected by the assay less than or equal to 35 was considered virus positive; (ii) the Ct value more than 40 was considered virus negative; (iii) the Ct value between 35 and 40 should be evaluated repeatedly. If Ct value was more than 40, the result was considered virus negative; otherwise, it was positive. In parallel, the detection results of RT-PCR assay with the conventional primers were used as method control.

3. Results

3.1. Priming specificity analysis of the DPO system

As shown in Fig. 1, we used the N gene of TGEV with three or five point mutations targeting the gene sequences designed for 5'- or 3'-segments of DPO primer as model (Fig. 1A) to evaluate the specificity of the DPO primer. In parallel, conventional PCR primer, designed from the same gene sequences as for DPO primer, was used as control. Following amplification, PCR products were respectively analyzed by agarose gel electrophoresis assay (Fig. 1B) and SYBR Green fluorescence assay (Fig. 1C). Our results showed that DPO primers had a higher priming specificity than that of conventional primer; moreover, when there were more than three nucleotides mismatched between the 3'- or 5'-segment of DPO primer and template, PCR amplification would be inhibited substantially and even the extension would be terminated, while the amplification efficiency of conventional PCR primers would not be affected.

3.2. Annealing temperature sensitivity analysis of the DPO system

To analyze the annealing temperature sensitivity of the DPO primer, the mixture of plasmids pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N was used as the template, and multiplex PCR was performed with DPO primers at the annealing temperature range from 40 °C to 65 °C, with conventional primers as control. As shown in Fig. 2, DPO primers allowed a wide range of annealing temperatures to efficiently amplify target genes with a nearly identical pattern, while conventional primers required a relatively rigid annealing temperature.




Fig. 3. Establishment of standard curves of DPO-based real-time PCR assay for TGEV, PEDV, PoRV, and PDCoV. A, B, and C for TGEV, and corresponding standard curve equation is $Y_{TGEV} = -3.7x + 36.14$, $R^2 = 0.992$; D, E, and F for PEDV, and corresponding standard curve equation is $Y_{PEDV} = -3.593x + 45.034$, $R^2 = 0.999$; G, H and I for PoRV, and corresponding standard curve equation is $Y_{PoRV} = -3.42x + 37.348$, $R^2 = 0.963$; J, K and L for PDCoV, and corresponding standard curve equation is $Y_{PDCoV} = -3.248x + 36.825$, $R^2 = 0.999$.

3.3. Standard curves

As shown in Fig. 3, using pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N, corresponding standard curves for viral quantification of TGEV, PEDV, PoRV, and PDCoV were established, respectively. T_m value (melting point) of the melting curve for TGEV, PEDV, PoRV, and PDCoV was 73.59 °C, 79.55 °C, 71.36 °C, and

75.28 °C, respectively.

3.4. Sensitivity, specificity and repeatability analysis of DPO-based real-time PCR assay

We used 10-fold serially diluted PEDV-, TGEV-, PoRV-, or PDCoV-positive feces sample to evaluate the limit of detection (LOD) of DPO-




Fig. 4. Limit of detection and specificity of DPO-based real-time PCR assay for TGEV, PEDV, PoRV, and PDCoV. (A) Limit of detection. 10-fold serially diluted PEDV-, TGEV-, PoRV-, or PDCoV-positive feces sample was used to evaluate the LOD of DPO-based real-time RT-PCR assay in clinical specimens, followed by calculation of the copy number of viral nucleic acids with standard curve equation established in this study. a: Amplification plot with 1–8: 10^1 , 10^2 , 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , and 10^8 dilution of feces sample; b: Melt curve. (B) Specificity result. a: Amplification plot; b: Melt curve. 1: target viruses; 2–12: other viruses and negative control.

based real-time RT-PCR assay for PEDV, TGEV, PoRV, and PDCoV in clinical specimens. This was followed by calculating the copy number of viral nucleic acids with corresponding standard curve equation established in this study. Our results showed that the LOD of DPO-based real-time RT-PCR assay for detecting PEDV, TGEV, PoRV, and PDCoV in clinical specimens was 8.63×10^2 copies/ μL , 1.92×10^2 copies/ μL , 1.74×10^2 copies/ μL , and 1.76×10^2 copies/ μL , respectively (Fig. 4A). In this work, the detection specificity of the assay for PEDV, TGEV, PoRV, and PDCoV was evaluated using TGEV, PEDV, PoRV, PDCoV, BVDV, BRV, IBPV, IBDV, BRSV, IHNV and FIPV, and our results showed that only target viruses were positive for the assay without non-specific priming, while other viruses were negative, indicating a strong specificity (Fig. 4B). Moreover, the intra-assay and inter-assay detection reproducibility was estimated using the plasmid standards pMD-TGEV-N, pMD-PEDV-N, pMD-PoRV-VP7, and pMD-PDCoV-N as templates,

and results showed that the coefficients of variation were all less than 1.0% (Table 2), indicating good repeatability.

3.5. Development of DPO-based multiplex real-time RT-PCR assay

In this work, DPO-based multiplex (duplex, triplex, and quadruplex) real-time RT-PCR assays were developed for the detection of different combinations of PEDV, TGEV, PoRV, and PDCoV, following the optimization of reaction system based on the single real-time RT-PCR assay. As shown in Fig. 5, using DPO-based multiplex real-time RT-PCR assays developed in this study, different target viral pathogens were accurately differentiated in a single reaction.

Table 2
Repeatability of DPO-based real-time RT-PCR assay.

Standards	Concentration (copies/ μ L)	n	Intra-assay variability		Inter-assay variability	
			X \pm SD (Ct)	CV (%)	X \pm SD (Ct)	CV (%)
pMD-TGEV-N	1.52×10^7	5	13.28 \pm 0.08	0.60	12.83 \pm 0.12	0.83
	1.52×10^5	5	21.43 \pm 0.13	0.22	21.19 \pm 0.13	0.52
	1.52×10^3	5	26.84 \pm 0.09	0.36	28.19 \pm 0.19	0.67
pMD-PEDV-N	1.64×10^7	5	14.13 \pm 0.09	0.63	13.73 \pm 0.15	0.83
	1.64×10^5	5	19.51 \pm 0.11	0.56	20.26 \pm 0.14	0.69
	1.64×10^3	5	26.16 \pm 0.08	0.31	27.54 \pm 0.16	0.58
pMD-PoRV-VP7	2.27×10^7	5	14.97 \pm 0.07	0.46	13.51 \pm 0.22	0.72
	2.27×10^5	5	19.23 \pm 0.12	0.62	20.79 \pm 0.29	0.89
	2.27×10^3	5	25.81 \pm 0.10	0.38	28.11 \pm 0.17	0.60
pMD-PDCoV-N	1.58×10^7	5	15.38 \pm 0.19	0.93	14.48 \pm 0.12	0.83
	1.58×10^5	5	19.96 \pm 0.23	0.85	20.76 \pm 0.25	0.97
	1.58×10^3	5	26.01 \pm 0.16	0.62	26.71 \pm 0.28	0.95




Fig. 5. DPO-based duplex, triplex, and quadruplex real-time RT-PCR assays.

3.6. Detection of clinical samples

A total of 672 piglet diarrhea samples collected from Heilongjiang, Jilin, and Liaoning provinces in Northeastern China from 2017 to 2018 were used to evaluate the feasibility of the assay developed in this study, using conventional RT-PCR assay as control. As shown in Table 3, the prevalence rate of PEDV, TGEV, PoRV, and PDCoV detected by the DPO-based real-time RT-PCR assay was 19.05% (128/672), 5.21% (35/672), 4.32% (29/672), and 3.87% (26/672), respectively, while that detected by conventional RT-PCR assay was 17.56% (118/

672), 4.46% (30/672), 3.87% (26/672), and 3.57% (24/672), respectively. The positive coincidence rate of these two detection methods for PEDV, TGEV, PoRV, and PDCoV was 92.18%, 85.71%, 89.66%, and 92.31%. We also detected mixed virus infections present in some samples. With the results of DPO-based real-time RT-PCR in terms of Ct value as shown in Fig. 6, it was clear that the quantities of the viruses detected in different virus-positive samples varied greatly with Ct values of 20–25, 25–30, and 30–35, respectively. Moreover, from the virus-positive results of the clinical samples, PEDV is the major cause of piglet viral diarrhea disease. Subsequently, to confirm the detection

Table 3
Detection results of clinical samples.

Year	n	Conventional RT-PCR assay				Prevalence rate TGEV/ PEDV/PoRV/PDCoV	DPO-based real-time RT-PCR assay				Prevalence rate TGEV/PEDV/PoRV/ PDCoV
		TGEV +/-	PEDV +/-	PoRV +/-	PDCoV +/-		TGEV +/-	PEDV +/-	PoRV +/-	PDCoV +/-	
2017	376	25/351	64/312	15/361	10/366	6.65%/17.02%/3.99%/ 2.66%	29/347	71/305	17/359	11/365	7.71%/18.88%/ 4.52%/2.93%
2018	296	5/291	54/242	11/285	14/282	1.69%/18.24%/3.72%/ 4.73%	6/290	57/239	12/284	15/281	2.03%/19.26%/ 4.05%/5.07%
Total	672	30/642	118/554	26/646	24/648	4.46%/17.56%/3.87%/ 3.57%	35/637	128/544	29/643	26/646	5.21%/19.05%/ 4.32%/3.87%
Coincidence rates ^a between conventional RT-PCR assay and DPO-based real-time RT-PCR assay						TGEV 85.71%	PEDV 92.19%	PoRV 89.66%	PDCoV 92.31%		

^a Coincidence rate was the positive results detected by conventional RT-PCR/the positive results detected by DPO-based real-time RT-PCR.




Fig. 6. Virus-positive results of the clinical samples and the rate of PEDV, TGEV, PoRV, and PDCoV in the positive samples.

accuracy of the assay developed in this study, the samples with different detection results by the two methods were subjected to sequence analysis of target genes. As shown in Fig. 7, the sequencing results verified that the detection results of DPO-based real-time RT-PCR assay were accurate.

4. Discussion

Porcine viral diarrhea disease still seriously endangers the development of the pig industry, and leads to significant economic losses for pig farmers worldwide. Clinically, PEDV, TGEV, PoRV, and PDCoV are the major causative agents of viral diarrhea in piglets. In some cases, mixed infections with two or more these viruses are common, which seriously interfere with clinical diagnosis [8,16,18,33,34]. Moreover, with the increasing growth of international trades and animal transports, the risk of transboundary spread of porcine viral diarrhea disease has also increased significantly. Therefore, to differentiate accurately infections of TGEV, PEDV, PoRV, and PDCoV in clinical specimens, and to prevent transboundary spread of porcine viral diarrhea disease, it is

necessary to develop a rapid, accurate, and multi-target diagnostic method for the differentiation of TGEV, PEDV, PoRV, and PDCoV.

Currently, PCR-based methods are commonly recognized as useful tools for the molecular diagnosis of various viral or bacterial pathogens. The most common PCR method used is the real-time PCR method, which includes probe assay such as Taqman-probe and fluorescence dye assay such as SYBR Green dye. Both assays have their advantages and disadvantages. Of them both Taqman-probe and SYBR Green fluorescence dye have high sensitivity, and TaqMan-probe has high specificity but poor versatility, which is only suitable for specific gene detection, while SYBR Green fluorescence dye has strong versatility. Moreover, the cost of TaqMan-probe is higher than that of SYBR Green fluorescence dye. Therefore, in order to reduce the cost of detection, we utilized SYBR Green fluorescence dye to develop the real-time PCR assay. Primer design is the key to determine the specificity of both probe- and fluorescence dye-based real-time PCR method, which requires very rigid primer parameters. Generally, to obtain reliable primers, there should be repeated optimization of primer parameters, such as specificity, length, melting temperature, GC content, less secondary




Fig. 7. Confirmation of detection results of DPO-based real-time RT-PCR assay by sequencing.

structure, and PCR annealing temperature, which is time-consuming, in particular the design of primers for developing a multiplex PCR assay. In this study, we introduced DPO primer that was different from the conventional primer to develop real-time RT-PCR assay with SYBR Green fluorescence dye for the detection of TGEV, PEDV, PoRV, and PDCoV. Compared with the conventional primer, DPO primer is easier because it does not require repeated optimization of primer parameters.

Moreover, DPO primer has 2 obvious advantages, compared with conventional primer. Firstly, the priming specificity of DPO primer is stronger than that of conventional primer. We used the TGEV N gene with point mutations targeting the gene sequences designed for the 3'- or 5'-segment of DPO primer as model to evaluate the priming specificity of the DPO primer. It was found that when five nucleotides mismatched with the template in the 5'-segment of the DPO primer, the PCR amplification efficiency decreased substantially, while the PCR products were still observed via agarose gel electrophoresis assay and SYBR Green fluorescence assay. Additionally, when five nucleotides mismatched with the template in the 3'-segment of DPO primer, the PCR amplification terminated and could not be analyzed by agarose gel electrophoresis assay, whereas a very weak fluorescence signal with Ct value of approximately 33 was observed via SYBR Green fluorescence assay. Our results were somewhat inconsistent with previous report [25], in which the researchers claimed that if there were 3 bp mismatched in the 3'-segment or 5'-segment of the DPO primer, the extension would not proceed. In any case, the priming specificity of DPO primer is stronger than that of conventional primers, which have a

single priming region, and extension might proceed even in the presence of mismatch between primer and template. Moreover, the high specificity of DPO primers could effectively reduce the competitive effect between different primers, promoting the amplification efficiency indirectly, and although the optimal PCR product size generally recommended for SYBR Green-based protocols was below 200 nucleotides, there was little effect on the amplification efficiency for TGEV (180 bp), PEDV (251 bp), PoRV (318 bp), and PDCoV (400 bp) designed in this work, showing 94.551%, 101.283%, 95.347%, and 102.305%, respectively. Secondly, due to the specific structure of DPO primer, there is wide range of annealing temperature to effectively amplify the target gene with a nearly identical pattern. However, the conventional primers required optimal annealing temperature to amplify effectively, suggesting that the DPO primer was a powerful tool to develop multiplex PCR assay.

In this study, we combined the DPO primer and SYBR Green fluorescence dye to develop the DPO-based real-time RT-PCR assay for clinically differentiating the infection of TGEV, PEDV, PoRV, and PDCoV, which showed high specificity, sensitivity, and detection repeatability. Generally, to demonstrate the new detection method, it is necessary to test the clinical samples and compare the results with that of control method. For this purpose, we collected 672 diarrhea samples of piglet from pig farms located in Northeastern China from 2017 to 2018 to evaluate the diagnostic capability of the assay developed in this study, using conventional RT-PCR assay as control. By contrast, the detection accuracy of DPO-based real-time RT-PCR assay is higher than

that of conventional RT-PCR assay, indicating a better diagnostic capability in clinical samples. Moreover, from the epidemiological investigation, we know that PEDV still is the major cause of piglet viral diarrhea disease in China, and PDCoV as a novel animal coronaviruses that emerged in China [35], has its infection on the rise.

In conclusion, DPO-based real-time RT-PCR assay for differentiating TGEV, PEDV, PoRV, and PDCoV developed in this study has both high specificity of the DPO primer and high sensitivity of SYBR Green fluorescence dye, and shows good diagnostic capability in practice, providing a useful tool for the rapid and accurate diagnosis and epidemiological investigation of these viral pathogens causing piglet diarrhea. Moreover, since SYBR Green-based PCR protocols rely on complex melting curves, they are no more sufficient for human diagnostics laboratories. Therefore, there protocols need further development to an internal probe-based detection system using different colors in order to comply with current diagnostic standards, even if this test is not being applied for humans.

Funding

This work was funded by the National Key Research and Development (R&D) Program of China (grant number 2016YFD0501003, 2016YFD0500704).

Conflicts of interest

The authors declare that they have no conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.mcp.2019.101435>.

References

- [1] J. Zhao, B.J. Shi, X.G. Huang, M.Y. Peng, X.M. Zhang, D.N. He, R. Pang, B. Zhou, P.Y. Chen, A multiplex RT-PCR assay for rapid and differential diagnosis of four porcine diarrhea associated viruses in field samples from pig farms in East China from 2010 to 2012, *J. Virol. Methods* 194 (2013) 107–112.
- [2] L. Rodak, L. Valicek, B. Smid, Z. Nevorankova, An ELISA optimized for porcine epidemic diarrhoea virus detection in faeces, *Vet. Microbiol.* 105 (2005) 9–17.
- [3] L. Wang, B. Byrum, Y. Zhang, Detection and genetic characterization of deltacoronavirus in pigs, Ohio, USA, 2014, *Emerg. Infect. Dis.* 20 (2014) 1227–1230.
- [4] Y. Ma, Y. Zhang, X. Liang, F. Lou, M. Oglesbee, S. Krakowka, J. Li, Origin, evolution, and virulence of porcine deltacoronaviruses in the United States, *mBio* 6 (2015) e00064-15.
- [5] J. Zhang, Porcine deltacoronavirus: overview of infection dynamics, diagnostic methods, prevalence and genetic evolution, *Virus Res.* 226 (2016) 71–84.
- [6] G. Jang, K.K. Lee, S.H. Kim, C. Lee, Prevalence, complete genome sequencing and phylogenetic analysis of porcine deltacoronavirus in South Korea, 2014–2016, *Transbound. Emerg. Dis.* 64 (2017) 1364–1370.
- [7] T. Ajayi, R. Dara, M. Misener, T. Pasma, L. Moser, Z. Poljak, Herd-level prevalence and incidence of porcine epidemic diarrhoea virus (PEDV) and porcine deltacoronavirus (PDCoV) in swine herds in Ontario, Canada, *Transbound. Emerg. Dis.* 65 (2018) 1197–1207.
- [8] M.C. Niederwerder, R.A. Hesse, Swine enteric coronavirus disease: a review of 4 years with porcine epidemic diarrhoea virus and porcine deltacoronavirus in the United States and Canada, *Transbound. Emerg. Dis.* 65 (2018) 660–675.
- [9] H. Zhang, Q. Liang, B. Li, X. Cui, X. Wei, Q. Ding, Y. Wang, H. Hu, Prevalence, phylogenetic and evolutionary analysis of porcine deltacoronavirus in Henan province, China, *Prev. Vet. Med.* 166 (2019) 8–15.
- [10] L.J. Saif, J.L. van Cott, T.A. Brim, Immunity to transmissible gastroenteritis virus and porcine respiratory coronavirus infections in swine, *Vet. Immunol. Immunopathol.* 43 (1994) 89–97.
- [11] M. Yu, L. Wang, S. Ma, X. Wang, Y. Wang, Y. Xiao, Y. Jiang, X. Qiao, L. Tang, Y. Xu, Y. Li, Immunogenicity of eGFP-marked recombinant *Lactobacillus casei* against transmissible gastroenteritis virus and porcine epidemic diarrhoea virus, *Viruses* 9 (2017) 274.
- [12] R.Q. Sun, R.J. Cai, Y.Q. Chen, P.S. Liang, D.K. Chen, C.X. Song, Outbreak of porcine epidemic diarrhoea in suckling piglets, China, *Emerg. Inf. Disp.* 18 (2012) 161–163.
- [13] W. Shi, S. Jia, H. Zhao, J. Yin, X. Wang, M. Yu, S. Ma, Y. Wu, Y. Chen, W. Fan, Y. Xu, Y. Li, Novel approach for isolation and identification of porcine epidemic diarrhoea virus (PEDV) strain NJ using porcine intestinal epithelial cells, *Viruses* 9 (2017) 19.
- [14] A.N. Vlasova, J.O. Amimo, L.J. Saif, Porcine rotaviruses: epidemiology, immune responses and control strategies, *Viruses* 9 (2017) 48.
- [15] X. Dou, Y. Li, J. Han, D.S. Zarlenga, W. Zhu, X. Ren, N. Dong, X. Li, G. Li, Cholesterol of lipid rafts is a key determinant for entry and post-entry control of porcine rotavirus infection, *BMC Vet. Res.* 14 (2018) 45.
- [16] K. Jung, H. Hu, B. Eyerly, Z. Lu, J. Chepngeno, L.J. Saif, Pathogenicity of 2 porcine deltacoronavirus strains in gnotobiotic pigs, *Emerg. Infect. Dis.* 21 (2015) 650–654.
- [17] S. Mesonero-Escuredo, K. Strutzberg-Minder, C. Casanovas, J. Segales, Viral and bacterial investigations on the aetiology of recurrent pig neonatal diarrhoea cases in Spain, *Porc. Heal. Manag.* 4 (2018) 5.
- [18] S. Tuantap, C. Phupolphan, S. Luengyosuechakul, A. Duangin, A. Theamboonlers, S. Wattanaphansak, S. Vongpunsawad, A. Amonsin, Y. Poovorawan, Porcine rotavirus C in pigs with gastroenteritis on Thai swine farms, 2011–2016, *PeerJ* 6 (2018) e4724.
- [19] S. Belak, The molecular diagnosis of porcine viral diseases: a review, *Acta Vet. Hung.* 53 (2005) 113–124.
- [20] A.N. Ben Salem, A.C. Sergei, P.B. Olga, G.A. Olga, A. Mahjoub, B.P. Larissa, Multiplex nested RT-PCR for the detection of porcine enteric viruses, *J. Virol. Methods* 165 (2010) 283–293.
- [21] G. Liu, Y. Jiang, T. Opriessnig, K. Gu, H. Zhang, Z. Yang, Detection and differentiation of five diarrhoea related pig viruses utilizing a multiplex PCR assay, *J. Virol. Methods* 263 (2019) 32–37.
- [22] S. Escutenaire, N. Mohamed, M. Isaksson, P. Thoren, B. Klingeborn, S. Belak, M. Berg, J. Blomberg, SYBR Green real-time reverse transcription-polymerase chain reaction assay for the generic detection of coronaviruses, *Arch. Virol.* 152 (2007) 41–58.
- [23] Y.H. Chun, Y.J. Jeong, S.I. Park, M. Hosmillo, D.J. Shin, H.J. Kwon, S.Y. Kang, S.K. Woo, M.I. Kang, S.J. Park, K.O. Cho, Development of one-step real-time reverse transcription polymerase chain reaction assays for rapid detection of porcine group C rotaviruses, *J. Vet. Diagn. Investig.* 22 (2010) 74–77.
- [24] H.Y. Han, H.H. Zheng, Y. Zhao, R.B. Tian, P.L. Xu, H.L. Hou, H.Y. Chen, M.F. Yang, Development of a SYBR green I-based duplex real-time fluorescence quantitative PCR assay for the simultaneous detection of porcine epidemic diarrhoea virus and porcine circovirus 3, *Mol. Cell. Probes* 44 (2019) 44–50.
- [25] J.Y. Chun, K.J. Kim, I.T. Hwang, Y.J. Kim, D.H. Lee, I.K. Lee, J.K. Kim, Dual priming oligonucleotide system for the multiplex detection of respiratory viruses and SNP genotyping of CYP2C19 gene, *Nucleic Acids Res.* 35 (2007) e40.
- [26] C.S. Lee, B.K. Kang, D.H. Lee, S.H. Lyou, B.K. Park, S.K. Ann, K. Jung, D.S. Song, One-step multiplex RT-PCR for detection and subtyping of swine influenza H1, H3, N1, N2 viruses in clinical samples using a dual priming oligonucleotide (DPO) system, *J. Virol. Methods* 151 (2008) 30–34.
- [27] S.R. Kim, C.S. Ki, N.Y. Lee, Rapid detection and identification of 12 respiratory viruses using a dual priming oligonucleotide system-based multiplex PCR assay, *J. Virol. Methods* 156 (2009) 111–116.
- [28] X. Ma, H. Xu, L. Shi, P. Yang, L. Zhang, X. Sun, W. Zhen, K. Hu, A multiplex PCR assay for the detection of five influenza viruses using a dual priming oligonucleotide system, *BMC Infect. Dis.* 15 (2015) 93.
- [29] J.Y. Yeh, J.H. Lee, H.J. Seo, J.Y. Park, J.S. Moon, I.S. Cho, I.S. Choi, S.Y. Park, C.S. Song, J.B. Lee, Simultaneous detection of rift valley fever, bluetongue, rinderpest, and peste des petits ruminants viruses by a single-tube multiplex reverse transcriptase-PCR assay using a dual-priming oligonucleotide system, *J. Clin. Microbiol.* 49 (2011) 1389–1394.
- [30] W.L. Fan, Z.W. Wang, Y. Qin, C. Sun, Z.M. Liu, Y.P. Jiang, X.Y. Qiao, L.J. Tang, Y.J. Li, Y.G. Xu, Use of dual priming oligonucleotide system-based multiplex RT-PCR combined with high performance liquid chromatography assay for simultaneous detection of five enteric viruses associated with acute enteritis, *J. Virol. Methods* 243 (2017) 80–82.
- [31] Y. Wang, B. Feng, C. Niu, S. Jia, C. Sun, Z. Wang, Y. Jiang, W. Cui, L. Wang, Y. Xu, Dendritic cell targeting of bovine viral diarrhoea virus E2 protein expressed by *Lactobacillus casei* effectively induces antigen-specific immune responses via oral vaccination, *Viruses* 11 (2019) 575.
- [32] X.W. Gao, Y.Y. Ma, Z. Wang, J. Bai, S. Jia, B.H. Feng, Y.P. Jiang, W. Cui, L.J. Tang, Y.J. Li, L. Wang, Y.G. Xu, Oral immunization of mice with a probiotic *Lactobacillus casei* constitutively expressing the α -toxin induces protective immunity against *Clostridium perfringens* α -toxin, *Virulence* 10 (2019) 166–179.
- [33] J.O. Amimo, A.N. Vlasova, L.J. Saif, Detection and genetic diversity of porcine group A rotaviruses in historic (2004) and recent (2011 and 2012) swine fecal samples in Ohio: predominance of the G9P 13 genotype in nursing piglets, *J. Clin. Microbiol.* 51 (2013) 1142–1151.
- [34] Q. Chen, P. Gauger, M. Stafne, J. Thomas, P. Arruda, E. Burrough, D. Madson, J. Brodie, D. Magstadt, R. Derscheid, M. Welch, J. Zhang, Pathogenicity and pathogenesis of a United States porcine deltacoronavirus cell culture isolate in 5-day-old neonatal piglets, *Virology* 482 (2015) 51–59.
- [35] B. Wang, Y. Liu, C.M. Ji, Y.L. Yang, Q.Z. Liang, P. Zhao, L.D. Xu, X.M. Lei, W.T. Luo, P. Qin, J. Zhou, Y.W. Huang, Porcine deltacoronavirus engages the transmissible gastroenteritis virus functional receptor porcine aminopeptidase N for infectious cellular entry, *J. Virol.* 92 (2018) e00318-18.