

## RESEARCH ARTICLE

# Association between vitamin D deficiency and allergic symptom in pregnant women

Kumiko T. Kanatani<sup>1,2\*</sup>, Yuichi Adachi<sup>3</sup>, Kei Hamazaki<sup>4</sup>, Kazunari Onishi<sup>5</sup>, Tohshin Go<sup>2</sup>, Kyoko Hirabayashi<sup>2</sup>, Motonobu Watanabe<sup>6</sup>, Keiko Sato<sup>2,7</sup>, Youichi Kurozawa<sup>5</sup>, Hidekuni Inadera<sup>4</sup>, Hiroshi Oyama<sup>8</sup>, Takeo Nakayama<sup>1</sup>, for the Japan Environment and Children's Study Group<sup>†</sup>

**1** Department of Health Informatics, Kyoto University School of Public Health, Kyoto, Japan, **2** Kyoto Unit Centre for Japan Environment and Children's Study, Graduate School of Medicine, Kyoto University, Kyoto, Japan, **3** Department of Pediatrics, Faculty of Medicine, University of Toyama, Toyama, Japan, **4** Department of Public Health, University of Toyama, Toyama, Japan, **5** Department of Public Health, Tottori University, Yonago, Tottori, Japan, **6** Doshisha University Center for Baby Science, Kizugawa, Kyoto, Japan, **7** Institute for Advancement of Clinical and Translational Science, Kyoto University Hospital, Kyoto, Japan, **8** Department of Clinical Information Engineering, Health Services Sciences, School of Public Health, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan

<sup>†</sup> Membership of the Japan Environment and Children's Study Group is provided in the Acknowledgments.

\* [kanatanik@mac.com](mailto:kanatanik@mac.com)



## OPEN ACCESS

**Citation:** Kanatani KT, Adachi Y, Hamazaki K, Onishi K, Go T, Hirabayashi K, et al. (2019) Association between vitamin D deficiency and allergic symptom in pregnant women. PLoS ONE 14(4): e0214797. <https://doi.org/10.1371/journal.pone.0214797>

**Editor:** Sakamuri V. Reddy, Charles P. Darby Children's Research Institute, UNITED STATES

**Received:** July 11, 2018

**Accepted:** March 20, 2019

**Published:** April 10, 2019

**Copyright:** © 2019 Kanatani et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data are unsuitable for public deposition due to ethical restrictions and legal framework of Japan. It is prohibited by the Act on the Protection of Personal Information (Act No. 57 of 30 May 2003, amendment on 9 September 2015) to publicly deposit the data containing personal information. Ethical Guidelines for Epidemiological Research enforced by the Japan Ministry of Education, Culture, Sports, Science and Technology and the Ministry of Health, Labour and Welfare also restricts the open sharing of the epidemiologic data. All inquiries about access to

## Abstract

### Background

Vitamin D has been reported to affect both innate, and acquired immunity with immune cells such as dendritic cells having the vitamin D receptors. The co-occurrence of the high prevalence of allergic diseases and vitamin D deficiency globally documented in recent decades, has prompted a hypothesis on whether there is a reasonable association between them.

### Objective

To investigate the association between serum vitamin D deficiency and allergic symptoms.

### Methods

Historical cohort. On a cohort study for the association between desert dust exposure and allergic symptoms in 3,327 pregnant women during spring and fall in 2011–2013 in Japan conducted as an adjunct study to the Japan Environment and Children's Study, we promptly acquired subjects' daily allergic symptom scores by sending a web-based questionnaire to each participant on some days. Of the 29,434 answers provided by 3,327 participating pregnant women, we extracted 13,356 answers from 1,475 pregnant women that were answered within a 3-month period after blood samplings. And we measured 25(OH)D levels on those samples to investigate the association between their vitamin D deficiency (serum 25(OH)D < 20ng/mL) and the occurrence of any allergic symptom (allergic symptom score > 0) within 3 months.

data should be sent to: [jecs-en@nies.go.jp](mailto:jecs-en@nies.go.jp). The person responsible for handling enquiries sent to this e-mail address is Dr Shoji F. Nakayama, JECS Programme Office, National Institute for Environmental Studies.

**Funding:** This work was supported by the Environment Research and Technology Development Fund from the Ministry of the Environment, Japan [5C-1152(2011-2013) 5-1555 (2015-2017)], and JSPS KAKENHI Grant Number 25350044 (2013-2015), and conducted as an adjunct study of Japan Environment and Children Study (JECS). The findings and conclusions of this article are solely the responsibility of the authors and do not represent the official views of the Japanese government. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Results

Serum 25(OH)D was less than 20ng/mL in 1,233 of 1,745 samples (70.7%). The adjusted odds ratio (aOR) for occurrence of any allergic symptom in deficient cases compared with non-deficient cases was 1.33 (95% CI: 1.07–1.64,  $p = 0.01$ ). Further, vitamin D deficiency significantly enhanced the risk increase at desert dust events and at pollen exposure ( $p$ -values for interaction  $<0.1$ ).

## Conclusion

We confirmed the association between serum vitamin D deficiency and allergic symptoms in Japanese pregnant women.

## Introduction

Allergic diseases are major public health problems worldwide with an increasing prevalence in the past 40 years[1,2], with their causes unknown. Many hypotheses link the allergy epidemic to stringent hygiene in early life, replacement of a traditional way of living with a westernized lifestyle, and a more accelerated pace of life. We have demonstrated that desert dust exposure is a risk factor for exacerbations of allergy symptoms[3]. However, some individuals were not affected at all by this extrinsic exposure, suggesting a partial role of the host, too. It would appear that these susceptible individuals, with intrinsic factors have increased in recent years.

As a consequence of indoor occupations followed by reduced exposure to sunlight, concerns have been raised that vitamin D deficiency is widespread in developed countries. Vitamin D has been reported to affect both innate, and acquired immunity through immune cells having the vitamin D receptors, such as dendritic cells and regulatory T cells[4].


In the present study, we hypothesized that vitamin D deficiency could be a cause of the excessive immune reactions at events such as desert dust exposure, and examined the association between serum vitamin D status and allergic symptom exacerbation, with respect to Asian dust events or pollen dispersion in pregnant women, by measuring frozen-stored serum 25 (OH)D levels of pregnant women who were in a panel study where they responded to questionnaires about their allergic symptoms on the day repeatedly during spring and fall.

The study protocol was registered at UMIN000010826, and was published[5].

## Methods

### Study design

Historical cohort. The study was started in 2011 as the first part of an adjunct study of the Japan Environment and Children's Study (JECS) to examine the effects of desert dust exposure on allergic symptoms in pregnant women, effect-modification potential of other pollutants and pollens, and to identify vulnerable populations for dust events in areas within Kyoto, Toyama, and Tottori, Japan[3,5–7]. The JECS participants from these regions who agreed to participate in the adjunct study were enrolled prior to delivery of their child. The details of the study design have been described elsewhere[3,5–7]. Briefly, questionnaires asking about allergic symptoms on the day were sent out during the Asian dust seasons (Feb-May and Oct-Nov) on Asian dust days and on several other randomly selected days for each participant during her pregnancy. The total number of questionnaires sent out was 41,020, of which 29,434



**Fig 1. Extraction of subjects in the adjunct study of the Japan Environment and Children's Study (JECS).** Of 7,829 JECS participants as of May 2013 in Kyoto, Toyama, and Tottori, 4,178 participants presented informed consent to the adjunct study, and 3,327 participants responded to symptom questionnaire during the Asian dust season. Of them, we extracted responses of 1,745 participants that were answered within 3 months after their blood-sampling.

<https://doi.org/10.1371/journal.pone.0214797.g001>


(71.8%) were completed within the set time limit of 28 hours after the questionnaire was sent out[3]. They also volunteered serum samples three times during pregnancy; once per pregnant trimester[7].

Of the 29,434 responses on 3,327 pregnant women, we extracted responses that were answered within 3 months after their blood-sampling (Figs 1 and 2). And we measured 25 (OH)D levels in those samples to investigate the association between their vitamin D deficiency (serum 25(OH)D < 20ng/mL) and the occurrence of any allergic symptoms (allergic symptom score > 0). The study-protocol was approved by the ethics committees of Kyoto University, University of Toyama, and Tottori University.

## Measurements

**Demographics.** Upon enrolment, baseline information such as history of diagnosis of asthma or other allergic diseases was collected via a web-based questionnaire[5]. In addition, various lifestyle parameters, including diet, housing environment, socioeconomic background, and serum IgE to common allergens, were obtained typically during their 1<sup>st</sup> trimesters[7]. The details of measured items are described elsewhere[3,5,7].

**Vitamin D levels.** Serum 25(OH)D was measured by a two-sandwich RIA method[8]. Biochemical analysis of blood was performed on serum samples at SRL laboratories. The serum samples were kept frozen (-30 °C) until analysis. 25(OH)D was measured using the 25-hydroxyvitamin D 125I RIA kit. The validated measurement range was 6–99900000 ng/ml.



**Fig 2. Study design.** Questionnaires asking about allergic symptoms on the day were sent out during the Asian dust seasons (Feb-May and Oct-Nov) on Asian dust days and on several other randomly selected days for each participant during her pregnancy. Serum samples were drawn three times during pregnancy; once per pregnant trimester. Responses that were answered within 3 months after their blood-sampling were extracted for the analysis.

<https://doi.org/10.1371/journal.pone.0214797.g002>

**Allergic symptoms.** A self-administered questionnaire including items from the Japanese version of the Allergy Control Score [9,10] was sent to participants via mobile phone and/or personal computer. The Allergy Control Score consists of items pertaining to daily nasal, ocular, and lung symptoms, which are measured on a 4-point scale (absent, mild, moderate, and severe) [9,10].

**Covariates.** We examined the following factors including the model based on our previous report[3]: Asian dust,  $SO_2$ , pollen count, lowest temperature on the day, temperature difference within the day, humidity, seasonality (spring or autumn), cedar-pollen-specific IgE class, and house-dust-mite (*Dermatophagoides pteronyssinus*)-specific IgE class. Details are shown elsewhere[3].

### Statistical analysis

Although there is little agreement on the definition of vitamin D insufficiency especially for non-musculoskeletal health, the Institute of Medicine panel concluded that circulating levels of 25(OH)D  $>20$  ng/mL are sufficient with respect to skeletal health[11]. *A priori* we defined vitamin D deficiency as serum 25(OH)D less than 20 ng/mL of serum 25(OH)D in the protocol based on previous studies conducted on the Japanese people[12,13]. For occurrences of any symptoms (symptom score exceeding 0), we estimated an odds ratio for an effect measure that compares the symptom occurrence between vitamin D deficient cases (25(OH)D

<20 ng/mL) and non-deficient cases, using the Generalized Estimating Equations (GEE) logistic regression analysis. We then included the interaction term of desert dust exposure and vitamin D deficiency or pollen exposure and vitamin D deficiency to investigate if the vitamin D sufficiency changes the risk of Asian dust or pollen exposure for the occurrence of allergic symptoms.

The robustness of the main results was tested by performing the following sensitivity analyses. First, we tried a cut-off value of 30 ng/mL serum 25(OH)D as vitamin D deficiency, and we also tried treating 25(OH)D as a continuous variable. Then we treated the outcome as a continuous variable (log-transformed symptom scores) using a linear mixed model.

All analysis were performed using SAS software, version 9.3 (SAS Institute), and 2-sided  $P < 0.05$  was considered statistically significant ( $P < 0.1$  for interaction terms).

## Results

### Subjects

Of the 29,434 answers on 3,327 pregnant women who participated in the study, we extracted 13,356 answers for 1,475 pregnant women, that were responded within 3 months after their blood-sample (Fig 1). Table 1 summarize the characteristics of extracted subjects. All were pregnant, in a wide age range from teens to over 45 years, and from various socioeconomic backgrounds. Approximately half exhibited positive IgE to cedar pollen and nearly half to dust mite allergen. Participant characteristics were similar to those reported by the Japanese government in 2012, except that the proportion of current smokers was lower among the study participants (1.6–2.7%) compared with those in the government report (12.8% for women in their 20s and 16.6% for women in their 30s) [14,15], and that the proportion of participants with an education level up to junior high school completion was lower (1.6–3.3%) compared to that in the government report (6.0%) [16,17]. Overall, we believe the extracted cohort was considered a good representation of pregnant women in Japan.

### Vitamin D sufficiency status

Serum 25(OH)D was less than 20ng/mL in 1,233 of 1,745 samples (70.7%), and 8.0% (139 of 1,745 samples) showed even less than 10 ng/mL which is characterized as severe vitamin D deficiency. Consistent with previous reports, there was a clear seasonal change with a peak at the end of summer and a trough in early spring [18] (Fig 3), both in allergic and non-allergic pregnant women (S1 and S2 Figs). The median level of serum 25(OH)D in each season was 15, 14, 19, and 20 ng/mL in winter (Dec-Feb), spring (Mar-May), summer (Jul-Aug) and autumn (Sep-Nov), respectively.

### Association between vitamin D deficiency and allergic symptoms

GEE logistic regression analysis with an exchangeable correlation matrix showed that pregnant women were more likely to develop allergic symptoms (symptom score >0) when they were vitamin D deficient (i.e. 25(OH)D < 20ng/mL) compared with non-deficient subjects (odds ratio [OR] 1.29; 95% confidence interval [95%CI]: 1.08–1.54,  $p = 0.005$ , adjusted OR 1.33 (95% CI: 1.07–1.64,  $p = 0.009$ ), Table 2). The increased odds of the vitamin D deficiency was observed independently of the pregnancy trimesters, that is, including the trimester in the adjusted model did not change the estimated OR (adjusted OR for allergic symptom development of vitamin D deficiency: 1.29 (95%CI: 1.05–1.59,  $p = 0.0167$ ); and the 25(OH)D deficiency was a consistent risk factor for the allergic symptoms through the trimesters (S1 and S2 Tables). Sensitivity analysis showed that this was true with a cut-off value of 30 ng/mL serum

Table 1. Characteristics of subjects.

	Kyoto n = 464	Toyama n = 744	Tottori n = 267
<b>Age (years)</b> <sup>a b</sup> n = 1457, 459, 734, 264	32.2±4.5	31.0±4.8	31.0±4.5
<b>Height (cm)</b> <sup>a b</sup> n = 455, 718, 257	159.0±5.5	158.7±5.2	157.6±5.0
<b>BMI before pregnancy</b> <sup>a b</sup> n = 453, 734, 262	20.8±2.9	21.0±2.8	20.7±2.7
<b>History of allergic diseases</b>	n (%)	n (%)	n (%)
Asthma	52 (11.2)	81 (10.9)	39 (14.6)
Allergic rhinitis	254 (45.3)	463 (37.8)	157 (41.2)
Atopic dermatitis	87 (18.8)	125 (16.8)	46(17.2)
<b>Serum positive IgE<sup>c</sup></b>	n (%)	n (%)	n (%)
Japanese Cedar Pollen positive	283 (61.0)	397 (53.4)	126 (47.2)
House Dust Mite positive	233 (50.2)	373 (50.1)	118 (44.2)
Sample not available	15 (3.2)	23 (3.1)	13 (4.9)
<b>Smoking history<sup>b</sup></b>	n (%)	n (%)	n (%)
Never smoked	297 (64.0)	432 (58.1)	163 (61.0)
Stopped before pregnancy	108 (23.3)	195 (26.2)	63 (23.6)
Stopped after pregnancy	34 (7.3)	69 (9.3)	24 (9.0)
Current smoker	7 (1.4)	18 (2.5)	6 (1.5)
Blank <sup>d</sup>	18 (3.9)	30 (4.0)	11 (4.1)
<b>Partner's smoking history<sup>b</sup></b>	n (%)	n (%)	n (%)
Never smoked	163 (35.1)	209 (28.1)	69 (25.8)
Stopped before pregnancy	129 (27.8)	183 (24.6)	68 (25.5)
Stopped after pregnancy	9 (1.9)	14 (1.9)	3 (1.1)
Current smoker	140 (31.5)	302 (40.8)	112 (45.0)
Blank <sup>d</sup>	23 (5.0)	36 (4.8)	15 (5.6)
<b>Education level<sup>b</sup></b>	n (%)	n (%)	n (%)
Junior high school	7 (1.5)	24 (3.2)	8 (3.0)
High school	84 (18.1)	172 (23.1)	79 (29.6)
Vocational school	65 (14.0)	167 (22.4)	61 (22.8)
College (2 years)	110 (23.7)	128 (17.2)	43 (16.1)
High vocational School	2 (1.1)	4 (2.7)	7 (3.0)
College (4 years)	165 (35.6)	214 (28.8)	57 (21.3)
Graduate school	26 (5.6)	15 (2.0)	4 (1.5)
Blank <sup>d</sup>	5 (1.1)	20 (2.7)	8 (3.0)
<b>Family income /year<sup>b</sup></b>	n (%)	n (%)	n (%)
Under \$20,000	11 (2.4)	11 (1.5)	15 (5.6)
\$20,000–40,000	125 (26.9)	209 (28.1)	99 (37.1)
\$40,000–60,000	157 (33.8)	261 (35.1)	67 (25.1)
\$60,000–80,000	85 (18.3)	117 (15.7)	41 (15.4)
\$80,000–100,000	39 (8.4)	62 (8.3)	12 (4.5)
Above \$100,000	27 (5.8)	26 (3.5)	13 (4.9)
Blank <sup>d</sup>	20 (4.3)	58 (7.8)	20 (7.5)
<b>Frequency exposed to sunlight</b>	n (%)	n (%)	n (%)
Rarely	43 (9.3)	161 (22.0)	51 (19.1)
1 to 2 days a week	94 (20.3)	192 (25.8)	68 (25.5)
3 to 4 days a week	117 (25.2)	165 (22.2)	55 (20.6)
More than 5 days a week	172 (37.1)	163 (21.9)	63 (23.6)
Blank <sup>d</sup>	38 (8.2)	60 (8.1)	30 (11.2)

(Continued)



Table 1. (Continued)

	Kyoto n = 464	Toyama n = 744	Tottori n = 267
UV block usage	n (%)	n (%)	n (%)
Never expose bare skin to sunlight	10 (2.2)	27 (3.6)	5 (1.9)
Sometimes use UV-blocks	275 (59.3)	362 (48.7)	123 (46.1)
Rarely use UV-blocks	120 (25.9)	262 (35.2)	104 (39.0)
Blank <sup>d</sup>	59 (12.7)	93 (12.5)	35 (13.1)


<sup>a</sup> Mean ± SD

<sup>b</sup> referred to JECS (tentative data)

<sup>c</sup> referred to JECS (fixed data)

<sup>d</sup> Blank includes those who did not return the answer sheet.

<https://doi.org/10.1371/journal.pone.0214797.t001>



**Fig 3. Serum 25(OH)D levels in relation to sampled months.** Serum 25(OH)D was less than 20ng/mL in 1,233 of 1,745 samples (70.7%). There was a clear seasonal change with a peak at the end of summer and a trough in early spring. The median level of serum 25(OH)D in each season was 15, 14, 19, and 20 ng/mL in winter (Dec-Feb), spring (Mar-May), summer (Jul-Aug) and autumn (Sep-Nov).

<https://doi.org/10.1371/journal.pone.0214797.g003>

**Table 2. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development.**

	OR	95% CI			P value
Vitamin D deficiency <sup>a</sup>	1.33	1.07	–	1.64	.009
IgE to cedar pollen (per class increase)	1.28	1.21	–	1.35	<.001
IgE to house dust mite (per class increase)	1.10	1.03	–	1.18	.008
Age (compared with >40 years)					
<20 years	0.81	0.19	–	3.55	.781
20–25 years	1.82	1.01	–	3.27	.046
25–30 years	1.49	0.95	–	2.35	.085
30–35 years	1.62	1.04	–	2.53	.034
35–40 years	1.47	0.93	–	2.32	.103
BMI before pregnancy (compared with >25)					
<18	1.00	0.68	–	1.47	.995
18–25	1.14	0.82	–	1.60	.436
Family income (per 20,000 increase)	0.95	0.87	–	1.01	.113
Smoking status of subjects (Compared with current smoker)					
Never smoker	0.98	0.54	–	1.79	.957
Stopped before pregnancy	1.12	0.61	–	2.07	.708
Stopped after pregnancy	1.04	0.54	–	1.97	.915
Smoking status of subjects' partners (Compared with current smoker)					
Never smoker	1.01	0.81	–	1.26	.940
Stopped before pregnancy	1.01	0.81	–	1.25	.956
Stopped after pregnancy	0.51	0.29	–	0.90	.020
Location (Compared with Kyoto)					
Toyama	0.99	0.81	–	1.21	.920
Tottori	1.03	0.80	–	1.34	.795

Adjusted by desert dust, pollen counts, SO<sub>2</sub>, humidity, lowest temperature of the day, temperature difference within the day, and month on the day.

Vitamin D deficiency was defined as serum 25(OH)D level < 20 ng/mL on serum samples taken within 3 months before the symptom response.

IgE class to cedar pollen and IgE class to house dust mite were treated as ordered categorical. The OR per one class increase for each is shown.

Age was categorized into 6 groups by 5 years and was treated as categorical.

BMI before pregnancy was categorized into 3 groups, and was treated as categorical.

Family income was categorized into 6 groups and was treated as ordered categorical.

<https://doi.org/10.1371/journal.pone.0214797.t002>

25(OH)D, which is often described as vitamin D insufficiency level, and the OR for symptom development decreased as the vitamin D level increased (OR 0.80; 95%CI: 0.64–1.00, p = 0.053 for those with 20–30 ng/mL, and OR 0.68; 95%CI: 0.47–0.99, p = 0.043 for those with <30 ng/mL, [S3 Table](#)).

### Vitamin D deficiency as an allergic-symptom risk modifier at events

The risk increases during Asian dust events appeared more evident in vitamin D deficient subjects compared with vitamin D non-deficient subjects (OR 1.15, 95%CI: 1.04–1.27 for deficient subjects, and OR 1.10, 95%CI: 0.88–1.36 for non-deficient subjects), and so did the one by pollen exposure (OR 1.45, 95%CI: 1.29–1.63 for deficient subjects, and OR 1.18, 95%CI: 0.90–1.54 for non-deficient subjects) ([S4](#) and [S5 Tables](#)). The models with interaction terms of ‘dust



**Table 3. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development with the interaction term of vitamin D deficiency and desert dust.**

	OR	95% CI			P value
Vitamin D deficiency <sup>a</sup>	1.03	0.97	–	1.09	.288
Desert dust <sup>b</sup>	1.04	1.01	–	1.06	.003
Interaction term	1.03	0.99	–	1.07	.098

Those who were seropositive for IgE to JCP and went outdoors on the day were included in the analysis.

Adjusted by pollen counts (log-transformed and standardized), humidity, lowest temperature of the day, temperature difference within the day, IgE to house dust mite, domestic income, smoking status, age, and BMI before pregnancy. IgE class to house dust mite was treated as ordered categorical.

Age was categorized into 6 groups by 5 years, and was treated as categorical.

BMI before pregnancy was categorized into 3 groups, and was treated as categorical.

Family income was categorized into 6 groups and was treated as ordered categorical.

<sup>a</sup> Vitamin D deficiency was defined as serum 25(OH)D level < 20 ng/mL on serum samples taken within 3 months before the symptom response.

<sup>b</sup> Desert dust level was log-transformed and standardized.

<https://doi.org/10.1371/journal.pone.0214797.t003>

**Table 4. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development with the interaction term between vitamin D deficiency and pollen exposure.**

	OR	95% CI			P value
Vitamin D deficiency <sup>a</sup>	1.05	0.99	–	1.11	.125
Pollen <sup>b</sup>	1.09	1.06	–	1.12	< .001
Interaction term	1.05	1.00	–	1.11	.059

Those who were seropositive for IgE to JCP and went outdoors on the day were included in the analysis.

Adjusted by desert dust (log-transformed and standardized), humidity, lowest temperature of the day, temperature difference within the day, IgE to house dust mite, family income, smoking status, age, and BMI before pregnancy. IgE class to house dust mite was treated as ordered categorical.

Age was categorized into 6 groups by 5 years, and was treated as categorical.

BMI before pregnancy was categorized into 3 groups, and was treated as categorical.

Family income was categorized into 6 groups and was treated as ordered categorical.

<sup>a</sup> Vitamin D deficiency was defined as serum 25(OH)D level < 20 ng/mL on serum samples taken within 3 months before the symptom response.

<sup>b</sup> Pollen count was log-transformed and standardized.

<https://doi.org/10.1371/journal.pone.0214797.t004>

level' and 'serum 25(OH)D deficiency' or 'pollen level' and 'serum 25(OH)D deficiency' showed statistically significant ( $p < 0.1$ ) interaction (Tables 3 and 4).

## Discussion

In this study, we showed that pregnant women are significantly more likely to have self-reported allergic symptoms when they have deficient level of 25(OH)D (< 20ng/mL) compared with non-deficient women. The increased risks of allergic symptoms during Asian dust exposure and pollen exposure were more evident in vitamin D deficient subjects compared with vitamin D sufficient subjects. Our finding supports the hypothesis that the recent epidemic of allergy could partly be caused by vitamin D deficiency, suggesting that we may be able to reduce the symptoms by approximately 30% by ensuring a level of 20 ng/mL of serum 25(OH)D level for pregnant women. The impact would be large because vitamin D deficiency

is observed wide-spread; Serum 25(OH)D was less than 20ng/mL in 1,233 of 1,745 samples (70.7%) in our study, and this was consistent with reports of pregnant women from Japan and other Asian countries[19,20].

Some epidemiologic studies suggest a protective effect of higher vitamin D exposure against allergic diseases, while other results point to increased risk or no associations with vitamin D exposure. The latest systematic review for vitamin D supplementation for pregnant women, breastfeeding women, and infants concluded that the supplementation did not decrease the risk of developing allergic diseases[21]. On the other hand, another latest systemic review revealed that vitamin D supplementation reduced the frequency of asthma exacerbations[22]. Besides of the heterogeneity in the definition of outcomes and timings of vitamin D measurements, there may be other potential reasons for the controversial findings. Hawrylowicz *et al.* recently showed an interesting hypothesis that the apparent beneficial association of vitamin D in allergies *in vivo*, which in many patients is a  $T_H2$ -type cytokine pathology, suggests that the direct action of vitamin D in promoting  $T_H2$  lymphocyte responses is less important *in vivo* than other vitamin D-mediated mechanisms (eg, vitamin D upregulated production of soluble decoy receptor, an inhibitor of IL-33 *in situ* in the airways mucosa, and the induction of regulatory mechanisms) in inhibiting  $T_H2$ -type cytokine responses[23]. Vitamin D supplementations may not reduce the risk of developing allergic diseases[21], but may reduce the inflammation (exacerbations of symptoms) in people with allergies. Our results that Asian dust, which includes silica and other possible danger signals[3] that may cause IL-33 release, exhibits adjuvant-like effects in pollen-sensitized pregnant women[3], and this was evident in vitamin D deficient women supports this hypothesis. Further work needs to be done in this field to elucidate the mechanisms.

The strengths of this study include a relatively large dataset on a population-based cohort resulting in high generalizability to pregnant women in Japan. Other strengths include the fact that the lists of symptoms were collected repeatedly during Asian dust/pollen season while their serum IgE levels were also measured. This enabled us to perform detailed analysis for symptom developments at events. And by sending a web-based questionnaire on the day, which required a response within a set time limit using a mobile phone or personal computer, we ensured a high response rate (72%) and reduced recall misclassifications of symptoms.

This study has three major limitations. The most notable is our inability to exclude the possibility of reverse causality for the association between allergic symptom and vitamin D deficiency, as is often the case with observational studies; that is, the pregnant women who had severe symptoms might not have dared to go outdoors to avoid pollen exposure, and this might have resulted in low 25(OH)D levels in the serum due to reduced sunlight exposure. Second, allergic symptoms were self-reported and not confirmed by medical professionals. In other words, symptoms were not distinguished by origin (i.e., allergy, infection, or other). Thus, our speculation regarding allergic symptoms is based on timing of the onset of symptoms (i.e., due to pollen or Asian dust) and vulnerability of participants (i.e., positive serum IgE to pollen). The third limitation relates to generalizability. We investigated pregnant women, whose immune responses as well as vitamin D status might have been altered due to pregnancy. Further studies should be performed to verify whether our findings also apply to the non-pregnant population.

In conclusion, pregnant women were significantly more likely to have allergic symptoms if their 25(OH)D level was < 20ng/mL, compared with non-deficient subjects (OR 1.28 (1.04–1.56)). We may be able to reduce the symptoms by approximately 30% by ensuring a level of 20 ng/mL of serum 25(OH)D level for pregnant women. The risk increase during Asian dust events or pollen exposure was more evident in vitamin D deficient pregnant women compared

with vitamin D non-deficient pregnant women. This adds to the growing evidence of the association between vitamin D insufficiency and allergic diseases.

## Supporting information

**S1 Fig. Serum 25(OH)D levels in relation to sampled months on subjects with/without history of allergic rhinitis.** There was a clear seasonal change with a peak at the end of summer and a trough in early spring regardless of past history of allergic rhinitis.

(TIFF)

**S2 Fig. Serum 25(OH)D levels in relation to sampled months on subjects seropositive/seronegative to Japanese Cedar Pollen (JCP).** There was a clear seasonal change with a peak at the end of summer and a trough in early spring regardless of the presence of serum IgE to JCP.

(TIFF)

**S1 Table. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development of vitamin D deficiency in their 1<sup>st</sup> trimester.**

(DOCX)

**S2 Table. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development of vitamin D deficiency in their 2<sup>nd</sup> trimester.**

(DOCX)

**S3 Table. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development on cases with sufficient in vitamin D ( $\geq 30$  ng/mL) and on cases with insufficient vitamin D (20–30 ng/mL) compared with cases deficient in vitamin D ( $< 20$  ng/mL).**

(DOCX)

**S4 Table. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development compared with cases deficient in vitamin D ( $< 20$  ng/mL).**

(DOCX)

**S5 Table. Odds ratio (OR) and its 95% Confidence Interval (95%CI) for allergic symptom development in Vitamin D non-deficient subjects (25(OH)D  $\geq 20$  ng/mL).**

(DOCX)

## Acknowledgments

Japan Environment and Children's Study Group; Kyoto Regional Centre: Ikuo Konishi, Toshio Heike, Takeo Nakayama, Yukuo Konishi, Tohshin Go, Keiko Sato, Miho Egawa, Yoshimitsu Takahashi, Motonobu Watanabe, Reiko Yasumi, Kyoko Hirabayashi, Masahito Morita, Kaori Konishi, Kumiko T Kanatani, Research Coordinators, and members of the JECS Kyoto Management Committee; Toyama Regional Centre: Hidekuni Inadera, Kei Hamazaki, Yuichi Adachi, Akiko Tsuchida, and Research Coordinators; Tottori Regional Centre: Soji Fukumoto, Ryota Teshima, Takao Inoue, Youichi Kurozawa, Tasuku Harada, Susumu Kan-zaki, Yoshihiro Maegaki, Kosaku Ohno, Tatsuya Koeda, Kazunari Onishi, Toshio Masumoto, and Research Coordinators.

The authors would like to thank all participants of the study who contributed to a very high response rate; Prof. Yoshio Tanigawa for fruitful discussions; Toyama Prefectural Environmental Science Research Centre, Kyoto Prefectural Government, Department of Culture and the Environment, Lake Biwa Environmental Research Institute, Kyoto City, Environmental Conservation Division, Toyama City Hall, Air and Water Environment Division, Tottori

Prefecture Environmental Sanitation Research Center for providing air pollution data; Japan Weather Association for pollen data; Japan Meteorological Agency for meteorological data; and personnel at the hospitals and clinics that collaborated in the JECS study.

## Author Contributions

**Conceptualization:** Kumiko T. Kanatani, Yuichi Adachi.

**Data curation:** Kumiko T. Kanatani, Kei Hamazaki, Kazunari Onishi, Tohshin Go, Kyoko Hirabayashi, Motonobu Watanabe, Keiko Sato.

**Formal analysis:** Kumiko T. Kanatani.

**Funding acquisition:** Kumiko T. Kanatani, Yuichi Adachi, Tohshin Go, Youichi Kurozawa, Hidekuni Inadera, Takeo Nakayama.

**Investigation:** Kumiko T. Kanatani, Yuichi Adachi, Tohshin Go.

**Methodology:** Kumiko T. Kanatani, Yuichi Adachi, Hiroshi Oyama.

**Project administration:** Kumiko T. Kanatani, Yuichi Adachi, Kei Hamazaki, Kazunari Onishi, Kyoko Hirabayashi, Motonobu Watanabe, Keiko Sato, Youichi Kurozawa, Hidekuni Inadera, Takeo Nakayama.

**Resources:** Kumiko T. Kanatani, Yuichi Adachi.

**Software:** Kumiko T. Kanatani.

**Supervision:** Yuichi Adachi, Tohshin Go, Youichi Kurozawa, Hidekuni Inadera, Hiroshi Oyama, Takeo Nakayama.

**Validation:** Kumiko T. Kanatani, Kei Hamazaki, Kazunari Onishi.

**Visualization:** Kumiko T. Kanatani, Hiroshi Oyama.

**Writing – original draft:** Kumiko T. Kanatani.

**Writing – review & editing:** Kumiko T. Kanatani, Yuichi Adachi, Kei Hamazaki, Kazunari Onishi, Tohshin Go, Kyoko Hirabayashi, Motonobu Watanabe, Keiko Sato, Youichi Kurozawa, Hidekuni Inadera, Hiroshi Oyama, Takeo Nakayama.

## References

1. The World Allergy Organization (WAO) White Book on Allergy. 2013. (Accessed Apr. 23, 2016, at [http://www.worldallergy.org/definingthespecialty/white\\_book.php](http://www.worldallergy.org/definingthespecialty/white_book.php).)
2. Rutkowski K, Sowa P, Rutkowska-Talipska J, Sulkowski S, Rutkowski R. Allergic diseases: the price of civilisational progress. *Postepy dermatologii i alergologii* 2014; 31:77–83. <https://doi.org/10.5114/pdia.2014.40936> PMID: 25097472
3. Kanatani KT, Hamazaki K, Inadera H, et al. Effect of desert dust exposure on allergic symptoms: A natural experiment in Japan. *Annals of allergy, asthma & immunology: official publication of the American College of Allergy, Asthma, & Immunology* 2016.
4. Chun RF, Liu PT, Modlin RL, Adams JS, Hewison M. Impact of vitamin D on immune function: lessons learned from genome-wide analysis. *Frontiers in physiology* 2014; 5:151. <https://doi.org/10.3389/fphys.2014.00151> PMID: 24795646
5. Kanatani KT, Adachi Y, Sugimoto N, et al. Birth cohort study on the effects of desert dust exposure on children's health: protocol of an adjunct study of the Japan Environment & Children's Study. *BMJ open* 2014; 4:e004863. <https://doi.org/10.1136/bmjopen-2014-004863> PMID: 24958210
6. Nitta H. Outline of the Japan Environment and Children's Study and the Framework of Genome Analysis. *Nihon eiseigaku zasshi Japanese journal of hygiene* 2016; 71:91–3. PMID: 26832622

7. Kawamoto T, Nitta H, Murata K, et al. Rationale and study design of the Japan environment and children's study (JECS). *BMC public health* 2014; 14:25. <https://doi.org/10.1186/1471-2458-14-25> PMID: [24410977](https://pubmed.ncbi.nlm.nih.gov/24410977/)
8. Hollis BW, Kamerud JQ, Selvaag SR, Lorenz JD, Napoli JL. Determination of vitamin D status by radioimmunoassay with an <sup>125</sup>I-labeled tracer. *Clinical chemistry* 1993; 39:529–33. PMID: [8448871](https://pubmed.ncbi.nlm.nih.gov/8448871/)
9. Tsuji Kanatani K, Slingsby BT, Mukaida K, et al. Translation and linguistic validation of the Allergy-CONTROL-Score for use in Japan. *Allergology international: official journal of the Japanese Society of Allergology* 2013; 62:337–41.
10. Hafner D, Reich K, Matricardi PM, Meyer H, Kettner J, Narkus A. Prospective validation of 'Allergy-Control-SCORE': a novel symptom-medication score for clinical trials. *Allergy* 2011; 66:629–36. <https://doi.org/10.1111/j.1398-9995.2010.02531.x> PMID: [21261656](https://pubmed.ncbi.nlm.nih.gov/21261656/)
11. Institute of Medicine Standing Committee on the Scientific Evaluation of Dietary Reference I. The National Academies Collection: Reports funded by National Institutes of Health. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. Washington (DC): National Academies Press (US) National Academy of Sciences.; 1997.
12. Kuroda T, Shiraki M, Tanaka S, Ohta H. Contributions of 25-hydroxyvitamin D, co-morbidities and bone mass to mortality in Japanese postmenopausal women. *Bone* 2009; 44:168–72. <https://doi.org/10.1016/j.bone.2008.03.023> PMID: [19010116](https://pubmed.ncbi.nlm.nih.gov/19010116/)
13. Suzuki T, Kwon J, Kim H, et al. Low serum 25-hydroxyvitamin D levels associated with falls among Japanese community-dwelling elderly. *Journal of bone and mineral research: the official journal of the American Society for Bone and Mineral Research* 2008; 23:1309–17.
14. Japan Statistical Yearbook 2013. First Editorial Section, Statistical Library, Statistical Research and Training Institute. 2013. (Accessed Apr. 19, 2016, at <http://www.stat.go.jp/data/nihon/18.htm>.)
15. The national health and nutrition survey in Japan, 2011. 2011. (Accessed Apr. 19, 2016, at <http://www.mhlw.go.jp/bunya/kenkou/eiyoudl/h24-houkoku-06.pdf>.)
16. The Population and Households of Japan 2010. at <http://www.stat.go.jp/data/kokusei/2010/pdf/waga3.pdf>.)
17. School Health Survey. at <http://www.e-stat.go.jp/SG1/estat/eStatTopPortalE.do>.)
18. Ratzinger F, Haslacher H, Stadlberger M, et al. 25(OH)D and 1,25(OH)D vitamin D fails to predict sepsis and mortality in a prospective cohort study. *Scientific reports* 2017; 7:40646. <https://doi.org/10.1038/srep40646> PMID: [28079172](https://pubmed.ncbi.nlm.nih.gov/28079172/)
19. Takaoka N, Nagao M, Umezawa M, Sairenchi T, Haruyama Y, Kobashi G. Distribution of serum 25-hydroxyvitamin D levels of reproductive age Japanese women. *Nihon Kosho Eisei Zasshi*. 2017; 64(3):133–142. Japanese. PMID: [28367934](https://pubmed.ncbi.nlm.nih.gov/28367934/)
20. Enkhmaa D, Tanz L, Ganmaa D, et al. Randomized trial of three doses of vitamin D to reduce deficiency in pregnant Mongolian women. *EBioMedicine*. 2019 Jan; 39:510–519. <https://doi.org/10.1016/j.ebiom.2018.11.060> PMID: [30552064](https://pubmed.ncbi.nlm.nih.gov/30552064/)
21. Yepes-Nunez JJ, Brozek JL, Fiocchi A, et al. Vitamin D supplementation in primary allergy prevention: Systematic review of randomized and non-randomized studies. *Allergy* 2018; 73:37–49. <https://doi.org/10.1111/all.13241> PMID: [28675776](https://pubmed.ncbi.nlm.nih.gov/28675776/)
22. Jolliffe DA, Greenberg L, Hooper RL, et al. Vitamin D supplementation to prevent asthma exacerbations: a systematic review and meta-analysis of individual participant data. *The Lancet Respiratory medicine* 2017; 5:881–90. [https://doi.org/10.1016/S2213-2600\(17\)30306-5](https://doi.org/10.1016/S2213-2600(17)30306-5) PMID: [28986128](https://pubmed.ncbi.nlm.nih.gov/28986128/)
23. Pfeffer PE, Chen YH, Woszczek G, et al. Vitamin D enhances production of soluble ST2, inhibiting the action of IL-33. *The Journal of allergy and clinical immunology* 2015; 135:824–7.e3. <https://doi.org/10.1016/j.jaci.2014.09.044> PMID: [25457999](https://pubmed.ncbi.nlm.nih.gov/25457999/)