

Research Article



Determination of the optimal diabetes duration for bone regeneration experiments in an alloxan-induced diabetic rabbit calvarial defect model

Sang-Hun Jeong ¹, Bo Hyun Jung ², Ki-Yeon Yoo ^{2,3}, Heung-Sik Um ¹,
Beom-Seok Chang ¹, Jae-Kwan Lee ^{1,3,*}, Won-Youl Choi ^{3,4,5,*}

¹Department of Periodontology, Gangneung-Wonju National University College of Dentistry, Gangneung, Korea

²Department of Anatomy, Gangneung-Wonju National University College of Dentistry, Gangneung, Korea

³Research Institute for Dental Engineering, Gangneung-Wonju National University, Gangneung, Korea

⁴Department of Metal and Materials Engineering, Gangneung-Wonju National University, Gangneung, Korea

⁵Wellnanos Co., Ltd., Gangneung, Korea

OPEN ACCESS

Received: Oct 18, 2018

Accepted: Dec 16, 2018

*Correspondence:

Jae-Kwan Lee

Department of Periodontology, Gangneung-Wonju National University College of Dentistry, Gangneung, Korea.

E-mail: periojk@gwnu.ac.kr

Tel: +82-33-640-3199

Fax: +82-33-640-3103

Won-Youl Choi

Department of Metal and Materials Engineering, Gangneung-Wonju National University and Wellnanos Co. Ltd., 24-3 Gwahakdangi-ro, Gangneung 25451, Korea.

E-mail: cwj@gwnu.ac.kr

cwj@wellnanos.com

Tel: +82-33-640-2483

Fax: +82-33-642-2245

Copyright © 2018. Korean Academy of Periodontology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>).

ORCID iDs

Sang-Hun Jeong

<https://orcid.org/0000-0003-0321-9705>

Bo Hyun Jung

<https://orcid.org/0000-0002-2283-6066>

Ki-Yeon Yoo

<https://orcid.org/0000-0002-0575-275X>

ABSTRACT

Purpose: The purpose of this study was to evaluate the optimal diabetes duration for bone regeneration experiments in an alloxan monohydrate (ALX)-induced diabetic rabbit calvarial defect model by evaluating the association between diabetes duration and bone healing capacity.

Methods: Twenty-four New Zealand white rabbits were used. Twenty-two rabbits were injected with 100 mg/kg of ALX to induce experimental diabetes. These rabbits were divided into 4 groups, including a control group and groups with diabetes durations of 1 week (group 1), 2 weeks (group 2), and 4 weeks (group 3). Calvarial defects were created at 1, 2, and 4 weeks after ALX injection and in the control rabbits. Cone-beam computed tomography (CBCT) scanning was performed on the day of surgery and at 2 and 4 weeks after surgery. The rabbits were sacrificed 4 weeks after surgery, followed by histological and immunofluorescence analysis.

Results: The diabetic state of all diabetic rabbits was well-maintained throughout the experiment. Reconstructed 3-dimensional CBCT imaging showed more rapid and prominent bone regeneration in the control group than in the experimental groups. Histological staining showed notable bone regeneration in the control group, in contrast to scarce bone formation in the experimental groups. The appearance and immunoreactivity of receptor activator of nuclear factor-kappa B and osteoprotegerin did not show notable differences among the groups.

Conclusion: ALX administration at 100 mg/kg successfully induced experimental diabetes in rabbits. The effect of diabetes on bone healing was evident when the interval between diabetes induction and the intervention was ≥ 1 week.

Keywords: Alloxan; Bone regeneration; Cone-beam computed tomography; Experimental diabetes mellitus; Rabbits

INTRODUCTION

The clinical significance of bone regeneration is rapidly expanding with the recent increase in patients requesting dental implant treatment. To assess regenerative materials such as

Heung-Sik Um 
<https://orcid.org/0000-0002-7986-1019>
 Beom-Seok Chang 
<https://orcid.org/0000-0002-5280-3249>
 Jae-Kwan Lee 
<https://orcid.org/0000-0003-1710-1580>
 Won-Youl Choi 
<https://orcid.org/0000-0002-2181-827X>

Funding

This research was financially supported by the Human Resource Training Program for Regional Innovation and Creativity and Local University Excellent Scientist Supporting Program through the Ministry of Education and National Research Foundation of Korea (Grant No. 2015H1C1A1035848 and 2017R1D1A3B03036068).

Author Contributions

Conceptualization: Jae-Kwan Lee, Ki-Yeon Yoo, Won-Youl Choi; Data curation: Sang-Hun Jeong, Jae-Kwan Lee, Bo Hyun Jung, Ki-Yeon Yoo, Won-Youl Choi; Formal analysis: Sang-Hun Jeong, Jae-Kwan Lee, Heung-Sik Um, Beom-Seok Chang; Funding acquisition: Won-Youl Choi; Investigation: Sang-Hun Jeong, Jae-Kwan Lee, Bo Hyun Jung; Methodology: Jae-Kwan Lee, Ki-Yeon Yoo, Heung-Sik Um, Beom-Seok Chang, Won-Youl Choi; Software: Sang-Hun Jeong, Jae-Kwan Lee, Ki-Yeon Yoo; Supervision: Jae-Kwan Lee, Heung-Sik Um, Beom-Seok Chang; Writing - original draft: Sang-Hun Jeong, Jae-Kwan Lee; Writing - review & editing: Jae-Kwan Lee, Ki-Yeon Yoo, Heung-Sik Um, Beom-Seok Chang, Won-Youl Choi.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

bone graft materials and barrier membranes after *ex vivo* testing, animal studies are usually performed [1]. Preclinical studies in animals can reproduce various conditions, including systemic diseases [2]. However, the exact protocol of such animal studies for systemic diseases such as diabetes mellitus (DM) has not yet been established.

DM is a chronic metabolic disorder that has been found to impair osseointegration and bone regeneration. It is characterized by hyperglycemia, resulting from defects in insulin secretion, insulin action, or both [3]. Type 1 DM is an absolute deficiency of insulin secretion, caused by autoimmune destruction of the insulin-producing β -cells of the pancreas. Type 2 DM involves impaired insulin function, such as resistance to insulin action, along with an insufficient compensatory insulin secretion response. In 2013, approximately 2.7 million Korean adults (8.03%) aged 30 years or more were found to have type 2 diabetes, and 25.0% of adults had prediabetes [4]. In addition, the prevalence of type 2 diabetes in people aged 60 years or older has been steadily increasing.

The complications of DM include problems associated with bone metabolism. Alteration of mineral homeostasis, osteoid production, and bone formation has been demonstrated in previous experimental diabetic models [5-7]. Goodman and Hori [5] described decreased osteoid formation in diabetic rats. Moreover, diabetic rats showed a longer osteoid mineralization lag time than control rats. The biological effects of DM on osseointegration have been evaluated in clinical studies [8,9]. In a prospective study to determine whether type 2 DM was a risk factor affecting the long-term clinical outcomes of dental implants [8], 255 implants were placed in type 2 DM patients, and a significantly high failure rate was found. Moy et al. [9] performed a retrospective cohort study to evaluate the hypothesis that coexisting conditions such as smoking, DM, and radiation therapy lead to increased rates of implant failure. Among 1,140 patients with 4,680 implants placed, 48 patients had DM, and they showed a failure rate of 31.25%. Thus, DM patients presented a higher relative risk of failure than healthy controls. In a preclinical study, Nevins et al. [10] observed less bone-implant contact in rats with chemically-induced diabetes than in the control group. Giglio et al. [11] reported retarded peri-implant bone healing in rats with chemically induced diabetes. Retrospective human studies also showed that DM affected implant osseointegration by impairing osseous wound and bone healing, decreasing bone density, and increasing fracture risk and susceptibility to periodontal disease [12,13].

The mechanism through which DM results in impaired osseous healing is not yet clear; however, some possible pathogenic pathways have been suggested, including hyperglycemia. In *ex vivo* studies, osteoblastic cells subjected to elevated glucose concentrations exhibited irregular bone mineral nodule formation and a suppressed cell-growth rate [14,15]. Non-enzymatic protein glycation occurs in diabetes, leading to the formation of various chemically modified proteins known as advanced glycation end products (AGEs) [16]. Santana et al. [17] found that the accumulation of AGEs in bone contributed to diminished bone healing in type 1 diabetes, possibly mediated by receptor for advanced glycation end products (RAGE). A plausible mechanism is that AGE-RAGE interactions may result in the activation of nuclear factor-kappa B (NF- κ B) in osteoblastic cells [18].

Many complications of diabetes, such as neuropathy and vascular diseases, are associated with the duration of DM [19-22]. Although a few animal studies on the effects of chemically-induced diabetes on bone defect healing have shown impaired healing of bone defects in diabetic animals [17,23,24], to our knowledge, none of those studies established a protocol

for assessing the association between the duration of diabetes and the extent of impairment of bone-healing capacity. Therefore, the purpose of this study was to examine the association between the duration of diabetes and its effect on bone-healing capacity in rabbits, and to establish an optimal experimental protocol for evaluating the bone-healing capacity of diabetic animals using cone-beam computed tomography (CBCT), histologic analysis, and immunofluorescence analysis.

MATERIALS AND METHODS

Experimental animals

Male New Zealand white rabbits (10–14 weeks, 2–3 kg) were purchased from Samtako (Osan, Korea). The animals were housed in conventional conditions, including room temperature (23°C), controlled humidity (60%), a 12-hour light/12-hour dark cycle, and *ad libitum* access to food and water. Twenty-four young male New Zealand white rabbits were divided into 4 groups, including a control group. Before any experimental manipulations, the rabbits were allowed to acclimate for at least 7 days. The study was conducted with approval of the Institutional Animal Care and Use Committee of Gangneung-Wonju National University (approval No. GWNVA-2015-37). All experiments were performed in a way that minimized the number of animals used and the distress caused by the procedures used in this study.

Study design

To evaluate the association between the duration of diabetes and bone-healing capacity, the experimental groups (groups 1, 2, 3) were organized according to the duration of diabetes. Bone defects were introduced at 1, 2, and 4 weeks after DM induction in groups 1, 2, and 3, respectively. The rabbits in the control group underwent bone defect surgery without DM induction. CBCT (Alphard Vega 3030, Asahi, Fukuoka, Japan) was performed in all groups on the day of surgery, 2 weeks after surgery, and 4 weeks after surgery. After the last CBCT scan, the rabbits were sacrificed.

Alloxan-induced diabetic rabbit model

For DM induction, alloxan monohydrate (ALX; Sigma Aldrich Co., St. Louis, MO, USA) was dissolved in 5 mL of sterile saline with a final concentration of 100 mg/kg, and was administered intravenously for 5 minutes via the marginal ear vein of rabbits in the experimental groups. The rabbits showing fasting blood sugar (FBS) levels higher than 200 mg/dL at 1 week after injection were classified as diabetic. FBS readings were taken 1 week later, using a blood glucose test meter (Roche, Basel, Switzerland) after 12 hours of fasting.

Surgical procedure

For bone defect induction, rabbits were anesthetized with isoflurane (Hana Pharm Co., Hwaseong, Korea). With a sharp blade and periosteal elevator, full dissection was done to expose the calvaria. Four circular bone defects with a diameter of 5 mm were created on frontal and parietal bone sites bilaterally in each rabbit using a trephine burr under saline irrigation. The surgical site was closed by bilayer suturing using 4-0 black silk (Ailee, Busan, Korea). All animals received an intramuscular injection of cefazolin (30 mg/kg) for 2 days to prevent postsurgical infection.

CBCT scanning and gray level unit (GLU) measurements

In the experimental and control groups, CBCT was performed under isoflurane anesthesia to evaluate the process of bone defect healing at 3-time points: just after surgery, 2 weeks later, and 4 weeks later. The exposure settings for the scans were as follows: voxel size, 100 μm ; tube current, 5 mA; tube voltage, 80 kV; and scan time, 17.0 seconds. To evaluate bone healing and mineralization, the GLUs of the bone defects were measured. The region of interest was set on the top of the bone defect margin in a circle with a diameter of 6 mm. The mean GLUs were recorded and statistically analyzed.

Tissue processing for histological analysis

Four weeks after surgery, the rabbits were sacrificed. For the histological analysis, the animals were anesthetized with isoflurane and perfused transcardially with 0.1 M phosphate-buffered saline (PBS; pH 7.4) followed by 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The frontal and parietal bones were retrieved and postfixed in the same fixative for 6 hours. The fixed tissues were decalcified with Decalcifying Solution-Lite (Sigma Aldrich Co.) for 12 hours at room temperature. The decalcifying process was completed after confirming that the bone was cut with a blade. The decalcified tissues were washed with running tap water and dehydrated by immersion for 2 hours in 50%, 70%, 80%, 90%, 95%, and 100% ethanol baths in succession at room temperature. Next, the tissues were placed twice for an hour each in fresh pure xylene and then in molds containing melted paraffin (Histowax, Leica, Wetzlar, Germany). Serial 5- μm -thick transverse sections were cut and mounted on saline-coated microscopy slides.

Histological analysis

Masson trichrome stain

A trichrome stain kit (Abcam, Cambridge, UK) was used. Specimens were washed twice with xylene for 10 minutes to remove the paraffin. Deparaffinized sections were treated with Bouin's fluid for 60 minutes and cooled for 10 minutes, followed by treatment with Weigert's hematoxylin for 5 minutes. Sections were washed in running water for 2 minutes. Then, sections were stained in Biebrich scarlet-acid fuchsin solution for 15 minutes and rinsed in distilled water (DW). Finally, the slides were placed in phosphomolybdic-phosphotungstic acid solution for 10 minutes, transferred into aniline blue solution for 5 minutes, and placed in 1% acetic acid solution. The slides were washed, dehydrated, and sealed in Canada balsam (Kanto Chemical, Tokyo, Japan) and then observed under a light microscope (Leica).

Hematoxylin/eosin staining

Sections were deparaffinized and hydrated using the conventional process. After hydration, slides were washed in DW for 10 minutes. The slides were then stained with Harris hematoxylin for 8 minutes, washed in running water, and placed in 1% acid alcohol for 30 seconds. After washing, the slides were placed in 0.5% ammonia in water and washed in water. Then, the slides were stained with eosin for 1 minute and placed in 95% alcohol for 10 minutes. The slides were placed in 100% ethanol twice for 10 minutes each, and then in xylene twice for 10 minutes each. Finally, the slides were sealed with Canada balsam and observed under a light microscope (Axio Imager A2, Zeiss, Jena, Germany).

Immunofluorescence analysis and optical density measurements

An immunofluorescence analysis was performed to identify the appearance of osteoprotegerin (OPG) and receptor activator of NF- κB (RANKL). Sections were placed in xylene twice for 10 minutes each. After deparaffinization, sections were placed in 100%

alcohol twice, and hydrated using the conventional hydration process. Sections were washed in PBS buffer thrice. Antigen retrieval was then performed by treating slides for 5 minutes in a microwave with citric acid buffer (pH 6.0). After sections were washed with PBS buffer, they were incubated at 4°C with goat anti-RANKL antibody (diluted 1:200) and mouse anti-OPG antibody (diluted 1:200) overnight. The slides were washed in PBS buffer and incubated in room temperature for 90 minutes with donkey anti-mouse 488 IgG for OPG and donkey anti-goat 546 IgG for RANKL. The slides were dried in an oven, placed in xylene for 1 minute, and sealed in the permount. The immunoreactions were observed under a light microscope (Axio Imager A2, Zeiss) attached to a fluorescent lamp (HBO 100 illuminator, Zeiss) and a charge-coupled device camera (Aziocam 506, Zeiss).

A densitometric analysis to quantify the immunoreactivity of OPG and RANKL was performed by determining the relative optical density (ROD) using Scion ImageJ (National Institutes of Health, Bethesda, MD, USA). The ROD was calculated as a percentage, and the ROD of the control group was designated as 100%.

Statistical analysis

The total sample size of 24 was calculated using the G*Power program on the basis of an effect size of 0.25, a type I error of 0.05, a power of 0.80, and the presence of 4 groups in 1-way repeated-measures analysis of variance. Six samples per group were required.

Non-parametric analysis of variance for repeated measures (the Friedman test) was performed to evaluate changes in GLUs among groups. The Wilcoxon signed-rank test was also performed for pairwise comparisons of time points within each group. The significance of multiple comparisons was adjusted using the Bonferroni method. Data are presented as the mean±standard deviation. *P* values less than 0.05 were considered to indicate statistical significance. All statistical analyses were performed using a software package for statistical analysis (SPSS version 23, IBM Corp., Armonk, NY, USA; Microsoft Excel, Microsoft, Redmond, WA, USA).

RESULTS

Alloxan-induced diabetic rabbits

Of the 22 experimental rabbits, 9 died within 7 days, 5 failed to show elevated blood glucose levels, and 2 died during bone defect surgery. Six rabbits showed elevated blood glucose levels, and were distributed into 3 experimental groups (groups 1, 2, and 3), each comprising 2 rabbits with DM. The rabbits underwent bone defect surgery 1, 2, and 4 weeks after DM induction, respectively. The control group comprised 2 rabbits without DM, which were subjected to bone defect surgery. Eight specimens per group were obtained. On the day of diabetes induction, the mean FBS levels were 149.63±19.38 mg/dL in the diabetic rabbits. FBS levels increased over time, reaching over 500 mg/dL by 4 weeks after the injection. The mean FBS level of diabetic rabbits at 4 weeks after DM induction was 554.75±29.94 mg/dL. The diabetic state of all diabetic rabbits was well-maintained throughout the experimental period (Figure 1).

CBCT scanning, GLU measurements, and 3-dimensional (3D) imaging

Table 1 shows the GLU levels of the groups. The control group showed a mean GLU of 326.25±64.27 at baseline, 685.88±175.56 at 2 weeks after surgery, and 915.25±274.37 at 4 weeks after surgery. Group 1 showed a mean GLU of 460.25±58.19 at baseline, 438.88±141.64




Figure 1. FBS levels of diabetic rabbits. FBS levels over 200 mg/dL were classified as indicating diabetes. All rabbits showed stable FBS throughout the experiment. No rabbits showed relapse of FBS. FBS: fasting blood sugar, W: weeks.

Table 1. Gray level units obtained from cone-beam computed tomography

Group	Baseline	2 weeks later	4 weeks later	P value
Control (n=8)	326.25±64.27 ^{a)}	685.88±175.56 ^{b)}	915.25±274.37 ^{c)}	<0.001
Group 1 (n=8)	460.25±58.19 ^{a)}	438.88±141.64 ^{a)}	463.00±215.74 ^{a)}	>0.05
Group 2 (n=8)	345.38±107.90 ^{a)}	322.25±151.61 ^{a)}	529.38±125.77 ^{a)}	>0.05
Group 3 (n=8)	331.00±45.41 ^{a)}	589.13±174.01 ^{a)}	530.88±197.48 ^{a)}	>0.05

Data are shown as mean±standard deviation.

^{a,b,c)}The same superscript letters indicate values that are not significantly different ($P>0.05$).

at 2 weeks after surgery, and 463.00±215.74 at 4 weeks after surgery. Group 2 showed a mean GLU of 345.38±107.90 at baseline, 322.25±151.61 at 2 weeks after surgery, and 529.38±125.77 at 4 weeks after surgery. Group 3 showed a mean GLU of 331.00±45.41 at baseline, 589.13±174.01 at 2 weeks after surgery, and 530.88±197.48 at 4 weeks after surgery. The change in GLU showed a significant relationship with the experimental group (i.e., to the time elapsed), and the GLU values of the control group were significantly different from those of the 3 experimental groups ($P<0.001$). However, no significant difference was observed among groups 1, 2, and 3. On reconstructed 3D imaging, the control group showed evident bone regeneration, while groups 1, 2, and 3 showed relatively slow bone healing (Figure 2).

Histological analysis

For a morphological analysis of bone defect areas, bone sections were stained with hematoxylin and eosin (H&E) and Masson trichrome. In the control group, the bone defect area was mostly filled with bone tissue (regenerated bone) stained with H&E or Masson trichrome. However, in groups 1, 2 and 3, there was significantly less regenerated bone in the bone defect areas than in the control group. In the experimental groups, the defect areas were mainly filled with connective tissue (Figure 3).

Immunofluorescence analysis

To identify the activated osteoclasts in the bone defect margin, double immunofluorescence staining for RANKL and OPG was performed. In all experimental groups, RANKL and OPG immunoreactivity was observed around the Howship lacuna. However, relative to the optical density of the control samples, RANKL, OPG, and RANKL/OPG immunoreactivity did not show any remarkable change (Figure 4).




Figure 2. Reconstructed 3-dimensional cone-beam computed tomography image. The control group (A) showed relatively rapid and prominent bone regeneration compared to groups 1 (B), 2 (C), and 3 (D).

DISCUSSION

We evaluated the association between the duration of DM and bone-healing capacity using an experimental diabetic rabbit model, and found that DM duration over 1 week could impair bone-healing capacity to some extent. However, we did not identify a link between the magnitude of bone healing impairment and the duration of DM. Since the levels of OPG and RANKL did not significantly differ among groups, osteoclast activity might not contribute to the alteration of bone metabolism in DM.

ALX is one of the most commonly used chemical agents for the induction of DM, and it is used for experimental DM in animals such as rabbits, rats, mice, and dogs. ALX has 2 distinct pathological effects: selective inhibition of glucose-induced insulin secretion through specific inhibition of glucokinase—the glucose sensor of β -cells—and induction of a state of insulin-dependent DM by the induction of reactive oxygen species formation, leading to selective necrosis of β -cells [25]. The major disadvantages of ALX are variability in the response to DM induction and systemic toxicity [26]. Susceptibility to both toxic and diabetogenic doses of ALX varies widely not only across different species, but also among animals of the same species [27]. In our study, the lethality of 100 mg/kg ALX was 52.6%, and the success rate of DM induction was 36.4%. In the literature, lethality rates of 25%–83% and success rates of 16%–50% have been demonstrated with the same dose of ALX [28,29]. When considering both the lethality rate and the DM induction success rate, the dose applied in this study can be regarded as acceptable. This study required 6 samples per group, but only 8 of the 24 rabbits survived. We created 4 bone defects per rabbit to obtain an adequate number of samples.




Figure 3. Hematoxylin and eosin staining and Masson trichrome staining of specimens (scale bar=1 mm). The control group (B) showed notable bone regeneration, whereas groups 1 (C), 2 (D), and 3 (E) showed scarce bone regeneration. (A) baseline. Black arrowheads indicate defect margin. CTL: control, DM: diabetes mellitus, W: weeks.

To evaluate bone healing in live diabetic rabbits in real time, we used GLUs obtained by CBCT measurements. A GLU is a value assigned to the voxels in CBCT. GLUs are not the same as Hounsfield units (HUs), which provide a quantitative assessment of bone density; however, Mah et al. [30] demonstrated that GLUs had a linear relationship with attenuation coefficients, and that HU values could be calculated from GLU values. The GLU values of the control group significantly differed from those of groups 1, 2, and 3. Moreover, since the appearance of GLUs at 4 weeks after surgery corresponded with the histological appearance to some extent, GLUs can be regarded as a practical means to quantify and assess bone healing.

Since an objective of this study was to establish an optimal protocol for the evaluation of the bone-healing capacity of diabetic animals, we studied the effect of varying intervals between DM induction and bone defect induction. As hyperglycemia is usually confirmed 1 week after DM induction, the shortest interval was set at 1 week. Two preclinical studies evaluated the effect of chemically-induced DM on bone-healing capacity [17,24]. Both studies evaluated




Figure 4. Immunofluorescence analysis of OPG and RANKL, as well as merged images (Scale bar = 25 μ m). There were no obvious differences among the groups. The relative optical density was derived from the immunofluorescence analysis. No statistically significant difference was shown among groups. (A) baseline, (B) control group, (C) group 1, (D) group 2, (E) group 3. OPG: osteoprotegerin, RANKL: receptor activator of nuclear factor-kappa B, CTL: control, DM: diabetes mellitus, IR: immunoreactivity, W: weeks.

the extent of damage to bone healing caused by DM depending on the size of cranial defects created about 3 weeks after DM induction, and demonstrated altered bone healing in diabetic animals. Designation of the precise duration of DM before surgical intervention that would be most suitable for expressing alterations in bone healing would be helpful for minimizing the experimental period and planning optimal schedules in future animal studies.

As can be seen in the 3D images obtained by CBCT and through the histological analysis, an interval of at least 1 week between DM induction and the intervention was sufficient for DM to exert a harmful effect on bone healing. Although there were some variations among the sites, bone regeneration was generally much more prominent in the control group than in groups 1, 2, and 3. Most of the regenerated bone was located on the top of the defect, and was misshapen owing to the protrusion of intracranial tissue into the bone defect [31]. Careful effort not to damage the dura mater during bone defect induction would minimize this phenomenon.

In the immunofluorescence analysis, the ROD of OPG and RANKL did not significantly differ among groups. OPG and RANKL have been identified as components of a ligand–receptor system that directly controls osteoclast differentiation and bone resorption. RANK is the cognate receptor of RANKL, and is expressed at high levels in osteoclast precursors [32]. The coupling of RANK and RANKL activates the formation and maturation of osteoclasts [33]. OPG works as a decoy receptor of RANKL and hinders the activation of RANK by selectively binding to RANKL, thereby protecting bone against destruction [34]. For these reasons, OPG and RANKL are considered good markers of osteoclast activity. Our results suggest that altered osteoclast activity does not contribute to altered bone metabolism in DM. In future studies, examination of other possible factors, like osteoblast activity, can be recommended.

New Zealand white rabbits were used in the present study. Many animal studies on DM have used rats as experimental animals. One disadvantage of rodents as experimental animals is that, due to their short life span, they may not develop the clinical complications that manifest in humans under long-term diabetic conditions [35]. In addition, calvarial bone and the other bones of rats are very small, making surgical interventions relatively difficult. Rabbits, however, are adaptable animals for prolonged-duration DM studies [36]. Rabbits are known to have suitable characteristics for experimental animals, such as appropriate size, a longer life span than small rodents, strain specificity, good temper, maneuverability, and relatively low cost [37]. In our study, all the diabetic rabbits survived and maintained elevated FBS throughout the experimental period. This demonstrates the viability and ease of handling of rabbits in DM animal models. Furthermore, the diabetic rabbit model can be used for implant studies due to the sufficient bone volume of the rabbit tibia.

We demonstrated that bone healing slowed when the interval between DM induction and surgical intervention was longer than 1 week. However, we failed to show a relationship between the extent of bone healing impairment and the duration of DM. Some limitations of this study, such as the small number of samples, relatively short period of the study, and high lethality and toxicity of ALX, might have been responsible for this outcome. The dose of ALX used in this study yielded relatively acceptable results; however, further research is needed to determine more precisely the dose of ALX that would maximize the success rate of DM induction and minimize lethality. A role of osteoclasts in altered bone metabolism was not evident. Other possible factors that would reveal the exact mechanisms affecting bone healing in DM should be considered in future studies.

Taken together, the dose of 100 mg/kg of ALX seems to have been an acceptable dose for DM induction in rabbits, and effects of DM on bone healing were demonstrated when the interval between DM induction and intervention was at least 1 week. However, osteoclast activity might not be responsible for altered bone metabolism in DM.

REFERENCES

1. Stavropoulos A, Sculean A, Bosshardt DD, Buser D, Klinge B. Pre-clinical *in vivo* models for the screening of bone biomaterials for oral/craniofacial indications: focus on small-animal models. *Periodontol 2000* 2015;68:55-65.
[PUBMED](#) | [CROSSREF](#)
2. Kantarci A, Hasturk H, Van Dyke TE. Animal models for periodontal regeneration and peri-implant responses. *Periodontol 2000* 2015;68:66-82.
[PUBMED](#) | [CROSSREF](#)
3. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997;20:1183-97.
[PUBMED](#) | [CROSSREF](#)
4. Korean Diabetes Association. Korean diabetes fact sheet 2015 [Internet]. Seoul: Korean Diabetes Association, c2011 [cited 2016 Aug 18]. Available from: <http://www.diabetes.or.kr/>.
5. Goodman WG, Hori MT. Diminished bone formation in experimental diabetes. Relationship to osteoid maturation and mineralization. *Diabetes* 1984;33:825-31.
[PUBMED](#) | [CROSSREF](#)
6. Nyomba BL, Verhaeghe J, Thomasset M, Lissens W, Bouillon R. Bone mineral homeostasis in spontaneously diabetic BB rats. I. Abnormal vitamin D metabolism and impaired active intestinal calcium absorption. *Endocrinology* 1989;124:565-72.
[PUBMED](#) | [CROSSREF](#)

7. Shires R, Teitelbaum SL, Bergfeld MA, Fallon MD, Slatopolsky E, Avioli LV. The effect of streptozotocin-induced chronic diabetes mellitus on bone and mineral homeostasis in the rat. *J Lab Clin Med* 1981;97:231-40.
[PUBMED](#)
8. Morris HF, Ochi S, Winkler S. Implant survival in patients with type 2 diabetes: placement to 36 months. *Ann Periodontol* 2000;5:157-65.
[PUBMED](#) | [CROSSREF](#)
9. Moy PK, Medina D, Shetty V, Aghaloo TL. Dental implant failure rates and associated risk factors. *Int J Oral Maxillofac Implants* 2005;20:569-77.
[PUBMED](#)
10. Nevins ML, Karimbux NY, Weber HP, Giannobile WV, Fiorellini JP. Wound healing around endosseous implants in experimental diabetes. *Int J Oral Maxillofac Implants* 1998;13:620-9.
[PUBMED](#)
11. Giglio MJ, Giannunzio G, Olmedo D, Guglielmotti MB. Histomorphometric study of bone healing around laminar implants in experimental diabetes. *Implant Dent* 2000;9:143-9.
[PUBMED](#) | [CROSSREF](#)
12. Lalla E, Lamster IB, Drury S, Fu C, Schmidt AM. Hyperglycemia, glycoxidation and receptor for advanced glycation endproducts: potential mechanisms underlying diabetic complications, including diabetes-associated periodontitis. *Periodontol* 2000 2000;23:50-62.
[PUBMED](#) | [CROSSREF](#)
13. Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M, Knowler WC, et al. Non-insulin dependent diabetes mellitus and alveolar bone loss progression over 2 years. *J Periodontol* 1998;69:76-83.
[PUBMED](#) | [CROSSREF](#)
14. Balint E, Szabo P, Marshall CF, Sprague SM. Glucose-induced inhibition of *in vitro* bone mineralization. *Bone* 2001;28:21-8.
[PUBMED](#) | [CROSSREF](#)
15. Terada M, Inaba M, Yano Y, Hasuma T, Nishizawa Y, Morii H, et al. Growth-inhibitory effect of a high glucose concentration on osteoblast-like cells. *Bone* 1998;22:17-23.
[PUBMED](#) | [CROSSREF](#)
16. Brownlee M, Cerami A, Vlassara H. Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 1988;318:1315-21.
[PUBMED](#) | [CROSSREF](#)
17. Santana RB, Xu L, Chase HB, Amar S, Graves DT, Trackman PC. A role for advanced glycation end products in diminished bone healing in type 1 diabetes. *Diabetes* 2003;52:1502-10.
[PUBMED](#) | [CROSSREF](#)
18. Lalla E, Lamster IB, Schmidt AM. Enhanced interaction of advanced glycation end products with their cellular receptor RAGE: implications for the pathogenesis of accelerated periodontal disease in diabetes. *Ann Periodontol* 1998;3:13-9.
[PUBMED](#) | [CROSSREF](#)
19. Brussee V, Guo G, Dong Y, Cheng C, Martinez JA, Smith D, et al. Distal degenerative sensory neuropathy in a long-term type 2 diabetes rat model. *Diabetes* 2008;57:1664-73.
[PUBMED](#) | [CROSSREF](#)
20. Diabetes Control and Complications Trial Research Group Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977-86.
[PUBMED](#) | [CROSSREF](#)
21. Kamiya H, Zhang W, Sima AA. Degeneration of the Golgi and neuronal loss in dorsal root ganglia in diabetic BioBreeding/Worcester rats. *Diabetologia* 2006;49:2763-74.
[PUBMED](#) | [CROSSREF](#)
22. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837-53.
[PUBMED](#) | [CROSSREF](#)
23. Follak N, Klötting I, Wolf E, Merk H. Histomorphometric evaluation of the influence of the diabetic metabolic state on bone defect healing depending on the defect size in spontaneously diabetic BB/OK rats. *Bone* 2004;35:144-52.
[PUBMED](#) | [CROSSREF](#)
24. Shyng YC, Devlin H, Sloan P. The effect of streptozotocin-induced experimental diabetes mellitus on calvarial defect healing and bone turnover in the rat. *Int J Oral Maxillofac Surg* 2001;30:70-4.
[PUBMED](#) | [CROSSREF](#)

25. Lenzen S. The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia* 2008;51:216-26.
[PUBMED](#) | [CROSSREF](#)
26. Lackey RW, Bunde CA, Gill AL, Harris LC. Glycogen in alloxan-treated rats. *Proc Soc Exp Biol Med* 1944;57:191-4.
[CROSSREF](#)
27. Zhao ZH, Watschinger B, Brown CD, Beyer MM, Friedman EA. Variations of susceptibility to alloxan induced diabetes in the rabbit. *Horm Metab Res* 1987;19:534-7.
[PUBMED](#) | [CROSSREF](#)
28. Nath MC, Gadgil JS, Hatwalne VG. Studies on alloxan diabetes. I. Increase of susceptibility caused by acetoacetate. *Biochem J* 1953;53:481-3.
[PUBMED](#) | [CROSSREF](#)
29. Hadour G, Ferrera R, Sebbag L, Forrat R, Delaye J, de Lorgeril M. Improved myocardial tolerance to ischaemia in the diabetic rabbit. *J Mol Cell Cardiol* 1998;30:1869-75.
[PUBMED](#) | [CROSSREF](#)
30. Mah P, Reeves TE, McDavid WD. Deriving Hounsfield units using grey levels in cone beam computed tomography. *Dentomaxillofac Radiol* 2010;39:323-35.
[PUBMED](#) | [CROSSREF](#)
31. Greenwald JA, Mehrara BJ, Spector JA, Chin GS, Steinbrech DS, Saadeh PB, et al. Biomolecular mechanisms of calvarial bone induction: immature versus mature dura mater. *Plast Reconstr Surg* 2000;105:1382-92.
[PUBMED](#)
32. Hsu H, Lacey DL, Dunstan CR, Solovyev I, Colombero A, Timms E, et al. Tumor necrosis factor receptor family member RANK mediates osteoclast differentiation and activation induced by osteoprotegerin ligand. *Proc Natl Acad Sci U S A* 1999;96:3540-5.
[PUBMED](#) | [CROSSREF](#)
33. Horowitz MC, Xi Y, Wilson K, Kacena MA. Control of osteoclastogenesis and bone resorption by members of the TNF family of receptors and ligands. *Cytokine Growth Factor Rev* 2001;12:9-18.
[PUBMED](#) | [CROSSREF](#)
34. Crotti T, Smith MD, Hirsch R, Soukoulis S, Weedon H, Capone M, et al. Receptor activator NF kappaB ligand (RANKL) and osteoprotegerin (OPG) protein expression in periodontitis. *J Periodontal Res* 2003;38:380-7.
[PUBMED](#) | [CROSSREF](#)
35. Rees DA, Alcolado JC. Animal models of diabetes mellitus. *Diabet Med* 2005;22:359-70.
[PUBMED](#) | [CROSSREF](#)
36. Wang J, Wan R, Mo Y, Zhang Q, Sherwood LC, Chien S. Creating a long-term diabetic rabbit model. *Exp Diabetes Res* 2010;2010:289614.
[PUBMED](#) | [CROSSREF](#)
37. Mir SH, Darzi MM. Histopathological abnormalities of prolonged alloxan-induced diabetes mellitus in rabbits. *Int J Exp Pathol* 2009;90:66-73.
[PUBMED](#) | [CROSSREF](#)