

Cost of a 5-year lung cancer survivor: symptomatic tumour identification vs proactive computed tomography screening

AW Castleberry¹, D Smith², C Anderson³, AJ Rotter² and FW Grannis Jr^{*,2}

¹University of Pittsburgh School of Medicine, Pittsburgh, PA, USA; ²Department of Thoracic Surgery, City of Hope National Medical Center, Duarte, CA, USA and ³University of California San Diego, San Diego, CA, USA

BACKGROUND: Our objective was to analyse the cost effectiveness of computed tomography (CT) screening for lung cancer in terms of the cost per long-term survivor, which has not been evaluated to date.

METHODS: Estimations were computed based on data from the Surveillance, Epidemiology, and End Results registries covering years 1999–2003. The design framework of our model allowed for the incorporation of multiple values taken from the epidemiological and clinical literature to be utilised for cost inputs, scope of patients screened, diagnostic staging, and survival percentages applied separately to two cohorts: age 40–79 and 60–79 years. This enabled the analysis of over 1400 scenarios, each containing a unique set of input values, for which the estimated cost per 5-year survivor (CP5YS) was compared between the symptom-detected and proactive screening approaches.

RESULTS: Estimated CP5YS were higher for the symptom-detected approach in all 729 scenarios analysed for the cohort ages 60–79 years, ranging from approximately \$5800 to \$116700 increased cost per 5-year survivor (CP5YS). For the cohort ages 40–79 years, 75% of the 729 scenarios analysed showed increased CP5YS for the symptom-detected approach ranging from \$5700 to \$110000 increased CP5YS. Total costs and total 5-year survivors were higher for the proactive screening method for all scenarios analysed across both cohorts with increases ranging from 50–256% and 98–309%, respectively.

CONCLUSION: The predicted increase in long-term survival with CT screening and the potential for better utilisation of health-care dollars in terms of CP5YS, particularly when screening patients over the age of 60 years, are critically important considerations in directing effective future lung cancer management strategy.

British Journal of Cancer (2009) **101**, 882–896. doi:10.1038/sj.bjc.6605253 www.bjcancer.com

Published online 18 August 2009

© 2009 Cancer Research UK

Keywords: lung cancer; screening; comparative effectiveness research; computerised tomography; cost-effectiveness; QALY

Lung cancer is the leading cause of cancer deaths nationwide. An estimated 162 000 people would have died of lung cancer in the United States in 2006 (American Cancer Society, 2006). Currently, only 16% of lung cancers are diagnosed when the disease is still localised and the 5-year survival for all stages combined is 15% (Jemal *et al*, 2005).

Because lung cancer is seldom symptomatic in early stage and treatment in advanced stage has very low survival (Mountain, 1997), current research has focused on methods to detect lung cancer at an earlier stage where curative treatment can be offered.

Results from the International Early Lung Cancer Action Project (I-ELCAP) demonstrating striking improvement in early stage detection of lung cancer by low-radiation-dose computed tomography (CT) screening (Henschke *et al*, 1999; Henschke *et al*, 2001) have sparked debate on the cost-effectiveness of a CT screening programme, with widely variant results reported (Marshall *et al*, 2001a; Marshall *et al*, 2001b; Chirikos *et al*, 2002; Mahadevia *et al*,

2003; Wisnivesky *et al*, 2003). In a systemic review published in January 2006 (Black *et al*, 2006), the authors conclude that many issues remain unresolved in the debate over the true cost-effectiveness of lung cancer screening programmes. Attention to this issue has further escalated following the reported results of over 30 000 persons screened in the large collaborative I-ELCAP study, which concluded that annual spiral CT screening can detect lung cancer that is curable in most cases (Henschke *et al*, 2006a). In addition, an analysis of the mean sojourn time and sensitivity of CT screening for lung cancer estimated approximately 23% mortality reduction possible by an annual CT screening programme opposed to observation (Chien and Chen, 2008). Knudsen *et al* (2007) have emphasised the need for mathematical models that examine the cost effectiveness of implementation of population CT screening.

Many investigations of cancer treatment cost-effectiveness utilise quality-adjusted life year (QALY) as the primary metric to measure the results of interventions that provide some improvement in quality and/or duration of life, but only for limited periods of time. For example, a cancer treatment that results in improvement in duration of survival by 3 months at cost of \$25 000 would result in a cost per year or QALY of approximately \$100 000, as it requires four patients, each experiencing a 3-month improvement in survival to add up to a year of life gained.

*Correspondence: FW Grannis Jr., Department of Thoracic Surgery, City of Hope National Medical Center, 1500 East Duarte Road, Duarte, CA 91010, USA; E-mail: fgrannis@coh.org
Received 7 January 2009; revised 8 July 2009; accepted 21 July 2009; published online 18 August 2009

Although this metric has the advantage of wide utilisation, in a certain sense, it is a misleading figure, as in this example no single patient would actually survive 1 year. Preventive screening measures, where the positive impact on health outcomes may not be measurable for many years, may be difficult to quantify using QALYs (Phillips and Thompson, 2003). Arguably, the most important consideration in determining cost effectiveness for differing approaches to lung cancer is the number of long-term survivors attained by a given strategy. Accordingly, we believe that the cost of a long-term survival is more meaningful to the reader, and is particularly more important to patients at high-risk of lung cancer. The purpose of this study is to evaluate the estimated cost-effectiveness of a proactive screening programme compared with current lung cancer strategies when applied to a representative subset of the US population. We compared the ratio of 5-year survivors to the total screening, diagnosis, and treatment costs for the current management algorithm that we will designate as the symptomatic tumour identification (STID) approach and for an early detection approach using CT screening (EDCTS). Second, we report the estimated total costs and total 5-year survivors associated with each strategy. The EDCTS protocol is based on I-ELCAP recommendations (Henschke, 2006). We separately applied this analysis to two cohorts distinguished by age. One cohort was in the age of 40–79 years and the second cohort was in the age of 60–79 years.

MATERIALS AND METHODS

We constructed a mathematical model based on multivariate analysis of empirical data to estimate the ratio of screening, diagnosis, and treatment costs per 5-year non-small-cell lung cancer (NSCLC) survivor for two different lung cancer management strategies, STID and EDCTS. These estimations were computed in 2005 USD based on the most recent 5 years of data available, extending from 1999 to 2003, within the 13 Surveillance, Epidemiology, and End Results (SEER) data registries (National Cancer Institute, 2005). With respect to selected demographic and epidemiologic factors, these areas are a reasonably representative subset of the US population (Ries *et al*, 2005). The scope of our analysis covered a population of approximately 15 million people.

The design framework of our model allowed for the incorporation of multiple values taken from the epidemiological and clinical literature to be utilised for cost inputs, scope of patients screened, diagnostic staging, and survival percentages. This methodology enabled the analysis of over 700 scenarios for each of the two cohorts (for a total of over 1400 scenarios), each containing a unique set of input values. For each of these scenarios, the estimated CP5YS was compared between the STID and EDCTS approaches. By analysing a broad range of parameter values, we were able to evaluate the sensitivity of selected variables by using a range of favourable and unfavourable estimates. In addition, this study design also enabled the evaluation of completely separate parameter estimates and calculations from different sources of published data. Our model assumes that the capital equipment and resources necessary for both management strategies are already in place.

Input variables and source data

Population screened. The initial I-ELCAP screening study published in 1999 screened at-risk persons aged 60 years or older (Henschke *et al*, 1999). The subsequent large collaborative I-ELCAP study published in 2006 screened at-risk persons aged 40 and older (Henschke *et al*, 2006a). Approximately 98.3% of the 31 567 persons screened in the large collaborative I-ELCAP study were under the age of 80 years. As such, we analysed two separate cohorts: one cohort was of the age 40–79 years and the second cohort was of the age 60–79 years. The population screened for

lung cancer in our analysis was based on inclusion criteria from I-ELCAP reported studies, which screened patients with a history of at least 10 pack-years of cigarette smoking, no history of cancer (other than non-melanoma skin cancer), and who were fit to undergo thoracic surgery (Henschke *et al*, 1999). We estimated this population using the following steps:

First, we determined the total population per the 13 SEER data registries from 1999–2003 for each of the two cohorts. This total population was reduced by 0.45% to reflect the incidence of cancer other than non-melanoma skin cancer per the SEER Cancer Statistics Review (Ries *et al*, 2005).

Second, we estimated the percentage of US adults in each cohort with at least a 10 pack-year history of smoking, regardless of current smoking status, using figures reported by the 2003 National Cancer Institute and Centers for Disease Control and Prevention Co-sponsored Tobacco Use Special Cessation Supplement to the Current Population Survey (CPS) (National Cancer Institute, 2003). To evaluate the sensitivity of this parameter and incorporate a range of variability in actual smoking history, a 10% increase/decrease in the CPS percentage was also analysed as upper and lower bound estimates, respectively (Table 1).

We then estimated the percent of this population that would be unfit for surgery by using SEER data to determine the percent of stage I tumours diagnosed in patients aged 40–79 and 60–79, respectively for each cohort, in whom surgery was not recommended or was contraindicated (Criteria: lung and bronchus NSCLC, invasive, microscopically confirmed, the patient's first primary cancer, and diagnosed between 1999 and 2003. Cases where surgery was recommended and the patient refused the procedure were counted the same as cases where surgery was performed. Cases where it was unknown whether surgery was recommended were omitted from the analysis). This resulted in an estimate of 14.7% for the cohort ages 40–79 years who are excluded from screening because they would not be fit for surgery and 17.0% for the cohort ages 60–79 years. It is important to emphasise that some individuals who would not be the candidates for standard surgical procedures might still receive potentially curative treatment with minimally invasive surgical approaches, radiation therapy, radiofrequency ablation, or other methods.

Using steps (1), (2), and (3) described above, we quantified an estimate of the total population that would be eligible for screening in each cohort in accordance with I-ELCAP inclusion criteria, with an upper and lower bound estimate for the percent of the population with a 10 pack-year history of smoking (Table 1). Statistics on compliance with breast cancer screening guidelines from three separate studies (Horton *et al*, 1996; Phillips *et al*, 1998; Rahman *et al*, 2003) were used to estimate the compliance with lung cancer screening recommendations (Table 1). Multiplying the SEER population meeting the inclusion criteria by the estimated compliance percent provides estimates of the total population that would receive the screening procedures (Table 2).

The choice of different inclusion criteria (e.g., older age at entry, or higher/lower pack-years requirement) would result in changes in number of individuals screened, costs, and results.

Baseline screening vs annual repeat screening. In the I-ELCAP studies, the percentage of patients requiring additional tests subsequent to the initial CT scan was different in the first year (baseline screen) than in annual repeat screening. In addition, the percent of cases where malignancy was found was also different for baseline screens vs annual repeat screens (Henschke *et al*, 1999; Henschke *et al*, 2001; Henschke *et al*, 2006a). For the purpose of our estimation of screening costs and malignancies detected, the 'Population Screened' described above for each year was divided into groups reflecting those receiving a baseline screen vs those receiving repeat annual screens. This was carried out based on the large collaborative I-ELCAP study, in which 31 567 persons received a baseline screen with 27 456 (87.0%) following up with

Table 1 Summary of input variables and source data^a

	Parameter values			Source		
	Input 1	Input 2	Input 3	Input 1	Input 2	Input 3
<i>Percent of US adults with a 10 pack-year smoking history %</i>						
Age 40–79 ^b	28.4	31.3	25.6	NCI (2003)	Estimate	Estimate
Age 60–79	32.3	35.5	29.1	NCI (2003)	Estimate	Estimate
	(A)	(B)	(C)			
Compliance with screening recommendations (%)	47.4	27.0	41.4	Horton et al (1996)	Phillips et al (1998)	Rahman et al (2003)
	(D)	(E)	(F)			
<i>Frequency (%) and type of screening procedure carried out</i>						
Baseline Screen						
MDCT		100.00				
Follow-up MDCT	—	8.40	—	—	Henschke et al. (1999, 2006b)	—
PET scan		2.20				
FNA biopsy		1.30				
<i>Annual repeat screen</i>						
MDCT		100.00				
Follow-up MDCT	—	2.96	—	—	Henschke et al. (2001, 2006b)	—
PET scan		0.59				
FNA biopsy		0.59				
<i>Percent of screened patients with lung cancer</i>						
<i>Baseline year</i>						
Age 40–79	—	1.3	—	—	Henschke et al (2006a)	—
Age 60–79	—	2.7	—	—	Henschke et al (1999)	—
<i>Years following baseline year</i>						
Age 40–79	—	0.3	—	—	Henschke et al (2006a)	—
Age 60–79	—	0.6	—	—	Henschke et al (2001)	—
<i>Screening procedure costs</i>						
MDCT	\$204	\$178	\$533			
Follow-up MDCT	\$204	\$178	\$533	Wisnivesky et al (2003)	Mahadevia et al (2003)	Mahadevia et al (2003)
FNA biopsy	\$620	\$355	\$498			
PET scan	\$1,150	\$1,150	\$3,018	2005 Medicare Rate	2005 Medicare Rate	Kelly et al (2004)
	(G)	(H)	(I)			
<i>Percent distribution of screened tumours by stage</i>						
Stage I	85	64	74			
Stage II	4	27	16			
Stage IIIA	7	9	8	Henschke et al (1999)	Swensen et al (2002)	Average
Stage IIIB	4	0	2			
Stage IV	0	0	0			
	(J)	(K)	(L)			
<i>Cost of diagnosis and treatment by stage (diagnosis to death)</i>						
Stage I	\$70 033	\$79 285	\$40 548			
Stage II	\$47 139	\$64 456	\$45 249			
Stage IIIA	\$47 139	\$64 456	\$41 933	Riley et al (1995)	Fireman et al (1997)	Evans et al (1995)
Stage IIIB	\$47 139	\$64 456	\$36 694			
Stage IV	\$31 814	\$49 500	\$31 265			
	(M)	(N)	(O)			
<i>Percent 5-year survivors by stage</i>						
Stage I	67	88	82		Henschke et al (2006a)	Patz et al (2000)
Stage II	55	23	23			
Stage IIIA	23	11	11	Mountain et al 1997	Fry et al (1999)	Fry et al (1999)
Stage IIIB	5	5	5			
Stage IV	1	1	1			
	(P)	(Q)	(R)			

Abbreviations: FNA = fine-needle aspiration; MDCT = multidetector computed tomography; PET = positron emission tomography. All costs in 2005 USD. ^aParentetical references are used to identify the combination of variables used for a given scenario. For example, ADGJMF would refer to a scenario in with the input values indicated by parenthetical reference (A), (D), (G), and so on. ^bThe data input listed above apply to both the cohort ages 40–79 years and the cohort aged 60–79 years unless specifically stated otherwise.

an annual repeat screen (Henschke et al, 2006a). This reflects a dropout rate (i.e., the percent of patients who receive a baseline scan and do not follow up with annual repeat scans) of 13%. This

information was used to estimate baseline and repeat screens as follows: (1) in year one, all screens are baseline screens; (2) in all subsequent years, 87.0% of the patients screened in the earlier year

Table 2 SEER population estimated to receive screening for lung cancer based on inclusion criteria from I-ELCAP reported studies, nationwide smoking history statistics, and estimated compliance with screening recommendations^a

	Year 1	Year 2	Year 3	Year 4	Year 5
SEER population age 40–79 years ^b	14 277 295	14 596 718	14 944 483	15 262 435	15 578 827
Less: cancer other than non-melanoma skin cancer (see text)	64 248	65 685	67 250	68 681	70 105
	14 213 047	14 531 033	14 877 233	15 193 754	15 508 722
Less: estimated patients unfit for surgery (see text)	2 089 318	2 136 062	2 186 953	2 233 482	2 279 782
	12 123 729	12 394 971	12 690 280	12 960 272	13 228 940
Less: <10 pack-year smoking history or nonsmoker (see text) ^a	8 675 948	8 870 053	9 081 381	9 274 592	9 466 855
	3 447 782	3 524 918	3 608 899	3 685 680	3 762 085
Less: non-compliance with screening recommendation (see text) ^a	1 813 533	1 854 107	1 898 281	1 938 668	1 978 857
Total screened population	1 634 248	1 670 811	1 710 618	1 747 012	1 783 228

^aThe values shown above for <10 pack-year smoking history and compliance with screening recommendations are based on the 'Input 1' column in Table 1. For simplicity, values generated using the columns labelled 'Input 2' and 'Input 3' values are not shown here. ^bFor simplicity, the values associated with the cohort ages 60–79 years are not shown here.

Table 3 Breakdown of patients receiving a baseline screen vs repeat annual screen in each year analysed

	Year 1	Year 2	Year 3	Year 4	Year 5
(A) Total screened population Age 40–79 (see Table 2) ^{ab}	1 634 248	1 670 811	1 710 618	1 747 012	1 783 228
<i>Breakdown of the above total population screened (A) into baseline screens and annual repeat screens based on assumptions outlined in the section 'Baseline screening vs annual repeat screening' (see text):</i>					
(B) Dropouts to screening in subsequent years (= 13% × (A))		212 452	222 380	227 112	231 820
(C) Patients who will get a repeat screen in the next year (= (A)–(B))	1 421 796	1 453 606	1 488 238	1 519 901	1 551 408
(D) Number of screens that are repeat annual screens (= (C) from earlier year)	NA	1 421 796	1 453 606	1 488 238	1 519 901
(E) Number of screens that represent new patients receiving baseline screens (= (A)–(D))	(All)	249 015	257 012	258 775	263 327
<i>Summary of the breakout of baseline vs annual repeat screens per year:</i>					
Patients receiving baseline screen (all in year 1; = (E) in subsequent years):	1 634 248	249 015	257 012	258 775	263 327
Patients receiving annual repeat screen (none in year 1; = (D) in subsequent years):	0	1 421 796	1 453 606	1 488 238	1 519 901
Total Screened Population (sum of baseline and annual repeat screens):	1 634 248	1 670 811	1 710 618	1 747 012	1 783 228

^aThe values shown above for total screened population are based on the 'Input 1' column in Table 1. For simplicity, values generated using the columns labelled 'Input 2' and 'Input 3' values are not shown here. Refer to Table 1 and Table 2. ^bFor simplicity, the values associated with the cohort ages 60–79 years are not shown here.

are assumed to receive repeat screens with the remaining patients screened in that year assumed to be new patients receiving a baseline screen (see Table 3).

Frequency and type of screening procedures carried out. The procedures carried out as part of the baseline and annual repeat screens were based on I-ELCAP protocol, revision dated 20 October 2006 (Henschke, 2006). The I-ELCAP recommendations for the diagnostic workup in participants with a positive result on the CT include multiple options for nodules meeting certain size criteria. The protocol used in the I-ELCAP research was reviewed semi-annually and modified over time and with increasing experience in the analysis of the study data, which likely impacted the volume of procedures indicated in years subsequent to the baseline year because of a temporal, learning effect. In addition, in the I-ELCAP studies, the decision regarding how to proceed is left to each participant and the referring physician (Henschke *et al*, 2006a). As such, the screening procedure protocol utilised in this analysis (referred to as 'EDCTS Screening Protocol') may not precisely reflect the I-ELCAP protocol or screening procedures presented in I-ELCAP studies. The EDCTS Screening Protocol identifies additional workup procedures which, based on the size of nodules identified in the initial CT scan, would be included in a patient's screening costs. We derived the percentage of total screened patients receiving each additional workup procedure based on the size distribution of screen tumours reported in the I-ELCAP baseline and repeat screening studies (Henschke *et al*, 1999; Henschke *et al*, 2001) (Tables 4 and 5). A summary of the percent of screening participants receiving each type of screening procedure is provided in Table 1. Workup procedures that also

involve treatment of the malignancy, such as video-assisted thoracoscopic surgical biopsy, are considered as treatment procedures and are not included as part of screening procedures in this analysis.

Screen-detected malignant tumours. For the cohort ages 60–79 years, the number of malignant tumours identified by screening in the baseline and annual repeat screens was derived by multiplying the percent of malignant tumours identified in the I-ELCAP studies published in 1999 and 2001 that screened patients aged 60 years and over (Henschke *et al*, 1999; Henschke *et al*, 2001) by the total population screened (Table 1). For the cohort ages 40–79 years, the percentage of malignant tumours identified in the large collaborative I-ELCAP study published in 2006 that screened patients aged 40 years and over was used (Henschke *et al*, 2006a) (Table 1).

Screening costs. For the EDCTS approach, our model incorporated three different screening procedure cost estimates for multidetector CT (MDCT) scans and fine-needle aspiration biopsies (FNAB), and two cost estimates for positron emission tomography (PET) scans taken from the clinical literature (published costs for conventional CT scans were used for MDCT costs, as there is typically no difference in the cost to the payor). The first set of estimates for CT and FNAB were based on the actual costs incurred by I-ELCAP volunteers, as recorded by the New York-Presbyterian Hospital's financial system cost database (Wisnivesky *et al*, 2003). The second and third sets of estimates were based on 2003 published figures from a cost effectiveness study by Mahadevia *et al* (2003). The Mahadevia *et al* (2003) study

Table 4 Percentage of patients receiving each screening procedure in the baseline screen

I-ELCAP baseline screening results ^a			EDCTS screening procedures that would be applied to the nodules sizes identified by I-ELCAP	
	Patients	% of total		
(A) Patients receiving initial CT scan:	1000	100	(A)	Receive MDCT
<i>Positive non-calcified nodules identified:</i>				
(B) Nodules <= 5 mm	136	13.6	(B)	MDCT in 1 year ^b
(C) Nodules 6–10 mm, negative upon workup	56	5.6	(C)	MDCT in 3 months, assume no growth and therefore additional MDCT in 1 year ^b
(D) Nodules 6–10 mm, positive upon workup	14	1.4	(D)	MDCT in 3 months, assume growth and therefore begin treatment ^b
(E) Nodules 11–20 mm, negative upon workup	14	1.4	(E)	Receive PET, assume negative and therefore additional MDCT in 3 months
(F) Nodules 11–20 mm, positive upon workup	8	0.8	(F)	Receive PET, assume positive and therefore perform FNAB
(G) Nodules >20 mm	5	0.5	(G)	All receive FNAB

Abbreviations: EDCTS = early detection approach using computed tomography screening; FNAB = fine-needle aspiration biopsy; MDCT = multidetector computed tomography; PET = positron emission tomography. ^aReference Henschke et al (1999). ^bFor purposes of this screening protocol, FNAB and PET are not utilised for nodules ≤ 10 mm in size.

Table 5 Percentage of patients receiving each screening procedure in the annual repeat screen

I-ELCAP annual repeat screening results ^a			EDCTS screening procedures that would be applied to the nodules sizes identified by I-ELCAP	
	Patients	% of total		
(A) Patients receiving initial CT scan:	1184	100	(A)	Receive MDCT
<i>Positive non-calcified nodules identified:</i>				
(B) Nodules ≤ 5 mm, negative upon workup	7	0.6	(B)	MDCT in 3 months, assume no growth ^b
(C) Nodules ≤ 5 mm, positive upon workup or not ruled out	9	0.8	(C)	MDCT in 3 months, assume growth and therefore patient receives treatment (no further screening procedures) ^b
(D) Nodules 6–10 mm, negative upon workup	7	0.6	(C)	MDCT in 3 months, assume no growth and therefore additional MDCT in 1 year ^b
(D) Nodules 6–10 mm, positive upon workup or not ruled out	9	0.8	(D)	MDCT in three months, assume growth and therefore begin treatment ^b
(E) Nodules 11–20 mm, negative upon workup	3	0.3	(F)	Receive PET, assume negative and therefore additional MDCT in 3 months
(F) Nodules 11–20 mm, positive upon workup or not ruled out	4	0.3	(E)	Receive PET, assume positive and therefore perform FNAB
(G) Nodules >20 mm	3	0.3	(G)	All receive FNAB

Abbreviations: EDCTS = early detection approach using computed tomography screening; FNAB = fine-needle aspiration biopsy; MDCT = multidetector computed tomography; PET = positron emission tomography. ^aReference Henschke et al (2001). ^bFor purposes of this screening protocol, FNAB and PET are not utilised for nodules ≤ 10 mm in size.

reported base-case costs for lung cancer screening procedures as well as favourable and unfavourable extremes. Values from both the favourable and unfavourable scenarios were utilised in our model (Table 1). The first estimate of PET scan costs was taken from 2005 Medicare Reimbursement rates, and the second cost estimate (unfavourable screening estimate) was taken from a 2004 published cost-effectiveness study by Kelly et al (2004) (Table 1). We did not consider the potential costs or implications of using PET-CT, as this was not included in the EDCTS Screening Protocol (see above). All costs were calibrated to 2005 US dollars using the medical component of the consumer price index. Screening costs for the STID method were assumed to be zero.

Quantity and distribution of tumours by stage. Tumour quantity and distribution by stage for the STID method was based on SEER data from 1999 to 2003, using lung and bronchus NSCLC that were invasive, microscopically confirmed, and the patient's first primary cancer. SEER tumours of unknown stage were allocated to stages based on the distribution percentages of the staged SEER tumours.

The quantity of tumours for the EDCTS method was comprised of two components: (1) screen-detected malignant tumours (quantification described above); and (2) non-screen-detected tumours ('symptomatic tumours'). For the first year of screening implementation it was assumed that the number of symptomatic tumours would include all SEER reported tumours for that year. For subsequent years, symptomatic tumours were calculated by

multiplying the percent of the population that did not comply with screening recommendations (based on estimates described for 'Population Screened' above) to the SEER tumour population. For example, if the SEER data identified 1000 tumours in a given year, and an estimated 60% of the population did not comply with screening recommendations, then it was assumed that 600 (= 1000 × 60%) symptomatic tumours would be recorded in that year in addition to screen-detected tumours. Components (1) and (2) were added together to determine the total EDCTS tumours identified each year.

The stage of EDCTS tumours was determined as follows: (1) Symptomatic tumours, as described above, maintain the SEER stage distribution, with the same treatment for tumours of unknown stage as described above for the STID method. (2) Three estimations were used to determine the number of remaining screen-detected malignant tumours allocated to each stage. The first estimation was based on I-ELCAP data (Henschke et al, 1999), the second was based on a prospective cohort CT screening study by Swensen et al (2002), and the final estimation was an average of the I-ELCAP and Swensen stage distribution percentages (Table 1).

Cost of treatment by stage. For this parameter, we included separate estimates and calculations from different sources of published data on the cost of diagnosis and treatment for lung cancer by stage. These sources included data based on Medicare payments for lung cancer patients in nine SEER registries (Riley et al, 1995), cost of medical care for patients of a Northern

California health maintenance organisation (Fireman *et al*, 1997), and costs under Canada's universal health-care system (Evans *et al*, 1995). A previous study by Wisnivesky *et al* (2003) noted that the overall distribution of the costs of lung cancer treatments by stage were similar in the United States and Canada. These estimates included the total lung cancer cost by stage from diagnosis to death, adjusted to 2005 US dollars (Table 1).

Total screening, diagnosis, and treatment costs. For the EDCTS approach, total costs were comprised of screening cost and the cost of treatment by stage. However, calculations for the cost of treatment by stage included diagnosis and staging procedures that are also covered in the screening protocol. To avoid double-counting these expenditures, the screening costs for patients diagnosed with malignant lung cancer were removed from the total costs, as these costs are already included in the cost of treatment by stage. As such, the resulting total costs for the EDCTS approach includes the sum total of screening costs for patients not diagnosed with malignant cancer (over 97% of total screening costs) and the cost of treatment by stage (including diagnostic costs) for the entire tumour population, including screen-detected and non-screen-detected tumours. Total costs for the STID approach were limited to the cost of treatment by stage with screening costs assumed to be zero.

Five-year survivors. Survival rates were applied to the population of tumours by stage to determine the estimated number of associated 5-year survivors. The same survival rates were used for both the STID and EDCTS calculations. Three different sets of survival rate estimates were included in our analysis. The first set of estimates was based on published data on over 5300 lung cancer cases compiled from The University of Texas MD Anderson Cancer Center and the classification research database from the Reference Center for Anatomic and Pathologic Classification of Lung Cancer (survival based on death from cancer or unknown cause) (Mountain, 1997). For the second set of estimates, stage I survival was based on I-ELCAP 10-year survival results (lung cancer-specific survival; 10-year survival data were used as a conservative estimate of 5-year survival) (Henschke *et al*, 2006a), with stage II through stage IV based on 713 043 primary lung malignancies submitted to the National Cancer Data Base (relative survival) (Fry *et al*, 1999). The final set of estimates used the same

National Cancer DataBase rates for stage II through stage IV disease, with the stage IA survival rate from a 2000 study published by Patz *et al* (2000), applied to the stage I tumour populations (all-cause survival) (Table 1).

Ratio of total CP5YS for 729 scenarios

Using the inputs described above, 729 scenarios were generated by combining all possible combinations of the variables containing multiple input values in Table 1. For example, the percent of the population eligible for screening (≥ 10 pack-year history of smoking) has three possible input values: (A), (B), and (C). Similarly, the estimated compliance percentage has three possible input values: (D), (E), and (F). Each combination of these values was used in a separate scenario examined (AD, AE, AF, BD, and so on). This method was applied to all six variables that contain multiple input values (three input values each) for a total of 729 ($= 3^6$) total scenarios for each cohort analysed (Figure 1). Each unique combination of input values was given a six letter 'scenario code' based on the combination of input identifiers utilised. For example, the scenario code of ADGJMP represents a scenario in which estimate (A) was used for the percent of US adults with a 10 pack-year history of smoking (28.4%), estimate (D) was used for the compliance with screening recommendations (27%), estimates (G) were used for screening procedure costs, and so on.

The total costs, total 5-year survivors, and the ratio of total cost to 5-year survivors were analysed for all 1458 scenarios (729 scenarios for each cohort). The CP5YS ratio was determined by first calculating the estimated number of NSCLC tumours distributed by stage for the 5-year period for both the STID and EDCTS approaches. The cost of diagnosis and treatment by stage were applied to these tumour populations, with screening cost also added to the EDCTS costs. The estimated percent of 5-year survivors for each stage was also applied to the tumour populations. The ratio of total CP5YS was then calculated (see Figure 2).

RESULTS

Note that the total survivors and total cost figures reported in this analysis refer only to the subset of the US population studied and are not extrapolated to represent nationwide estimates.




Figure 1 Illustration of the method of using multiple input value combinations used to generate 729 scenarios for each cohort. Identification codes (A), (B), and (C) represent the three input values for the percent of US adult smokers from Table 1. Similarly, (D), (E), and (F) represent three input values for percent complying with screening recommendations, and (G), (H), and (I) represent the three input values for screening procedure costs. 'Scenario code' refers to the combination of input value identifiers for multiple variables.




Figure 2 Schematic illustration of the calculation of costs and 5-year survivors for the early detection computed tomography screening (EDCTS) method. Costs and 5-year survivors for the symptomatic tumour identification (STID) method follow the path of the 'un-screened population' above. †The calculation exemplified by the above schematic was separately carried out for age range 40–79 years and age range 60–79 years.




Figure 3 Increase in STID costs per 5-year survivor (CP5YS) compared with the EDCTS method (STID CP5YS–EDCTS CP5YS). The scenario codes represent the combination of input variables and associated codes from Table 1. All 729 scenario codes are represented in each graph, though not all Y axis values are labelled. Note that the data are not sorted alphabetically by the X axis scenario codes. STID = symptomatic tumour identification; EDCTS = early detection approach using computed tomography screening. Note: Across all scenarios, the total number of 5-year survivors was higher for the EDCTS method, ranging from approximately 13 400 to 39 300 increased 5-year survivors for the population aged 40–79 years and 9300–27 400 for the population aged 60–79 years.

Cohort ages 40–79 years

Across all 729 scenarios analysed for the cohort ages 40–79 years, the total CP5YS ranged from \$157 300 to \$293 800 for the STID method, and from \$105 100 to \$288 100 for the EDCTS method. The costs per 5-year survivor were higher for the STID approach in 75% of the scenarios analysed, ranging from approximately \$5700 to \$110 000 increased CP5YS under the STID method for those scenarios (Figure 3) (See the Appendix Table A1 for a listing of the CP5YS for all 729 scenarios). The total number of 5-year survivors over the 5-year period analysed was higher for the EDCTS method for all 729 scenarios, ranging from approximately 13 400 to 39 300 increased 5-year survivors under the EDCTS method, which is a 112–309% increase. The total screening, diagnosis, and treatment costs were also higher for the EDCTS method in all 729 scenarios, ranging from approximately \$1.5B to \$8.7B in increased total costs, which is a 77–256% increase (Figure 4).

The EDCTS method resulted in a 27–66% increase in the number of tumours identified when compared with the STID method (Table 6).

Scenario code BDHJOQ was associated with the lowest CP5YS of approximately \$105 100 per 5-year survivor for the EDCTS method and \$157 300 for the STID method. Scenario code CEIKNP was associated with the highest CP5YS of approximately \$288 100 per 5-year survivor for the EDCTS method and \$293 800 for the STID method.

The scenario most favourable towards the EDCTS method in terms of the comparative decrease in costs per 5-year survivor was scenario code BDHJNR, in which the EDCTS CP5YS was approximately \$110 000 less than the STID ratio for that same scenario (305% more EDCTS 5-year survivors in that scenario). The scenario least favourable towards the EDCTS method in terms of the comparative increase in costs per 5-year survivor was scenario code BDIKOR, in which the EDCTS CP5YS was




Figure 4 Comparison of the total screening, diagnosis, and treatment costs between the STID and EDCTS methods. The scenario codes represent the combination of input variables and associated codes from Table 1. All 729 scenario codes are represented in each graph, though not all Y axis values are labelled. Note that the data are not sorted alphabetically by the X axis scenario codes. STID = symptomatic tumour identification, EDCTS = early detection approach using computed tomography screening.

approximately \$53 400 more than the STID ratio for that same scenario (248% more EDCTS 5-year survivors in that scenario).

Cohort ages 60–79 years

Across all 729 scenarios analysed for the cohort ages 60–79 years, the total CP5YS ranged from \$149 400 to \$282 100 for the STID method, and from \$86 400 to \$233 300 for the EDCTS method. The costs per 5-year survivor were higher for the STID approach in all scenarios analysed, ranging from approximately \$5800 to \$116 700 increased CP5YS under the STID method for those scenarios (Figure 3) (See the Appendix Table A2 for a listing of the CP5YS for all 729 scenarios). The total number of 5-year survivors over the 5-year period analysed was higher for the EDCTS method for all 729 scenarios, ranging from approximately 9000 to 26 600 increased 5-year survivors under the EDCTS method, which is a 98–272% increase. The total screening, diagnosis, and treatment costs were also higher for the EDCTS method in all 729 scenarios, ranging from approximately \$728 M to \$4.0B in increased total costs, which is a 50–159% increase (Figure 4).

The EDCTS method resulted in a 24–60% increase in the number of tumours identified when compared with the STID method (Table 6).

Scenario code BDHJQJ was associated with the lowest CP5YS of approximately \$86 400 per 5-year survivor for the EDCTS method and

\$149 400 for the STID method. Scenario code CEIJNP was associated with the highest CP5YS of approximately \$233 300 per 5-year survivor for the EDCTS method and \$282 000 for the STID method.

The scenario most favourable towards the EDCTS method in terms of the comparative decrease in costs per 5-year survivor was scenario code BDHJNR, in which the EDCTS CP5YS was approximately \$116 700 less than the STID ratio for that same scenario (269% more EDCTS 5-year survivors in that scenario). The scenario least favourable towards the EDCTS method in terms of the comparative decrease in costs per 5-year survivor was scenario code CEIKOQ, in which the EDCTS CP5YS was approximately \$5800 less than the STID ratio for that same scenario (98% more EDCTS 5-year survivors in that scenario).

DISCUSSION

The advisability of population screening for lung cancer is currently a topic of active debate. Because of the potential high cost of lung cancer screening programmes, cost-effectiveness questions are an important component to be evaluated. This article attempts to inform the debate by a multivariable analysis model. We incorporated a range of optimistic and pessimistic input variables taken from literature sources on both sides of this debate. In addition, this analysis presents an important method for evaluating a lung cancer-screening programme in contrast to

Table 6 Comparison of the total number of diagnosed tumours between the STID and EDCTS methods

Scenario code (includes all scenario codes beginning with the first two letters indicated) ^b	Cohort ages 40–79 years ^a			
	Total number of diagnosed tumours (5-year period)		Increase in EDCTS tumours	% Increase in EDCTS tumours
	STID	EDCTS ^c		
AD...		86 818	31 357	57
AE...		73 323	17 862	32
AF...		82 849	27 388	49
BD...	55 461	92 044	36 583	66
BE...	(Same for all scenarios)	76 300	20 839	38
BF...		87 414	31 953	58
CD...		81 592	26 131	47
CE...		70 346	14 885	27
CF...		78 284	22 823	41

Abbreviations: EDCTS = early detection approach using computed tomography screening; STID = symptomatic tumour identification. ^aApplying the same analysis to the cohort ages 60–79 years (not shown here), the increase in tumours identified under the EDCTS method ranged from 24 to 60%. ^bThe scenario codes represent the combination of input variables and associated codes from Table 1. ^cThe estimated number of EDCTS tumour varies depending on the following input parameters: (1) Estimated percentage of patients with a 10 pack-year history of smoking; and (2) the estimated percent of patients complying with screening recommendations. As such, the EDCTS number of diagnosed tumours will vary for scenario codes representing different values for these input parameters

analyses using QALYs, which can be misleading in terms of the efficacy of treatment regimens, particularly when there is little chance of long-term survival, as in the treatment of stage IIIB and IV NSCLC with chemotherapy. Owing to the frequent use of the QALY metric in the literature, we have provided for the reader a cost-per-year of life calculation for all 729 scenarios in the Appendix Table A3 and Table A4 to allow a rough comparison to other studies evaluating cost-effectiveness on a per-year basis. It should be emphasised, however, that the data in Table A3 and Table A4 is calculated assuming that these lung cancer survivors live only 5 years. It is therefore a conservative estimate and overstates the cost-per-year of life, as many lung cancer survivors detected by screening will live longer than 5 years (Marcus *et al*, 2000; Sobue *et al*, 2002; Henschke *et al*, 2006a).

The combination of decreased CP5YS and increased 5-year survivors in all 729 scenarios for the cohort ages 60–79 years and in the majority of scenarios for the cohort ages 40–79 years suggests that health-care dollars spent on proactively screening for lung cancer may achieve higher cost-effectiveness. These data may also suggest advantages in focusing a screening programme on patients aged 60 years or older as the marginal costs for implementing screening in the cohort ages 60–79 years were between 46 and 60% of the marginal cost for the cohort ages 40–79 years, with greater cost effectiveness in the 60–79 age group when compared with the STID method (see Figure 3). These findings are consistent with a 2008 study by Whynes (2008), analysing the potential cost effectiveness of lung cancer screening in the United Kingdom.

The I-ELCAP study results have shown the ability of CT scanning to detect asymptomatic lung cancers. By definition, a lung cancer programme that detects symptomatic and asymptomatic malignancies will diagnose more tumours at the onset of the programme than a system that detects symptomatic tumours alone, as illustrated in Figure 5. The magnitude of this increase is unknown, and sufficient information to evaluate the reasonableness of the 24–66% increase in diagnosed tumours under the EDCTS method presented in this model is currently unavailable. Additional long-term data on the prevalence of lung cancer in an asymptomatic at-risk population annually screened over multiple years are needed to enable further estimates in this area.

Increased total costs associated with a successful proactive screening programme are to be expected, not only due to the cost of screening procedures, but also more importantly due to the increased patient survival and related increase in long-term treatment costs and follow-up care. The available data for cost of treatment utilised in this study (see Table 1, inputs M, N, and O) all indicate that the treatment

for early stage cancer is more expensive than cancers detected in stages IIIB and IV. This is counter-intuitive, as stage I NSCLC is treated by surgery alone, whereas advanced stage lung cancers require multimodality treatment, adding the cost of expensive radiation and chemotherapy. Furthermore, treatment of advanced stage lung cancer is changing, which may potentially shift this balance, making the treatment of late-stage cancers increasingly more expensive. The percentage of patients in higher stages who are receiving expensive radiation therapy and chemotherapy appears to be increasing (Ramsey *et al*, 2004; Langer *et al*, 2005). In addition, new molecular 'targeted' medications used in second- and third-line treatment may markedly increase treatment costs (Adis International Ltd. Erlotinib, 2003). As I-ELCAP data report increased actuarial 10-year survival of 80% with screen-detected tumours (Henschke *et al*, 2006a), which is dramatically different than the survival of symptom-detected tumours which are disproportionately detected in later stages, it would follow that increased costs of long-term care would be expected. The cost of follow-up and care in the majority of survivors, however, adds little to routine health-care expenditures, and only a small minority would require downstream salvage or palliative radiation therapy and chemotherapy. In addition, the cost of screening programmes and treatment may reasonably be expected to decrease as a result of economies of scale, if lung cancer screening were implemented at a state or nationwide level.

It is important to consider several potential sources of bias when evaluating lung cancer screening:

(1) 'Overdiagnosis bias': small, slow-growing lesions are detected by screening for intervention that would never become symptomatic within a patient's lifetime in the absence of screening (Black *et al*, 2006). This could be caused by an improper pathological diagnosis; however, all lung cancers in the I-ELCAP study are vetted by a panel of prominent international pathologists (Flieder *et al*, 2006). With regard to the theoretical possibility of screening the detection of very slow growing malignant neoplasms that do not cause symptoms during the patient's anticipated normal lifespan, for ethical reasons, a randomised trial comparing surgery with no surgery for stage I NSCLC is not possible. However, data on untreated screen-detected NSCLC from screening studies including the Johns Hopkins study, the Memorial Sloan-Kettering study, the Mayo Clinic study, and the I-ELCAP study indicate that almost all of these untreated patients die within 5 years (Fleehinger *et al*, 1992; Henschke *et al*, 2006a). A study by Raz *et al* (2007), examined the natural history of patients with stage I NSCLC and concluded that long-term survival with untreated stage I NSCLC is uncommon with an overall survival of 6% for untreated stage I




Figure 5 Illustration demonstrating the theory for an increase in the number of tumours identified by screening for asymptomatic malignancies in addition to identifying symptomatic tumours. 'Tumour crop' refers to a cohort of lung cancers that clinically manifested at a point in time (i.e., a group of tumours for which the tumours' inception is at the same time point represented on the X axis).

NSCLC. In addition, a 2003 study by Henschke *et al* analysed data from 885 cases of stage IA lung cancer and concluded that almost all diagnosed cases of Stage IA lung cancer have a malignant natural course, fatal if not treated, thus representing genuine cancer. Although some speculate that the explanation for the paradoxical result in the Mayo Lung Trial of increased survival but no reduction in mortality with chest radiograph screening was due to, in part, overdiagnosis bias (Patz, 2006), no empirical evidence to support this theory exists. Furthermore, Yankelevitz *et al*, (2003) analysed the doubling time of stage I tumours in the Mayo Lung Trial and concluded that 'the hypothesis that early-stage lung tumours diagnosed on chest radiography during lung carcinoma screening may frequently be overdiagnosed, indolent cases needs to be rejected'.

Another form of overdiagnosis could exist in instances where comorbid disease would kill the patient before symptoms of lung cancer were experienced. This form of overdiagnosis bias can be reasonably assumed to be rendered largely irrelevant in medical environments, where patients are managed with good, sensible clinical judgment by physicians adept in identifying comorbid disease and reasonably accurately estimating the anticipated survival of their patients. Furthermore, in this study we have carefully taken into account the percentage of patients eligible for screening who have serious comorbid disease and excluded them from our analysis (refer to the 'Population Screened' section of the Materials and Methods above). We have been conservative in this approach by additionally excluding patients aged 80 years and above from screening, although many 80-year-olds are perfectly capable of undergoing minimally invasive surgical resection of lung cancers or being treated with radiotherapy.

Finally, overdiagnosis could result if spontaneous remission of a preclinical cancer were to occur. Case reports of this phenomenon are extremely rare (Kappauf *et al*, 1997).

(2) 'Length bias': detection of more patients with less aggressive disease and fewer of those with more aggressive disease, because the duration of asymptomatic disease is longer in less aggressive

tumours (Black *et al*, 2006). This also could result in 'overdiagnosis', discussed above. The baseline round of screening is inherently different from the repeat rounds because cancers with a longer latent (asymptomatic) phase are more frequently identified in the baseline round, whereas cancers found in repeat rounds are found earlier in their latent phase than in the baseline round (Morrison, 1982; Henschke and Yankelevitz, 2008). Cancers that are diagnosed at baseline, thus, tend to grow more slowly than does the subtype in general; they also grow more slowly than do tumours that are diagnosed in repeated screenings. As noted by the I-ELCAP researchers, this fact does not introduce a bias, but it may call for making a distinction between baseline screening and repeated screening (Henschke *et al*, 2007). 'Length bias' also implies that the faster growing tumours may present symptomatically between screening exams, however, the I-ELCAP data showed a very low incidence of such cases.

(3) 'Lead-time bias': screening-detected patients are accorded extended survival times solely because cancer was detected earlier owing to screening, although death occurred at the same time as would have happened without screening (Black *et al*, 2006). To address this potential source of bias, the estimated percentage of stage I 5-year survivors for all scenarios using input variable 'Q' (see Table 1) are based on 10-year survival percentages reported by I-ELCAP (Henschke *et al*, 2006a). If lead-time bias was evident in this study, a large number of individuals who survived 5 years would be expected to die before 10 years; the I-ELCAP survival curve shows no decrement in survival between 5 and 10 years (Henschke *et al*, 2006a).

Although some emphasise that the risks and complications of lung cancer screening may be considerable (Bach *et al*, 2007), the minutes of National Cancer Institute's National Lung Screen Trial (NLST) Data Safety Monitoring Board (DSMB), which we have reviewed, indicate that during the first 5 years of the study, which began in 2002, no unanticipated complications or risks have been recognised. The DSMB, which is responsible for the safety of NLST

research subjects, has neither terminated the study nor reported information on new complications to study subjects.

Limitations

Input variables based on SEER data may not be representative of actual nationwide lung cancer incidence or staging. Our model does not account for costs associated with complications from biopsies or other screening procedures, nor does it account for the increase in capital equipment and resources necessary to implement large-scale comprehensive CT screening programmes. In addition, our study design does not consider the indirect cost of lost productivity attributable to lung cancer morbidity and mortality. Of the estimated \$167 billion costs of all diseases caused by tobacco products, indirect costs (\$92 billion) are substantially higher than direct costs (Centers for Disease Control and Prevention, 2005). As survival increases, indirect costs attributable to loss of patient income, spousal income, and other factors may reasonably be expected to diminish substantially. This study is based on a representative subset of the US population and it is beyond the scope of this analysis to extrapolate the results to a state or nationwide level. The model does not factor in any additional benefit conferred by

survival beyond 5 years. On the basis of the results from the Mayo Lung Trial (Marcus *et al*, 2000), Japanese Anti-Lung Cancer Association (ALCA) trial (Sobue *et al*, 2002), and I-ELCAP (Henschke *et al*, 2006a), there is a strong evidence to suggest that the majority of 5-year survivors will continue to survive for 10, 15, and even 20 years following diagnosis and treatment. Finally, there is no accurate method to calculate a dollar value for not dying of lung cancer in an individual or group of individuals with lung cancer.

CONCLUSION

Our analytical model offers an innovative tool that provides data estimates that contribute insights into the continuing debate on the wisdom and advisability of implementing state and/or nationwide CT screening programmes. The predicted increase in long-term survival with CT screening and the potential for better utilisation of health-care dollars in terms of CP5YS, particularly when screening patients over the age of 60 years, are critically important considerations in directing effective future lung cancer management strategy.

REFERENCES

- Adis International Ltd. Erlotinib (2003) CP 358774, NSC 718781, OSI 774, R 1415. *Drugs R D* 4(4): 243–248
- American Cancer Society (2006) *Cancer Facts and Figures 2006*. Atlanta: American Cancer Society
- Bach PB, Jett JR, Pastorino U, Tockman MS, Swensen SJ, Begg CB (2007) Computed tomography screening and lung cancer outcomes. *JAMA* 297(9): 953–961
- Black C, Bagust A, Boland A, Walker S, McLeod C, De Verteuil R, Ayres J, Bain L, Thomas S, Godden D, Waugh N (2006) The clinical effectiveness and cost-effectiveness of computed tomography screening for lung cancer: systematic reviews. *Health Technol Assess* 10(3): 8, 38
- Centers for Disease Control and Prevention (2005) Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses – United States, 1997–2001. *MMWR* 54: 625–628
- Chien CR, Chen TH (2008) Mean sojourn time and effectiveness of mortality reduction for lung cancer screening with computed tomography. *Int J Cancer* 122(11): 2594–2599
- Chirikos TN, Hazelton T, Tockman M, Clark R (2002) Screening for lung cancer with CT: a preliminary cost-effectiveness analysis. *Chest* 121: 1507–1514
- Evans WK, Will BP, Berthelot JM, Wolfson MC (1995) Estimating the cost of lung cancer diagnosis and treatment in Canada: the POHEM model. *Can J Oncol* 5: 408–419
- Fireman BH, Quesenberry CP, Somkin CP, Jacobson AS, Baer D, West D, Potosky AL, Brown ML (1997) Cost of care for cancer in a health maintenance organization. *Health Care Financ Rev* 18: 51–76
- Fleehinger BJ, Kimmel M, Melamed MR (1992) The effect of surgical treatment on survival from early lung cancer. Implications for screening. *Chest* 101(4): 1013–1018
- Flieder DB, Vazquez M, Carter D, Brambilla E, Gazdar A, Noguchi M, Travis WD, Kramer A, Yankelevitz DF, Henschke CI (2006) Pathologic findings of lung tumors diagnosed on baseline CT screening. *Am J Surg Pathol* 30(5): 606–613
- Fry WA, Phillips JL, Menck HR (1999) Ten-year survey of lung cancer treatment and survival in hospitals in the United States: a national cancer data base report. *Cancer* 86: 1867–1876
- Henschke CI (2006) International early lung cancer action program: protocol. <http://www.ielcap.org/ielcap.pdf>, 10/20/2006. Accessed 26 December 2006.
- Henschke CI, McCauley DI, Yankelevitz DF, Naidich DP, McGuinness G, Miettinen OS, Libby DM, Pasmantier MW, Koizumi J, Altorki NK, Smith JP (1999) Early Lung Cancer Action Project: overall design and findings from baseline screening. *Lancet* 354: 99–105
- Henschke CI, Naidich DP, Yankelevitz DF, McGuinness G, McCauley DI, Smith JP, Libby D, Pasmantier M, Vazquez M, Koizumi J, Flieder D, Altorki N, Miettinen OS (2001) Early lung cancer action project. Initial findings on repeat screenings. *Cancer* 92: 153–159
- Henschke CI, Smith JP, Miettinen OS (2007) Response to letters to the editor. *N Engl J Med* 356: 743–747
- Henschke CI, Wisnivesky JP, Yankelevitz DF, Miettinen OS (2003) Small stage I cancers of the lung: genuineness and curability. *Lung Cancer* 39(3): 327–330
- Henschke CI, Yankelevitz DF, Libby DM, Pasmantier MW, Smith JP, Miettinen OS (2006a) Survival of patients with stage I lung cancer detected on CT screening. *N Engl J Med* 355(17): 1763–1771
- Henschke LI, Yankelevitz DF (2008) CT screening for lung cancer: Update 2007. *Oncologist* 13: 65–78
- Horton JA, Cruess DF, Romans MC (1996) Compliance with mammography screening guidelines: 1995 mammography attitudes and usage study report. *Womens Health Issues* 6: 239–245
- Jemal A, Murray T, Ward E, Samuels A, Tiwari RC, Ghafoor A, Feuer EJ, Thun MJ (2005) Cancer statistics, 2005. *CA Cancer J Clin* 55: 10–30
- Kappauf H, Gallmeier WM, Wunsch PH (1997) Complete spontaneous remission in a patient with metastatic non-small-cell lung cancer. Case report, review of literature, and discussion of possible biological pathways involved. *Ann Oncol* 8(10): 1031–1039
- Kelly RF, Tran T, Holmstrom A, Murar J, Seguro Jr RJ (2004) Accuracy and cost-effectiveness of [18F]-2-fluoro-deoxy-D-glucose-positron emission tomography scan in potentially resectable non-small cell lung cancer. *Chest* 125: 1413–1423
- Knudsen AB, McMahon PM, Gazelle GS (2007) Use of modeling to evaluate the cost-effectiveness of cancer screening programs. *J Clin Oncol* 25(2): 203–208
- Langer CJ, Moughan J, Movsas B, Komaki R, Ettinger D, Owen J, Wilson JF (2005) Patterns of care survey (PCS) in lung cancer: how well does current U.S. practice with chemotherapy in the non-metastatic setting follow the literature? *Lung Cancer* 48(1): 93–102
- Mahadevia PJ, Fleisher LA, Frick KD, Eng J, Goodman SN, Powe NR (2003) Lung cancer screening with helical computed tomography in older adult smokers: a decision and cost-effectiveness analysis. *JAMA* 289: 313–322
- Marcus PM, Bergstralh EJ, Fagerstrom RM, Williams DE, Fontana R, Taylor WF, Prorok PC (2000) Lung cancer mortality in the Mayo Lung Project: impact of extended follow-up. *J Natl Cancer Inst* 92: 1308–1316
- Marshall D, Simpson KN, Earle CC, Chu CW (2001a) Economic decision analysis model of screening for lung cancer. *Eur J Cancer* 37: 1759–1767
- Marshall D, Simpson KN, Earle CC, Chu C (2001b) Potential cost-effectiveness of one-time screening for lung cancer (LC) in a high risk cohort. *Lung Cancer* 32: 227–236
- Morrison AS (1982) The effects of early treatment, lead time and length bias on the mortality experienced by cases detected by screening. *Int J Epidemiol* 11: 261–267
- Mountain CF (1997) Revisions in the International System for staging lung cancer. *Chest* 111(6): 1710–1717
- National Cancer Institute (2005) Surveillance, Epidemiology, and End Results (SEER) Program, (www.seer.cancer.gov) Public-Use Data (1973–2003), DCCPS, Surveillance Research Program, Cancer Statistics Branch, released April 2006, based on the November 2005 submission.

National Cancer Institute (2003) Centers for Disease Control and Prevention Co-sponsored Tobacco Use Special Cessation Supplement to the Current Population Survey. <http://riskfactor.cancer.gov/studies/tus-cps/>.

Patz Jr EF (2006) Lung cancer screening, overdiagnosis bias, and reevaluation of the Mayo Lung Project. *J Natl Cancer Inst* 98(11): 724–725

Patz Jr EF, Rossi S, Harpole Jr DH, Herndon JE, Goodman PC (2000) Correlation of tumor size and survival in patients with pathologic stage IA nonsmall cell lung cancer. *Chest* 117: 1568–1571

Phillips C, Thompson G (2003) What is a QALY? Newmarket, United Kingdom: Hayward Medical Communications; 2003. www.evidence-based-medicine.co.uk. Accessed 8 September 2006.

Phillips KA, Kerlikowske K, Baker LC, Chang SW, Brown ML (1998) Factors associated with women's adherence to mammography screening guidelines. *Health Serv Res* 33: 29–53

Rahman SM, Dignan MB, Shelton BJ (2003) Factors influencing adherence to guidelines for screening mammography among women aged 40 years and older. *Ethn Dis* 13: 477–484

Ramsey SD, Howlader N, Etzioni RD, Donato B (2004) Chemotherapy use, outcomes, and costs for older persons with advanced non-small-cell lung cancer: evidence from surveillance, epidemiology and end results-Medicare. *J Clin Oncol* 22(24): 4971–4978

Raz DJ, Zell JA, Ou SH, Gandara DR, Anton-Culver H, Jablons DM (2007) Natural history of stage I non-small cell lung cancer: implications for early detection. *Chest* 132(1): 193–199

Ries LAG, Harkins D, Krapcho M, Mariotto A, Miller BA, Feuer EJ, Clegg L, Eisner MP, Horner MJ, Howlader N, Hayat M, Hankey BF, Edwards BK

(eds) (2005) SEER Cancer Statistics Review, 1975–2003, National Cancer Institute. Bethesda, MD. http://seer.cancer.gov/csr/1975_2003/, based on November 2005 SEER data submission, posted to the SEER web site, 2006.

Riley GF, Potosky AL, Lubitz JD, Kessler LG (1995) Medicare payments from diagnosis to death for elderly cancer patients by stage of diagnosis. *Med Care* 33: 828–841

Sobue T, Moriyama N, Kaneko M, Kusumoto M, Kobayashi T, Tsuchiya R, Kakinuma R, Ohmatsu H, Nagai K, Nishiyama H, Matsui E, Eguchi K (2002) Screening for lung cancer with low-dose helical computed tomography: anti-lung cancer association project. *J Clin Oncol* 20: 911–920

Swensen SJ, Jett JR, Sloan JA, Midthun DE, Hartman TE, Sykes AM, Aughenbaugh GL, Zink FE, Hillman SL, Noetzel GR, Marks RS, Clayton AC, Pairolo PC (2002) Screening for lung cancer with low-dose spiral computed tomography. *Am J Respir Crit Care Med* 165: 508–513

Whynes DK (2008) Could CT screening for lung cancer ever be cost effective in the United Kingdom? *Cost Eff Resour Alloc* 6: 5

Wisnivesky JP, Mushlin AI, Sicherman N, Henschke C (2003) The cost-effectiveness of low-dose CT screening for lung cancer: preliminary results of baseline screening. *Chest* 124: 614–621

Yankelevitz DF, Kostis WJ, Henschke CI, Heelan RT, Libby DM, Pasmantier MW, Smith JP (2003) Overdiagnosis in chest radiographic screening for lung carcinoma: frequency. *Cancer* 97(5): 1271–1275

Appendix

See Tables A1 to A4

Table A1 Summary of the cost per 5-year survivor for all 729 scenarios^a analysed for the cohort ages 40–79 years (2005 USD, in thousands)

		EDCTS																															
		A									B									C													
		D			E			F			D			E			F			D			E			F							
STID		G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I		
P	219	180	173	259	187	181	251	181	175	257	179	172	259	186	180	252	180	174	258	180	174	258	188	183	250	182	176	256					
M	Q	199	146	141	210	155	151	208	148	143	210	145	140	211	154	149	209	147	142	210	147	142	210	156	152	208	149	144	210				
	R	211	156	151	225	165	161	223	158	153	224	155	150	225	164	159	223	157	152	225	157	152	225	167	162	222	159	154	224				
	P	294	207	201	286	223	218	288	211	205	287	206	199	286	221	216	288	209	203	287	209	203	286	226	221	288	213	207	287				
J	N	Q	267	169	164	233	185	181	239	172	167	234	167	162	232	183	178	238	171	166	234	170	165	234	188	184	240	174	169	235			
	R	283	180	175	249	198	193	255	184	179	250	179	173	249	195	191	254	183	177	250	182	177	250	201	196	256	186	181	251				
	P	173	136	130	215	143	138	208	138	132	214	136	129	216	142	137	209	137	131	215	137	131	215	144	139	206	139	133	213				
O	Q	157	111	106	175	119	114	172	113	108	175	110	105	176	118	113	173	112	107	175	112	107	175	120	116	172	114	109	174				
	R	167	119	113	187	127	122	184	121	115	187	118	112	188	126	121	185	120	114	187	120	114	187	128	124	183	122	116	186				
	P	219	176	169	256	184	179	249	178	171	254	175	169	257	183	177	250	177	170	255	177	170	255	188	180	248	179	173	254				
M	Q	199	164	158	239	171	166	232	166	160	238	164	158	240	170	165	233	165	159	238	165	159	239	172	167	230	167	161	237				
	R	211	175	168	255	182	177	246	176	170	253	174	168	255	181	175	248	176	169	254	176	169	254	183	178	245	177	171	252				
	P	294	206	200	287	223	217	288	210	204	287	205	198	287	220	215	288	208	202	287	208	202	287	225	220	288	212	206	287				
K	N	Q	267	193	187	268	207	202	268	196	190	268	192	186	268	205	200	268	195	189	268	195	189	268	209	204	268	198	192	268			
	R	283	205	199	285	220	215	285	209	202	285	204	197	285	218	213	285	207	201	285	207	201	285	222	217	285	210	204	285				
	P	173	140	134	221	146	141	212	142	135	219	140	133	221	146	140	213	141	135	219	141	135	220	147	142	210	142	136	217				
O	Q	157	131	125	206	136	131	197	132	127	204	131	125	207	135	130	198	132	126	205	132	126	205	137	132	195	133	127	203				
	R	167	140	133	219	145	140	209	141	135	217	139	133	220	144	139	211	140	134	218	140	134	218	146	141	208	141	135	216				
	P	219	178	171	257	185	180	250	179	173	256	177	170	258	184	179	251	179	172	257	179	172	257	186	181	249	180	174	255				
M	Q	199	155	149	224	163	158	220	157	151	223	154	149	225	162	157	221	156	150	224	156	150	224	164	160	219	158	152	223				
	R	211	165	159	239	173	169	234	167	161	238	165	159	240	172	167	235	166	160	239	166	160	239	175	170	234	168	163	238				
	P	294	207	201	287	223	218	288	211	204	287	205	199	286	221	215	288	209	203	287	209	202	287	225	220	288	213	207	287				
L	N	Q	267	181	175	250	196	191	253	184	179	251	179	173	250	194	189	253	182	177	250	182	177	250	198	194	253	186	181	251			
	R	283	193	187	267	209	204	270	196	190	267	191	185	266	207	201	269	195	189	267	194	188	267	211	207	270	198	193	268				
	P	173	139	132	218	145	140	210	140	134	216	138	131	219	144	139	211	139	133	217	139	133	217	146	141	209	141	135	215				
O	Q	157	121	115	190	127	123	184	122	117	189	120	115	191	126	122	185	122	116	190	122	116	190	128	124	183	123	118	188				
	R	167	129	123	203	136	131	197	130	125	202	128	122	204	135	130	197	130	124	202	130	124	202	137	132	195	131	126	201				

□ Early detection approach using computed tomography screening (EDCTS).

■ Symptomatic tumour identification (STID) method.

^aRow and column headings represent input variable identification codes corresponding to Table 1. Each figure within the table above represents a unique combination of input variables with the six-letter combination of row and column headings corresponding to the 'scenario code' for that set of variables (see Figure 1 for further explanation of scenario codes). Note that the STID costs per 5-year survivor do not change across variables A through I.

Table A2 Summary of the cost per 5-year survivor for all 729 scenarios^a analysed for the cohort ages 60–79 years (2005 USD, in thousands)

		EDCTS																											
		A									B									C									
		D			E			F			D			E			F			D			E			F			
STID	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I		
M	P	211	154	151	191	165	162	195	156	153	192	152	149	191	163	161	195	155	152	192	155	152	192	167	164	196	158	155	193
	Q	191	125	123	156	137	135	162	128	125	157	124	121	155	135	133	161	126	124	156	126	124	156	139	137	163	129	127	158
	R	202	134	131	167	146	144	173	137	134	168	133	130	166	144	142	172	135	133	167	135	132	167	148	146	174	138	136	169
J	P	282	181	178	219	201	199	231	186	183	222	180	177	218	198	196	230	184	181	221	184	181	220	204	202	233	188	186	223
	Q	254	148	145	178	167	165	192	152	150	181	146	143	177	164	162	190	150	148	180	150	147	180	170	168	194	154	152	183
	R	270	158	155	191	178	176	205	163	160	194	156	153	189	176	173	203	160	158	193	160	158	192	182	180	207	165	163	196
O	P	166	110	107	148	121	119	151	113	110	149	109	106	148	120	117	151	112	109	148	112	109	148	123	121	152	114	111	149
	Q	149	90	87	121	101	99	126	92	90	122	89	86	120	99	97	125	91	89	121	91	89	121	102	100	127	94	91	122
	R	159	96	94	129	107	105	134	99	96	130	95	92	128	106	104	134	98	95	130	97	95	130	109	107	135	100	98	131
M	P	211	150	146	188	162	159	192	152	149	189	148	145	187	160	158	192	151	148	188	151	148	188	164	161	193	154	151	189
	Q	191	140	137	175	150	148	178	142	139	176	139	136	175	148	146	178	141	138	176	141	138	175	151	149	179	143	141	176
	R	202	148	145	186	159	157	190	151	148	187	147	144	186	158	155	189	150	147	187	150	147	187	161	159	190	152	149	187
K	P	282	180	177	218	200	198	231	185	182	221	178	175	217	198	195	229	183	180	220	182	179	220	203	201	233	187	184	223
	Q	254	168	165	204	186	183	214	172	169	206	166	163	203	183	181	213	170	167	205	170	167	205	188	186	216	174	172	207
	R	270	179	176	217	197	195	228	183	180	219	177	174	216	195	192	226	181	178	218	181	178	218	200	198	229	185	183	221
O	P	166	114	111	152	124	122	155	116	113	153	113	110	152	123	120	154	115	112	152	115	112	152	126	123	155	117	115	153
	Q	149	106	103	142	115	113	143	108	105	142	105	102	142	114	111	143	107	104	142	107	104	142	116	114	144	109	107	142
	R	159	113	110	151	122	120	152	115	112	151	112	109	151	121	118	152	114	111	151	114	111	151	124	121	153	116	113	151
M	P	211	152	148	189	163	161	194	154	151	190	150	147	189	162	159	193	153	150	190	153	150	190	165	163	194	156	153	191
	Q	191	132	129	165	143	141	170	135	132	166	131	128	165	142	140	169	133	131	166	133	131	166	145	143	171	136	134	167
	R	202	141	138	176	153	150	181	144	141	177	140	137	176	151	149	181	142	140	177	142	139	177	155	152	182	145	142	178
L	P	282	181	178	219	201	198	231	185	182	222	179	176	218	198	196	230	183	180	220	183	180	220	204	202	233	188	185	223
	Q	254	158	155	191	176	174	203	162	159	194	156	153	190	174	171	201	160	157	192	160	157	192	179	177	205	164	162	195
	R	270	168	165	203	188	186	216	173	170	206	166	163	202	185	183	215	171	168	205	170	168	205	191	189	218	175	172	208
O	P	166	112	109	150	123	120	153	115	112	151	111	108	150	121	119	153	113	110	151	113	110	150	124	122	154	116	113	151
	Q	149	98	95	131	108	106	134	100	98	132	97	94	131	106	104	134	99	96	131	99	96	131	109	107	135	101	99	132
	R	159	104	101	140	115	112	143	107	104	140	103	100	139	113	111	143	106	103	140	105	103	140	116	114	144	108	105	141

□ Early detection approach using computed tomography screening (EDCTS).

■ Symptomatic tumour identification (STID) method.

^aRow and column headings represent input variable identification codes corresponding to Table 1. Each figure within the table above represents a unique combination of input variables with the six-letter combination of row and column headings corresponding to the 'scenario code' for that set of variables (see Figure 1 for further explanation of scenario codes). Note that the STID costs per five-year survivor do not change across variables A through I.

Table A3 Summary of the cost per lung cancer survivor year of life for all 729 scenarios^a analysed for the cohort ages 40–79 years (2005 USD, in thousands)^b

		EDCTS																											
		A									B									C									
		D			E			F			D			E			F			D			E			F			
		STID	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I
M	P	44	36	35	52	37	36	50	36	35	51	36	34	52	37	36	50	36	35	52	36	35	52	38	37	50	36	35	51
	Q	40	29	28	42	31	30	42	30	29	42	29	28	42	31	30	42	29	28	42	29	28	42	31	30	42	30	29	42
	R	42	31	30	45	33	32	45	32	31	45	31	30	45	33	32	45	31	30	45	31	30	45	33	32	44	32	31	45
J	P	59	41	40	57	45	44	58	42	41	57	41	40	57	44	43	58	42	41	57	42	41	57	45	44	58	43	41	57
	Q	53	34	33	47	37	36	48	34	33	47	33	32	46	37	36	48	34	33	47	34	33	47	38	37	48	35	34	47
	R	57	36	35	50	40	39	51	37	36	50	36	35	50	39	38	51	37	35	50	36	35	50	40	39	51	37	36	50
O	P	35	27	26	43	29	28	42	28	26	43	27	26	43	28	27	42	27	26	43	27	26	43	29	28	41	28	27	43
	Q	31	22	21	35	24	23	34	23	22	35	22	21	35	24	23	35	22	21	35	22	21	35	24	23	34	23	22	35
	R	33	24	23	37	25	24	37	24	23	37	24	22	38	25	24	37	24	23	37	24	23	37	26	25	37	24	23	37
M	P	44	35	34	51	37	36	50	36	34	51	35	34	51	37	35	50	35	34	51	35	34	51	37	36	50	36	35	51
	Q	40	33	32	48	34	33	46	33	32	48	33	32	48	34	33	47	33	32	48	33	32	48	34	33	46	33	32	47
	R	42	35	34	51	36	35	49	35	34	51	35	34	51	36	35	50	35	34	51	35	34	51	37	36	49	35	34	50
K	P	59	41	40	57	45	43	58	42	41	57	41	40	57	44	43	58	42	40	57	42	40	57	45	44	58	42	41	57
	Q	53	39	37	54	41	40	54	39	38	54	38	37	54	41	40	54	39	38	54	39	38	54	42	41	54	40	38	54
	R	57	41	40	57	44	43	57	42	40	57	41	39	57	44	43	57	41	40	57	41	40	57	44	43	57	42	41	57
O	P	35	28	27	44	29	28	42	28	27	44	28	27	44	29	28	43	28	27	44	28	27	44	29	28	42	28	27	43
	Q	31	26	25	41	27	26	39	26	25	41	26	25	41	27	26	40	26	25	41	26	25	41	27	26	39	27	25	41
	R	33	28	27	44	29	28	42	28	27	43	28	27	44	29	28	42	28	27	44	28	27	44	29	28	42	28	27	43
M	P	44	36	34	51	37	36	50	36	35	51	35	34	52	37	36	50	36	34	51	36	34	51	37	36	50	36	35	51
	Q	40	31	30	45	33	32	44	31	30	45	31	30	45	32	31	44	31	30	45	31	30	45	33	32	44	32	30	45
	R	42	33	32	48	35	34	47	33	32	48	33	32	48	34	33	47	33	32	48	33	32	48	35	34	47	34	33	48
L	P	59	41	40	57	45	44	58	42	41	57	41	40	57	44	43	58	42	41	57	42	40	57	45	44	58	43	41	57
	Q	53	36	35	50	39	38	51	37	36	50	36	35	50	39	38	51	36	35	50	36	35	50	40	39	51	37	36	50
	R	57	39	37	53	42	41	54	39	38	53	38	37	53	41	40	54	39	38	53	39	38	53	42	41	54	40	39	54
O	P	35	28	26	44	29	28	42	28	27	43	28	26	44	29	28	42	28	27	43	28	27	43	29	28	42	28	27	43
	Q	31	24	23	38	25	25	37	24	23	38	24	23	38	25	24	37	24	23	38	24	23	38	26	25	37	25	24	38
	R	33	26	25	41	27	26	39	26	25	40	26	24	41	27	26	39	26	25	40	26	25	40	27	26	39	26	25	40

□ Early detection approach using computed tomography screening (EDCTS).

■ Symptomatic tumour identification (STID) method.

^aRow and column headings represent input variable identification codes corresponding to Table 1. Each figure within the table above represents a unique combination of input variables with the six-letter combination of row and column headings corresponding to the 'scenario code' for that set of variables (see Figure 1 for further explanation of scenario codes). Note that the STID costs per 5-year survivor do not change across variables A through I. ^bAs a conservative estimate, this calculation assumes that the lung cancer survivors live no longer than 5 years.

Table A4 Summary of the cost per lung cancer survivor year of life for all 729 scenarios^a analysed for the cohort ages 60–79 years (2005 USD, in thousands)^b

		EDCTS																												
		A									B									C										
		D			E			F			D			E			F			D			E			F				
		STID	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	G	H	I	
Clinical Studies	J	P	42	31	30	38	33	32	39	31	31	38	30	30	38	33	32	39	31	30	38	31	30	38	33	33	39	32	31	39
		Q	38	25	25	31	27	27	32	26	25	31	25	24	31	27	27	32	25	25	31	25	25	31	28	27	33	26	25	32
		R	40	27	26	33	29	29	35	27	27	34	27	26	33	29	28	34	27	27	33	27	26	33	30	29	35	28	27	34
	M	P	56	36	36	44	40	40	46	37	37	44	36	35	44	40	39	46	37	36	44	37	36	44	41	40	47	38	37	45
		Q	51	30	29	36	33	33	38	30	30	36	29	29	35	33	32	38	30	30	36	30	29	36	34	34	39	31	30	37
		R	54	32	31	38	36	35	41	33	32	39	31	31	38	35	35	41	32	32	39	32	32	38	36	36	41	33	33	39
	K	P	33	22	21	30	24	24	30	23	22	30	22	21	30	24	23	30	22	22	30	22	22	30	25	24	30	23	22	30
		Q	30	18	17	24	20	20	25	18	18	24	18	17	24	20	19	25	18	18	24	18	18	24	20	20	25	19	18	24
		R	32	19	19	26	21	21	27	20	19	26	19	18	26	21	21	27	20	19	26	19	19	26	22	21	27	20	20	26
	L	P	42	30	29	38	32	32	38	30	30	38	30	29	37	32	32	38	30	30	38	30	30	38	33	32	39	31	30	38
		Q	38	28	27	35	30	30	36	28	28	35	28	27	35	30	29	36	28	28	35	28	28	35	30	30	36	29	28	35
		R	40	30	29	37	32	31	38	30	30	37	29	29	37	32	31	38	30	29	37	30	29	37	32	32	38	30	30	37
M	P	56	36	35	44	40	40	46	37	36	44	36	35	43	40	39	46	37	36	44	36	36	44	41	40	47	37	37	45	
	Q	51	34	33	41	37	37	43	34	34	41	33	33	41	37	36	43	34	33	41	34	33	41	34	33	41	38	37	43	
	R	54	36	35	43	39	39	46	37	36	44	35	35	43	39	38	45	36	36	44	36	36	44	40	40	46	37	37	44	
N	P	33	23	22	30	25	24	31	23	23	31	23	22	30	25	24	31	23	22	30	23	22	30	25	25	31	23	23	31	
	Q	30	21	21	28	23	23	29	22	21	28	21	20	28	23	22	29	21	21	28	21	21	28	23	23	29	22	21	28	
	R	32	23	22	30	24	24	30	23	22	30	22	22	30	24	24	30	23	22	30	23	22	30	23	22	30	25	24	31	
O	P	42	30	30	38	33	32	39	31	30	38	30	29	38	32	32	39	31	30	38	31	30	38	33	33	39	31	31	38	
	Q	38	26	26	33	29	28	34	27	26	33	26	26	33	28	28	34	27	26	33	27	26	33	29	29	34	27	27	33	
	R	40	28	28	35	31	30	36	29	28	35	28	27	35	30	30	36	28	28	35	28	28	35	31	30	36	29	28	36	
P	P	56	36	36	44	40	40	46	37	36	44	36	35	44	40	39	46	37	36	44	37	36	44	41	40	47	38	37	45	
	Q	51	32	31	38	35	35	41	32	32	39	31	31	38	35	34	40	32	31	38	32	31	38	36	35	41	33	32	39	
	R	54	34	33	41	38	37	43	35	34	41	33	33	40	37	37	43	34	34	41	34	34	41	34	34	41	38	38	44	
Q	P	33	22	22	30	25	24	31	23	22	30	22	22	30	24	24	31	23	22	30	23	22	30	25	24	31	23	23	30	
	Q	30	20	19	26	22	21	27	20	20	26	19	19	26	21	21	27	20	19	26	20	19	26	20	19	26	22	21	27	
	R	32	21	20	28	23	22	29	21	21	28	21	20	28	23	22	29	21	21	28	21	21	28	23	23	29	22	21	28	

□ Early detection approach using computed tomography screening (EDCTS).

■ Symptomatic tumour identification (STID) method.

^aRow and column headings represent input variable identification codes corresponding to Table I. Each figure within the table above represents a unique combination of input variables with the six-letter combination of row and column headings corresponding to the 'scenario code' for that set of variables (see Figure 1 for further explanation of scenario codes). Note that the STID costs per 5-year survivor do not change across variables A through I. ^bAs a conservative estimate, this calculation assumes that the lung cancer survivors live no longer than 5 years.