

RAPID COMMUNICATION

Low-Level Stress Induces Production of Neuroprotective Factors in Wild-Type but Not BDNF^{+/-} Mice: Interleukin-10 and Kynurenic Acid

Allison M. Dugan, BS; Jennifer M. Parrott, BS; Laney Redus, BS;
Julie G. Hensler, PhD; Jason C. O'Connor, PhD

Department of Pharmacology, University of Texas Health Science Center at San Antonio, San Antonio, TX (Ms Dugan, Ms Parrott, Ms Redus, Dr Hensler, and Dr O'Connor); Audie L Murphy VA Hospital, South Texas Veterans Health System, San Antonio, TX (Dr O'Connor).

Correspondence: Jason C. O'Connor, PhD, Department of Pharmacology, MSC-7764, The University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900 (occonnorj@uthscsa.edu).

Abstract

Background: Brain-derived neurotrophic factor (BDNF) deficiency confers vulnerability to stress, but the mechanisms are unclear. BDNF^{+/-} mice exhibit behavioral, physiological, and neurochemical changes following low-level stress that are hallmarks of major depression. After immune challenge, neuroinflammation-induced changes in tryptophan metabolism along the kynurenine pathway mediate depressive-like behaviors.

Methods: We hypothesized that BDNF^{+/-} mice would be more susceptible to stress-induced neuroinflammation and kynurenine metabolism, so BDNF^{+/-} or wild-type littermate mice were subject to repeated unpredictable mild stress. Proinflammatory cytokine expression and kynurenine metabolites were measured.

Results: Unpredictable mild stress did not induce neuroinflammation. However, only wild-type mice produced the neuroprotective factors interleukin-10 and kynurenic acid in response to repeated unpredictable mild stress. In BDNF^{+/-} mice, kynurenine was metabolized preferentially to the neurotoxic intermediate 3-hydroxykynurenine following repeated unpredictable mild stress.

Conclusions: Our data suggest that BDNF may modulate kynurenine pathway metabolism during stress and provide a novel molecular mechanism of vulnerability and resilience to the development of stress-precipitated psychiatric disorders.

Keywords: Kynurenine, neuroinflammation, vulnerability

Introduction

In the adult brain, brain-derived neurotrophic factor (BDNF) is essential to the survival function of neurons and has profound effects on the plasticity and structural stability of synapses. Decreased levels of BDNF, particularly in the hippocampus and frontal cortex, occur with severe or chronic stress and are mechanistically linked with depressive phenotypes (Altar, 1999; Duman and Monteggia, 2006; Calabrese et al., 2009). While

conventional BDNF^{-/-} mice do not typically survive beyond post-natal day 14, heterozygous (BDNF^{+/-}) mice exhibit a 50% to 60% reduction in BDNF expression in both the frontal cortex and hippocampus (Chourbaji et al., 2004). We and others have shown that mice deficient in BDNF are more sensitive to stress than wild-type mice and exhibit behavioral, physiological, and neurochemical changes following low-level stress that are hallmarks

Received: June 23, 2015; Revised: July 23, 2015; Accepted: July 28, 2015

© The Author 2015. Published by Oxford University Press on behalf of CINP.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

of major depression (Duman et al., 2007; Advani et al., 2009; Burke et al., 2013). However, the mechanism mediating this vulnerability remains unclear.

A growing literature indicates that neuroinflammation contributes to the development of major depression (Raison et al., 2006; Capuron and Miller, 2011). We and others have found that neuroinflammation upregulates the kynurenine pathway of tryptophan metabolism and that inflammation-induced depressive-like behaviors in mice are dependent on this pathway (O'Connor et al., 2009; Walker et al., 2013). Microglia, innate immune cells of the brain, produce proinflammatory cytokines interleukin (IL)-1 β , IL-6, and tumor necrosis factor (TNF)- α when activated, and these cytokines stimulate the induction of the kynurenine pathway. Chronic or severe stress results in increased production of proinflammatory cytokines by activated microglia (Wohleb et al., 2011; Patki et al., 2013). Thus, central inflammation and kynurenine metabolism may be one of the factors linking stress with depression.

We have reported that BDNF^{+/-} mice exhibit an exaggerated immune response and more severe depressive-like behavior following acute peripheral immune challenge compared with wild-type mice (Parrott et al., 2012). These observations led us to hypothesize that low-level stress results in an exaggerated increase in proinflammatory cytokines in the brain under conditions of BDNF deficiency. We report here no change in IL-1 β , IL-6 or TNF- α mRNA or protein in the forebrain of wild-type or BDNF^{+/-} mice following repeated unpredictable mild stress (UMS). We did, however, observe an increase in antiinflammatory cytokine IL-10 mRNA in wild-type mice, suggestive of a protective mechanism that is absent in BDNF^{+/-} mice. Given the role of IL-10 in activating astrocytes and negatively modulating microglia, we focused our attention on distinct branches of kynurenine pathway localized in these nonneuronal cells (Guillemin et al., 2001; Schwarcz and Pellicciari, 2002). Here we report that measurement of kynurenine pathway enzymes and metabolites indicated that in wild-type mice, UMS induced changes in the kynurenine pathway that resulted in increased production of kynurenic acid, an NMDA receptor antagonist. By contrast, in stress-sensitive BDNF^{+/-} mice, the production of the neurotoxic kynurenine metabolite 3-hydroxykynurenine predominated following UMS. Our data provide an innovative perspective on stress resilience and vulnerability and suggest a mechanism by which decreased levels of BDNF confer vulnerability to stress-precipitated psychiatric disorders such as major depression.

Methods

BDNF heterozygous (BDNF^{+/-}) and wild-type (BDNF^{+/+}) mice were bred at the University of Texas Health Science Center-San Antonio. Breeding pairs consisted of wild-type female (C57BL/6J) and heterozygote male (B6.129S4-Bdnf^{tm1.1ae}/J) mice (The Jackson Laboratory). Mice were genotyped as recommended by The Jackson Laboratory. Mice were group-housed and maintained on a 12-hour-day/-night cycle (lights on 7:00 AM) with constant access to food and water. All procedures were approved by the University of Texas Health Science Center-San Antonio Institutional Animal Care and Use Committee and carried out in accordance with the Guide for the Care and Use of Laboratory Animals as adopted by the National Institutes of Health.

Male wild-type C57BL/6J and BDNF^{+/-} mice, 4 to 5 months of age, were subjected to UMS, which consisted of: Day 1, cage shaking (2 hours); Day 2, switching cages with other mice in the cohort (1 hour); Day 3, 45° cage tilt (8 hours); Day 4, wet bedding

(3 hours); Day 5, exposure to rat odor (3 hours); Day 6, crowding; Day 7, strobe light and loud noises in dark (1hr); Day 8, restraint (2 hrs). Mice were sacrificed 24 hours after the last stress and forebrain collected for determination of cytokine mRNA by real time reverse transcription-polymerase chain reaction (RT-PCR) using the 2^{- $\Delta\Delta$ Ct} method with glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as an endogenous control gene as we have previously described (Lawson et al., 2013) and protein analysis by Luminex-based multiplex array according to the manufacturer's instructions (ProcartaPlex Mouse Cytokine/Chemokine Panel 1, eBioscience; intra-assay CV <5%). For RT-PCR experiments, inventoried gene assays were purchased from Life Technologies (Grand Island, NY; indoleamine 2,3-dioxygenase1, Mm00492586; tryptophan 2,3-dioxygenase, mM00451266; kynurenine monooxygenase, Mm00505511; kynurenine aminotransferase 2, Mm00496169). In experiments examining the effect of UMS on the kynurenine pathway, mRNA for kynurenine pathway enzymes was determined by RT-PCR as previously described (Heisler et al., 2015), and metabolite levels were determined by liquid chromatography/mass spectrometry as described (Walker et al., 2013; Mass Spectrometry Core Facility, University of Texas Health Science Center-San Antonio).

Results

We found no change in proinflammatory cytokine IL-1 β , IL-6, or TNF- α mRNA or protein in forebrain of wild-type or BDNF^{+/-} mice following the UMS procedure. We did, however, observe an increase in antiinflammatory cytokine IL-10 mRNA in wild-type mice following UMS, but IL-10 mRNA expression in BDNF^{+/-} mice was not different that wild-type controls following the UMS procedure (data expressed as fold change relative to the wild-type no-stress control: wild-type = 1.10 \pm 0.21, BDNF^{+/-} = 1.72 \pm 0.58; mean \pm SD, n = 6 per genotype; P = .034). We found no differences in baseline expression of IL-10 between wild-type and BDNF^{+/-} mice in forebrain IL-10 mRNA (data not shown).

In naïve mice, we found no difference between wild-type and BDNF^{+/-} mice in forebrain mRNA levels of kynurenine pathway enzymes. Eight days of UMS, however, resulted in increased expression of the 2 enzymes responsible for the metabolism of tryptophan to kynurenine, indoleamine 2,3-dioxygenase (IDO), and tryptophan 2,3-dioxygenase (TDO), in hippocampus of wild-type mice (Figure 1A-B, E). Concurrently, the expression of kynurenine monooxygenase (KMO) was downregulated (Figure 1C), which would be expected to decrease kynurenine metabolism in the neurotoxic branch of the kynurenine pathway in microglia (Figure 1E). Kynurenine aminotransferase (KAT) expression was not altered (Figure 1D). Collectively, these stress-induced changes in the expression of enzymes of the kynurenine pathway in wild-type mice favor the formation of kynurenic acid (Figure 1E).

In naïve mice, we found no difference between wild-type and BDNF^{+/-} mice in forebrain levels of kynurenine pathway metabolites, suggesting that there is no difference in activation of the kynurenine pathway between wild-type and BDNF^{+/-} mice at baseline. In wild-type mice, 8 days of UMS robustly increased kynurenine levels in brain, which is reflected in a significant increase in the kynurenine/tryptophan ratio (Figure 2A). The increase in the kynurenine/tryptophan ratio is indicative of increased metabolism of tryptophan to kynurenine and is consistent with the upregulation of IDO and TDO in wild-type mice following UMS (Figure 1). Kynurenine is not neurochemically active per se, but serves as a precursor for the synthesis of neurochemically active metabolites. Importantly, kynurenic acid levels were increased in wild-type mice following UMS




Figure 1. In wild-type (WT) mice, low-level stressor induced changes in the expression of enzymes in the kynurenine pathway that favor the formation of kynurenic acid. WT or brain-derived neurotrophic factor (BDNF^{-/-}) mice were subjected to 8 days of unpredictable mild stress (UMS) and sacrificed 24 hours after the last stress. RNA was isolated from forebrain. Real time reverse transcription (RT)-PCR data are expressed as relative fold change using the $2^{-\Delta\Delta Ct}$ calculation method. Mean \pm SEM, $n=7$ /group. One-way ANOVA, * $P < .05$ posthoc Tukey's Multiple Comparisons Test.

(Figure 2B). This may be due in part to the downregulation of KMO in wild-type mice following UMS (Figure 1C). In striking contrast to what was observed in wild-type mice, 8 days of UMS resulted in a marked increase in 3-hydroxykynurenine in BDNF^{-/-} mice (Figure 2C).

Discussion

In the present study, we found that low-level stress resulted in a marked increase in the expression of the antiinflammatory cytokine IL-10 in wild-type mice. Interestingly, this was not

observed in BDNF-deficient mice following UMS. IL-10, secreted by both microglia and astrocytes (Ledebuer et al., 2002; Henry et al., 2009), negatively modulates microglia activation (Norden et al., 2014). Our data suggest that this protective mechanism present in wild-type mice is absent in BDNF^{-/-} mice.

Both microglia and astrocytes express receptors for IL-10 (Ledebuer et al., 2002). While microglia activation is attenuated by IL-10, astrocytes are redirected to secrete antiinflammatory factors, which further inhibit microglia activity (Norden et al., 2014). In addition to providing negative feedback to activated microglia, astrocytes produce many regulatory factors that are




Figure 2. Low-level stressor resulted in a marked increase in (A) kynurenine and (B) kynurenic acid in wild-type (WT) mice. By contrast, low-level stressor resulted in a marked increase in (C) 3-hydroxykynurenine in brain-derived neurotrophic factor (BDNF)^{+/-} mice. Mice were subjected to 8 days of unpredictable mild stress (UMS) and sacrificed 24 hours after the last stress. Forebrain samples were prepared as described (Walker et al., 2013) and kynurenine metabolites analyzed by liquid chromatography/mass spectrometry. n = 7/group. One-way ANOVA, *P < .05 posthoc Tukey's Multiple Comparisons Test.

neuroprotective. Astrocytes express KAT, an enzyme of the kynurenine pathway that produces kynurenic acid, an ionotropic glutamate receptor antagonist (Guillemin et al., 2001; Schwarcz and Pellicciari, 2002). By contrast, microglia do not express KAT but do express KMO and therefore preferentially produce excitotoxic metabolites 3-hydroxykynurenine and quinolinic acid when activated (Guillemin et al., 2001; Schwarcz and Pellicciari, 2002). Thus, the neuroprotective and neurotoxic branches of the kynurenine pathway are compartmentalized in astrocytes and microglia, respectively.

In wild-type mice, UMS resulted in an upregulation of IDO and TDO expression and an increase in the kynurenine/tryptophan ratio, indicating stimulation of the kynurenine pathway. UMS also resulted in the downregulation of KMO expression in wild-type mice, potentially through IL-10-mediated inhibition of microglia (Norden et al., 2014). Of note, systemic administration of kynurenine to naïve mice, without the concurrent downregulation or inhibition of KMO, produces a depressive-like behavioral phenotype (O'Connor et al., 2009; Salazar et al., 2012). Our data indicate that in wild-type mice, low-level stress results in increased kynurenine metabolism through the neuroprotective branch of the kynurenine pathway in astrocytes. Consistent with this, UMS resulted in a marked increase in the formation of kynurenic acid.

In BDNF^{+/-} mice, where UMS did not upregulate IL-10 expression, we found no change in IDO or TDO expression and no increase in the kynurenine/tryptophan ratio. Following UMS, KMO was not downregulated, and we observed a marked increase in the excitotoxic metabolite 3-hydroxykynurenine. The neurotoxic metabolite 3-hydroxykynurenine contributes to oxidative stress and is a precursor of the NMDA receptor agonist quinolinic acid. Thus, in stress-sensitive BDNF^{+/-} mice (Duman et al., 2007; Advani et al., 2009; Burke et al., 2013), low-level stress results in kynurenine metabolism through the neurotoxic branch of the kynurenine pathway in microglia.

A single nucleotide polymorphism in the human *bdnf* gene (rs6265; val66met) is a risk factor associated with stress-precipitated psychiatric disorders, such as major depression (Hosang et al., 2014). The met allele results in impaired cellular processing and activity-dependent secretion of BDNF (Egan et al., 2003; Chen et al., 2004). Transgenic mice expressing the met allele exhibit a depressive-like behavioral phenotype with exposure to low-level stress (Yu et al., 2012). Further mechanistic studies are necessary to firmly establish the relationship between the kynurenine pathway and stress-precipitated depressive-like phenotypes in mice as well as the role BDNF may play in

the modulation of the kynurenine pathway. Such studies are necessary to increase our understanding of the cellular and molecular mechanisms by which individuals may be at risk for stress-precipitated mental illness. This work may inform clinical practice by leading to the identification of peripheral biomarkers for stress-related mental disorders and of novel therapeutic targets to restore equilibrium in the brain kynurenine pathway.

Acknowledgments

The authors gratefully acknowledge the excellent technical support of the Mass Spectrometry Core Facility, University of Texas Health Science Center-San Antonio.

This work was supported by the Institute for Integration of Medicine and Science at the University of Texas Health Science Center-San Antonio (J.G.H.), the Norman Hackerman Advanced Research Program (J.C.O.), and the National Institutes of Health (T32 NS 0821451 [A.M.D.], F31MH102070-01 [J.M.P.], and MH090127 [JCO]).

Statement of Interest

None.

References

- Advani T, Koek W, Hensler JG (2009) Gender differences in the enhanced vulnerability of BDNF^{+/-} mice to mild stress. *Int J Neuropsychopharmacol* 12:583–588.
- Altar CA (1999) Neurotrophins and depression. *Trends Pharmacol Sci* 20: 59–61.
- Burke TF, Advani T, Adachi M, Monteggia LM, Hensler JG (2013) Sensitivity of hippocampal serotonin_{1A} receptors to stress in mice deficient in brain-derived neurotrophic factor. *Int J Neuropsychopharm* 16:631–645.
- Calabrese F, Molteni R, Racagni G, Riva MA (2009) Neuronal plasticity: a link between stress and mood disorders. *Psychoneuroendocrinology* 34 [Suppl 1]:S208–S216.
- Capuron L, Miller AH (2011) Immune system to brain signaling: neuropsychopharmacological implications. *Pharmacol Ther* 130:226–238.
- Chen ZY, Patel PD, Sant G, Meng CX, Teng KK, Hempstead BL, Lee FS (2004) Variant brain-derived neurotrophic factor (BDNF) (Met66) alters the intracellular trafficking and activity-dependent secretion of wild-type BDNF in neurosecretory cells and cortical neurons. *J Neurosci* 24:4401–4411.

- Chourbaji S, Hellweg R, Brandis D, Zorner B, Zacher C, Lang UE, Henn FA, Hortnagl H, Gass P (2004) Mice with reduced brain-derived neurotrophic factor expression show decreased choline acetyltransferase activity, but regular brain monoamine levels and unaltered emotional behavior. *Brain Res Mol Brain Res* 121:28–36.
- Deak T, Bordner KA, McElderry NK, Barnum CJ, Blandino P Jr, Deak MM, Tammariello SP (2005) Stress-induced increases in hypothalamic IL-1: a systematic analysis of multiple stressor paradigms. *Brain Res Bull* 64:541–556.
- Duman CH, Schlesinger L, Kodama M, Russell DS, Duman RS (2007) A role for MAP kinase signaling in behavioral models of depression and antidepressant treatment. *Biol Psychiatry* 61:661–670.
- Duman RS, Monteggia LM (2006) A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 59:1116–1127.
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Koachana BS, Bertolino A, Zaitsev E, Gold B, Goldman D, Dean M, Lu B, Weinberger DR (2003) The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 112:257–269.
- Guillemin GJ, Kerr SJ, Smythe GA, Smith DG, Kapoor V, Armati PJ, Croitoru, J, Brew BJ (2001) Kynurenine pathway metabolism in human astrocytes: a paradox for neuronal protection. *J Neurochem* 78:842–853.
- Heisler JM, O'Connor JC (2015) Indoleamine 2,3-dioxygenase-dependent neurotoxic kynurenine metabolism mediates inflammation induced deficit in recognition memory. *Brain Behav Immun* June 27 Epub. doi: 10.1016/j.bbi.2015.06.022.
- Henry CJ, Huang Y, Wynne AM, Godbout JP (2009) Peripheral lipopolysaccharide (LPS) challenge promotes microglial hyperactivity in aged mice that is associated with exaggerated induction of both pro-inflammatory IL-1 β and anti-inflammatory IL-10 cytokines. *Brain Behav Immun* 23:309–317.
- Hosang GM1, Shiles C, Tansey KE, McGuffin P, Uher R (2014) Interaction between stress and the BDNF Val66Met polymorphism in depression: a systematic review and meta-analysis. *BMC Med* 12:7.
- Lawson MA, Parrott JM, McCusker RH, Dantzer R, Kelley KW, O'Connor JC (2013) Intracerebroventricular administration of lipopolysaccharide induces indoleamine-2,3-dioxygenase-dependent depression-like behaviors. *J Neuroinflammation* 10:87.
- Ledeboer A, Brevé JJ, Wierinckx A, van der Jagt S, Bristow AF, Leysen JE, Tilders FJ, Van Dam AM (2002) Expression and regulation of interleukin-10 and interleukin-10 receptor in rat astroglial and microglial cells. *Eur J Neurosci* 16:1175–1185.
- Norden D M, Fenn AM, Dugan A, Godbout JP (2014) TGF β produced by IL-10 redirected astrocytes attenuates microglial activation. *Glia* 62:881–895.
- O'Connor JC, Lawson MA, Andre C, Moreau M, Lestage J, Castanon N, Kelley KW, Dantzer R (2009). Lipopolysaccharide-induced depressive-like behavior is mediated by indoleamine 2,3-dioxygenase activation in mice. *Mol Psychiatry* 14:511–522.
- Parrott JM, Redus L, O'Connor JC (2012) Brain-derived neurotrophic factor deficient mice exhibit vulnerability to lipopolysaccharide-induced neuropsychiatric-like behaviors. Program No. 660.13/R7. Society for Neuroscience, 2012.
- Patki G, Solanki N, Atrooz F, Allam F, Salim S (2013) Depression, anxiety-like behavior and memory impairment are associated with increased oxidative stress and inflammation in a rat model of social stress. *Brain Res* 1539:73–86.
- Raison CL, Capuron L, Miller AH (2006) Cytokines sing the blues: inflammation and the pathogenesis of depression. *Trends Immunol* 27:24–31.
- Salazar A, Gonzalez-Rivera BL, Redus L, Parrott JM, O'Connor JC (2012) Indoleamine 2,3-dioxygenase mediates anhedonia and anxiety-like behaviors caused by peripheral lipopolysaccharide immune challenge. *Horm Behav* 62:202–209.
- Schwarcz R, Pellicciari R (2002) Manipulation of brain kynurenes: glial targets, neuronal effects, and clinical opportunities. *J Pharmacol Exp Ther* 303:1–10.
- Walker AK, Budac DP, Bisulco S, Lee AW, Smith RA, Beenders, B, Kelley KW, Dantzer R (2013) NMDA receptor blockade by ketamine abrogates lipopolysaccharide-induced depressive-like behavior in C57BL/6J mice. *Neuropsychopharmacology* 38:1609–1616.
- Wohleb ES, Hanke ML, Corona AW, Powell ND, Stiner LM, Bailey MT, Nelson RJ, Godbout JP, Sheridan JF (2011) β -Adrenergic receptor antagonism prevents anxiety-like behavior and microglial reactivity induced by repeated social defeat. *J Neurosci* 31:6277–6288.
- Yu H, Wang DD, Wang Y, Liu T, Lee FS, Chen ZY (2012) Variant brain-derived neurotrophic factor Val66Met polymorphism alters vulnerability to stress and response to antidepressants. *J Neurosci* 32:4092–4101.