#### **REVIEW**



# Barbaloin: an amazing chemical from the 'wonder plant' with multidimensional pharmacological attributes

Shreya Sikdar Mitra<sup>1</sup> · Mimosa Ghorai<sup>1</sup> · Samapika Nandy<sup>1</sup> · Nobendu Mukherjee<sup>2</sup> · Manoj Kumar<sup>3</sup> · Radha<sup>4</sup> · Arabinda Ghosh<sup>5</sup> · Niraj Kumar Jha<sup>6,7,8</sup> · Jarosław Proćków<sup>9</sup> · Abhijit Dey<sup>1</sup>

Received: 17 April 2022 / Accepted: 9 September 2022 / Published online: 29 September 2022 © The Author(s) 2022

#### **Abstract**

Aloe vera (L.) Burm.f. is nicknamed the 'Miracle plant' or sometimes as the 'Wonder plant'. It is a plant that has been used since ancient times for the innumerable health benefits associated with it. It is one of the important plants that has its use in conventional medicinal treatments. It is a perennial succulent, drought-tolerant member of the family Asphodelaceae. There are scores of properties associated with the plant that help in curing various forms of human ailments. Extracts and gels obtained from plants have been shown to be wonderful healers of different conditions, mainly various skin problems. Also, this plant is popular in the cosmetics industry. The underlying properties of the plant are now mainly associated with the natural phytochemicals present in the plant. Diverse groups of phytoingredients are found in the plant, including various phenolics, amino acids, sugars, vitamins, and different other organic compounds, too. One of the primary ingredients found in the plant is the aloin molecule. It is an anthraquinone derivative and exists as an isomer of Aloin A and Aloin B. Barbaloin belonging to the first group is a glucoside of the aloe-emodin anthrone molecule. Various types of pharmacological properties exhibited by the plant can be attributed to this chemical. Few significant ones are antioxidant, anti-inflammatory, anti-diabetic, anti-cancer, anti-microbial, and anti-viral, along with their different immunity-boosting actions. Recently, molecular coupling studies have also found the role of these molecules as a potential cure against the ongoing COVID-19 disease. This study comprehensively focuses on the numerous pharmacological actions of the primary compound barbaloin obtained from the Aloe vera plant along with the mechanism of action and the potent application of these natural molecules under various conditions.

**Keywords** Aloe vera · Barbaloin · Aloins · Anthraquinone · Pharmacological actions · Medicinal uses

- Abhijit Dey abhijit.dbs@presiuniv.ac.in
- Department of Life Sciences, Presidency University, 86/1 College Street, Kolkata 700073, West Bengal, India
- Department of Health Sciences, Novel Global Community Educational Foundation, Hebersham, Australia
- Ohemical and Biochemical Processing Division, ICAR-Central Institute for Research On Cotton Technology, Mumbai 400019, Maharashtra, India
- School of Biological and Environmental Sciences, Shoolini University of Biotechnology and Management Sciences, Solan 173229, Himachal Pradesh, India

- Department of Botany, Gauhati University, 781014 Guwahati, Assam, India
- Department of Biotechnology, School of Engineering & Technology, Sharda University, Greater Noida, Uttar Pradesh 201310, India
- Department of Biotechnology Engineering and Food Technology, Chandigarh University, Mohali 140413, Punjab, India
- Department of Biotechnology, School of Applied & Life Sciences, Uttaranchal University, Dehradun 248007, Uttarakhand, India
- Department of Plant Biology, Institute of Environmental Biology, Wrocław University of Environmental and Life Sciences, Kożuchowska 5b, 51-631 Wrocław, Poland



#### Introduction

Aloe vera, a drought-tolerant, perennial succulent member of the family Xanthorrhoeaceae/Asphodelaceae/Liliaceae family having various similarities with Cactus sp., is one plant of its type that can be considered a blessing for humankind for centuries to date (Añibarro-Ortega et al. 2019). A native plant species of the hot and dry regions of North Africa and different Mediterranean countries has yet to be widely cultivated in the Indian subcontinent as well and has become a naturalized plant species of the region. Around the world, almost more than 300 varieties of the plant have been identified. Since time immemorial it has been used in alternative treatment approaches for its multifarious medicinal benefits and at times also finds use as an ornamental plant (Khare et al. 2021; Anand et al. 2022a). It is often called the 'Miracle Plant' and also at times the 'wound healer' due to its amazing ability to counter various human ailments and in turn heal and comfort man, and hence the names (Habeeb et al. 2007). Derived from the Arabic word 'Alloeh', the name Aloe vera means 'bitter substance with shine'. Two thousand years back, Greek scientists considered this plant to be the universal panacea. The Egyptians named it 'the plant of immortality'. The mucilage gel-like tissue obtained from the leaves of the Aloe plant has wide uses in the cosmetic industry as the Aloe gel and also has potent therapeutic and medicinal applications. Planned, detailed scientific investigations on various medical applications of A. vera as a medicinal plant have gained significant concentration, and research is carried out all over the globe on the different phytochemicals, their isolation, characterization, analysis, and estimation for their diverse use as the pharmaceutical as well as nutraceutical agent. This particular study aims at providing an overview of the multifaceted attributes of one of the chief chemical components of Aloe vera, Barbaloin that revolves primarily around the biological, medicinal, and phytochemical parameters. Focus is given on the effect of chemicals against various diseases and conditions, with some emphasis on the mode of action as well.

#### Taxonomic position of *Aloe vera*

Kingdom: Plantae

Clade: Tracheophytes Clade: Angiosperms Clade: Monocots Order: Asparagales

Family: Xanthorrhoeaceae/Asphodelaceae/Liliaceae

Subfamily: Asphodeloideae



Genus: Aloe

Species: vera (L.) Burm.f.

# Chemical components of Aloe vera

The primary chemical present in the Aloe vera plant is a bitter chemical that when dried produces a yellow color belonging to the class called Aloin, which, as per historical shreds of evidence, shows the potency to be used as the cathartic medicines. Aloe vera is made up of 99 to 99.5% water and has a pH of around 4.5. Aloins can again be classified into two groups according to the chemical compositions: (1) Nataloins that react with nitric acid produce oxalic acid and picric acid and do not produce any red colour; (2) Barbaloin that in reaction with nitric acid forms aloetic acid (C7H2N3O5), chrysammic acid (C7H2N2O6), picric and oxalic acids and forms reddish colouration with the acid (Rajeshwari et al. 2012). Bright yellowish scale-like substances are formed by Nataloins, whereas yellowish prismatic crystals are formed by Barbaloin (Rajeshwari et al. 2012).

Different other phytochemicals were also identified from other parts of Aloe vera plants, viz. Anthraquinones/ anthrones (barbaloin, aloetic acid, aloe-emodin, anthranol, emodin, isobarbaloin, cinnamic acid esters); chromones [8-C-glusoly-(2'-O-cinnamoly)-7-omethlyaloediol A, 8-C-glucosyl-noreugenin, 8-Cglucosyl-7-O-methylaloediol A, 8-C-glucosyl-(S)-aloesol, 8-C-glucosyl-7-0-methylaloediol, isorabaichromone, neoalosin A, isoaloeresin D]; lipids and other organic compounds (Y-linolenic acid, arachidonic acid); steroids (cholesterol, campesterol, β-sitosterol); triterpenoids; triglycerides; lignins; gibberellin; salicylic acid; potassium sorbate; and uric acid (Minwuyelet et al. 2017). The plant houses vitamins A, B, B2, choline, niacin, C, F, and folic acid with traces of vitamin B12. Enzymes such as alkaline phosphatase, acid phosphatase, lactic dehydrogenase, amylase, and lipase were also present. Table 1 represents the main chemical ingredients of the Aloe vera plant, as well as the individual chemicals that are present.

#### Comprehensive applications of *Aloe vera*

There are numerous scientific studies that focus on the multiple uses of *Aloe vera* as a medicinal and cosmetic agent. There are different properties associated with the plant such as laxative, anti-microbial, analgesic, antioxidant, anti-inflammatory, anti-diabetic, anti-proliferative, anti-tumor, anti-mutagenic, wound healing, neuroprotective, hepatoprotective, immunoregulative, anti-aging, and radioprotective activities. Table 2 makes a comprehensive listing of a few

Table 1 The main chemical constituents found in the Aloe vera plant with the specific compounds present

Types of compounds	Name of compounds	Reference
Phenolic bitter compounds made of free anth- raquinones/anthrones and their derivative compounds	Barbaloin, aloe-emodin, isobarbaloin, aloin A and B, anthrone C-glycosides, chromones	Rodríguez et al. 2010; Quispe et al. 2018; Shakib et al. 2019
Vitamins	A, B <sub>1</sub> , B <sub>2</sub> , B <sub>6</sub> , B <sub>9</sub> , B <sub>12</sub> , C, E	Maan et al. 2018; Kahramanoğlu et al. 2019
Minerals	Potassium, sodium, calcium, magnesium, manganese, copper, zinc, iron, selenium	Rahmani et al. 2015; Sánchez et al. 2020
Sterols	Campesterol, sitosterol, lupeol	Ahlawat and Khatkar, 2011; Bawankar et al. 2013
Enzymes	Amylase, lipase, catalase, peroxidase, carboxy- peptidase, cyclo-oxidase	Ray et al. 2013; Radha and Laxmipriya 2015; Hęś et al. 2019
Sugars	Long-chain polysaccharides containing glucose and mannose combining together to form gluco-mannans	Hamman, 2008; Zhang et al. 2018
Essential and non-essential amino acids	Arginine, aspartic acid, alanine, glutamic acid, histidine, glycine, leucine, lysine, proline, methionine, tyrosine, threonine	Quispe et al. 2018; Martínez-Sánchez et al. 2020

Table 2 Multiple properties and uses of Aloe vera

Properties	Functions	Reference
Laxative effect	Effective as a cure for constipation, irritable bowel syndrome, indigestion, and other intestinal disorders	Foster et al. 2011; Hęś et al. 2019
Anti-microbial property	Effective against various classes of microorganisms such as virus and bacteria	Goudarzi et al. 2015; Forno-Bell et al. 2019; Arbab et al. 2021
Anti-inflammatory action	Effective as a cure for arthritis and other inflammations such as different types of wounds	Klaikeaw et al. 2020; Farid et al. 2021
Immuno-boosting action	Enhances immunity through various modes of actions that involve different immune cells and other associated chemicals	Gao et al. 2019; Mehrabi et al. 2019; Darzi et al. 2021
Antioxidant	Prevents cell damage by destroying unwanted free radi- cals and inhibits different harmful oxidation reactions within the cell	de Oliveira et al. 2018; López-Cervantes et al. 2018; Bendjedid et al. 2021; Kaparakou et al. 2021; Reza Nazifi et al. 2019
Anti-tumor	Inhibits tumor formation and potential malignant tumors	Majumder et al. 2019; Medina-Cruz et al. 2021; Xie et al. 2021
Anti-diabetic	Hypoglycemic effect by lowering insulin levels in the blood	Aldayel et al. 2020; Muñiz-Ramirez et al. 2020
Cosmetic uses	Moisturizing and anti-aging properties help in the use for skin care and various beauty related products	Pressman et al. 2019; Martínez-Burgos et al. 2022

of the well-known properties. Figure 1 presents the diverse pharmacological attributes of barbaloin.

## **Barbaloin**

One of the principal ingenious phytoconstituents of the *Aloe vera* plant is barbaloin (10-beta-D-glucopyranosyl-1, 8-dihydroxy-3-hydroxymethyl-9(10H)-anthracenone) (Groom and Reynolds, 1987). A yellowish-brown colour latex-based chemical with a persistent taste and having  $C_{21}H_{22}O_9$  as the molecular formula. It is the C-glucoside of the aloe-emodin anthrone, located mainly in the outer part of the plant. According to historical evidence, this compound found its

use as a laxative before World War II. A salient feature of this compound is the production of yellow fluorescence. The content of barbaloin is generally higher in the younger parts of the *Aloe* plant mainly in the leaves than in the older leaves. The diverse range of medicinal properties is associated with this particular compound having universal distribution and isolated from the *Aloe vera* plant.

This particular ingredient is responsible for the typical bitterness of bitter aloes and helps protect the plant against an attack by herbivores and acts as the defense molecule for the plants. According to a report, 30% of the dried leaf extract of the *Aloe vera* plant contains barbaloin and aloins. Barbaloin is found to occur mainly in two forms: the pure form and the impure form, which is available on the market



**Fig. 1** The diverse pharmacological attributes of barbaloin



as various types of formulations. Conventionally having its use as cathartics but also possessing several other properties such as antioxidant, anti-cancer, anti-inflammatory, antimicrobial, and anti-fungal (Zhang et al. 2017a; Brilhante et al. 2021).

Chemical structure of barbaloin

Aloin isolated from different natural sources is found to exist as a combination of two diastereomers, namely Aloin A (or barbaloin) and Aloin B (or isobarbaloin), with similar chemical attributes. Aloin, being an anthraquinone glycosyl is an anthraquinone containing a skeletal structure with the addition of one sugar moiety. Anthraquinones are naturally occurring phenolic compounds containing a 9,10-anthraquinone skeleton and are the common and largest family of indigenously occurring quinones that contain some of the significant chemicals such as emodin, chrysophanol, aloe-emodin, physcion, and rhein, that are found in various types of natural orange, yellow, and red pigments, and many possess the purgative action similar to that of aloins. Aloin was found to share

similarities with aloe-emodin that has similar biological characteristics as aloins but does not contain the sugar group. The chemical structures of some of the main bioactive compounds found in *Aloe yera* leaf extracts are listed below in Table 3.

# Methodology

The data presented in this manuscript were obtained from different available international databases and electronic resources such as Google Scholar, PubMed, Science Direct, Scopus, Wiley, etc. as well as from certain textbooks. The main keywords such as *Aloe vera*, Barbaloin, medicinal values of Barbaloin, pharmacological activity, chemical components, treatment, prevention, pharmacological activities, nutraceutical uses, health, clinical trials, and other related words were used singly or in combination. A thorough and careful reading of all sources of the available literature was carried out to obtain the best possible information.



Table 3 Chemical structures of the chief aloins present in the Aloe leaf extract

Name of the bioactive	Class of the compound	Structure of the compound
compound		
Barbaloin	Anthraquinone	HO OH OH OH OH Aloin A (Barbaloin)
Isobarbaloin	Anthraquinone	OH OH OH OH OH O OH Aloin B (Isobarbaloin)
Aloe emodin	Dihydroxanthraquinone	HO OH HO
Chromones	Phenylpropanoids	0

## **Barbaloin isolation and extraction methods**

Fresh leaves from the *Aloe* plant are collected mainly from some local nurseries, and then the outer portion of the leaves is peeled off and the leaves are cleaned by placing

them under clean water and thoroughly washing them. The inner gel obtained after the process can be scrapped and then smaller pieces cut from it. Various standardized protocols are available for barbaloin extraction, such as the Soxhlet extraction procedure, ultrasound-assisted



extraction methods, batch extraction experiments (Singh et al. 2020). An elaborate drying process is followed using the solar drying method for 3 weeks at 30–45 °C. Standard barbaloin was retrieved with the Fluka method and ultra-assisted sonication for the calibration process. Analysis by HPLC involves a diverse range of solvents of different grades manufactured by S.D. Fine Chemicals, India. Taiwan-made ultrasonic probes were used in the sonication technique and the probes were 20 kHz and had dimensions of 1.5 cm diameter or 30 cm length. Maintaining a constant temperature for the process is important. Subsequently, the samples were collected at certain time gaps to estimate the barbaloin quantity from the *Aloe vera* sample.

# Medicinal importance and pharmacological activities of barbaloin

As already mentioned above, Barbaloin is one of the primary and functional phytocomponents of the *Aloe vera* plant for its innumerable benefits. Due to the multifarious and beneficial properties of this compound, it is in high demand as a remedy for treatment of various diseases. Various research studies point to the efficacy of barbaloin due to its varied pharmacological and medicinal aspects. The usefulness of the compound against different ailments because of its various properties like antioxidant, anti-inflammatory, anti-cancer, anti-diabetic, and anti-microbial is discussed through this study.

#### **Antioxidant property**

Barbaloin has been found to possess an antioxidant property by helping in the process of destroying free radicals produced in the process of oxidative stresses taking place within the cell. It exhibited its antioxidant properties in the regions of free radical induced with the deoxyribonucleic acid (DNA) component (Singh et al. 2020). Barbaloin also offered protection against the OH-induced components of DNA and broke it compared to another control group. At lower concentrations, it improved DNA damage or indicated its antioxidant attributes. The highest effect of barbaloin is reduced in a minimum ratio. On the other hand, there was a gradual decrease in the power of the free radical effects of barbaloin, and this may have led to results to protect the component of deoxyribonucleic acid (Singh et al. 2020). The presence of enzymes glutathione peroxidase, superoxide dismutase enzymes, and phenolic antioxidants were responsible for the antioxidant property of Aloe vera and its phytocomponents (Sharma et al. 2014).



Various scientific reports are available that pointed to some of the anti-cancer activities of the aloins, especially the barbaloin. A study by Zhang et al. (2017a, b) showed that different types of conventional treatment approaches for non-small cell lung carcinoma (NSCLC), one of the most common types of lung cancer, had their own set of side effects (Mitra et al. 2022c; Anand et al. 2022b). However, in contrast, cells treated with barbaloin because of its anti-cancer potential were found to inhibit metastasis and cell growth both in vitro and in vivo and without any side effects as traditional treatment methods (Anand et al. 2020; Halder et al. 2021). An increase in the levels of different proteins related to apoptosis such as caspase-3, -8, and an -9 and alteration in the concentrations of various cyclin A, p53, p27, and other such checkpoint proteins associated with the cell cycle process gave convincing evidence in A549 cells regarding the antiviability effect of barbaloin. In addition, barbaloin was found to significantly cause a reduction in invasion and migration of A549 cells and showed controlled expression of proteins involved in the tumor metastasis process. Further studies were also conducted investigating the initiation of various pro-survival or pro-metastasis signaling pathways such as AKT, nuclear factor kappa B (NF-κB), mitogen-activated protein kinase (MAPK) and β-catenin. The results showed that barbaloin hindered the p38MAPK/Cdc25B/Hsp27 pathway by blocking nuclear translocation of p38, but no other notable effects were found in other pathways. Lastly, this compound was also found to inhibit in vivo liver metastases and growth of A549 cells in vivo. The results of these studies, if combined, showed the inhibiting action of barbaloin in vitro and in vivo of NSCLC cells that point to safer and more efficient treatment approaches of NSCLC (Zhang et al. 2017a).

A research study by El-Shemy et al. (2010) tried to analyze the possible anti-cancer and immune modulatory effects of different phytochemicals isolated from the Aloe vera plant. The primary anthraquinone molecules were selected and verectin, a N-terminal octapeptide derived from a biologically active 14 kDa glycoprotein present in A. vera, was also examined. In vivo active chemicals demonstrated a notable expansion of longevity of the tumortransplanted animals in the order: barbaloin > octapeptide > aloesin > aloe-emodin. The active principles of A. vera also showed substantial inhibition of the number of Ehrlich ascites carcinoma cells (EACC) in the order: barbaloin > aloe-emodin > octapeptide > aloesin. Furthermore, in the trypan blue cell viability assay, the active constituents exhibited remarkable concentration-dependent cytotoxicity against acute myeloid leukemia (AML) and acute lymphocyte leukemia (ALL) cell lines (El-Shemy et al. 2010). The data was strong enough to indicate the chemopreventive



activity of *A. vera* phytochemicals and that they can be potential, efficient, safe, specific, and economical treatment strategies. Therefore, because of the anti-neoplastic properties of barbaloin, aloe-emodin, these chemicals have been found to be used against the growth of malignant cells, and hence they have been used as a treatment for different types of cancers (Mitra et al. 2022a).

#### **Anti-inflammatory property**

Different studies aimed to show the anti-inflammatory action of barbaloin. These compounds had been shown to gradually retard the synthesis of nitric oxide by showing a negative effect on the expression of nitric oxide mRNA (Singh et al. 2020). Also, the effect was shown with respect to the manufacture of Prostaglandin E2. Studies showed that the inflammatory action of barbaloin was exhibited by reducing and restricting the mRNA expression of the nitric oxide synthase effect and the effect of cytooxygenase-2 mRNA in the body (Singh et al. 2020). The anti-inflammatory action of barbaloin was also found to contribute to its use in the case of inflammatory bowel disorder (Langmead et al. 2004). Due to its anti-inflammatory properties, Aloe vera has already been used in different types of inflammatory disorders such as gout, arthritis, etc. It is also used to strengthen joint muscles and reduces pain, inflammation, and redness. Often Aloebased phytochemicals have been used to reduce different types of arthritis (Nandal and Bhardwaj, 2012).

#### Anti-diabetic and hypoglycemic effect

As already discussed, *Aloe vera* extract is known to contain a high quantity of polyphenols with the highest ratio of barbaloin. Barbaloin administered by oral route in rats for a long time up to 4 weeks was effective in controlling blood glucose levels and reduced the weight of mice compared to the control group (Singh et al. 2020).

#### **Anti-microbial property**

Barbaloin was found to possess anti-microbial activities against different classes of microorganisms ranging from bacteria, protozoa, viruses, etc. The phytoconstituent barbaloin from the *Aloe* extract showed high anti-microbial efficacy against the bacterium *Neisseria gonorrhoeae*. Pure barbaloin had been found to inhibit the growth of the bacteria *Neisseria gonorrhoeae* responsible for causing sexually transmitted disease gonorrhoeae (Kambiz and Afolayan 2008; Khare et al. 2021). Barbaloin, when administered with phyto-extracts from *Withania somnifera*, was found to show anti-viral properties against the herpes simplex type 1 virus (Singh et al. 2020). Also, a study by Van et al. (2002) pointed at the anti-plasmodial property

of Barbaloin (Van et al. 2002). Aloe extract containing Aloins was found to exert an anti-mycobacterial effect on tuberculosis-causing bacteria Mycobacterium tuberculosis and a strong anti-bacterial effect against bacteria, namely Pseudomonas aeruginosa, Escherichia coli, Staphylococcus aureus, and Salmonella typhi (Banik and Sharangi, 2019). When taken in large quantities, the compounds were found to show a purgative effect, but when taken in little concentration they helped in the absorption process in the gut and act as anti-microbial agents (Banik and Sharangi, 2019).

#### **Anti-viral actions**

There is clear evidence in favor of the anti-viral actions of Aloe vera and its different components. There are several documentations regarding the anti-viral properties of Aloe vera and its associated phytochemicals. The gel fractions comprising lectins had shown a direct inhibitory effect on cytomegalovirus (CMC) in cell cultures (Saoo et al. 1996). Alves et al. (2004) reported that a chemical called chrysophanic acid obtained from A. vera at concentrations of 0.21 and 0.02 µg/mL demonstrated 50% inhibition of viral replication in type 2 and 3 polioviruses. Aloin had been found to exert an anti-viral rather virostatic effect on an RNAnegative enveloped virus HSV (hemorrhagic rhabdovirus septicemia virus) (Alves et al. 2004). Aloe vera was found to exert an inhibitory effect on a negative RNA enveloped viral hemorrhagic septicemia rhabdovirus (VHSV) (Alves et al. 2004). Vlietinck et al. (1998), showed the anti-viral effect of Aloe compounds like lectins on different viral groups including the human immunodeficiency virus (HIV).

#### In the treatment of various skin diseases

Skin is the largest, most widely, and most varied organ of the human body. It is present as the external barrier of the body that acts as a defense organ against various deleterious external factors found in the environment. Often, various harmful overlying symptoms are found to affect the skin mainly caused by some microbial activities and other modes of action too. Products obtained from the *Aloe* plant have diverse beneficial effects on the skin as well as the discussed compound barbaloin. Table 4 focuses on all such skin disorders as well as the preventive mode of action shown by the compound barbaloin, which may play a significant role in developing treatment strategies against such skin ailments.

#### **Immunity-boosting property**

One property that has gained high attention in modern pandemic times is the property of immunity boost. Humans are trying to consume different types of foods that can function as immunity and energy boosters (Paul et al. 2021; Mandal



Table 4 The use of barbaloin against a variety of skin disorders

Skin ailments	Signs and symptoms	Responsible property	References
Acne	The formation of closed and open plugged pores in the skin along with the development of pus-filled papules on skin	Anti-bacterial property	Akhoondinasab et al. 2014
Wounds and burns	Redness of the skin, blister formation, peeling of the skin, mild pain	Anti-inflammatory property	Moriyama et al. 2016
Cold sores	Itching, tingling, and burning sensation around the lips, formation of blisters	Anti-inflammatory property	Rajeshwari et al. 2012
Dandruff	Small white flakes on the scalp mainly around the hairlines, itching, irritation, inflammation	Anti-inflammatory, antioxidant, moisturizing properties	Mazzarello et al. 2018
Eczema	Itching, dryness, red to brownish-gray patches around eyelids, hands, and feet	Anti-microbial and anti-inflammatory properties	Klein and Penneys, 1988
Psoriasis	Red patches and silvery scale-like structures that cause inflammation, itching, and pain	Anti-bacterial and anti-inflammatory property	Laxmi et al. 2013
Rashes	Red in colour, inflammation, itching on the skin, dryness, formation of minute blisters filled with fluid	Anti-inflammatory and anti-microbial property	Singh et al. 2020
Sunburns	Pinkish to redness of the skin with irritation, pain, headaches, and at times fever also	Anti-inflammatory and moisturizing property	Qadir, 2009

et al. 2022). The *Aloe vera* plant has long been found to possess this quality and this feature can be associated with the principal phenolic compound barbaloin. The bitter aloes found in the *Aloe vera* plant are found to contain various types of free anthraquinones and their derivatives such as barbaloin, isobarbaloin, anthrone-C-glycosides, aloe-emodin-9-anthrone, and chromones as discussed previously. Studies suggest that barbaloin has immunity-boosting properties. It is found to enhance the immune system. Shreds of evidence suggest the use of these phytoingredients as immunomodulators and can hence be found effective against cold and cough conditions, fatigue, and other such related diseases associated with low immunity such as cold, fatigue, and other diseases caused by a low immune system. In larger quantities, these chemicals are found to possess purgative actions, but in minute amounts they act as a powerful antimicrobial agent and help in the process of absorption in the gut and many other related functions. Madan et al. (2008) in their study showed that mice when administered with Aloe vera extract (150 mg/kg and 300 mg/kg) respectively for 5 days showed a substantial increase in the total count of WBC and the number of macrophages count, indicating the immunomodulatory action of these compounds.

# **Barbaloin and COVID-19 disease**

As already discussed above, barbaloin is found to contain anti-viral properties, and in modern times this word virus is the most discussed one. Although vaccination and different forms of medications are already available against the pandemic disease (Anand et al. 2021a, b; Anand et al.

2021b; Iyer et al. 2021), the role of different nutraceuticals and various phyto-based foods is not less, too. Often, such phytochemicals are used in complementary alternative treatment approaches as prophylactic measures against this disease (Banerjee et al. 2021; Dutta et al. 2021). Various study reports indicate the efficacy of barbaloin against COVID-19 disease. A study conducted by Ho et al. (2007) pointed to the use of aloe-emodin against the novel coronavirus (SARS-CoV) and its ability to inhibit the SARS-CoV spike protein and the ACE-2 receptor protein. Abd-Alla et al. (2012) showed the use of phytoconstituents such as barbaloin and aloe-emodin against various hemagglutinating viruses. Molecular coupling studies identified various Aloe-based compounds as potential inhibitors of the main protease enzyme of SARS-CoV-2 in the SARS-CoV-2 virus (Mpiana et al. 2020a, b). Several other studies also pointed to the usage of aloin compounds as a potential candidate against COVID-19 disease, which can also be of great use in this time of pandemic in the near future. But a more detailed study about the mode of action and other areas is needed in this field.

#### **Discussion**

Through this review, we can find the multidimensional usage of the *Aloe* plant with special reference to one of its chief phyto-components Barbaloin. This single phenolic compound, like several other phytoingredients, is found to possess diverse properties, ranging from antioxidant, anti-inflammatory, antiseptic, anti-tumor, hypoglycemic, to moistening anti-aging properties as well. As a result, it is



very clear that if this single chemical can be given more focus and importance in terms of both research and commercialization, it can serve so many ailments and disorders (Anand et al. 2019). Although there is a significant amount of studies on this chemical, much more needs to be done in the field of isolation, purification, availability, and tissue culture techniques to grow the plant, as well as to harvest the particular compound and commercialization of this product. In the current perspective, this chemical is found to have anti-viral properties as well as immunity-boosting properties, too. Hence, it can be used in alternative treatment approaches too against viral diseases, COVID-19 disease being the most focused of the present time. Furthermore, the inclusion of *Aloe*-based products containing these aloins can also improve our immune system and help boost our immunity protecting us from various foreign attacks. Therefore, more research work is required on this wonder chemical from the miracle plant.

The entire process of research based on natural products-based research surrounds the applicability of different naturally obtained molecules (Mitra et al. 2022b). But in reality, it calls for a lot of patience and endurance. The entire process of natural products-based research is in itself time-consuming, and the desired results call for perseverance compared to those confined mainly to the synthetic molecules (Biswas et al. 2021). Although the quality associated with these products and their environmentally friendly impacts are much higher compared to synthetic ones. Hence, the speed of generating the molecules is a major constraint. But there is definitely a surplus of global pharmaceutical companies that are also focused on this avenue of research. Furthermore, the phytochemical purification of the obtained phytochemical is another important step that requires a lot of standardizations, and any error in the steps can disrupt the entire process of isolation and the obtaining of the natural compound. There are many more protocols based on tissue culture to cultivate plants and isolate the compounds. Therefore, more cutting-edge research needs to be done in this field to utilize this astounding chemical to its full potential.

#### **Conclusions**

Numerous applications of the miracle herb have been made possible in different cosmetic products, as well as a potential drug for the cure of different diseases along with traditional drugs (Bandopadhyay et al. 2022). Although there is evidence in respect of its uses, more and more controlled trials have to be carried out to find out the true potency along with the interaction between the herb and the drug. Several preclinical studies indicate the great potential of *Aloe*-based compounds as curative remedies against various diseases. However, more randomized clinical trials

are needed for the firm establishment of these efficacies and the realization of the utmost potential of the drug and its derived phytochemicals. To conclude, it can be said that Aloe vera is a very potent plant that produces various types of an extremely useful plant-based drug against various diseases including COVID-19 disease, also. Through discussion of various study results, the broad-range antiviral activity of the drug has been highlighted. It has been found to show efficacy against both DNA and RNA viruses. Anthraquinone-based phytochemicals such as barbaloin, isobarbaloin, and aloe-emodin are found to have anti-viral actions comparable with conventional anti-viral drugs such as Lopinavir and Ritonavir. The mechanism of action mainly centers on inhibition or binding to a variety of different viral proteins such as ACE-2 receptors, spike proteins, and 3CLPro. However, more research is needed in this field, and an elaborate mode of action must be deciphered to strongly establish these Aloe-based phytochemicals against various kinds of viruses. Therefore, the diverse range of activities of the principal phytoingredients of the Aloe vera plant truly justifies its name as 'wonder plant' or 'miracle plant'.

**Acknowledgements** The authors acknowledge respective departments and institutions for providing facilities and support.

Authors' contributions Shreya Sikdar Mitra came up with the study idea, planned and designed the review structure, wrote the first draft of the manuscript, and prepared the tables and final draft. Mimosa Ghorai and Samapika Nandy contributed to writing—review and editing, arranged the references, and prepared the tables. Nobendu Mukherjee participated in discussion, manuscript revision, and revision of the tables. Manoj Kumar contributed to overall revision of the manuscript, discussion, and suggestions. Radha revised the manuscript and response. Arabinda Ghosh and Niraj Kumar Jha revised the tables and thoroughly revised the manuscript and suggestions. Jarosław Proćków helped in formal interpretation, completed the critical revision of the entire manuscript and supervised the drafting process of the review, response, resources, final draft, project administration, and funding acquisition. Abhijit Dey came up with the study idea, revised the review structure and suggestions, completed the critical revision of the entire manuscript and formal interpretation, and supervised the drafting process of the review, resources, and final draft. All authors have read and approved the final version of the manuscript for submission to this journal.

**Funding** The Article Publishing Charge is financed by the Ministry of Education and Science, Warsaw, Poland.

Data availability Not applicable.

#### **Declarations**

**Ethics approval and consent to participate** Not applicable.

**Consent for publication** All authors have read and approved the final manuscript.



Competing interests The authors declare that they have no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

#### References

- Abd-Alla HI, Abu-Gabal NS, Hassan AZ, El-Safty MM, Shalaby NM (2012) Antiviral activity of Aloe hijazensis against some haemagglutinating viruses infection and its phytoconstituents. Arch Pharmacal Res 35(8):1347–1354
- Ahlawat KS, Khatkar BS (2011) Processing, food applications and safety of aloe vera products: a review. J Food Sci Technol 48(5):525–533
- Akhoondinasab MR, Akhoondinasab M, Saberi M (2014) Comparison of healing effect of *Aloe vera* extract and silver sulfadiazine in burn injuries in experimental rat model. World J Plast Surg 3(1):29
- Aldayel TS, Alshammari GM, Omar UM, Grace MH, Lila MA, Yahya MA (2020) Hypoglycaemic, insulin releasing, and hepatoprotective effect of the aqueous extract of Aloe perryi Baker resin (Socotran Aloe) in streptozotocin-induced diabetic rats. J Taibah Univ Sci 14(1):1671–1685
- Alves DS, Pérez-Fons L, Estepa A, Micol V (2004) Membrane-related effects underlying the biological activity of the anthraquinones emodin and barbaloin. Biochem Pharmacol 68(3):549–561
- Anand U, Jacobo-Herrera N, Altemimi A, Lakhssassi N (2019) A comprehensive review on medicinal plants as antimicrobial therapeutics: potential avenues of biocompatible drug discovery. Metabolites 9(11):258. https://doi.org/10.3390/metabo9110258
- Anand U, Nandy S, Mundhra A, Das N, Pandey DK, Dey A (2020) A review on antimicrobial botanicals, phytochemicals and natural resistance modifying agents from Apocynaceae family: Possible therapeutic approaches against multidrug resistance in pathogenic microorganisms. Drug Resist Updat 51:100695. https://doi.org/10. 1016/j.drup.2020.100695
- Anand U, Cabreros C, Mal J, Ballesteros F Jr, Sillanpää M, Tripathi V, Bontempi E (2021a) Novel coronavirus disease 2019 (COVID-19) pandemic: from transmission to control with an interdisciplinary vision. Environ Res 197:111126
- Anand U, Jakhmola S, Indari O, Chandra Jha H, Chen ZS, Tripathi V, Pérez de la Lastra JM (2021b) Potential therapeutic targets and vaccine development for COVID-19 management: A review on the recent update. Front Immunol 12:2454
- Anand U, Bianco F, Suresh S, Tripathi V, Núñez-Delgado A, Race M (2021) SARS-CoV-2 and other viruses in soil: An environmental outlook. Environ Res. 198:111297. https://doi.org/10.1016/j. envres.2021.111297
- Anand U, Tudu CK, Nandy S, Sunita K, Tripathi V, Loake GJ, Proćków J (2022a) Ethnodermatological use of medicinal plants in India: From Ayurvedic formulations to clinical perspectives—A review. J Ethnopharmacol 284:114744

- Anand U, Biswas P, Kumar V, Ray D, Ray P, Loake VIP, Kandimalla R, Chaudhary A, Singh B, Routhu NK, Chen ZS, Procków J, Dey A (2022b) Podophyllum hexandrum and its active constituents: Novel radioprotectants. Biomed Pharmacother. 146:112555. https://doi.org/10.1016/j.biopha.2021.112555
- Añibarro-Ortega M, Pinela J, Barros L, Ćirić A, Silva SP, Coelho E ... and Ferreira IC (2019) Compositional features and bioactive properties of Aloe vera leaf (Fillet, mucilage, and rind) and flower. Antioxidants 8(10):444
- Arbab S, Ullah H, Weiwei W, Wei X, Ahmad SU, Wu L, Zhang J (2021) Comparative study of antimicrobial action of Aloe vera and antibiotics against different bacterial isolates from skin infection. Vet Med Sci 7(5):2061–2067
- Bandopadhyay S, Anand U, Gadekar VS, Jha NK, Gupta PK, Behl T, Kumar M, Radha SMS, Dey A (2022) Dioscin: A review on pharmacological properties and therapeutic values. BioFactors 48(1):22–55. https://doi.org/10.1002/biof.1815
- Banerjee S, Anand U, Ghosh S, Ray D, Ray P, Nandy S, Deshmukh GD, Tripathi V, Dey A (2021) Bacosides from Bacopa monnieri extract: an overview of the effects on neurological disorders. Phytother Res 35(10):5668–5679. https://doi.org/10.1002/ptr.7203
- Banik S, Sharangi AB (2019) Phytochemistry, health benefits and toxicological profile of Aloe. J Pharmacog Phytochem 8(3):4499–4506
- Bawankar R, Deepti VC, Singh P, Subashkumar R, Vivekanandhan G, Babu S (2013) Evaluation of bioactive potential of an Aloe vera sterol extract. Phytother Res 27(6):864–868
- Bendjedid S, Lekmine S, Tadjine A, Djelloul R, Bensouici C (2021) Analysis of phytochemical constituents, antibacterial, antioxidant, photoprotective activities and cytotoxic effect of leaves extracts and fractions of Aloe vera. Biocatal Agric Biotechnol 33:101991
- Biswas P, Anand U, Ghorai M, Pandey DK, Jha NK, Behl T, Kumar M, Radha SMS, Dey A (2021) Unraveling the promise and limitations of CRISPR/Cas system in natural product research: Approaches and challenges. Biotechnol J 9:e2100507. https://doi.org/10.1002/biot.202100507
- Brilhante RSN, dos Santos Araújo G, Fonseca XMQC, de Melo Guedes GM, de Aguiar L, de SouzaCollares Maia Castelo-Branco D, de Aguiar Cordeiro R, Sidrim JJC, Pereira Neto WA, Rocha MFG (2021) Antifungal effect of anthraquinones against Cryptococcus neoformans: detection of synergism with amphotericin B. Medical Mycology 59(6):564–570
- Darzi S, Paul K, Leitan S, Werkmeister JA, Mukherjee S (2021) Immunobiology and application of aloe vera-based scaffolds in tissue engineering. Int J Mol Sci 22(4):1708
- de Oliveira ACL, Tabrez S, Shakil S, Khan MI, Asghar MN, Matias BD, da Silva Batista JMA, Rosal MM, de Lima MMDF, Gomes SRF, de Carvalho RM (2018) Mutagenic, antioxidant and wound healing properties of Aloe vera. J Ethnopharmacol 227:191–197
- Dutta T, Anand U, Saha SC, Mane AB, Prasanth DA, Kandimalla R, Procków J, Dey A (2021) Advancing urban ethnopharmacology: a modern concept of sustainability, conservation and cross-cultural adaptations of medicinal plant lore in the urban environment. Conserv Physiol. 9(1):coab073. https://doi.org/10.1093/conphys/coab073
- El-Shemy HA, Aboul-Soud MAM, Nassr-Allah AA, Aboul-Enein KM, Kabash A, Yagi A (2010) Antitumor properties and modulation of antioxidant enzymes' activity by Aloe vera leaf active principles isolated via supercritical carbon dioxide extraction. Curr Med Chem 17(2):129–138
- Farid A, Tawfik A, Elsioufy B, Safwat G (2021) In vitro and in vivo anti-Cryptosporidium and anti-inflammatory effects of Aloe vera gel in dexamethasone immunosuppressed mice. Int J Parasitol Drugs Drug Resist 17:156–167
- Forno-Bell N, Bucarey SA, García D, Iragüen D, Chacón O, San Martín B (2019) Antimicrobial effects caused by aloe barbadensis



- miller on bacteria associated with mastitis in dairy cattle. Nat Prod Commun 14(12):p.1934578X19896670
- Foster M, Hunter D, Samman S (2011) Evaluation of the nutritional and metabolic effects of Aloe vera. In: Benzie IFF, Wachtel-Galor S (eds.) Herbal Medicine: Biomolecular and Clinical Aspects (2nd ed.). CRC, Boca Raton
- Gao Y, Kuok KI, Jin Y, Wang R (2019) Biomedical applications of Aloe vera. Crit Rev Food Sci Nutr 59(sup1):S244–S256
- Goudarzi M, Fazeli,M, Azad M, Seyedjavadi SS and Mousavi R (2015) Aloe vera gel: effective therapeutic agent against multidrug-resistant Pseudomonas aeruginosa isolates recovered from burn wound infections. Chemother Res Pract 2015
- Groom QJ, Reynolds T (1987) Barbaloin in Aloe species. Planta Med 53(04):345–348
- Habeeb F, Shakir E, Bradbury F, Cameron P, Taravati MR, Drummond AJ ... and Ferro VA (2007) Screening methods used to determine the anti-microbial properties of Aloe vera inner gel. Methods 42(4), 315-320
- Halder S, Anand U, Nandy S, Oleksak P, Qusti S, Alshammari EM, El-Saber Batiha G, Koshy EP, Dey A (2021) Herbal drugs and natural bioactive products as potential therapeutics: A review on pro-cognitives and brain boosters perspectives. Saudi Pharm J 29(8):879–907. https://doi.org/10.1016/j.jsps.2021.07.003
- Hamman JH (2008) Composition and applications of Aloe vera leaf gel. Molecules 13(8):1599–1616
- Hęś M, Dziedzic K, Górecka D, Jędrusek-Golińska A, Gujska E (2019) Aloe vera (L.) Webb.: natural sources of antioxidants–a review. Plant Foods Hum Nutr 74(3):255–265
- Ho TY, Wu SL, Chen JC, Li CC, Hsiang CY (2007) Emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction. Antiviral Res 74(2):92–101
- Iyer M, Tiwari S, Renu K, Pasha MY, Pandit S, Singh B, Raj N, Krothapalli S, Kwak HJ, Balasubramanian V, Jang SB, G DK, Uttpal A, Narayanasamy A, Kinoshita M, Subramaniam MD, Nachimuthu SK, Roy A, Valsala Gopalakrishnan A, Ramakrishnan P, Cho SG, Vellingiri B (2021) Environmental survival of SARS-CoV-2 A solid waste perspective. Environ Res. 197:111015. https://doi.org/10.1016/j.envres.2021.111015
- Kahramanoğlu İ, Chen C, Chen J, Wan C (2019) Chemical constituents, antimicrobial activity, and food preservative characteristics of Aloe vera gel. Agronomy 9(12):831
- Kambiz L, Afolayan AJ (2008) Extracts from Aloe ferox and Withania somnifera inhibit Candida albicans and Neisseria gonorrhoea. Afr J Biotechnol 7(1):12–15
- Kaparakou EH, Kanakis CD, Gerogianni M, Maniati M, Vekrellis K, Skotti E, Tarantilis PA (2021) Quantitative determination of aloin, antioxidant activity, and toxicity of Aloe vera leaf gel products from Greece. J Sci Food Agric 101(2):414–423
- Khare T, Anand U, Dey A, Assaraf YG, Chen ZS, Liu Z, Kumar V (2021) Exploring Phytochemicals for Combating Antibiotic Resistance in Microbial Pathogens. Front Pharmacol 12:720726. https://doi.org/10.3389/fphar.2021.720726
- Klaikeaw N, Wongphoom J, Werawatganon D, Chayanupatkul M, Siriviriyakul P (2020) Anti-inflammatory and anti-oxidant effects of aloe vera in rats with non-alcoholic steatohepatitis. World J Hepatol 12(7):363
- Klein AD, Penneys NS (1988) Aloe vera. J Am Acad Dermatol 18(4):714–720
- Langmead L, Makins RJ, Rampton DS (2004) Anti-inflammatory effects of Aloe vera gel in human colorectal mucosa in vitro. Aliment Pharmacol Ther 19(5):521–527
- Laxmi RJ, Karthikeyan R, Babu PS, Babu RN (2013) Formulation and evaluation of antipsoriatic gel using natural excipients. J Acute Dis 2(2):115–121
- López-Cervantes J, Sánchez-Machado DI, Cruz-Flores P, Mariscal-Domínguez MF, de la Mora-López GS, Campas-Baypoli ON

- (2018) Antioxidant capacity, proximate composition, and lipid constituents of Aloe vera flowers. J App Res Med Aromat Plants 10:93–98
- Maan AA, Nazir A, Khan MKI, Ahmad T, Zia R, Murid M, Abrar M (2018) The therapeutic properties and applications of Aloe vera: A review. J Herb Med 12:1–10
- Madan J, Sharma AK, Inamdar N, Rao HS and Singh R (2008) Immunomodulatory properties of Aloe vera gel in mice. IJGP 2(3)
- Mandal S, Ghorai M, Anand U, Roy D, Kant N, Mishra T, Mane AB, Jha NK, Lal MK, Tiwari RK, Kumar M, Radha GA, Bhattacharjee R, Proćków J, Dey A (2022) Cytokinins: a genetic target for increasing yield potential in the CRISPR era. Front Genet 13:883930. https://doi.org/10.3389/fgene.2022.883930
- Majumder R, Das CK, Mandal M (2019) Lead bioactive compounds of Aloe vera as potential anticancer agent. Pharmacol Res 148:104416
- Martínez-Burgos WJ, Serra JL, MarsigliaF RM, Montoya P, Sarmiento-Vásquez Z, Marin O, Gallego-Cartagena E, Paternina-Arboleda CD (2022) Aloe vera: From ancient knowledge to the patent and innovation landscape—A review. S Afr J Bot 147:993–1006
- Martínez-Sánchez A, López-Cañavate ME, Guirao-Martínez J, Roca MJ, Aguayo E (2020) Aloe vera flowers, a byproduct with great potential and wide application, depending on maturity stage. Foods 9(11):1542
- Mazzarello V, Donadu MG, Ferrari M, Piga G, Usai D, Zanetti S, Sotgiu MA (2018) Treatment of acne with a combination of propolis, tea tree oil, and Aloe vera compared to erythromycin cream: two double-blind investigations. Clin Pharmacol Adv App 10:175
- Medina-Cruz D, Vernet-Crua A, Mostafavi E, González MU, Martínez L, Jones AAD, Kusper M, Sotelo E, Gao M, Geoffrion LD, Shah V (2021) Aloe vera-mediated Te nanostructures: highly potent antibacterial agents and moderated anticancer effects. Nanomaterials 11(2):514
- Mehrabi Z, Firouzbakhsh F, Rahimi-Mianji G, Paknejad H (2019) Immunostimulatory effect of Aloe vera (Aloe barbadensis) on non-specific immune response, immune gene expression, and experimental challenge with Saprolegnia parasitica in rainbow trout (Oncorhynchus mykiss). Aquaculture 503:330–338
- Minwuyelet T, MogesSewalem M, Gashe M (2017) Review on therapeutic uses of Aloe vera. Global J Pharmacol 11(2):14–20
- Mitra S, Anand U, Jha NK, Shekhawat MS, Saha SC, Nongdam P, Rengasamy KRR, Proćków J, Dey A (2022) Anticancer Applications and Pharmacological Properties of Piperidine and Piperine: A Comprehensive Review on Molecular Mechanisms and Therapeutic Perspectives. Front Pharmacol. 12:772418. https://doi.org/10.3389/fphar.2021.772418
- Mitra S, Anand U, Ghorai M, Vellingiri B, Jha NK, Behl T, Kumar M, Radha Shekhawat MS, Proćków J, Dey A (2022) Unravelling the therapeutic potential of botanicals against chronic obstructive pulmonary disease (COPD): molecular insights and future perspectives. Front Pharmacol. 13:824132. https://doi.org/10.3389/fphar.2022.824132
- Mitra S, Anand U, Sanyal R, Jha NK, Behl T, Mundhra A, Ghosh A, Radha Kumar M, Proćków J, Dey A (2022) Neoechinulins: Molecular, cellular, and functional attributes as promising therapeutics against cancer and other human diseases. Biomed Pharmacother. 145:112378. https://doi.org/10.1016/j.biopha. 2021.112378
- Moriyama M, Moriyama H, Uda J, Kubo H, Nakajima Y, Goto A, Hayakawa T (2016) Beneficial effects of the genus Aloe on wound healing, cell proliferation, and differentiation of epidermal keratinocytes. PLoS ONE 11(10):e0164799
- Mpiana PT, Tshibangu DS, Kilembe JT, Gbolo BZ, Mwanangombo DT, Inkoto CL and Tshilanda DD (2020a) Aloe vera (L.) Burm.



- f. as a potential Anti-COVID-19 plant: a mini-review of its antiviral activity. European J Med Plants 86–93
- Mpiana PT, Tshibangu DS, Kilembe JT, Gbolo BZ, Mwanangombo DT, Inkoto CL, Lengbiye EM, Mbadiko CM, Matondo A, Bongo GN, Tshilanda DD (2020b) Identification of potential inhibitors of SARS-CoV-2 main protease from Aloe vera compounds: a molecular docking study. Chem Phys Letters 754:137751
- Muñiz-Ramirez A, Perez RM, Garcia E and Garcia FE (2020) Antidiabetic activity of Aloe vera leaves. Evid Based Complementary Altern Med 2020
- Nandal U, Bhardwaj RL (2012) Aloe vera: A valuable wonder plant for food, medicine and cosmetic use-a review. Int J Pharm Sci Rev Res 13(1):59–67
- Paul S, Chakraborty S, Anand U, Dey S, Nandy S, Ghorai M, Saha SC, Patil MT, Kandimalla R, Proćków J, Dey A (2021) Withania somnifera (L.) Dunal (Ashwagandha): A comprehensive review on ethnopharmacology, pharmacotherapeutics, biomedicinal and toxicological aspects. Biomed Pharmacother 143:112175. https://doi.org/10.1016/j.biopha.2021.112175
- Pressman P, Clemens R, Hayes AW (2019) Aloe vera at the frontier of glycobiology and integrative medicine: health implications of an ancient plant. SAGE Open Medicine 7:2050312119875921
- Qadir MI (2009) Medicinal and cosmetological importance of Aloe vera. Int J Nat Ther 2:21–26
- Quispe C, Villalobos M, Bórquez J, Simirgiotis M (2018) Chemical composition and antioxidant activity of Aloe vera from the Pica oasis (Tarapacá, Chile) by UHPLC-Q/Orbitrap/MS/MS. J Chemother 6123850:1–12
- Radha MH, Laxmipriya NP (2015) Evaluation of biological properties and clinical effectiveness of Aloe vera: a systematic review. J Tradit Complement Med 5(1):21–26
- Rahmani AH, Aldebasi YH, Srikar S, Khan AA, Aly SM (2015) Aloe vera: Potential candidate in health management via modulation of biological activities. Pharmacogn Rev 9(18):120
- Rajeswari R, Umadevi M, Rahale CS, Pushpa R, Selvavenkadesh S, Kumar KS, Bhowmik D (2012) Aloe vera: the miracle plant its medicinal and traditional uses in India. J Pharmacog Phytochem 1(4):118–124
- Ray A, Gupta SD, Ghosh S (2013) Evaluation of anti-oxidative activity and UV absorption potential of the extracts of Aloe vera L. gel from different growth periods of plants. Ind Crops Prod 49:712–719

- Reza Nazifi SM, Asgharshamsi MH, Dehkordi MM, Zborowski KK (2019) Antioxidant properties of Aloe vera components: a DFT theoretical evaluation. Free Radical Res 53(8):922–931
- Rodríguez ER, Martín JD, Romero CD (2010) Aloe vera as a functional ingredient in foods. Crit Rev Food Sci Nutr 50(4):305–326
- Sánchez M, González-Burgos E, Iglesias I, Gómez-Serranillos MP (2020) Pharmacological update properties of Aloe vera and its major active constituents. Molecules 25(6):1324
- Saoo K, Miki H, Ohmori M, Winters WD (1996) Antiviral activity of Aloe extracts against cytomegalovirus. Phytother Res 10(4):348-350
- Shakib Z, Shahraki N, Razavi BM, Hosseinzadeh H (2019) Aloe vera as an herbal medicine in the treatment of metabolic syndrome: A review. Phytother Res 33(10):2649–2660
- Sharma P, Kharkwal AC, Kharkwal H, Abdin MZ, Varma A (2014) A review on pharmacological properties of Aloe vera. Int J Pharm Sci Rev Res 29(2):31–37
- Singh N, Goyal K, Sondhi S, Jindal S (2020) Traditional and medicinal use of Barbaloin: potential for the management of various diseases. J Appl Pharma Res 8(3):21–30
- Van Zyl RL, Viljoen AM, Jäger AK (2002) In vitro activity of Aloe extracts against Plasmodium falciparum. S Afr J Bot 68(1):106-110
- Vlietinck AJ, De Bruyne T, Apers S, Pieters LA (1998) Plant-derived leading compounds for chemotherapy of human immunodeficiency virus (HIV) infection. Planta Med 64(02):97–109
- Xie J, Wu J, Yang S and Zhou H (2021) Network pharmacology-based study on the mechanism of aloe vera for treating cancer. Evid Based Complement Alternat Med 2021:8
- Zhang Y, Bao Z, Ye X, Xie Z, He K, Mergens B, Li W, Yatcilla M, Zheng Q (2018) Chemical investigation of major constituents in Aloe vera leaves and several commercial Aloe juice powders. J AOAC Int 101(6):1741–1751
- Zhang Z, Rui W, Wang ZC, Liu DX, Du L (2017) Anti-proliferation and anti-metastasis effect of barbaloin in non-small cell lung cancer via inactivating p38MAPK/Cdc25B/Hsp27 pathway. Oncol Rep 38(2):1172–1180

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

