

Review

Distribution in Different Organisms of Amino Acid Oxidases with FAD or a Quinone As Cofactor and Their Role as Antimicrobial Proteins in Marine Bacteria

Jonatan C. Campillo-Brocal, Patricia Lucas-Elío and Antonio Sanchez-Amat *

Received: 4 November 2015; Accepted: 8 December 2015; Published: 16 December 2015

Academic Editor: Vassilios Roussis

Department of Genetics and Microbiology, Faculty of Biology, University of Murcia, Murcia 30100, Spain;

jonatancristian.campillo@um.es (J.C.C.-B.); patlucel@um.es (P.L.-E.)

* Correspondence: antonio@um.es; Tel. +34-868-884-955; Fax +34-868-883-963

Abstract: Amino acid oxidases (AAOs) catalyze the oxidative deamination of amino acids releasing ammonium and hydrogen peroxide. Several kinds of these enzymes have been reported. Depending on the amino acid isomer used as a substrate, it is possible to differentiate between L-amino acid oxidases and D-amino acid oxidases. Both use FAD as cofactor and oxidize the amino acid in the alpha position releasing the corresponding keto acid. Recently, a novel class of AAOs has been described that does not contain FAD as cofactor, but a quinone generated by post-translational modification of residues in the same protein. These proteins are named as LodA-like proteins, after the first member of this group described, LodA, a lysine epsilon oxidase synthesized by the marine bacterium *Marinomonas mediterranea*. In this review, a phylogenetic analysis of all the enzymes described with AAO activity has been performed. It is shown that it is possible to recognize different groups of these enzymes and those containing the quinone cofactor are clearly differentiated. In marine bacteria, particularly in the genus *Pseudoalteromonas*, most of the proteins described as antimicrobial because of their capacity to generate hydrogen peroxide belong to the group of LodA-like proteins.

Keywords: amino acid oxidases; quinone cofactor; flavoprotein; antimicrobial

1. Introduction

In a broad sense, amino acid oxidases (AAOs) can be described as enzymes that oxidize amino acids releasing ammonium and hydrogen peroxide. Two big groups of these enzymes are recognized depending on the chirality of the amino acid used as substrate. L-Amino acid oxidases, EC 1.4.3.2, (commonly abbreviated as LAAOs or LAOs) are flavoenzymes that oxidize L-amino acids releasing the corresponding α -keto acid in addition to ammonium and hydrogen peroxide. LAAOs are distributed in many biological groups. The best characterized members of this family have been studied in snake venoms [1]. Additionally, several enzymes with this activity have also been described in many other groups including bacteria [2]. D-Amino acid oxidases (DAAOs or DAOs), EC 1.4.3.3, are flavoproteins showing strict specificity for D-amino acids [3,4]. These enzymes are also broadly distributed in different groups of organisms with distinct physiological roles. Glycine is the unique amino acid that has no enantiomers. In *Bacillus*, a glycine oxidase has been described [5]. Recently, a novel class of AAOs has been reported. The marine bacterium *Marinomonas mediterranea* synthesizes LodA, a L-lysine epsilon-oxidase (EC 1.4.3.20) [6,7]. The cofactor of LodA is cysteine tryptophylquinone (CTQ) (Figure 1), a type of quinone cofactor generated by the post-translational modification of two residues in the same protein [8,9]. GoxA is an enzyme with sequence similarity

to LodA and the same kind of cofactor, but it shows glycine oxidase activity [10,11]. The analysis of sequenced microbial genomes has revealed that, approximately, 1% of them contain genes encoding proteins similar to LodA and GoxA [12].




Figure 1. Cysteine tryptophylquinone (CTQ) cofactor.

Enzymes with AAO activity are of great scientific interest because of the number of physiological activities displayed in different organisms. For example, the generation of hydrogen peroxide has been described from a physiological point of view as involved in antimicrobial processes. Among these processes, bacterial biofilm development [13], microbial competition related to biocontrol processes in fungi [14], protection of fish skin bacterial infections [15] and participation in the human immune system [16] are included. AAOs are also of interest in relation to their biotechnological applications [4,17]. Recently, their potential interest in medicine as antitumorals and as antimicrobials has increased [18,19]. In both cases, the hydrogen peroxide generated as product of the reaction plays a very important role.

LAAOs have been known for more than half a century. In many cases, when an enzyme showed that activity it was assumed to be a flavoprotein. However, it is important to point out that not in all the cases it has been demonstrated that the activity is due to an enzyme with a flavin cofactor. It is also important to bear in mind that the enzymatic activity of a protein may depend on the presence of other compounds, as it has been described for an enzyme whose activity can change from L-amino acid oxidase to monooxygenase by either mutation or change in the conditions of the assay [20]. Besides, it is well known that many enzymes may show latent promiscuous activities, and that the actual enzymatic activity could be different to the one initially described. As an example, an enzyme from *Streptococcus oligofermentans* was initially described as an L-amino acid oxidase [21] and lately reclassified as an aminoacetone oxidase involved in antioxidant mechanisms [22].

The aim of this review has been to study the phylogenetic relationship of all the enzymes described as amino acid oxidases, including the new family with quinone cofactor, in order to facilitate future work on novel enzymes with that activity. In the last section, the enzymes synthesized by marine microorganisms described as antimicrobial proteins will be discussed.

2. Phylogenetic Analysis of Proteins with Amino Acid Oxidase Activity

A bibliographic search has been performed in order to obtain representative sequences of proteins with amino acid oxidase activity. All the microbial proteins considered are shown in supplementary Table 1. Table S1 includes proteins characterized at the molecular level, as well as other enzymes in which only the enzymatic activity has been reported. In order to shed light on the evolutionary relationships of AAOs, a phylogenetic analysis has been performed that includes the microbial proteins for which the encoding gene has been cloned, plus other representative proteins from higher organisms (Figure 2). Those AAOs from higher organisms are listed in supplementary Table 2.




Figure 2. Phylogenetic relationships of enzymes with amino acid oxidase activity. The tree was created by the Neighbor-Joining method integrated in the program MEGA6 [23]. Sequences were aligned using the program MUSCLE built in MEGA6. The evolutionary distances were computed using the *p*-distance method and are in the units of the number of amino acid differences per site. Numbers at branches indicate bootstrap values higher than 70% for both Neighbor-Joining and Maximum Likelihood trees. The colored groups are detailed in Figures 3 and 4. LAAOs, L-amino acid oxidases; DAAO, D-amino acid oxidases; LASPOs, L-aspartate oxidases.

It has been possible to recognize several closely related groups which are named after their common characteristic. The proteins similar to LodA containing a quinone cofactor form a phylogenetic group clearly differentiated from all the other enzymes. Among the rest of enzymes with AAO activity, DAAOs from different organisms constitute a well-defined cluster. Another group clearly differentiated is the one containing L-aspartate oxidases. Regarding LAAOs, they constitute a broad group distributed in different organisms where they have evolved to meet different physiological functions. In the next sections, the different groups will be discussed separately.

2.1. AAOs with a Quinone Cofactor (LodA-Like Proteins)

L-Lysine ϵ -oxidase (LodA) synthesized by the melanogenic marine bacterium *M. mediterranea* was the first quinoprotein reported with LAAO activity [7]. LodA received a new number by the Enzyme Commission (EC 1.4.3.20) since it catalyzes the oxidative deamination of the amine group in epsilon position of L-Lys [6]. It has been demonstrated that the cofactor of LodA is cysteine tryptophylquinone (CTQ) (Figure 1), which is generated by post-translational modification of residues in the same protein [9,11]. In the post-translational modification of LodA, the flavoprotein LodB, encoded in the same operon, plays an important role [8,24]. Regarding its physiological

function, it has been shown that LodA plays a role, mediated by the hydrogen peroxide generated, in the development and differentiation of microbial biofilms [13].

Another quinoprotein with AAO activity was identified in *M. mediterranea*. This is a novel glycine oxidase (GoxA) which shows important differences in terms of substrate range and sensitivity to inhibitors to other glycine oxidases previously described [10]. GoxA is more specific for glycine and structurally different since it also contains the cofactor CTQ [11]. Genome mining revealed genes encoding proteins similar to LodA/GoxA in a number of microbial genomes. They are mostly present in *Bacteria*, absent in *Archaea* and, in *Eukarya*, only detected in a small group of fungi of the class *Agaromycetes* [12]. This new family of proteins has been named as the LodA-like family of proteins. Sequence alignment of the LodA-like proteins allowed the detection of several conserved residues. All these proteins showed a Cys and a Trp that aligned with the residues that are forming part of the CTQ cofactor in LodA and GoxA. Thus, this observation strongly suggests that proteins of the LodA-like family may contain a CTQ quinone cofactor [12]. Apart from LodA and GoxA, various oxidases of this group have been characterized in marine bacteria as discussed later (Section 3).

2.2. D-Amino Acid Oxidases

D-Amino acid oxidases (EC 1.4.3.3) are FAD-containing proteins with a strict stereospecificity. They catalyze the oxidative deamination of neutral and basic D-amino acids to give α -keto acids, ammonium and hydrogen peroxide [3,25]. However, DAAOs show a negligible or no activity towards acidic D-amino acids, which are substrates for the closely related animal flavoproteins D-aspartate oxidases (DDO, also abbreviated as DASPO; EC 1.4.3.1) [26] (Figure 3B). DAAOs also cluster close to the glycine oxidases group (Figure 2). DAAO activity has been described in a wide variety of organisms such as bacteria [27], fungi [28], plants [29], nematodes [30], fishes [31], and mammals, including humans [32]. DAAOs participate in different physiological functions [33]. In yeast, they have a catabolic role since DAAOs allow yeast cells to grow on D-amino acids. In humans, DAAOs manage the levels of the neuromodulator D-serine of NMDA (*N*-methyl-D-aspartate) receptors, which are related with many pathological processes. However, the physiological role of bacterial DAAOs is still largely unknown [25]. DAAOs can be utilized for a broad range of applications, such as the determination of D-amino acids, production of building blocks for pharmaceuticals, synthesis of α -keto acids, and diagnosis and treatment of certain diseases [4,34].

2.3. L-Aspartate Oxidases

L-Aspartate oxidases (LASPOs; EC 1.4.3.16) are flavoproteins encoded by the gene *nadB* that play an important role in NAD⁺ synthesis in the cells, as they catalyze the formation of iminoaspartate from L-aspartate, the first step in the *de novo* biosynthesis of NAD⁺ in many bacterial, archaeal and plant cells [35]. NadB forms a reversible multienzyme complex with NadA, which carries on the transformation leading to quinolinic acid, an intermediary in the NAD⁺ synthetic pathway. The high activity of NadB seems to imply an oxidative stress for the cells, due to the generation of hydrogen peroxide [36]. It is noteworthy that, under physiological conditions, LASPOs do not produce ammonia, since the final product of the pathway conserves the original N-atom of L-aspartic acid. In this feature, they do not conform to the classical definition of amino acid oxidases. Interestingly, NadB function is substituted by an NADP-dependent L-aspartate dehydrogenase in *Thermotoga maritima* which has no sequence similarities with any amino acid oxidase. Moreover, the gene encoding that enzyme is next to *nadA* in the genome, as described in *Bacillus* [37].

Sequence analysis reveals that LASPOs have some similarities with fumarate reductases and succinate dehydrogenases in terms of folding topology and conformation of the FAD-binding site and active center, suggesting a similar catalytic mechanism [38,39] and a common evolutionary origin [40]. This implies that L-aspartate oxidases are different from other flavin-dependent amino acid oxidases, such as the D-amino acid oxidases. This fact is reflected as a separate and defined cluster in the phylogenetic tree (Figures 2 and 3C).

One of the similarities between LASPOs and the succinate dehydrogenase/fumarate reductase oxidoreductase family is their ability to use different electron acceptors [41]. Accordingly, NadB can use either molecular oxygen or fumarate as electron acceptors of the reaction, permitting aspartate oxidation in anaerobic conditions [42]. Other aspartate oxidase, such as those from archaea, are being widely proposed for bioanalytical methods due to their high thermostability [43,44]. For instance, the LASPO from *Sulfolobus tokodaii*, which has been recombinantly expressed, has a broad pH range of activity and shows a weak inhibition pattern. This enzyme not only has a high thermal stability but also binds tightly the FAD cofactor [43].

Although aspartate oxidases were proposed as potential drug targets, since they are absent in mammals [38], this research has not been followed, probably because prokaryotes may have alternative routes for NAD⁺ production [45]. On the other hand, the gene is considered as an antivirulence loci (AVL), either inactivated or lost in all pathogenic *Shigella* strains studied, as the presence of quinolinic acid is detrimental to its invasive process [46]. This pathoadaptive evolution is accompanied by a nicotinic acid auxotrophy that bypasses the need of quinolinic acid production in the NAD⁺ synthetic pathway.




Figure 3. Phylogenetic relationships of representative LodA-like proteins (A), D-amino acid oxidases (B) and L-aspartate oxidases (C); The tree was created by the Neighbor-Joining method integrated in the program MEGA6 [23]. Sequences were aligned using the program MUSCLE built in MEGA6. The evolutionary distances were computed using the *p*-distance method and are in the units of the number of amino acid differences per site. Numbers at branches indicate bootstrap values higher than 70% for both Neighbor-Joining and Maximum Likelihood trees. An asterisk indicates that this branch was not detected, or it had a value lower than 70%.

2.4. LAAOs in Animals

The LAAOs from animals clusters in two different groups. One of them includes enzymes from vertebrates, and the other group includes gastropod enzymes (Figure 2). This observation is in agreement with previous studies that indicated that those enzymes may have evolved separately in the innate immune system of both groups [18,47].

2.4.1. LAAOs from Vertebrates

The phylogenetic analysis of AAOS revealed a cluster including enzymes from snake venoms, fishes and mammals (Figures 2 and 4A). Venom LAAOs have been subject of study for many years and several reviews have been published dealing with their biochemical properties and their interest in pharmacology [1,48]. Venom LAAOs show preference towards hydrophobic amino acids such as L-Leu, L-Phe and L-Met [48]. Numerous biological effects, such as antiparasitic, antimicrobial, apoptotic, etc., have been reported for those enzymes [1,49].

The skin of the fish is a barrier preventing the infection by microorganisms. The mucus layer contains many peptides and proteins that play a role in that process. In this layer, LAAOs specific for L-lysine have been detected in different fishes such as the rockfish *Sebastes schlegelii* [15] and the great sculpin *Myoxocephalus polyacanthocephalus* [50]. The *Sebastes* LAAO has been used for the design of a biosensor for L-Lys [51]. LAAOs are not only present in the skin mucus but also in other tissues and the serum of fishes [52,53]. The induction of the LAAO of the Atlantic cod *Gadus morhua* after exposure of the fish to pathogenic bacteria supports its involvement in antibacterial defense [54].

Regarding mammals, the interleukin 4-induced gene (IL4I1) codes for an L-amino acid oxidase with strong preference for L-phenylalanine as substrate, which seems to act as a regulator of the immune system [55]. IL4I1 shows antimicrobial properties that have been associated to the hydrogen peroxide generated, the basification of the medium due to ammonia accumulation and the deprivation of the amino acid [16]. A LAAO has been detected in the milk of the mouse with substrate similarity to the human enzyme that has been proposed to participate in the protection against bacterial infections [56].

2.4.2. LAAOs in Gastropods

Several gastropods are known to produce LAAOs (Figure 4B). The sea hare *Aplysia californica*, synthesizes a LAAO named escatin that is present in a separate gland from its substrates (Lys and Arg). The mixing of the secretion of both glands at the time of the attack of the predator generates a defensive mechanism against them [57]. Escatin shows antimicrobial activity against a range of bacteria, even when it is recombinantly expressed [58]. It has been shown that the antimicrobial activity of escatin depends not only on the hydrogen peroxide generated, but also on chemical compounds generated during the oxidation of L-lysine [59,60]. Another sea hare (*Aplysia punctata*) synthesizes a very similar LAAO (93% identical to escatin) whose antitumoral activity has been characterized [61]. Achacin is a LAAO isolated and characterized as an antimicrobial protein from the skin mucus of the giant snail *Achatina fulica* [62]. The cytotoxicity of achacin is mediated by two different mechanisms: hydrogen peroxide generation and apoptosis induction mediated by L-amino acids depletion [63].

2.5. Fungal LAAOs

Different flavoproteins with L-amino acid oxidase activities have been described in fungi. The *Neurospora crassa* and *Aspergillus nidulans* enzymes do not cluster in the same branch as the others, in spite of being close to them in the phylogenetic tree (Figure 4C). It is proposed that the utilization of amino acids as nitrogen source is their major physiological function. This role is also assumed for LAAOs in other eukaryotic microorganisms such as the single-cell alga *Chlamydomonas reinhardtii* [64,65]. Accordingly, fungal LAAOs usually have a broad substrate range to be able to use distinct amino acids for growing. This is the case of the LAAO from *Aspergillus nidulans* and *Neurospora crassa*, which are induced under nitrogen-limiting conditions [66,67]. Similarly, LAAO synthesized by *Laccaria bicolor* and *Hebeloma* sp. are also involved in amino acid catabolism and over-expressed under limiting concentrations of nitrogen. In addition, these enzymes seem to perform an important ecological role since they catalyze nitrogen mineralization from amino acids [68].

Aspergillus fumigatus produces another LAAO of wide substrate specificity, although it shows a certain degree of substrate preference. The enzyme is more specific for hydrophobic aromatic amino acids, namely L-tyrosine and L-phenylalanine [69]. *A. fumigatus* LAAO can be applied in the resolution of some racemic DL-amino acids, yielding optically pure D-amino acids [70].

Not all LAAOs reported in fungi are of broad substrate range and play a metabolic function. The L-lysine α -oxidase from *Trichoderma viride* (LysOX) exhibits high substrate specificity for L-lysine [71]. Recently, the crystal structure of LysOX has revealed similar overall structure to that of snake venom LAAOs. However, some residues in the funnel to the active site as well as the residues involved in the substrate side-chain recognition are distinct [72]. LysOX is expected to be a potential anti-cancer agent since it inhibits growth of cancer cells but shows relatively low cytotoxicity for normal cells [72,73]. Furthermore, LysOX is stable over a wide range of pH and temperature, so it is an attractive candidate for an enzyme-based L-lysine sensor [74,75]. More recently, an extracellular LAAO with antimicrobial activity has been isolated from *Trichoderma harzianum* ETS 323. This oxidase shows the best substrate specificity constant for L-phenylalanine, although it is also active on L-Lys, L-Glu and L-Ala. It is proposed that membrane permeabilization and reactive oxidative species production are involved in the mechanism responsible for its antibacterial activity [76]. Notably, LysOX and *T. harzianum* LAAO may be related with the capability of different *Trichoderma* species as biocontrol agents.

In addition, other fungal LAAOs with substrate specificity have been described. For instance, L-lysine oxidase from induced *Saccharomyces cerevisiae* has been reported as potential biosensor for L-lysine [77]. *Coprinus* sp. SF-1 LAAO catalyzes specifically the oxidative deamination of L-tryptophan, although it can also catalyze its decarboxylation depending on conditions. This oxidase is supposed to be involved in the L-tryptophan metabolism of the fungus [78].




Figure 4. Phylogenetic relationships of representative L-amino acid oxidases in vertebrates (A), gastropods (B) and fungi (C). The tree was created by the Neighbor-Joining method integrated in the program MEGA6 [23]. Sequences were aligned using the program MUSCLE built in MEGA6. The evolutionary distances were computed using the p -distance method and are in the units of the number of amino acid differences per site. Numbers at branches indicate bootstrap values higher than 70% for both Neighbor-Joining and Maximum Likelihood trees.

2.6. Other Bacterial LAAOs

There are many bacterial enzymes showing LAAO activity that do not form a defined clustered or cannot be included in any of the previously described groups. In many cases, bacterial enzymes are more similar to proteins from eukaryotes than to other bacterial LAOs. An example would be glutamate oxidases from actinobacteria which are closely related to LAAOs from animals. This observation suggests that these enzymes have an old evolutionary origin and that they have evolved to meet different physiological functions in distinct groups. In this section, we will discuss separately the most relevant features of the enzymes selected.

The flavoprotein glycine oxidase (GO, EC 1.4.3.19) catalyzes the oxidative deamination of some primary and secondary small-sized amines and D-amino acids to yield the corresponding α -keto acid, ammonium and hydrogen peroxide [79,80]. Glycine oxidase in *Bacillus subtilis* (ThiO), encoded by the *yjbR* gene, is involved in the biosynthesis of the thiazole moiety of thiamine [5]. In addition, another glycine oxidase has been described in the marine thermophilic bacterium *Geobacillus kaustophilus* (GoxK) [81]. This protein shares a similar substrate range and structure with the glycine oxidase from *B. subtilis*. Regarding applications, glycine oxidase can be used as a biosensor for glycine in biological samples [82], and in the degradation of the herbicide glyphosate [83,84]. Flavoproteins with sarcosine oxidase activity (SOX, EC 1.5.3.1), like the one from *Bacillus* sp. B-0618 (MSOX), exhibit substrate similarity with glycine oxidases since it catalyzes the oxidative demethylation of sarcosine (N-methylglycine) and other secondary amino acids (N-methyl-L-alanine, N-ethylglycine, and L-proline) [85]. However, MSOX does not cluster in the same group with glycine oxidases (Figure 2).

The hydrogen peroxide generated by a LAAO activity was described as a factor responsible for the competition of the oral *Streptococcus oligofermentans*, involved in the initial steps of dental biofilm formation, against *Streptococcus mutans*, one of the main caries-causing pathogens. The gene responsible for this action was cloned and its activity characterized as being held on 7 different amino acids, L-aspartic acid being the one with a higher rate of hydrogen peroxide production [21]. However, latter studies revealed that this flavoprotein is an aminoacetone oxidase (Aao) with a role in the antioxidant defence of *S. oligofermentans* against cellular ROS generation [22]. Phylogenetic analyses of several *Streptococcus* strains revealed that the gene in *S. oligofermentans* was acquired by horizontal gene transfer from a source closely related to *Streptococcus sanguinis* as a mechanism of adaptation to the aerobic conditions in the oral cavity [86]. In fact, BLAST analysis of *S. oligofermentans* *aao* gene product reveals that the most similar proteins are annotated as hypothetical or as pyridine nucleotide-disulfide oxidoreductases, such as in the case of *S. sanguinis*. In agreement with that, structure-function relationships of SoAAO set apart this protein from the oxidase/dehydrogenase class of flavoproteins [87]. These enzymes have been investigated for their role in diverse processes such as epoxyalkanes degradation in *Xanthobacter* [88], as part of a redox system based on coenzyme A in *Staphylococcus aureus* [89] or as catalyzing the disulfide bond formation in *Chromobacterium violaceum* FK228 anticancer depsipeptide [90].

Rhodococcus opacus synthesizes a LAAO with broad substrate range that makes it interesting in some biotechnological applications such as the solving of racemic mixtures of amino acids [91]. The structure of the enzyme has been determined, what has contributed to a better understanding of its molecular mechanism of action [92]. Other LAAOs with broad substrate range have been cloned from different *Rhodococcus* strains [93]. As far as we know, no physiological function has been proposed for these enzymes, since they have been characterized in terms of their possible biotechnological applications. Other enzymes with broad substrate range were described in *Bacillus carotarum* [94], *Cellulomonas* [95], *Morganella morganii* [96] and *Corynebacterium* [97]. An example of LAAO with high substrate specificity is the glutamate oxidase synthesized by *Streptomyces* [98,99]. The physiological function of this extracellular enzyme is unknown. From the biotechnological point of view it is used as part of biosensors for the determination of glutamate [100].

Several LAAOs have been described in bacteria of the genus *Pseudomonas*. The strain *Pseudomonas* sp. AIU 813 synthesizes an enzyme with oxidase activity on L-Lys, ornithine and L-Arg [101]. This flavoprotein was lately reclassified as an L-amino acid/monooxygenase since the second activity was higher than the first [20]. Interestingly, the capacity to act as an oxidase or as a decarboxilating monooxygenase is shared by other enzymes such as the PAO (L-phenylalanine oxidase) from *Pseudomonas* P501 which is able to show different activities depending on the substrate [102,103]. *Pseudomonas savastanoi* TMO (tryptophan monooxygenase) does not show oxidase activity, but it can be revealed by mutagenesis [104].

VioA from *Chromobacterium violaceum* encodes an enzyme with L-tryptophan 2',3'-oxidase activity that is involved in the synthesis of violacein [105]. An enzyme with the same enzymatic activity also participates in the synthesis of other secondary metabolites such as rebeccamycin and staurosporine produced by actinomycetes [106,107]. Interestingly the enzymes from the actinobacteria are not phylogenetically close to the one synthesized by *Chromobacterium* (Figure 2).

LAAOs with preference for basic amino acids have been described in cyanobacteria [108]. In *Synechococcus* PCC 7942 that enzyme has been related to the use of Arg as nitrogen source [109].

3. Amino Acid Oxidases with Antimicrobial Activity in Marine Bacteria

The colonization of submerged surfaces is of great importance in marine environments. On the one hand, it can generate the biofouling of surfaces. On the other hand, bacteria associated with surfaces are important factors inducing the settlement and metamorphosis of larvae of higher organisms [110]. The genus *Pseudoalteromonas* has been recognized as an important component of the surface of many different marine surfaces [111,112]. *Marinomonas mediterranea* is a bacterium that forms part of the microbiota of the seagrass *Posidonia oceanica* [113]. From both, *Pseudoalteromonas* and *Marinomonas*, proteins with antibacterial activity were reported [114,115]. As described above, LodA is an amino acid oxidase with quinone cofactor. The autolytic protein of the marine bacterium *Pseudoalteromonas tunicata* (AlpP) also possesses lysine oxidase activity and high sequence similarity with LodA [13]. LodA and AlpP are involved in biofilm formation and mediate differentiation, dispersal, and phenotypic variation among the dispersed cells [13]. In two non-marine bacteria, *Caulobacter crescentus* and *Chromobacterium violaceum*, LodA-like proteins also generate hydrogen peroxide related with biofilm differentiation in those microorganisms, but the substrate of the activity was not reported [13].

Interestingly, other LodA-like proteins in the genus *Pseudoalteromonas* have been described as having antimicrobial activity. Two different strains of *Pseudoalteromonas flavigipulchra* synthesize antimicrobial proteins of the Lod-A like group. *P. flavigipulchra* JG1 produces an extracellular antibacterial protein (PfaP) with high similarity to LodA and AlpP [116]. The AAO of the strain C2 was reported to have a broad substrate range oxidizing L-Lys, L-Met, L-Glu, L-Leu, L-Gln, L-Tyr and L-Phe (Table S1.). The hydrogen peroxide generated by its catalysis mediates its antibacterial activity [117]. Another antibacterial LAAO with similar broad substrate range (L-Met, L-Gln, L-Leu, L-Phe, L-Glu, L-Trp, etc.) has been described in several *Pseudoalteromonas luteoviolacea* strains but the genes encoding those enzymes have not been reported yet [118]. As far as we know, the only case of LAAO activity not related to a quinoprotein in the genus *Pseudoalteromonas* has been described in the strain B3, in which it has been associated to a flavoprotein with similarity to LASPOs [119,120] (Figure 3C). Genome mining of the genomes of *Pseudoalteromonas* strains, and other marine bacteria (data not shown) has revealed that they contain homologues to the gene encoding LASPO (*nadB*) as well as homologues to *nadA*. It remains to be determined whether these LASPOs participate in NAD synthesis or have evolved to meet an antimicrobial role. In the genomes of marine bacteria it is also possible to detect other genes encoding flavoproteins, some of them with similarity to amino oxidases. However, to the best of our knowledge, it has not been determined the actual enzymatic activity of the product of those genes or their possible antimicrobial activity.

The data available indicate that in the genomes of marine bacteria of the genus *Pseudoalteromonas* and *Marinomonas* it is common the presence of genes encoding proteins similar to LodA [12]. Although LodA-like proteins have been mainly described in marine bacteria, genes encoding proteins similar to them are present in approximately 1% of the microbial genomes sequenced, including microorganisms present in different environments. In the case of the freshwater *Rheinheimera aquatica* GR5, a protein with a peptide fragment with high similarity to LodA and AlpP was identified. This is an antimicrobial protein due to the hydrogen peroxide generation in the specific oxidation of L-Lys [121]. Nevertheless, it is important to bear in mind that in some microorganisms it is possible to detect several *lodA*-like genes [12]. Thus far, it has been shown that two of those genes in *Marinomonas* code for proteins with, respectively, L-lysine ε-oxidase and glycine oxidase activity. This observation suggests that they could have different, perhaps complementary physiological functions. LodA-like proteins may also constitute a reservoir of novel enzymatic activities of potential biotechnological interest.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Izidoro, L.F.; Sobrinho, J.C.; Mendes, M.M.; Costa, T.R.; Grabner, A.N.; Rodrigues, V.M.; da Silva, S.L.; Zanchi, F.B.; Zuliani, J.P.; Fernandes, C.F.; et al. Snake venom L-amino acid oxidases: Trends in pharmacology and biochemistry. *Biomed. Res. Int.* **2014**, *2014*, 196754. [[CrossRef](#)] [[PubMed](#)]
- Yu, Z.; Qiao, H. Advances in non-snake venom L-amino acid oxidase. *Appl. Biochem. Biotechnol.* **2012**, *167*, 1–13. [[CrossRef](#)] [[PubMed](#)]
- Pollegioni, L.; Molla, G.; Sacchi, S.; Rosini, E.; Verga, R.; Pilone, M. Properties and applications of microbial D-amino acid oxidases: Current state and perspectives. *Appl. Microbiol. Biotechnol.* **2008**, *78*, 1–16. [[CrossRef](#)] [[PubMed](#)]
- Pollegioni, L.; Molla, G. New biotech applications from evolved D-amino acid oxidases. *Trends Biotechnol.* **2011**, *29*, 276–283. [[CrossRef](#)] [[PubMed](#)]
- Settembre, E.C.; Dorrestein, P.C.; Park, J.H.; Augustine, A.M.; Begley, T.P.; Ealick, S.E. Structural and mechanistic studies on ThiO, a glycine oxidase essential for thiamin biosynthesis in *Bacillus subtilis*. *Biochemistry* **2003**, *42*, 2971–2981. [[CrossRef](#)] [[PubMed](#)]
- Gomez, D.; Lucas-Elio, P.; Sanchez-Amat, A.; Solano, F. A novel type of lysine oxidase: L-Lysine-epsilon-oxidase. *Biochim. Biophys. Acta* **2006**, *1764*, 1577–1585. [[CrossRef](#)] [[PubMed](#)]
- Lucas-Elio, P.; Gomez, D.; Solano, F.; Sanchez-Amat, A. The antimicrobial activity of marinocine, synthesized by *Marinomonas mediterranea*, is due to hydrogen peroxide generated by its lysine oxidase activity. *J. Bacteriol.* **2006**, *188*, 2493–2501. [[CrossRef](#)] [[PubMed](#)]
- Chacon-Verdu, M.; Gomez, D.; Solano, F.; Lucas-Elio, P.; Sanchez-Amat, A. LodB is required for the recombinant synthesis of the quinoprotein L-lysine-ε-oxidase from *Marinomonas mediterranea*. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 2981–2989. [[CrossRef](#)] [[PubMed](#)]
- Okazaki, S.; Nakano, S.; Matsui, D.; Akaji, S.; Inagaki, K.; Asano, Y. X-ray crystallographic evidence for the presence of the cysteine tryptophylquinone cofactor in L-lysine epsilon-oxidase from *Marinomonas mediterranea*. *J. Biochem.* **2013**, *154*, 233–236. [[CrossRef](#)] [[PubMed](#)]
- Campillo-Brocal, J.C.; Lucas-Elio, P.; Sanchez-Amat, A. Identification in *Marinomonas mediterranea* of a novel quinoprotein with glycine oxidase activity. *Microbiologyopen* **2013**, *2*, 684–694. [[CrossRef](#)] [[PubMed](#)]
- Chacon-Verdu, M.; Campillo-Brocal, J.C.; Lucas-Elio, P.; Davidson, V.L.; Sanchez-Amat, A. Characterization of recombinant biosynthetic precursors of the cysteine tryptophylquinone cofactors of L-lysine-epsilon-oxidase and glycine oxidase from *Marinomonas mediterranea*. *Biochim. Biophys. Acta* **2015**, *1854*, 1123–1131. [[CrossRef](#)] [[PubMed](#)]
- Campillo-Brocal, J.C.; Chacon-Verdu, M.D.; Lucas-Elío, P.; Sanchez-Amat, A. Distribution in microbial genomes of genes similar to *lodA* and *goxA* which encode a novel family of quinoproteins with amino acid oxidase activity. *BMC Genomics* **2015**, *16*, 231. [[CrossRef](#)] [[PubMed](#)]

13. Mai-Prochnow, A.; Lucas-Elio, P.; Egan, S.; Thomas, T.; Webb, J.S.; Sanchez-Amat, A.; Kjelleberg, S. Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several Gram-negative bacteria. *J. Bacteriol.* **2008**, *190*, 5493–5501. [[CrossRef](#)] [[PubMed](#)]
14. Yang, C.A.; Cheng, C.H.; Lo, C.T.; Liu, S.Y.; Lee, J.W.; Peng, K.C. A novel L-amino acid oxidase from *Trichoderma harzianum* ETS 323 associated with antagonism of *Rhizoctonia solani*. *J. Agric. Food Chem.* **2011**, *59*, 4519–4526. [[CrossRef](#)] [[PubMed](#)]
15. Kitani, Y.; Kikuchi, N.; Zhang, G.; Ishizaki, S.; Shimakura, K.; Shiomi, K.; Nagashima, Y. Antibacterial action of L-amino acid oxidase from the skin mucus of rockfish *Sebastodes schlegelii*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2008**, *149*, 394–400. [[CrossRef](#)] [[PubMed](#)]
16. Puiffe, M.L.; Lachaise, I.; Molinier-Frenkel, V.R.; Castellano, F. Antibacterial properties of the mammalian L-amino acid oxidase IL4I1. *PLoS ONE* **2013**, *8*. [[CrossRef](#)] [[PubMed](#)]
17. Pollegioni, L.; Motta, P.; Molla, G. L-Amino acid oxidase as biocatalyst: A dream too far? *Appl. Microbiol. Biotechnol.* **2013**, *97*, 9323–9341. [[CrossRef](#)] [[PubMed](#)]
18. Kasai, K.; Ishikawa, T.; Nakamura, T.; Miura, T. Antibacterial properties of L-amino acid oxidase: Mechanisms of action and perspectives for therapeutic applications. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 1–11. [[CrossRef](#)] [[PubMed](#)]
19. Rosini, E.; Pollegioni, L.; Ghisla, S.; Orru, R.; Molla, G. Optimization of D-amino acid oxidase for low substrate concentrations—Towards a cancer enzyme therapy. *FEBS J.* **2009**, *276*, 4921–4932. [[CrossRef](#)] [[PubMed](#)]
20. Matsui, D.; Im, D.H.; Sugawara, A.; Fukuta, Y.; Fushinobu, S.; Isobe, K.; Asano, Y. Mutational and crystallographic analysis of L-amino acid oxidase/monooxygenase from *Pseudomonas* sp. AIU 813: Interconversion between oxidase and monooxygenase activities. *FEBS Open Bio.* **2014**, *4*, 220–228. [[CrossRef](#)] [[PubMed](#)]
21. Tong, H.; Chen, W.; Shi, W.; Qi, F.; Dong, X. SO-LAAO, a novel L-amino acid oxidase that enables *Streptococcus oligofermentans* to outcompete *Streptococcus mutans* by generating H₂O₂ from peptone. *J. Bacteriol.* **2008**, *190*, 4716–4721. [[CrossRef](#)] [[PubMed](#)]
22. Zhou, P.; Liu, L.; Tong, H.; Dong, X. Role of operon *aaoSo-mutT* in antioxidant defense in *Streptococcus oligofermentans*. *PLoS ONE* **2012**, *7*. [[CrossRef](#)] [[PubMed](#)]
23. Tamura, K.; Stecher, G.; Peterson, D.; Filipski, A.; Kumar, S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* **2013**, *30*, 2725–2729. [[CrossRef](#)] [[PubMed](#)]
24. Sehanobish, E.; Chacon-Verdu, M.D.; Sanchez-Amat, A.; Davidson, V.L. Roles of active site residues in LodA, a cysteine tryptophylquinone dependent epsilon-lysine oxidase. *Arch. Biochem. Biophys.* **2015**, *579*, 26–32. [[CrossRef](#)] [[PubMed](#)]
25. Takahashi, S.; Abe, K.; Kera, Y. Bacterial D-amino acid oxidases: Recent findings and future perspectives. *Bioengineered* **2015**, *6*, 237–241. [[CrossRef](#)] [[PubMed](#)]
26. D'Aniello, A.; Vetere, A.; Petruccelli, L. Further study on the specificity of D-amino acid oxidase and of D-aspartate oxidase and time course for complete oxidation of D-amino acids. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **1993**, *105*, 731–734. [[CrossRef](#)]
27. Takahashi, S.; Furukawara, M.; Omae, K.; Tadokoro, N.; Saito, Y.; Abe, K.; Kera, Y. A highly stable D-amino acid oxidase of the thermophilic bacterium *Rubrobacter xylanophilus*. *Appl. Environ. Microbiol.* **2014**, *80*, 7219–7229. [[CrossRef](#)] [[PubMed](#)]
28. Caligiuri, A.; DArrigo, P.; Rosini, E.; Pedrocchi-Fantoni, G.; Tessaro, D.; Molla, G.; Servi, S.; Pollegioni, L. Activity of yeast D-amino acid oxidase on aromatic unnatural amino acids. *J. Mol. Catal. B Enzym.* **2008**, *50*, 93–98. [[CrossRef](#)]
29. Gholizadeh, A.; Kohnehrouz, B.B. Molecular cloning and expression in *Escherichia coli* of an active fused *Zea mays* L. D-amino acid oxidase. *Biochemistry (Mosc.)* **2009**, *74*, 137–144. [[CrossRef](#)] [[PubMed](#)]
30. Katane, M.; Saitoh, Y.; Seida, Y.; Sekine, M.; Furuchi, T.; Homma, H. Comparative characterization of three D-aspartate oxidases and one D-amino acid oxidase from *Caenorhabditis elegans*. *Chem. Biodivers.* **2010**, *7*, 1424–1434. [[CrossRef](#)] [[PubMed](#)]
31. Sarower, G.M.; Okada, S.; Abe, H. Molecular characterization of D-amino acid oxidase from common carp *Cyprinus carpio* and its induction with exogenous free D-alanine. *Arch. Biochem. Biophys.* **2003**, *420*, 121–129. [[CrossRef](#)] [[PubMed](#)]

32. Molla, G.; Sacchi, S.; Bernasconi, M.; Pilone, M.S.; Fukui, K.; Pollegioni, L. Characterization of human D-amino acid oxidase. *FEBS Lett.* **2006**, *580*, 2358–2364. [[CrossRef](#)] [[PubMed](#)]
33. Pollegioni, L.; Piubelli, L.; Sacchi, S.; Pilone, M.S.; Molla, G. Physiological functions of D-amino acid oxidases: From yeast to humans. *Cell. Mol. Life Sci.* **2007**, *64*, 1373–1394. [[CrossRef](#)] [[PubMed](#)]
34. Khoronenkova, S.V.; Tishkov, V.I. D-Amino acid oxidase: Physiological role and applications. *Biochemistry (Mosc)* **2008**, *73*, 1511–1518. [[CrossRef](#)] [[PubMed](#)]
35. Nasu, S.; Wicks, F.D.; Ghelton, R.K. L-Aspartate oxidase, a newly discovered enzyme of *Escherichia coli*, is the B protein of quinolinate synthetase. *J. Biol. Chem.* **1982**, *257*, 626–632. [[PubMed](#)]
36. Korshunov, S.; Imlay, J.A. Two sources of endogenous hydrogen peroxide in *Escherichia coli*. *Mol. Microbiol.* **2010**, *75*, 1389–1401. [[CrossRef](#)] [[PubMed](#)]
37. Yang, Z.; Savchenko, A.; Yakunin, A.; Zhang, R.; Edwards, A.; Arrowsmith, C.; Tong, L. Aspartate dehydrogenase, a novel enzyme identified from structural and functional studies of TM1643. *J. Biol. Chem.* **2003**, *278*, 8804–8808. [[CrossRef](#)] [[PubMed](#)]
38. Bossi, R.T.; Negri, A.; Tedeschi, G.; Mattevi, A. Structure of FAD-bound L-aspartate oxidase: Insight into substrate specificity and catalysis. *Biochemistry* **2002**, *41*, 3018–3024. [[CrossRef](#)] [[PubMed](#)]
39. Mattevi, A.; Tedeschi, G.; Bacchella, L.; Coda, A.; Negri, A.; Ronchi, S. Structure of L-aspartate oxidase: Implications for the succinate dehydrogenase/fumarate reductase oxidoreductase family. *Structure* **1999**, *7*, 745–756. [[CrossRef](#)]
40. Tedeschi, G.; Nonnis, S.; Strumbo, B.; Cruciani, G.; Carosati, E.; Negri, A. On the catalytic role of the active site residue E121 of *E. coli* L-aspartate oxidase. *Biochimie* **2010**, *92*, 1335–1342.
41. Sakuraba, H.; Satomura, T.; Kawakami, R.; Yamamoto, S.; Kawarabayashi, Y.; Kikuchi, H.; Ohshima, T. L-Aspartate oxidase is present in the anaerobic hyperthermophilic archaeon *Pyrococcus horikoshii* OT-3: Characteristics and role in the de novo biosynthesis of nicotinamide adenine dinucleotide proposed by genome sequencing. *Extremophiles* **2002**, *6*, 275–281. [[CrossRef](#)] [[PubMed](#)]
42. Marinoni, I.; Nonnis, S.; Monteferrante, C.; Heathcote, P.; Hartig, E.; Bottger, L.H.; Trautwein, A.X.; Negri, A.; Albertini, A.M.; Tedeschi, G. Characterization of L-aspartate oxidase and quinolinate synthase from *Bacillus subtilis*. *FEBS J.* **2008**, *275*, 5090–5107. [[CrossRef](#)] [[PubMed](#)]
43. Bifulco, D.; Pollegioni, L.; Tessaro, D.; Servi, S.; Molla, G. A thermostable L-aspartate oxidase: A new tool for biotechnological applications. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 7285–7295. [[CrossRef](#)] [[PubMed](#)]
44. Mutaguchi, Y.; Ohmori, T.; Sakuraba, H.; Yoneda, K.; Doi, K.; Ohshima, T. Visible wavelength spectrophotometric assays of L-aspartate and D-aspartate using hyperthermophilic enzyme systems. *Anal. Biochem.* **2011**, *409*, 1–6. [[CrossRef](#)] [[PubMed](#)]
45. Lin, H.; Kwan, A.L.; Dutcher, S.K. Synthesizing and salvaging NAD: Lessons learned from *Chlamydomonas reinhardtii*. *PLoS Genet.* **2010**, *6*. [[CrossRef](#)] [[PubMed](#)]
46. Di Martino, M.L.; Fioravanti, R.; Barbabella, G.; Prosseda, G.; Colonna, B.; Casalino, M. Molecular evolution of the nicotinic acid requirement within the *Shigella*/EIEC pathotype. *Int. J. Med. Microbiol.* **2013**, *303*, 651–661. [[CrossRef](#)] [[PubMed](#)]
47. Hughes, A.L. Origin and diversification of the L-amino oxidase family in innate immune defenses of animals. *Immunogenetics* **2010**, *62*, 753–759. [[CrossRef](#)] [[PubMed](#)]
48. Du, X.Y.; Clemetson, K.J. Snake venom L-amino acid oxidases. *Toxicon* **2002**, *40*, 659–665. [[CrossRef](#)]
49. Calderon, L.A.; Sobrinho, J.C.; Zaqueo, K.D.; de Moura, A.A.; Grabner, A.N.; Mazzi, M.C.V.; Marcussi, S.; Nomizo, A.; Fernandes, C.F.C.; Zuliani, J.P.; et al. Antitumoral activity of snake venom proteins: New trends in cancer therapy. *Biomed. Res. Int.* **2014**, *2014*, 203639. [[CrossRef](#)] [[PubMed](#)]
50. Nagashima, Y.; Tsukamoto, C.; Kitani, Y.; Ishizaki, S.; Nagai, H.; Yanagimoto, T. Isolation and cDNA cloning of an antibacterial L-amino acid oxidase from the skin mucus of the great sculpin *Myoxocephalus polyacanthocephalus*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2009**, *154*, 55–61. [[CrossRef](#)] [[PubMed](#)]
51. Endo, H.; Hayashi, Y.; Kitani, Y.; Ren, H.; Hayashi, T.; Nagashima, Y. Optical enzyme sensor for determining L-lysine content using L-lysine oxidase from the rockfish *Sebastodes schlegeli*. *Anal. Bioanal. Chem.* **2008**, *391*, 1255–1261. [[CrossRef](#)] [[PubMed](#)]
52. Kitani, Y.; Ishida, M.; Ishizaki, S.; Nagashima, Y. Discovery of serum L-amino acid oxidase in the rockfish *Sebastodes schlegeli*: Isolation and biochemical characterization. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2010**, *157*, 351–356. [[CrossRef](#)] [[PubMed](#)]

53. Wang, F.; Li, R.; Xie, M.; Li, A. The serum of rabbitfish (*Siganus oramin*) has antimicrobial activity to some pathogenic organisms and a novel serum L-amino acid oxidase is isolated. *Fish. Shellfish Immunol.* **2011**, *30*, 1095–1108. [[CrossRef](#)] [[PubMed](#)]
54. Kitani, Y.; Fernandes, J.M.O.; Kiron, V. Identification of the atlantic cod L-amino acid oxidase and its alterations following bacterial exposure. *Dev. Comp. Immunol.* **2015**, *50*, 116–120. [[CrossRef](#)] [[PubMed](#)]
55. Boulland, M.L.; Marquet, J.; Molinier-Frenkel, V.R.; Moller, P.; Guiter, C.; Lasoudris, F.; Copie-Bergman, C.; Baia, M.; Gaulard, P.; Leroy, K.; et al. Human IL4I1 is a secreted L-phenylalanine oxidase expressed by mature dendritic cells that inhibits T-lymphocyte proliferation. *Blood* **2007**, *110*, 220–227. [[CrossRef](#)] [[PubMed](#)]
56. Sun, Y.; Nonobe, E.; Kobayashi, Y.; Kuraishi, T.; Aoki, F.; Yamamoto, K.; Sakai, S. Characterization and expression of L-amino acid oxidase of mouse milk. *J. Biol. Chem.* **2002**, *277*, 19080–19086. [[CrossRef](#)] [[PubMed](#)]
57. Johnson, P.M.; Kicklighter, C.E.; Schmidt, M.; Kamio, M.; Yang, H.; Elkin, D.; Michel, W.C.; Tai, P.C.; Derby, C.D. Packaging of chemicals in the defensive secretory glands of the sea hare *Aplysia californica*. *J. Exp. Biol.* **2006**, *209*, 78–88. [[CrossRef](#)] [[PubMed](#)]
58. Yang, H.; Johnson, P.M.; Ko, K.C.; Kamio, M.; Germann, M.W.; Derby, C.D.; Tai, P.C. Cloning, characterization and expression of escapin, a broadly antimicrobial FAD-containing L-amino acid oxidase from ink of the sea hare *Aplysia californica*. *J. Exp. Biol.* **2005**, *208*, 3609–3622. [[CrossRef](#)] [[PubMed](#)]
59. Ko, K.C.; Wang, B.; Tai, P.C.; Derby, C.D. Identification of potent bactericidal compounds produced by escapin, an L-amino acid oxidase in the ink of the sea hare *Aplysia californica*. *Antimicrob. Agents Chemother.* **2008**, *52*, 4455–4462. [[CrossRef](#)] [[PubMed](#)]
60. Ko, K.C.; Tai, P.C.; Derby, C.D. Mechanisms of action of escapin, a bactericidal agent in the ink secretion of the sea hare *Aplysia californica*: Rapid and long-lasting DNA condensation and involvement of the OxyR-regulated oxidative stress pathway. *Antimicrob. Agents Chemother.* **2012**, *56*, 1725–1734. [[CrossRef](#)] [[PubMed](#)]
61. Butzke, D.; Hurwitz, R.; Thiede, B.; Goedert, S.; Rudel, T. Cloning and biochemical characterization of APIT, a new L-amino acid oxidase from *Aplysia punctata*. *Toxicon* **2005**, *46*, 479–489. [[CrossRef](#)] [[PubMed](#)]
62. Ehara, T.; Kitajima, S.; Kanzawa, N.; Tamiya, T.; Tsuchiya, T. Antimicrobial action of achacin is mediated by L-amino acid oxidase activity. *FEBS Lett.* **2002**, *531*, 509–512. [[CrossRef](#)]
63. Kanzawa, N.; Shintani, S.; Ohta, K.; Kitajima, S.; Ehara, T.; Kobayashi, H.; Kizaki, H.; Tsuchiya, T. Achacin induces cell death in HeLa cells through two different mechanisms. *Arch. Biochem. Biophys.* **2004**, *422*, 103–109. [[CrossRef](#)] [[PubMed](#)]
64. Perez-Alegre, M.; Franco, A.R. Resistance to L-methionine-S-sulfoximine in *Chlamydomonas reinhardtii* is due to an alteration in a general amino acid transport system. *Planta* **1998**, *207*, 20–26. [[CrossRef](#)]
65. Vallon, O.; Bulte, L.; Kuras, R.; Olive, J.; Wollman, F.A. Extensive accumulation of an extracellular L-amino-acid oxidase during gametogenesis of *Chlamydomonas reinhardtii*. *Eur. J. Biochem.* **1993**, *215*, 351–360. [[CrossRef](#)] [[PubMed](#)]
66. Davis, M.A.; Askin, M.C.; Hynes, M.J. Amino acid catabolism by an *areA*-regulated gene encoding an L-amino acid oxidase with broad substrate specificity in *Aspergillus nidulans*. *Appl. Environ. Microbiol.* **2005**, *71*, 3551–3555. [[CrossRef](#)] [[PubMed](#)]
67. Sikora, L.; Marzluf, G.A. Regulation of L-amino acid oxidase and of D-amino acid oxidase in *Neurospora crassa*. *Mol. Gen. Genet.* **1982**, *186*, 33–39. [[CrossRef](#)] [[PubMed](#)]
68. Nuutinen, J.T.; Timonen, S. Identification of nitrogen mineralization enzymes, L-amino acid oxidases, from the ectomycorrhizal fungi *Hebeloma* spp. and *Laccaria bicolor*. *Mycol. Res.* **2008**, *112*, 1453–1464. [[CrossRef](#)] [[PubMed](#)]
69. Singh, S.; Gogoi, B.K.; Bezbaruah, R.L. Optimization of medium and cultivation conditions for L-amino acid oxidase production by *Aspergillus fumigatus*. *Can. J. Microbiol.* **2009**, *55*, 1096–1102. [[CrossRef](#)] [[PubMed](#)]
70. Singh, S.; Gogoi, B.; Bezbaruah, R. Racemic resolution of some DL-amino acids using *Aspergillus fumigatus* L-amino acid oxidase. *Curr. Microbiol.* **2011**, *63*, 94–99. [[CrossRef](#)] [[PubMed](#)]
71. Kusakabe, H.; Kodama, K.; Kuninaka, A.; Yoshino, H.; Misono, H.; Soda, K. A new antitumor enzyme, L-lysine alpha-oxidase from *Trichoderma viride*. Purification and enzymological properties. *J. Biol. Chem.* **1980**, *255*, 976–981. [[PubMed](#)]

72. Amano, M.; Mizuguchi, H.; Sano, T.; Kondo, H.; Shinyashiki, K.; Inagaki, J.; Tamura, T.; Kawaguchi, T.; Kusakabe, H.; Imada, K.; *et al.* Recombinant expression, molecular characterization and crystal structure of antitumor enzyme, L-lysine- α -oxidase from *Trichoderma viride*. *J. Biochem.* **2015**, *157*, 549–559. [[CrossRef](#)] [[PubMed](#)]
73. Lukasheva, E.V.; Berezov, T.T. L-Lysine alpha-oxidase: Physicochemical and biological properties. *Biochemistry (Mosc)* **2002**, *67*, 1152–1158. [[CrossRef](#)] [[PubMed](#)]
74. Chauhan, N.; Narang, J.; Sunny; Pundir, C.S. Immobilization of lysine oxidase on a gold-platinum nanoparticles modified Au electrode for detection of lysine. *Enzyme Microb. Technol.* **2013**, *52*, 265–271. [[CrossRef](#)] [[PubMed](#)]
75. Pohlmann, A.; Stamm, W.W.; Kusakabe, H.; Kula, M.R. Enzymatic determination of L-lysine by flow-injection techniques. *Anal. Chim. Acta* **1990**, *235*, 329–335. [[CrossRef](#)]
76. Yang, C.A.; Cheng, C.H.; Liu, S.Y.; Lo, C.T.; Lee, J.W.; Peng, K.C. Identification of antibacterial mechanism of L-amino acid oxidase derived from *Trichoderma harzianum* ETS 323. *FEBS J.* **2011**, *278*, 3381–3394. [[CrossRef](#)] [[PubMed](#)]
77. Akyilmaz, E.; Erdogan, A.; Ozturk, R.; Yasa, I. Sensitive determination of L-lysine with a new amperometric microbial biosensor based on *Saccharomyces cerevisiae* yeast cells. *Biosens. Bioelectron.* **2007**, *22*, 1055–1060. [[CrossRef](#)] [[PubMed](#)]
78. Furuya, Y.; Sawada, H.; Hirahara, T.; Ito, K.; Ohshiro, T.; Izumi, Y. A novel enzyme, L-tryptophan oxidase, from a basidiomycete, *Coprinus* sp. SF-1: Purification and characterization. *Biosci. Biotechnol. Biochem.* **2000**, *64*, 1486–1493. [[CrossRef](#)] [[PubMed](#)]
79. Nishiya, Y.; Imanaka, T. Purification and characterization of a novel glycine oxidase from *Bacillus subtilis*. *FEBS Lett.* **1998**, *438*, 263–266. [[CrossRef](#)]
80. Job, V.; Marcone, G.L.; Pilone, M.S.; Pollegioni, L. Glycine oxidase from *Bacillus subtilis*. Characterization of a new flavoprotein. *J. Biol. Chem.* **2002**, *277*, 6985–6993. [[CrossRef](#)] [[PubMed](#)]
81. Martinez-Martinez, I.; Navarro-Fernandez, J.; Garcia-Carmona, F.; Takami, H.; Sanchez-Ferrer, A. Characterization and structural modeling of a novel thermostable glycine oxidase from *Geobacillus kaustophilus* HTA426. *Proteins* **2008**, *70*, 1429–1441. [[CrossRef](#)] [[PubMed](#)]
82. Rosini, E.; Piubelli, L.; Molla, G.; Frattini, L.; Valentino, M.; Varriale, A.; D'Auria, S.; Pollegioni, L. Novel biosensors based on optimized glycine oxidase. *FEBS J.* **2014**, *281*, 3460–3472. [[CrossRef](#)] [[PubMed](#)]
83. Nicolia, A.; Ferradini, N.; Molla, G.; Biagetti, E.; Pollegioni, L.; Veronesi, F.; Rosellini, D. Expression of an evolved engineered variant of a bacterial glycine oxidase leads to glyphosate resistance in alfalfa. *J. Biotechnol.* **2014**, *184*, 201–208. [[CrossRef](#)] [[PubMed](#)]
84. Yao, P.; Lin, Y.; Wu, G.; Lu, Y.; Zhan, T.; Kumar, A.; Zhang, L.; Liu, Z. Improvement of glycine oxidase by DNA shuffling, and site-saturation mutagenesis of F247 residue. *Int. J. Biol. Macromol.* **2015**, *79*, 965–970. [[CrossRef](#)] [[PubMed](#)]
85. Trickey, P.; Wagner, M.A.; Jorns, M.S.; Mathews, F.S. Monomeric sarcosine oxidase: Structure of a covalently flavyliylated amine oxidizing enzyme. *Structure* **1999**, *7*, 331–345. [[CrossRef](#)]
86. Boggs, J.M.; South, A.H.; Hughes, A.L. Phylogenetic analysis supports horizontal gene transfer of L-amino acid oxidase gene in *Streptococcus oligofermentans*. *Infect. Genet. Evol.* **2012**, *12*, 1005–1009. [[CrossRef](#)] [[PubMed](#)]
87. Molla, G.; Nardini, M.; Motta, P.; D'Arrigo, P.; Panzeri, W.; Pollegioni, L. Aminoacetone oxidase from *Streptococcus oligofermentans* belongs to a new three-domain family of bacterial flavoproteins. *Biochem. J.* **2014**, *464*, 387–399. [[CrossRef](#)] [[PubMed](#)]
88. Swaving, J.; de Bont, J.A.; Westphal, A.; de, K.A. A novel type of pyridine nucleotide-disulfide oxidoreductase is essential for NAD⁺- and NADPH-dependent degradation of epoxyalkanes by *Xanthobacter* strain Py2. *J. Bacteriol.* **1996**, *178*, 6644–6646. [[PubMed](#)]
89. delCardayre, S.B.; Davies, J.E. *Staphylococcus aureus* coenzyme A disulfide reductase, a new subfamily of pyridine nucleotide-disulfide oxidoreductase. Sequence, expression, and analysis of cdr. *J. Biol. Chem.* **1998**, *273*, 5752–5757. [[CrossRef](#)] [[PubMed](#)]
90. Wang, C.; Wesener, S.R.; Zhang, H.; Cheng, Y.Q. An FAD-dependent pyridine nucleotide-disulfide oxidoreductase is involved in disulfide bond formation in FK228 anticancer depsipeptide. *Chem. Biol.* **2009**, *16*, 585–593. [[CrossRef](#)] [[PubMed](#)]

91. Geueke, B.; Hummel, W.; Bommarius, A. L-Amino Acid Oxidase from *Rhodococcus* Species. U.S. Patent 6461841 B2, 8 October 2002.
92. Faust, A.; Niefind, K.; Hummel, W.; Schomburg, D. The structure of a bacterial L-amino acid oxidase from *Rhodococcus opacus* gives new evidence for the hydride mechanism for dehydrogenation. *J. Mol. Biol.* **2007**, *367*, 234–248. [CrossRef] [PubMed]
93. Isobe, K.; Satou, S.; Matsumoto, E.; Yoshida, S.; Yamada, M.; Hibi, M.; Ogawa, J. Characterization and application of a L-specific amino acid oxidase from *Rhodococcus* sp. AIU LAB-3. *J. Biosci. Bioeng.* **2013**, *115*, 613–617. [CrossRef] [PubMed]
94. Bearley, G.M.; Price, C.P.; Atkinson, T.; Hammond, P.M. Purification and partial characterisation of a broad-range L-amino acid oxidase from *Bacillus carotarum* 2Pfa isolated from soil. *Appl. Microbiol. Biotechnol.* **1994**, *41*, 670–676. [CrossRef]
95. Braun, M.; Kim, J.M.; Schmid, R.D. Purification and some properties of an extracellular L-amino acid oxidase from *Cellulomonas cellulans* AM8 isolated from soil. *Appl. Microbiol. Biotechnol.* **1992**, *37*, 594–598. [CrossRef]
96. Bouvrette, P.; Luong, J.H. T. Purification and further characterization of an L-phenylalanine oxidase from *Morganella morganii*. *Appl. Biochem. Biotechnol.* **1994**, *48*, 61–74. [CrossRef]
97. Coudert, M. Charcterization and physiological function of a soluble L-amino acid oxidase in *Corynebacterium*. *Arch. Microbiol.* **1975**, *102*, 151–153. [CrossRef] [PubMed]
98. Arima, J.; Sasaki, C.; Sakaguchi, C.; Mizuno, H.; Tamura, T.; Kashima, A.; Kusakabe, H.; Sugio, S.; Inagaki, K. Structural characterization of L-glutamate oxidase from *Streptomyces* sp. X-119-6. *FEBS J.* **2009**, *276*, 3894–3903. [CrossRef] [PubMed]
99. Arima, J.; Tamura, T.; Kusakabe, H.; Ashiuchi, M.; Yagi, T.; Tanaka, H.; Inagaki, K. Recombinant expression, biochemical characterization and stabilization through proteolysis of an L-glutamate oxidase from *Streptomyces* sp. X-119-6. *J. Biochem.* **2003**, *134*, 805–812. [CrossRef] [PubMed]
100. Batra, B.; Pandir, C.S. An amperometric glutamate biosensor based on immobilization of glutamate oxidase onto carboxylated multiwalled carbon nanotubes/gold nanoparticles/chitosan composite film modified Au electrode. *Biosens. Bioelectron.* **2013**, *47*, 496–501. [CrossRef] [PubMed]
101. Isobe, K.; Sugawara, A.; Domon, H.; Fukuta, Y.; Asano, Y. Purification and characterization of an L-amino acid oxidase from *Pseudomonas* sp. AIU 813. *J. Biosci. Bioeng.* **2012**, *114*, 257–261. [CrossRef] [PubMed]
102. Koyama, H. Oxidation and oxygenation of L-amino acids catalyzed by a L-phenylalanine oxidase (Deaminating and decarboxylating) from *Pseudomonas* Sp. P-501. *J. Biochem.* **1984**, *96*, 421–427. [PubMed]
103. Suzuki, H.; Higashi, Y.; Asano, M.; Suguro, M.; Kigawa, M.; Maeda, M.; Katayama, S.; Mukouyama, E.B.; Uchiyama, K. Sequencing and expression of the L-phenylalanine oxidase gene from *Pseudomonas* sp. P-501. Proteolytic activation of the proenzyme. *J. Biochem.* **2004**, *136*, 617–627. [CrossRef] [PubMed]
104. Sobrado, P.; Fitzpatrick, P.F. Analysis of the role of the active site residue Arg98 in the flavoprotein tryptophan 2-monooxygenase, a member of the L-amino oxidase family. *Biochemistry* **2003**, *42*, 13826–13832. [CrossRef] [PubMed]
105. Genet, R.; Benetti, P.H.; Hammadi, A.; Menez, A. L-Tryptophan 2',3'-oxidase from *Chromobacterium violaceum*. Substrate specificity and mechanistic implications. *J. Biol. Chem.* **1995**, *270*, 23540–23545. [CrossRef] [PubMed]
106. Kameya, M.; Onaka, H.; Asano, Y. Selective tryptophan determination using tryptophan oxidases involved in bis-indole antibiotic biosynthesis. *Anal. Biochem.* **2013**, *438*, 124–132. [CrossRef] [PubMed]
107. Nishizawa, T.; Aldrich, C.; Sherman, D.H. Molecular analysis of the rebeccamycin L-amino acid oxidase from *Lechevalieria aerocolonigenes* ATCC 39243. *J. Bacteriol.* **2005**, *187*, 2084–2092. [CrossRef] [PubMed]
108. Gau, A.E.; Heindl, A.F.; Nodop, A.F.; Kahmann, U.F.; Pistorius, E.K. L-Amino acid oxidases with specificity for basic L-amino acids in cyanobacteria. *Z. Naturforsch. C Biosci.* **2007**, *62*, 273–284. [CrossRef]
109. Bockholt, R.; Scholten-Beck, G.F.; Pistorius, E.K. Construction and partial characterization of an L-amino acid oxidase-free *Synechococcus* PCC 7942 mutant and localization of the L-amino acid oxidase in the corresponding wild type. *Biochim. Biophys. Acta* **1996**, *1307*, 111–121. [CrossRef]
110. Qian, P.Y.; Lau, S.C.K.; Dahms, H.U.; Dobretsov, S.; Harder, T. Marine biofilms as mediators of colonization by marine macroorganisms: Implications for antifouling and aquaculture. *Mar. Biotechnol.* **2007**, *9*, 399–410. [CrossRef] [PubMed]

111. Holmstrom, C.; Egan, S.; Franks, A.; McCloy, S.; Kjelleberg, S. Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. *FEMS Microbiol. Ecol.* **2002**, *41*, 47–58. [[CrossRef](#)]
112. Shikuma, N.J.; Pilhofer, M.; Weiss, G.L.; Hadfield, M.G.; Jensen, G.J.; Newman, D.K. Marine tubeworm metamorphosis induced by arrays of bacterial phage tail-like structures. *Science* **2014**, *343*, 529–533. [[CrossRef](#)] [[PubMed](#)]
113. Celdran, D.; Espinosa, E.; Sanchez-Amat, A.; Atucha, A. Effects of epibiotic bacteria on leaf growth and epiphytes of seagrass *Posidonia oceanica*. *Mar. Ecol. Prog. Ser.* **2012**, *456*, 21–27. [[CrossRef](#)]
114. James, S.G.; Holmstrom, C.; Kjelleberg, S. Purification and characterization of a novel antibacterial protein from the marine bacterium D2. *Appl. Environ. Microbiol.* **1996**, *62*, 2783–2788. [[PubMed](#)]
115. Lucas-Elio, P.; Hernandez, P.; Sanchez-Amat, A.; Solano, F. Purification and partial characterization of marinocine, a new broad-spectrum antibacterial protein produced by *Marinomonas mediterranea*. *Biochim. Biophys. Acta* **2005**, *1721*, 193–203. [[CrossRef](#)] [[PubMed](#)]
116. Yu, M.; Wang, J.; Tang, K.; Shi, X.; Wang, S.; Zhu, W.M.; Zhang, X.H. Purification and characterization of antibacterial compounds of *Pseudoalteromonas flavipulchra* JG1. *Microbiology* **2012**, *158*, 835–842. [[CrossRef](#)] [[PubMed](#)]
117. Chen, W.M.; Lin, C.Y.; Chen, C.A.; Wang, J.T.; Sheu, S.Y. Involvement of an L-amino acid oxidase in the activity of the marine bacterium *Pseudoalteromonas flavipulchra* against methicillin-resistant *Staphylococcus aureus*. *Enzyme Microb. Technol.* **2010**, *47*, 52–58. [[CrossRef](#)]
118. Gomez, D.; Espinosa, E.; Bertazzo, M.; Lucas-Elio, P.; Solano, F.; Sanchez-Amat, A. The macromolecule with antimicrobial activity synthesized by *Pseudoalteromonas luteoviolacea* strains is an L-amino acid oxidase. *Appl. Microbiol. Biotechnol.* **2008**, *79*, 925–930. [[CrossRef](#)] [[PubMed](#)]
119. Yu, Z.; Zhou, N.; Qiao, H.; Qiu, J. Identification, cloning, and expression of L-amino acid oxidase from marine *Pseudoalteromonas* sp. B3. *Sci. World J.* **2014**, *2014*, 979858. [[CrossRef](#)] [[PubMed](#)]
120. Yu, Z.; Wang, J.; Lin, J.; Zhao, M.; Qiu, J. Exploring regulation genes involved in the expression of L-amino acid oxidase in *Pseudoalteromonas* sp. Rf-1. *PLoS ONE* **2015**, *10*. [[CrossRef](#)] [[PubMed](#)]
121. Chen, W.M.; Lin, C.Y.; Sheu, S.Y. Investigating antimicrobial activity in *Rheinheimera* sp. due to hydrogen peroxide generated by L-lysine oxidase activity. *Enzyme Microb. Technol.* **2010**, *46*, 487–493. [[CrossRef](#)] [[PubMed](#)]



© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).