


# Endogenous expression and localization of CAV-1::GFP in *C. elegans*

Dillon E. Sloan<sup>1</sup> and Joshua N. Bembenek<sup>1§</sup><sup>1</sup>University of Michigan

§To whom correspondence should be addressed: bembenek@umich.edu



**Figure 1. Endogenous CAV-1::GFP:** (A) Schematic of *cav-1* edit (B-G) Single-plane images of CAV-1::GFP localization: at the bend of the gonad arm (B), in the -1 oocyte (C), in a prometaphase I embryo (D), in an anaphase I embryo (E), in a bean stage embryo (F), and in a fold-stage embryo (G). (H) CAV-1::GFP in adult worm. Numbers correspond with number of cells in embryo.

## Description

Caveolins are integral membrane proteins responsible for the formation of caveolae, invaginations of the plasma membrane linked to various disease states (Parton *et al.* 2020). In *C. elegans*, there are two caveolin proteins, CAV-1 and CAV-2. The *cav-1* gene shares homology with all three mammalian caveolin genes (Tang *et al.* 1997). *C. elegans* CAV-1 protein does not appear to form caveolae, but a double-knockout mutant of *cav-1* and *cav-2* affects egg laying, and knockdown of *cav-1* affects locomotion in a dynamin mutant background (Parker *et al.* 2007, Kirkham *et al.* 2008, Sato *et al.* 2008). Based on exogenous expression, CAV-1::GFP is known to localize to cortical granules and the plasma membrane in oocytes and the early embryo, to the plasma membrane in later embryos, as well as to the neuromuscular system in larvae and adult worms (Sato *et al.* 2006, Bembenek *et al.* 2007, Parker *et al.* 2007).

Using CRISPR/Cas9, we generated an endogenous CAV-1::GFP fluorescent fusion strain, primarily to create a cortical granule marker (Figure 1A). Imaging endogenous CAV-1::GFP revealed similar patterns as previously reported in the germ line (Sato *et al.* 2006). However, unlike the *pie-1* driven exogenous line, endogenous CAV-1::GFP was not clearly detected in the distal gonad until the bend region of the gonad arm, where it localized to the plasma membrane, and vesicle-shaped structures (Figure 1B). Oocytes had slightly higher expression showing enrichment at plasma membrane localization but faint cytoplasmic signal, and no obvious enrichment to vesicles (Figure 1C). After ovulation, CAV-1::GFP remained on the plasma membrane during meiosis I and was no longer detected by meiosis II (Figure 1D,E). In 5/5

anaphase I embryos, CAV-1::GFP was not significantly localized to cortical granules at a time when separase could be observed on vesicles. Overall, endogenous CAV-1::GFP signal is less intense in the germline than exogenous *pie-1*-driven lines. However, endogenous CAV-1::GFP signal became increasingly intense at the plasma membrane sometime after the 16-cell embryo stage (Figure 1H). The membrane signal remains intense through later stages of embryonic development (Figure 1F,G). This potentially indicates a dynamic role for *cav-1* in embryonic development. Finally, adult somatic expression appeared to follow previously observed patterns (Figure 1H, Parker *et al.* 2007).

In summary, we have developed an endogenous CAV-1::GFP line by CRISPR, which can be of use to researchers interested in CAV-1.

## Methods

[Request a detailed protocol](#)

### CRISPR/Cas9 Gene editing

We followed the CRISPR/Cas9 protocol generated by Seydoux lab for C-terminal GFP tagging of the *C. elegans cav-1* gene (Figure 1A, Paix *et al.* 2015). The repair template was amplified from the pDD282 plasmid. All guide RNAs and oligos were obtained commercially.

The primer sequences are listed below:

*cav-1::GFP* Forward (termed DS\_027):

ACGGAATCAATCAAGAACTACTGCTCCATGCGTCATGAGTAAAGGAGAAGAATTGTTCACTG

*cav-1::GFP* Reverse (termed DS\_026):

AGTAAAATGAATTTGAGATAAATTAATAAATTTACTTGTAGAGCTCGTCCATTCC

The crRNA sequence recognizing the spacer upstream of NGG is as follows:

(Termed DS\_g003): AUUAAAUAAAUUUAGACGCA

### Microscopy

Imaging was done as indicated in Bai *et al.* 2020. Embryos were staged using the localization of separase fused with a red fluorescent protein as well as embryo position and age in the animal.

## Reagents

JAB192: *cav-1(erb78[cav-1::GFP])*

This strain is available upon request.

**Acknowledgments:** We thank members of the Bembenek and Csankovszki labs, especially Chris Turpin, for useful feedback.

## References

- Bai X., Melesse M., Sorensen Turpin C.G., Sloan D.E., Chen C.Y., Wang W.C., Lee P.Y., Simmons J.R., Nebenfuhr B., Mitchell D., Klebanow L.R., Mattson N., Betzig E., Chen B.C., Cheerambathur D., Bembenek J.N. (2020). Aurora B functions at the apical surface after specialized cytokinesis during morphogenesis in *C. elegans*. *Development* 147(1), dev181099. PMID: 31806662.
- Bembenek, J.N., Richie, C.T., Squirrel, J.M., Campbell, J.M., Eliceiri, K.W., Poteryaev, D., Spang, A., Golden, A., and White, J.G. (2007). Cortical granule exocytosis in *C. elegans* is regulated by cell cycle components including separase. *Development* 134, 3837–3848. PMID: 17913784.
- Kirkham, M., Nixon, S.J., Howes, M.T., Abi-Rached, L., Wakeham, D.E., Hanzal-Bayer, M., Ferguson, C., Hill, M.M., Fernandez-Rojo, M., Brown, D.A., Hancock, J.F., Brodsky, F.M., Parton, R.G. (2008). Evolutionary analysis and molecular dissection of caveola biogenesis. *J. Cell Sci.* 121, 2075–2086. PMID: 18505796.
- Paix, A., Folkmann, A., Rasoloson, D., Seydoux, G. (2015). High efficiency, homology-directed genome editing in *Caenorhabditis elegans* using CRISPR–Cas9 ribonucleoprotein complexes. *Genetics* 201, 47–54. PMID: 26187122.
- Parker, S., Peterkin, H. S., Baylis, H. A. (2007). Muscular dystrophy associated mutations in caveolin-1 induce neurotransmission and locomotion defects in *Caenorhabditis elegans*. *Invert. Neurosci* 7, 157–164. PMID: 17629760.
- Parton R.G., McMahon K.A., Wu Y. (2020). Caveolae: Formation, dynamics, and function. *Curr Opin Cell Biol* 65, 8–16. PMID: 32146331.

Sato, K., Sato, M., Audhya, A., Oegema, K., Schweinsberg, P., and Grant, B. D. (2006). Dynamic regulation of caveolin-1 trafficking in the germ line and embryo of *Caenorhabditis elegans*. *Mol. Biol. Cell* 17, 3085-3094. PMID: 16672374.

Sato, M., Grant, B.D., Harada, A., Sato, K. (2008). Rab11 is required for synchronous secretion of chondroitin proteoglycans after fertilization in *Caenorhabditis elegans*. *J Cell Sci* 121, 3177–3186. PMID: 18765566.

Tang, Z., Okamoto, T., Boontrakulpoontawee, P., Katada, T., Otsuka, A. J., and Lisanti, M. P. (1997). Identification, sequence, and expression of an invertebrate caveolin gene family from the nematode *Caenorhabditis elegans*. Implications for the molecular evolution of mammalian caveolin genes. *J. Biol. Chem.* 272, 2437–2445. PMID: 8999956.

**Funding:** National Institutes of Health (R01 GM114471)

**Author Contributions:** Dillon E. Sloan: Data curation, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review and editing, Software. Joshua N. Bembenek: Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Validation, Writing - review and editing.

**Reviewed By:** Sara Olson

**History:** **Received** August 6, 2020 **Revision received** September 18, 2020 **Accepted** September 18, 2020 **Published** September 21, 2020

**Copyright:** © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Sloan, DE; Bembenek, JN (2020). Endogenous expression and localization of CAV-1::GFP in *C. elegans*. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000311>