

Research Article

The Effect of Corneal Refractive Power Area Changes on Myopia Progression during Orthokeratology

Minfeng Chen, Xinting Liu, Zhu Xie, Pengqi Wang, Miaoran Zheng, and Xinjie Mao 

School of Ophthalmology and Optometry, Wenzhou Medical University, Wenzhou, Zhejiang 325000, China

Correspondence should be addressed to Xinjie Mao; mxj@mail.eye.ac.cn

Received 21 February 2022; Accepted 28 May 2022; Published 16 June 2022

Academic Editor: Kofi Asiedu

Copyright © 2022 Minfeng Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. To investigate the effect of corneal refractive power area changes on myopia progression during orthokeratology. **Methods.** One hundred and sixteen children who met the inclusion criteria and insisted on wearing orthokeratology lenses for two years were retrospectively assessed. Seventy-two children with the orthokeratology lens decentration distance more than 0.5 mm but less than 1.5 mm were in the decentered group, and forty-four children with the orthokeratology lens decentration distance less than 0.5 mm were in the centric group. The orthokeratology decentration via tangential difference topography was analyzed. This study calculated the different power areas in the central 4 mm pupillary area by axial-difference corneal topography, compared the differences of the different power areas between these two groups, and evaluated the relationships between corneal positive-power area, orthokeratology decentration, and AL changes. **Results.** The axial length changes of the centric group presented a statistical difference with the decentered group (0.52 ± 0.37 mm vs. 0.38 ± 0.26 mm; $t = 2.403$, $p = 0.018$). For all children, both the AL changes (0.43 ± 0.31 mm) and decentration distance (0.64 ± 0.33 mm) showed a significant correlation with the positive-power area ($r = -0.366$, $p < 0.001$ and $r = 0.624$, $p < 0.001$); AL changes also presented a statistical correlation with decentration distance ($r = -0.343$, $p < 0.001$), baseline age ($r = -0.329$, $p < 0.001$), and baseline spherical equivalent refractive power ($r = 0.335$, $p < 0.001$). In the centric group and decentered group, the AL changes (centric group: $r = -0.319$, $p = 0.035$; decentered group: $r = -0.332$, $p = 0.04$) and decentration distance (centric group: $r = 0.462$, $p = 0.002$; decentered group: $r = 0.524$, $p < 0.001$) had a significant correlation with the positive-power area yet. In the multiple regression analysis, AL changes were increased with less baseline age (beta, 0.015; $p < 0.001$), positive-power area (beta, 0.021; $p = 0.002$), and larger SER (beta, 0.025; $p = 0.018$). **Conclusions.** The corneal positive-power area had a positive impact on affirming AL changes during orthokeratology. This area might be formed by lens decentration to provide an additional myopia-defocusing influence on the retina to achieve better myopia control.

1. Introduction

Currently, there are around 1.4 billion myopic people in the world, nearly half of whom live in China. Without effective intervention, it is estimated that by 2050, there will be 4.758 billion children with myopia, accounting for about half of the total population [1, 2]. Thus, myopia is a global public health concern. Many studies have confirmed that orthokeratology is a safe, effective, and reversible method for slowing axial length (AL) changes [3–5]. After orthokeratology treatment, AL changes are reduced by 30–50% [3–6], effectively mitigating the development of myopia.

Orthokeratology changes the corneal surface by flattening the central cornea and steepening the mid-peripheral cornea

to provide a clear unaided distance vision and myopia control; this is achieved through the application of gas-permeable contact lenses [6–8]. However, the mechanism underlying the effect of orthokeratology on the control of myopia progression remains unknown. Moreover, the effects of orthokeratology on mitigating AL changes vary significantly among children. Many factors [8–10] influence the level of myopia control achieved with orthokeratology, including baseline age, baseline spherical equivalent refractive power (SER), and pupil diameter. Zhong et al. also found that changes in corneal refractive power significantly affected AL changes; specifically, subjects with a larger magnitude of corneal refractive power change after orthokeratology treatment typically experienced slower AL changes [11–16].

Following orthokeratology treatment, the cornea becomes a multifocal cornea [6]. Hiraoka et al. found that corneal multifocality was negatively correlated with the AL change, whereas patients with increased corneal multifocality showed less AL elongation [13–17]. Hiraoka et al. suggested that corneal multifocality might be affected by lens decentration. Recent studies [11, 12] have also revealed that lens decentration affects myopia control during orthokeratology. Wang and Yang [11] found that acceptable lens decentration could delay the development of myopia more effectively than when the lens is in a centric position.

In previous studies [11–20], both corneal refractive power change and lens decentration have been shown to have an effect on AL changes, and lens decentration may have an effect on changes in the corneal refractive power [11, 12, 20, 21]. Thus, it is worth investigating whether there might be relationships between decentration, corneal refractive power changes, and AL changes.

To this end, the current study calculated lens decentration using tangential corneal topography and analyzed the areas of corneal refractive power change during orthokeratology using axial corneal topography [18–23]. Then, the relationships between lens decentration, AL changes, and corneal refractive power changes were investigated.

2. Methods

2.1. Study Subjects. A retrospective analysis was performed on 116 children (66 males and 50 females), ranging in age from 8 to 14 years (mean age, 11.19 ± 1.63 years). The subjects were all treated at Wenzhou Medical University Eye Hospital between June 2016 and October 2018 and were asked to wear orthokeratology lenses for no less than 8 hours per night over a two-year period. The inclusion criteria were as follows: age 8–15 years; spherical refractive power (SR) of -1.00 to -6.00 diopters (D); astigmatism of 0.00 to -1.50 D; binocular best-corrected visual acuity of 20/20 or greater; intraocular pressure of 10 to 21 mmHg; lens-fitting decentration less than 1.5 mm, to prevent the sclera from being covered [6, 11]; no other eye diseases; no history of surgery or use of atropine or contact lenses to control myopia progression; no systemic or ocular conditions that might affect vision. This study was performed in accordance with the tenets of the Declaration of Helsinki and was approved by the Ethics Committee of Wenzhou Medical University Eye Hospital.

2.2. Instrumentation. The orthokeratology lenses used in this study were four-zoned reverse-geometry lenses (Euclid Systems Corporation, USA). These lenses range in diameter from 10.2 mm to 11.2 mm and have a central thickness ranging from 0.22 mm to 0.23 mm. Each subject underwent a comprehensive baseline eye examination, including slit-lamp examination, refraction, uncorrected visual acuity, best-corrected visual acuity, AL (ZEISS IOLMaster; Carl Zeiss AG, Oberkochen, Germany), corneal topography (E-300; Medmont International Pty. Ltd.), corneal endothelium

microscopy, and intraocular pressure (NT-2000; NIDEK CO., LTD., Gamagori, Japan).

All children were treated by doctors who had worked in the field of orthokeratology treatment at the eye hospital for more than 10 years. Corneal topography was measured with a Medmont E300 (Medmont International Pty. Ltd.); measurement was performed by a specialized technician within one hour of removal of the orthokeratology lenses. The topographic images used for analysis were each subject's best focus image (with an accuracy greater than 95%) from the four frames that were captured automatically. Based on each subject's corneal topography and the fitting evaluation based on corneal fluorescein pattern analysis, the doctor ordered lenses for the subject that were the most suitable based on their corneal parameters, according to the manufacturer's guidelines.

2.3. Measurements. Through corneal topography, the areas of different refractive powers within the central 4 mm pupillary area were calculated and recorded as the refractive distribution of the cornea; this is used as an index of corneal multifocality [6, 13, 20–24]. In all subjects, the right eye was selected for analysis.

Based on the recorded power areas of the pupillary area obtained from axial difference topography, the following measurements were made: positive-power area (positive D value), corrected area (D value = baseline SER \pm 0.25 D), overcorrected area (D value > baseline SER $-$ 0.25 D), and undercorrected area (D value < baseline SER $+$ 0.25 D). To calculate the different refractive powers in the central 4 mm pupillary area, axial difference topography was performed, based on the axial topography [25], at the baseline and after three months of continuous orthokeratology lens wear. According to the axial difference topography, the refractive power parameter was set as the individual baseline SER power (Figure 1).

To assess the optical zone decentration distance, the tangential difference topography was constructed through tangential topography measurements collected at the baseline and after three months of continuous orthokeratology. According to the tangential difference topography, the optical zones ranged from the corneal vertex to (1) where the keratometry values changed within 1 D and (2) there were less than two colors (red and blue) in the refractive power parameter [18–21].

After importing the tangential difference topography into MATLAB (MathWorks, Inc), 16 points were selected in a clockwise fashion, on the red-blue transition zone, to delineate the margins of the optical zone [18, 26, 27]. The pupil center was determined by the Medmont E300. The decentration distance of orthokeratology was defined from the center of the optical zone to the pupil center [12, 13, 28, 29] and was measured using ImageJ software (National Institutes of Health, Bethesda, MD, USA) (Figure 2). Based on the decentration distance, participants were divided into a centric group (decentration distance less than 0.5 mm) and decentered group (decentration distance larger than 0.5 mm).




FIGURE 1: Areas of different refractive powers in the central 4 mm pupillary area ((a) positive-power area, (b) overcorrected area, (c) corrected area, and (d) undercorrected area).




FIGURE 2: Optical zone decentration distance measurement (decentration distance was measured from the center of the optical zone to the pupil center).

2.4. Statistical Analysis. The degree of myopia was expressed by the SER. Measurement data are expressed as mean \pm standard deviation (SD). Intragroup comparisons were performed using the paired *t*-test and intergroup comparisons were performed using analysis of variance or the independent *t*-test; the Wilcoxon rank-sum test for two related samples was used to compare data across time points. Pearson correlation analysis was performed to examine the relationships between the variables. A *p* value less than 0.05 was considered a statistically significant difference. Statistical Package for the Social Sciences (SPSS) version 25 software (IBM Corporation, Armonk, NY, USA) was used for the statistical analyses.

3. Results

3.1. Study Subjects. A total of 116 myopic children (mean age, 11.19 ± 1.63 years, range 8 to 14 years) were included in the sample. Before orthokeratology, the SER, SR, and regular astigmatism values were -3.56 ± 1.21 D (range -6.00 to -1.00 D), -3.41 ± 1.17 D (range -6.00 to -1.00 DS), and -0.36 ± -0.40 D (range -1.50 to 0.00 DC), respectively. The baseline parameters for both the centric group (decentration

distance: 0.31 ± 0.13) and decentered group (decentration distance: 0.84 ± 0.25) are shown in Table 1.

Among the total sample, significant differences were observed between the baseline and two-year AL change (25.07 ± 0.81 mm vs. 25.50 ± 0.76 mm; $t = -12.547$, $p < 0.001$). The AL change in the centric group was significantly different from that of the decentered group (0.52 ± 0.37 mm vs. 0.38 ± 0.26 mm; $t = 2.403$, $p = 0.018$). After orthokeratology, the distributions of the different refractive power areas in the pupillary areas of these two groups are shown in Figure 2. There were no statistically significant differences in the following parameters between the centric group and decentered group: undercorrected area (8.49 ± 3.56 mm² vs. 7.37 ± 3.20 mm², $t = 1.745$, $p = 0.084$), corrected area (1.72 ± 1.71 mm² vs. 1.66 ± 1.50 mm², $t = 0.190$, $p = 0.850$), and overcorrected area (1.79 ± 3.01 mm² vs. 1.90 ± 2.49 mm², $t = -0.215$, $p = 0.830$). However, the positive-power area (0.51 ± 0.86 mm² vs. 1.61 ± 1.27 mm², $t = -5.087$, $p < 0.001$) significantly differed between the centric group and the decentered group, as shown in Figure 3.

Among the 116 children, both the AL change (0.43 ± 0.31 mm) and decentration distance (0.64 ± 0.33 mm) were significantly correlated with the positive-power area ($r = -0.366$, $p < 0.001$ and $r = 0.624$, $p < 0.001$, respectively). Further, AL change was significantly correlated with the decentration distance ($r = -0.343$, $p < 0.001$), baseline age ($r = -0.329$, $p < 0.001$), and baseline SER ($r = 0.335$, $p < 0.001$). In the centric group and decentered group, the AL change (centric group: $r = -0.319$, $p = 0.035$; decentered group: $r = -0.332$, $p = 0.04$) and decentration distance (centric group: $r = 0.462$, $p = 0.002$; decentered group: $r = 0.524$, $p < 0.001$) were significantly correlated with the positive-power area, as shown in Figure 4.

The relationships between baseline age, baseline SER, decentration distance, different power areas, and AL change among the total sample of 116 children are shown in Table 2. In the multiple regression analysis, AL change was associated with younger baseline age (beta, 0.015; $p < 0.001$), positive-power area (beta, 0.021; $p = 0.002$), and larger SER (beta, 0.025; $p = 0.018$). However, the decentration distance, overcorrected area, and corrected area were not significantly associated with AL change in the multiple linear regression analysis, as shown in Table 2.

TABLE 1: The baseline parameters between the centric group and decentered group (mean \pm SD).

Groups	Centric position ($n = 44$)	Decentered position ($n = 72$)	t	p
Age (y)	10.86 \pm 1.84	11.39 \pm 1.47	-1.697	0.092
SR (DS)	-3.27 \pm 1.22	-3.49 \pm 1.15	0.967	0.336
Astigmatism (DC)	0.37 \pm 0.43	0.36 \pm 0.39	-0.151	0.880
SER (D)	-3.41 \pm 1.31	-3.64 \pm 1.14	0.985	0.327
AL (mm)	25.14 \pm 0.83	25.03 \pm 0.81	0.573	0.569
Flat-K (D)	43.02 \pm 1.24	42.63 \pm 1.24	1.546	0.125
Steep-K (D)	44.22 \pm 1.23	44.08 \pm 2.07	0.378	0.706
ΔK (D)	1.20 \pm 0.44	1.30 \pm 0.63	-0.847	0.399

Note. SR: spherical refractive power; SER: spherical equivalent refractive power; AL: axial length; flat-K: flat radial refractive power; steep-K: steep radial refractive power; ΔK : the difference between flat-K and steep-K.




FIGURE 3: Comparison of the distribution of different refractive power areas between the centric group and decentered group (** $p < 0.001$).




FIGURE 4: Correlation analysis of AL change and decentration distance with the positive-power area in the centric group and decentered group ((a) correlation between AL change and positive-power area in the centric group and decentered group; (b) correlation between decentrated distance and positive-power area in the centric group and decentered group).

4. Discussion

Orthokeratology is a safe and effective method for slowing changes in the AL [3–5], and more and more children and parents are choosing orthokeratology to prevent increases in

myopia. However, the effect of orthokeratology on myopia control is influenced by eyelid tension, lens movement, and corneal asymmetry, among other factors. As a result, decentration [11, 12, 20, 21] has become a common phenomenon and cannot be completely avoided during

TABLE 2: Univariate and multiple linear regression analysis to determine the correlations between AL change and baseline age, baseline SER, decentration distance, and the different power areas.

	Univariate linear regression			Multiple linear regression		
	B	β	<i>p</i>	B	β	<i>p</i>
Age (y)	-0.063	-0.329	<0.001	-0.055	0.015	<0.001
SER (D)	0.086	0.335	<0.001	0.066	0.021	0.002
Decentration distance (mm)	-0.325	-0.343	<0.001			
Overcorrected area (mm ²)	0.022	0.194	0.037			
Corrected area (mm ²)	0.046	0.234	0.011			
Positive-power area (mm ²)	-0.091	-0.366	<0.001	-0.060	0.025	0.018

Note. AL: axial length; SER: spherical equivalent refractive power.

orthokeratology treatment. Recently, studies [11, 12] have found that decentration might also have an effect on myopia control, in addition to the above previously known factors.

In the study by Wang and Yang [11], after orthokeratology, the AL change in the lens decentered group was smaller than that in the lens centric group. Moreover, Chen et al. [12] found a negative correlation between the lens decentration distance and AL change. In the current study, the myopic children wore orthokeratology lenses for two years and did not experience any serious complications that required them to stop wearing the lenses. In the sample of the present study, the mean decentration distance was 0.64 ± 0.33 mm (range: 0.38–1.40 mm), similar to the distances of 0.73 ± 0.25 mm (range: 0.5–1.50 mm) and 0.64 ± 0.38 mm (range: 0.13–1.78 mm) reported in the studies conducted by Wang and Yang [11], and Chen et al. [12]. Furthermore, consistent with other recent studies [11, 12], in the current study, in addition to the correlations between AL change and both baseline age and baseline SER, children with a greater decentration distance exhibited less change in AL after orthokeratology.

Modern orthokeratology is a process that uses reverse-geometry-designed rigid contact lenses to reshape the eye over time. Each lens had a back optical zone diameter of 6.0 mm, a reverse curve of 0.6 mm in width, an alignment curve of 1.25 mm in width, and a peripheral curve of 0.4 mm in width [14, 25, 28]. After the base curve flattens the cornea, ideally, a flat zone is formed to correct the refractive error. Unfortunately, after orthokeratology, the cornea becomes a multifocal cornea with many different refractive power areas in this zone, including an overcorrected area, corrected area, and undercorrected area. Hiraoka et al. [14] showed that greater decentration caused a greater change in corneal multifocality.

According to previous studies, the 4 mm area in the center of the pupil is the effective applanation zone. To investigate the relationships between the decentration distance and AL change, the areas of different refractive power in the central 4 mm pupil area were investigated using corneal topography. The results indicated that, compared with the centric group, the decentered group had a greater positive-power area in the central 4 mm pupil area, and the positive-power area was significantly negatively correlated with the decentration distance in both the centric group and decentered group. However, there were no significant differences in the undercorrected area, corrected area, and overcorrected area

between the centric group and decentered group. The reversal curve, which is next to the base curve, forms the positive-power area in the cornea; thus, when the position of the orthokeratology lens is decentered, there is a greater positive-power area in the central 4 mm pupil area.

Correlation analysis indicated that the positive-power area was negatively correlated with the AL change, with a greater positive-power area associated with less AL elongation. This is consistent with the relationship between the decentration distance and AL change reported in previous studies [11, 12], where a larger decentration distance was associated with a greater positive-power area. In the univariate linear regression analysis, the decentration distance and positive-power area were significantly associated with AL change, but in the multiple linear regression analysis, the positive-power area, and not the decentration distance, was significantly associated with AL change. Thus, it appears that decentration changes the corneal refractive power of the pupil area, impacting the positive-power area and, in turn, influencing AL change.

Lam et al. [29] found that a custom-made spectacle lens (composed of a central optical zone for correcting the distance refractive error and an annular multifocal zone with multiple segments having a relative positive power) could effectively slow myopia progression. Lam et al.'s study confirmed that paracentral myopia defocus is effective for myopia control. Today, children have to work at a close range to their viewing target for long durations, due to the increasing burden of schoolwork. In this context, although myopia may be corrected with orthokeratology, these children may still experience a certain degree of adjustment lag [30–32], which might cause the fundus image to form behind the retina instead of on the retina, leading to hyperopia defocus [33, 34] on the retina; this would, in turn, accelerate the progression of myopia. However, among children with positive-power areas in the pupil zone, it appears that there is a positive-power lens in front of the cornea, providing paracentral myopia defocus on the retina; this would move the image from behind the retina to onto the retina or even in front of the retina, turning the hyperopia defocus into myopia defocus [33, 34]. This would result in an additional degree of myopia defocus on the paracentral retina, and thus, would provide a better myopia-control effect.

One limitation of the current study is that the measurement of lens decentration was only performed after the treatment had begun. Although previous studies have

confirmed that there are no significant changes in decentration after two years of treatment, it would be valuable to know the magnitudes of decentration and positive-power area changes during the course of orthokeratology. Another limitation is that the degree of defocus could not be measured according to the positive-power area. This makes it difficult to decide how large the positive-power area should be in order to provide the best myopia control treatment effect. Additional studies should be performed to clarify these points.

5. Conclusions

In summary, after orthokeratology lens wear, optical zone decentration is common and unavoidable. A greater decentration distance might change the corneal refractive power of the pupil area, resulting in a certain degree of positive-power area. This might, in turn, change hyperopia defocus into myopia defocus, resulting in an additional degree of myopia defocus on the paracentral retina. This would be helpful for children when performing their schoolwork at a close range, thus providing a better myopia-control effect.

Data Availability

The data used to support the findings of this study are available from the corresponding author (Xinjie Mao) upon request.

Consent

The authors approved the final manuscript for submission.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Xinjie Mao, the co-corresponding author, contributed to project design and the revision of this manuscript. Minfeng Chen, the first author, contributed to data collection and analysis, and writing of the manuscript. Xinting Liu, Zhu Xie, Pengqi Wang, and Miaoran Zheng contributed to data collection and analysis. All authors read and approved the final manuscript.

Acknowledgments

The authors are grateful for the support and help of the Optometry Center of Wenzhou Medical University Eye Hospital. This study was supported by the National Key Research and Development Project (2019YFC1710204).

References

- [1] E. Dolgin, "The myopia boom," *Nature*, vol. 519, no. 7543, pp. 276–278, 2015.
- [2] B. A. Holden, T. R. Fricke, D. A. Wilson et al., "Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050," *Ophthalmology*, vol. 123, no. 5, pp. 1036–1042, 2016.
- [3] T. Hiraoka, Y. Sekine, F. Okamoto, T. Mihashi, and T. Oshika, "Safety and efficacy following 10-years of overnight orthokeratology for myopia control," *Ophthalmic and Physiological Optics*, vol. 38, no. 3, pp. 281–289, 2018.
- [4] J. Huang, D. Wen, Q. Wang et al., "Efficacy comparison of 16 interventions for myopia control in children: a network meta-analysis," *Ophthalmology*, vol. 123, no. 4, pp. 697–708, 2016.
- [5] Y. M. Liu and P. Xie, "The safety of orthokeratology--a systematic review," *Eye and Contact Lens: Science and Clinical Practice*, vol. 42, no. 1, pp. 35–42, 2016.
- [6] J. Chen, W. Huang, R. Zhu, J. Jiang, and Y. Li, "Influence of overnight orthokeratology lens fitting decentration on corneal topography reshaping," *Eye and Vision*, vol. 5, no. 1, p. 5, 2018.
- [7] A. Queirós, A. Amorim-De-Sousa, D. Lopes-Ferreira, C. Villa-Collar, ÁR. Gutiérrez, and J. M. González-Méijome, "Relative peripheral refraction across 4 meridians after orthokeratology and LASIK surgery," *Eye and Vision*, vol. 5, no. 1, p. 12, 2018.
- [8] P. Cho, S. W. Cheung, and M. Edwards, "The longitudinal orthokeratology research in children (LORIC) in Hong Kong: a pilot study on refractive changes and myopic control," *Current Eye Research*, vol. 30, no. 1, pp. 71–80, 2005.
- [9] P. Cho and S. W. Cheung, "Retardation of myopia in orthokeratology (ROMIO) study: a 2-year randomized clinical trial," *Investigative Ophthalmology and Visual Science*, vol. 53, no. 11, p. 7077, 2012.
- [10] J. Santodomingo-Rubido, C. Villa-Collar, B. Gilmartin, and R. Gutiérrez-Ortega, "Factors preventing myopia progression with orthokeratology correction," *Optometry and Vision Science*, vol. 90, no. 11, pp. 1225–1236, 2013.
- [11] A. Wang and C. Yang, "Influence of overnight orthokeratology lens treatment zone decentration on myopia progression," *Journal of Ophthalmology*, vol. 2019, Article ID 2596953, 7 pages, 2019.
- [12] R. Chen, Y. Chen, M. Lipson et al., "The effect of treatment zone decentration on myopic progression during orthokeratology," *Current Eye Research*, vol. 45, no. 5, pp. 645–651, 2020.
- [13] T. Hiraoka, T. Kakita, F. Okamoto, and T. Oshika, "Influence of ocular wavefront aberrations on axial length elongation in myopic children treated with overnight orthokeratology," *Ophthalmology*, vol. 122, no. 1, pp. 93–100, 2015.
- [14] T. Hiraoka, T. Mihashi, C. Okamoto, F. Okamoto, Y. Hirohara, and T. Oshika, "Influence of induced decentered orthokeratology lens on ocular higher-order wavefront aberrations and contrast sensitivity function," *Journal of Cataract and Refractive Surgery*, vol. 35, no. 11, pp. 1918–1926, 2009.
- [15] T. Hiraoka, T. Kakita, F. Okamoto, H. Takahashi, and T. Oshika, "Long-term effect of overnight orthokeratology on axial length elongation in childhood myopia: a 5-year follow-up study," *Investigative Ophthalmology and Visual Science*, vol. 53, no. 7, p. 3913, 2012.
- [16] Y. Zhong, Z. Chen, F. Xue, J. Zhou, L. Niu, and X. Zhou, "Corneal power change is predictive of myopia progression in orthokeratology," *Optometry and Vision Science*, vol. 91, no. 4, pp. 404–411, 2014.
- [17] J. Santodomingo-Rubido, C. Villa-Collar, B. Gilmartin, and R. Gutiérrez-Ortega, "Short-term and long-term changes in corneal power are not correlated with axial elongation of the eye induced by orthokeratology in children," *Eye and Contact*

- Lens: Science and Clinical Practice*, vol. 44, no. 4, pp. 260–267, 2018.
- [18] J. Santodomingo-Rubido, C. Villa-Collar, B. Gilmartin, and R. Gutiérrez-Ortega, “Myopia control with orthokeratology contact lenses in Spain: refractive and biometric changes,” *Investigative Ophthalmology and Visual Science*, vol. 53, no. 8, p. 5060, 2012.
- [19] J. Santodomingo-Rubido, C. Villa-Collar, B. Gilmartin, R. Gutiérrez-Ortega, and A. Suzuki, “The effects of entrance pupil centration and coma aberrations on myopic progression following orthokeratology,” *Clinical and Experimental Optometry*, vol. 98, no. 6, pp. 534–540, 2015.
- [20] X. Yang, X. Zhong, X. Gong, and J. Zeng, “Topographical evaluation of the decentration of orthokeratology lenses,” *Yan Ke Xue Bao*, vol. 21, no. 3, 2005.
- [21] Y. Y. Tsai and J. M. Lin, “Ablation centration after active eye-tracker-assisted photorefractive keratectomy and laser in situ keratomileusis,” *Journal of Cataract and Refractive Surgery*, vol. 26, no. 1, pp. 28–34, 2000.
- [22] J. Li, C. Yang, W. Xie et al., “Predictive role of corneal Q-value differences between nasal-temporal and superior-inferior quadrants in orthokeratology lens decentration,” *Medicine (Baltimore)*, vol. 96, no. 2, Article ID e5837, 2017.
- [23] V. K. Maseedupally, P. Gifford, E. Lum et al., “Treatment zone decentration during orthokeratology on eyes with corneal toricity,” *Optometry and Vision Science*, vol. 93, no. 9, pp. 1101–1111, 2016.
- [24] Y. Hu, C. Wen, Z. Li, W. Zhao, X. Ding, and X. Yang, “Areal summed corneal power shift is an important determinant for axial length elongation in myopic children treated with overnight orthokeratology,” *British Journal of Ophthalmology*, vol. 103, no. 11, pp. 1571–1575, 2019.
- [25] T. Gu, B. Gong, D. Lu et al., “Influence of corneal topographic parameters in the decentration of orthokeratology,” *Eye and Contact Lens: Science and Clinical Practice*, vol. 45, no. 6, pp. 372–376, 2019.
- [26] Z. Chen, F. Xue, J. Zhou et al., “Prediction of orthokeratology lens decentration with corneal elevation,” *Optometry and Vision Science*, vol. 94, no. 9, pp. 903–907, 2017.
- [27] Y. Y. Tsai, S. H. Tseng, and J. M. Lin, “Comparison of ablation centration in initial and retreatment active eye-tracker-assisted laser in situ keratomileusis and the effect on visual outcome,” *Journal of Cataract and Refractive Surgery*, vol. 30, no. 7, pp. 1521–1525, 2004.
- [28] L. H. Yu, W. Q. Jin, X. J. Mao, and J. Jiang, “Effect of orthokeratology on axial length elongation in moderate myopic and fellow high myopic eyes of children,” *Clinical and Experimental Optometry*, vol. 104, no. 1, pp. 22–27, 2021.
- [29] C. S. Y. Lam, W. C. Tang, D. Y. Y. Tse et al., “Defocus incorporated multiple segments (DIMS) spectacle lenses slow myopia progression: a 2-year randomised clinical trial,” *British Journal of Ophthalmology*, vol. 104, no. 3, pp. 363–368, 2020.
- [30] D. O. Mutti, G. L. Mitchell, J. R. Hayes et al., “Accommodative lag before and after the onset of myopia,” *Investigative Ophthalmology and Visual Science*, vol. 47, no. 3, p. 837, 2006.
- [31] N. S. Logan and J. S. Wolffsohn, “Role of un-correction, under-correction and over-correction of myopia as a strategy for slowing myopic progression,” *Clinical and Experimental Optometry*, vol. 103, no. 2, pp. 133–137, 2020.
- [32] J. J. Walline, A. Gaume Giannoni, L. T. Sinnott et al., “A randomized trial of soft multifocal contact lenses for myopia control: baseline data and methods,” *Optometry and Vision Science*, vol. 94, no. 9, pp. 856–866, 2017.
- [33] D. Moderiano, M. Do, S. Hobbs et al., “Influence of the time of day on axial length and choroidal thickness changes to hyperopic and myopic defocus in human eyes,” *Experimental Eye Research*, vol. 182, pp. 125–136, 2019.
- [34] E. P. Tarutta, O. V. Proskurina, N. A. Tarasova, S. V. Milash, and G. A. Markosyan, “Otdalennyye rezul’taty ochkovoï korrektsii s perifokal’nym defokusom u detei s progressivnoy miopiei [Long-term results of perifocal defocus spectacle lens correction in children with progressive myopia],” *Vestnik Oftalmologii*, vol. 135, no. 5, pp. 46–53, 2019.