

RESEARCH ARTICLE

Open Access

Risk Factors for HSV-2 Infection among Sexual Partners of HSV-2/HIV-1 Co-Infected Persons

Andrew Mujugira^{1*}, Amalia S Magaret², Jared M Baeten^{1,3,4}, Connie Celum^{1,3,4}, Jairam Lingappa^{1,3,5}

Abstract

Background: Herpes simplex virus type 2 (HSV-2) is the most frequent cause of genital ulcer disease worldwide and has been associated with increased risk for HIV-1 acquisition and transmission. We conducted a cross-sectional analysis of risk factors for HSV-2 infection among HIV-1 uninfected partners, whose partners were co-infected with HIV-1 and HSV-2.

Methods: Between November 2004 and April 2007, 3408 HIV-discordant couples, in which the HIV-1 infected partners were HSV-2 seropositive with CD4 250 cells/mm³ or greater, were enrolled in an HSV-2 suppression trial to prevent HIV-1 transmission at 14 sites in 7 African countries. Clinical & behavioral data, HSV-2 and HIV-1 testing were conducted at enrolment. Univariate and multivariate Poisson regression analyses were performed separately, by gender of the HIV-1 infected partner.

Results: Among 3354 HIV-1 uninfected participants, 32% were female and overall 71% were HSV-2 seropositive. Among couples with female HIV-1 infected partners, HIV-1 plasma RNA [aPR 1.03; 95% CI: 0.99 to 1.06; p = 0.11] and CD4 count [aPR 1.00; 95% CI: 0.98 to 1.01; p = 0.48] in the HSV-2/HIV-1 dually infected female and circumcision in the HIV-1 uninfected male partner [aPR 0.94; 95% CI: 0.88 to 1.00; p = 0.06] were not associated with reduced risk of HSV-2 seropositivity, after adjusting for other factors.

Conclusions: In this cross-sectional analysis of African HIV-1 serodiscordant heterosexual couples with prevalent HSV-2 infection in the HIV-1 infected partner, HIV-1 plasma RNA and CD4 count in the dually-infected partner and male circumcision in the HIV-1 uninfected partner were not associated with HSV-2 concordance.

Trial Registration: ClinicalTrials.gov NCT00194519

Background

Herpes simplex virus type 2 (HSV-2) is the most common cause of genital ulcer disease (GUD) worldwide. HSV-2 prevalence in sub-Saharan Africa ranges from 27-57% in men to 30-74% in women [1,2] and is higher in HIV-1 infected persons [3]. Epidemiologic studies suggest synergy between HIV-1 and HSV-2 that facilitates the spread of both viruses, with HSV-2 increasing HIV-1 susceptibility and infectiousness [4] and HIV-1 infection increasing HSV-2 reactivation frequency [5].

The increased risk of HIV-1 transmission and acquisition due to HSV-2 infection in the context of an overall high risk of HIV-1 transmission among stable, heterosexual HIV-1 serodiscordant couples (one partner HIV-

1 infected and the other HIV-1 uninfected) [6], which are highly prevalent in Africa [7], make evaluation of risk factors for HSV-2 within HIV-1 serodiscordant couples from diverse parts of Africa of interest.

We conducted a cross-sectional analysis of risk factors for HSV-2 seropositivity in >3300 HIV-1 uninfected members of African HIV-1 serodiscordant couples in which the HIV-1 infected partner was dually-infected with HSV-2. In particular, we sought to assess whether HIV-1 plasma RNA level or CD4 count in the HIV-1 infected partner, as indicators of higher HIV-1 infectiousness and more advanced immunosuppression, modifies HSV-2 infectiousness, and if male circumcision in the HIV-uninfected partners of women with HSV-2/HIV-1 dual infection protected them from HSV-2 acquisition.

* Correspondence: mujugira@uw.edu

¹Department of Global Health, University of Washington, Seattle, WA, USA
Full list of author information is available at the end of the article

Methods

Study Population

From November 2004 to April 2007, 3408 heterosexual HIV-1 discordant couples from 14 sites in Botswana, Kenya, Rwanda, South Africa, Tanzania, Uganda and Zambia were enrolled into the Partners in Prevention HSV/HIV Transmission Study, a randomized clinical trial of suppressive acyclovir therapy to reduce sexual HIV-1 transmission from HSV-2/HIV-1 dually-infected women and men to their HIV-1 uninfected partners [7]. HIV-1 discordant couples were eligible for enrolment into the trial if they were sexually active and planning to remain partnered for at least 24 months and if the HIV-1-infected participant was HSV-2 infected, had a CD4 count 250 cells/mm³ or greater, was not on antiretroviral therapy, and was not pregnant. HIV-1 uninfected members of the discordant couples could be either HSV-2 seropositive or seronegative. Clinical and behavioral data collected at study enrolment were used for this analysis.

Laboratory Methods

HIV-1 testing was done using 2 rapid assays in parallel, most commonly Determine HIV-1/2 (Abbott Diagnostics), Uni-Gold HIV-1 (Trinity Biotech), or Capillus HIV-1/HIV-2 (Trinity Biotech) and paired positive rapid assay results were confirmed by HIV-1 EIA [7]. At all study sites, HSV-2 serostatus was determined using HerpeSelect HSV-2 gG2-based EIA (Focus Technologies). Given the results of previous studies using this assay in African cohorts, a cut-off value of ≥ 3.5 was chosen to define seropositivity, in order to improve test specificity [8-10]. CD4 quantification was performed for HIV-1 infected participants at 6-month intervals using standard flow cytometry. Plasma HIV-1 RNA was quantified using the COBAS Ampliprep/COBAS TaqMan real-time HIV-1 RNA assay, version 1.0 (Roche Diagnostics, Indianapolis, IN). All HIV-1 and HSV-2 serologic results were confirmed by Western blot, performed at the University of Washington [11].

Statistical Analysis

Data were entered onto standard case report forms and faxed to a central database at the University of Washington. Analysis was done using SAS version 9.1 (SAS Institute, Cary, NC). Univariate and multivariate analyses were performed using Poisson regression with HSV-2 status of the partner participant as the outcome variable. In order to examine male circumcision and to allow for other possible gender differences in transmission risk, separate analyses were performed for male and female HIV-1 infected participants. Backward elimination, starting with all variables significant at $p < 0.1$ in univariate regression, was performed to select a final multivariate model.

Ethical Approval

Institutional review boards at the University of Washington and at all collaborating site organizations approved study procedures. All participants provided written informed consent. The trial was registered through ClinicalTrials.gov (NCT00194519).

Results

Characteristics of study population

Of 3408 enrolled HIV-1-uninfected participants, 3354 were eligible for this analysis (54 were excluded because either they or their partner were ineligible for the study, based on confirmatory HIV-1 or HSV-2 testing). HIV-1 seronegative partners had a median age of 35 years and 1084 (32%) were female. Couples reported a median of 4 sex acts in the month prior to enrollment (IQR 2-8). The seroprevalence of HSV-2 among HIV-1 seronegative partners was 71% overall; 63% in men versus 89% in women ($p < 0.001$). By study design, all HIV-1 infected partners were co-infected with HSV-2. Median CD4 count among HIV-1 infected partners was 462 cells/mm³ (IQR 347-631) and median plasma HIV-1 RNA was 4.1 log₁₀ copies/mL (IQR 3.3-4.7).

Risk factors for HSV-2 infection among HIV-1 uninfected partners

Among male HIV-1 uninfected partners of HIV-1/HSV-2 dually infected females, older age (adjusted prevalence ratio [aPR] 1.11, $p < 0.001$), a greater number of children (aPR 1.02 per 1 child increase, $p < 0.001$) and greater number of years living together (aPR 1.07, per 1 year increase, $p = 0.009$) were significantly associated with HSV-2 infection (Table 1). Characteristics of the female HIV-1 infected partner associated with HSV-2 infection in their male HIV-1 uninfected partners included having herpetic lesions observed on genital exam at enrollment (aPR 0.92, $p = 0.04$), other sexual partners (aPR 1.28, $p = 0.05$), and using vaginal drying agents (aPR 1.08, $p = 0.05$). Circumcised HIV-1 uninfected men were at slightly decreased risk for HSV-2 (aPR 0.94, $p = 0.06$), although this did not achieve statistical significance.

Among female HIV-1 uninfected partners of HIV-1/HSV-2 dually-infected males, only older age (aPR 1.08, $p < 0.0001$) and greater number of years of education (aPR 0.90, $p = 0.001$) of the HIV-1 infected partner were associated with HSV-2 seropositivity.

Notably, neither HIV-1 plasma RNA level [aPR 1.03; 95% CI: 0.99 to 1.06; $p = 0.11$] nor CD4 count [aPR 1.00; 95% CI: 0.98 to 1.01; $p = 0.48$], factors that might indicate increased HSV-2 infectiousness [12], were associated with increased risk of HSV-2 infection in HIV-1 uninfected partners.

Table 1 Population characteristics and risk factors for HSV-2 seropositivity among HIV-1 uninfected members of African HIV-1 serodiscordant partnerships

	HIV-1 uninfected female partners of males dually-infected with HSV-2 and HIV-1 (N = 1084)					HIV-1 uninfected male partners of females dually-infected with HSV-2 and HIV-1 (N = 2270)						
	HSV-2 negative (n = 123) # (%) or median (IQR)	HSV-2 positive (n = 961) # (%) or median (IQR)	Prevalence ratio (95% CI)	p-value	Adjusted Prevalence Ratio [#] (95% CI)	p-value	HSV-2 negative (n = 841) # (%) or median (IQR)	HSV-2 positive (n = 1429) # (%) or median (IQR)	Prevalence ratio (95% CI)	p-value	Adjusted Prevalence ratio [#] (95% CI)	p-value
Characteristics of HIV-1/HSV-2 dually-infected partner												
Genital lesions at enrollment	27 (22%)	233 (24.2%)	1.01 (0.97, 1.07)	0.6			202 (24%)	287 (20.1%)	0.92 (0.85, 0.99)	0.028	0.92 (0.85, 0.99)	0.04
CD4 count	424 (335-566)	424 (333-571)	1.00 (0.99, 1.01)*	0.66			484 (358-669)	482 (354-659)	1.00 (0.98, 1.01)*	0.48		
Plasma HIV-1 RNA	4.35 (3.71-4.95)	4.32 (3.67-4.87)	0.99 (0.97, 1.02)	0.64			3.92 (3.13-4.50)	3.96 (3.25-4.55)	1.03 (0.99, 1.06)	0.11		
Has any other sexual partners	9 (7.3%)	78 (8.1%)	1.01 (0.94, 1.09)	0.76			7 (0.8%)	26 (1.8%)	1.26 (0.99, 1.59)	0.058	1.28 (1.00, 1.63)	0.047
Characteristics of HIV-1 uninfected partner												
Age	26 (22-30)	31 (26-39)	1.09 (1.06, 1.12) [†]	<.001	1.08 (1.06, 1.11) [†]	<.001	32 (28-39)	37 (31-44)	1.17 (1.14, 1.21) [†]	<.001	1.11 (1.06, 1.15)	<.001
Years of education	8 (7-12)	8 (6-10)	0.88 (0.83, 0.93) [†]	<.0001	0.90 (0.85, 0.96) [†]	0.0011	9 (7-12)	9 (7-12)	0.89 (0.82, 0.97) [†]	0.01	!	
Has any other sexual partners	1 (1.5%)	4 (0.8%)	0.91 (0.65, 1.28)	0.59			31 (7.9%)	67 (10.0%)	1.09 (0.94, 1.27)	0.27		
Gender-specific characteristics												
Male partner is circumcised (physical exam)	37 (30.1%)	329 (34.3%)	1.02 (0.98, 1.07)	0.36			483 (57.4%)	760 (53.2%)	0.94 (0.88, 1.00)	0.049	0.94 (0.88, 1.00)	0.0614
Female partner uses hormonal contraceptives	18 (14.6%)	157 (16.3%)	1.01 (0.96, 1.07)	0.63			160 (19.0%)	279 (19.5%)	1.01 (0.93, 1.10)	0.77		
Female uses vaginal drying	24 (19.5%)	196 (20.4%)	1.01 (0.95, 1.06)	0.82			156 (18.5%)	322 (22.5%)	1.09 (1.01, 1.18)	0.025	1.08 (1.00, 1.17)	0.051
Couple characteristics[‡]												
Couple is married	94 (76.4%)	776 (80.7%)	1.03 (0.98, 1.09)	0.26			589 (70.0%)	1078 (75.4%)	1.11 (1.03, 1.20)	0.005		
Duration of sexual relationship	4 (2-8)	7 (3-15)	1.06 (1.04, 1.09) [†]	<.001			4 (2-7)	6 (3-10)	1.18 (1.13, 1.24) [†]	<.001		

Table 1 Population characteristics and risk factors for HSV-2 seropositivity among HIV-1 uninfected members of African HIV-1 serodiscordant partnerships (Continued)

Years living together	3 (1-6)	6 (3-13)	1.06 (1.03, 1.09) [†]	<.001	3 (1-6)	5 (2-10)	1.20 (1.15, 1.26) [†]	<.001	1.07 (1.02, 1.13)	0.009
Number of children	1 (1-3)	3 (1-4)	1.03 (1.02, 1.04)	<.001	2 (1-3)	2 (1-4)	1.05 (1.04, 1.07)	<.001	1.02 (1.01, 1.04)	0.001
Informal housing	50 (41.3%)	348 (37.5%)	0.98 (0.94, 1.03)	0.41	346 (42.5%)	540 (38.9%)	0.95 (0.89, 1.01)	0.099		
Number of sex acts with study part-ner in previous month (median, IQR)	3 (2-7)	4 (2-8)	1.02 (0.99, 1.05) [†]	0.26	4 (2-8)	4 (2-8)	0.96 (0.92, 1.01) [†]	0.13		
Any sex acts with partner in previous month	114 (92.7%)	885 (92.1%)	0.99 (0.92, 1.07)	0.82	796 (94.6%)	1342 (93.9%)	0.95 (0.83, 1.09)	0.47		
Any unprotected sex with study partner in previous month	32 (26.0%)	259 (27.0%)	1.01 (0.96, 1.05)	0.83	245 (29.1%)	451 (31.6%)	1.04 (0.97, 1.12)	0.23		
Enrollment site in Eastern (vs. southern) Africa	78 (63.4%)	593 (61.7%)	0.99 (0.95, 1.04)	0.71	509 (60.5%)	874 (61.2%)	1.01 (0.95, 1.08)	0.76		

[#] Multivariate risk ratios, adjusted for other factors in the model as shown.

* Risk ratio of HSV2 positivity for every 100 ct increase in CD4.

[†] Risk ratio of HSV2 positivity for every increase of 10.

[‡] As reported by HIV-1 uninfected participant.

(Note: Length of sexual relationship strongly associated with age of partner, rho = 0.66, p < .0001.)

! Years of education was not a significant risk factor for HSV-2 seropositivity in male HIV-1 uninfected partners

Discussion

In this multi-national study of 3354 African HIV-1 serodiscordant couples in which one partner was dually-infected with HIV-1 and HSV-2, HIV-1 plasma RNA and CD4 count of the HIV-1 infected partner were not significantly associated with HSV-2 infection concordance in the HIV-1 uninfected partner. Furthermore, despite a recent prospective clinical trial reporting association of male circumcision with reduced HSV-2 acquisition [13], male circumcision in male HIV-1 uninfected partner was only marginally protective against HSV-2 infection in our cross-sectional multivariate analysis.

Prevalence of HSV-2 in the HIV-1 uninfected partner was high in our study (63% in men and 89% in women) compared to other African cohorts [14,15], possibly due to the high rate of HSV-2 transmission reported in stable, long-term partnerships [16,17]. Also consistent with other studies [18-20], HSV-2 infection at enrollment in the HIV-1 uninfected partner was significantly associated with older age and was higher among women compared to men. The former may indicate that age is a surrogate for duration of sexual exposure in these couples [21].

A limitation of this cross-sectional study was the inability to determine temporal relationships of HIV-1 and HSV-2 infection in the dually-infected partner or recency of HSV-2 infection in the HIV-uninfected partners, thus limiting our analysis of behavioral correlates with HSV-2 infection. HSV-2 concordance may reflect HSV-2 transmission within the couple or acquisition outside the partnership. Our study eligibility criteria included that couples be sexually active and intend to remain together; as such we may have selected for older, more stable couples.

Conclusions

In summary, among African HIV-1 serodiscordant couples in which all the HIV-infected partners were dually infected with HSV-2, 71% of their HIV-1 uninfected partners had HSV-2 infection, and HSV-2 infection in the HIV-uninfected partner was not associated with HIV-1 plasma RNA and CD4 count in the HIV-infected partner, or male circumcision in the HIV-1 uninfected partner. Given the need to identify interventions to reduce HSV-2 transmission, future research should focus on prospective analysis of risk factors for HSV-2 transmission, which will be conducted in this cohort.

Acknowledgements

We are grateful to the study participants for their participation and dedication. We thank the study team members at the research sites and at the University of Washington for their contributions to study implementation and data collection. This study was supported through a research grant from the Bill & Melinda Gates Foundation (grant ID # 26469)

Partners in Prevention HSV/HIV Transmission Study Team:

University of Washington Coordinating Center and Central Laboratories,

Seattle, USA: Connie Celum (principal investigator), Anna Wald (protocol co-chair), Jairam Lingappa (medical director), Jared M. Baeten, Mary Campbell, Lawrence Corey, Robert W. Coombs, James P. Hughes, Amalia Magaret, M. Juliana McElrath, Rhoda Morrow, James I. Mullins

Study sites and site principal investigators:

Cape Town, South Africa (University of Cape Town): David Coetzee; Eldoret, Kenya (Moi University, Indiana University): Kenneth Fife, Edwin Were; Gaborone, Botswana (Botswana Harvard Partnership): Max Essex, Joseph Makhema; Kampala, Uganda (Infectious Disease Institute, Makerere University): Elly Katabira, Allan Ronald; Kigali, Rwanda (Rwanda Zambia HIV Research Group, and Emory University): Susan Allen, Kayitesi Kayitenkore, Etienne Karita; Kisumu, Kenya (Kenya Medical Research Institute, University of California San Francisco): Elizabeth Bukusi, Craig Cohen; Kitwe, Zambia (Rwanda Zambia HIV Research Group, and Emory University): Susan Allen, William Kanweka; Lusaka, Zambia (Rwanda Zambia HIV Research Group, and Emory University): Susan Allen, Bellington Vwalika; Moshi, Tanzania (Kilimanjaro Christian Medical College, Harvard University): Saidi Kapiga, Rachel Manongi; Nairobi, Kenya (University of Nairobi, University of Washington): Carey Farquhar, Grace John-Stewart, James Kiarie; Ndola, Zambia (Rwanda Zambia HIV Research Group, and Emory University): Susan Allen, Mubiana Inambao; Orange Farm, South Africa (Reproductive Health Research Unit, University of the Witwatersrand): Sinead Delany-Moretlwe, Helen Rees; Soweto, South Africa (Perinatal HIV Research Unit, University of the Witwatersrand): Guy de Bruyn, Glenda Gray, James McIntyre; Thika, Kenya (University of Nairobi, University of Washington): Nelly Rwamba Mugo
Data management was provided by DF/Net Research, Inc. (Seattle, USA) and site laboratory oversight was provided by Contract Lab Services (University of the Witwatersrand, Johannesburg, South Africa).

Author details

¹Department of Global Health, University of Washington, Seattle, WA, USA.

²Department of Laboratory Medicine, University of Washington, Seattle, WA, USA. ³Department of Medicine, University of Washington, Seattle, WA, USA.

⁴Department of Epidemiology, University of Washington, Seattle, WA, USA.

⁵Department of Pediatrics, University of Washington, Seattle, WA, USA.

Authors' contributions

AM wrote the first draft along with JL and JMB. ASM analyzed the data. All authors contributed to the gathering of data, interpretation of results, and writing of the manuscript, and all approved the final draft.

Competing interests

The authors declare that they have no competing interests.

Received: 1 March 2011 Accepted: 15 March 2011

Published: 15 March 2011

References

1. Looker KJ, Garnett GP, Schmid GP: An estimate of the global prevalence and incidence of herpes simplex virus type 2 infection. *Bull World Health Organ* 2008, **86**(10):805-12, A.
2. Corey L, Wald A, Celum CL, et al: The Effects of Herpes Simplex Virus-2 on HIV-1 Acquisition and Transmission: A Review of Two Overlapping Epidemics. *J Acquir Immune Defic Syndr* 2004, **35**(5):435-445.
3. Robinson NJ: Age-specific prevalence of infection with herpes simplex virus types 2 and 1: a global review. *J Infect Dis* 2002, **186**(Suppl 1):S3-28.
4. Freeman EE, Weiss HA, Glynn JR, et al: Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *AIDS* 2006, **20**(1):73-83.
5. Van de Perre P, Segondy M, Foulongne V, et al: Herpes simplex virus and HIV-1: deciphering viral synergy. *Lancet Infect Dis* 2008, **8**(8):490-7.
6. Dunkle KL, Stephenson R, Karita E, et al: New heterosexually transmitted HIV infections in married or cohabiting couples in urban Zambia and Rwanda: an analysis of survey and clinical data. *Lancet* 2008, **371**(9631):2183-91.
7. Lingappa JR, Lambdin B, Bukusi EA, et al: Regional Differences in Prevalence of HIV-1 Discordance in Africa and Enrollment of HIV-1 Discordant Couples into an HIV-1 Prevention Trial. *PLoS ONE* 2008, **3**(1): e1411.

8. Laeyendecker O, Henson C, Gray RH, *et al*: Performance of a commercial, type-specific enzyme-linked immunosorbent assay for detection of herpes simplex virus type 2-specific antibodies in Ugandans. *J Clin Microbiol* 2004, **42**(4):1794-6.
9. Gamiel JL, Tobian AA, Laeyendecker OB, *et al*: Improved performance of enzyme-linked immunosorbent assays and the effect of human immunodeficiency virus coinfection on the serologic detection of herpes simplex virus type 2 in Rakai, Uganda. *Clin Vaccine Immunol* 2008, **15**(5):888-90.
10. Golden MR, Ashley-Morrow R, Swenson P, *et al*: Herpes simplex virus type 2 (HSV-2) Western blot confirmatory testing among men testing positive for HSV-2 using the focus enzyme-linked immunosorbent assay in a sexually transmitted disease clinic. *Sex Transm Dis* 2005, **32**(12):771-7.
11. Celum C, Wald A, Lingappa JR, *et al*: Acyclovir and Transmission of HIV-1 from Persons Infected with HIV-1 and HSV-2. *N Engl J Med* 2010, **362**(5):427-439.
12. Schacker TW, Zeh JE, Hu H, *et al*: Changes in plasma human immunodeficiency virus type 1 RNA associated with herpes simplex virus reactivation and suppression. *J Infect Dis* 2002, **186**(12):1718-1725.
13. Tobian AA, Serwadda D, Quinn TC, *et al*: Male circumcision for the prevention of HSV-2 and HPV infections and syphilis. *N Engl J Med* 2009, **360**(13):1298-309.
14. Tobian AA, Charvat B, Ssempijja V, *et al*: Factors Associated with the Prevalence and Incidence of Herpes Simplex Virus Type 2 Infection among Men in Rakai, Uganda. *J Infect Dis* 2009.
15. Freeman EE, Glynn JR: Factors affecting HIV concordancy in married couples in four African cities. *AIDS* 2004, **18**(12):1715-21.
16. Halton K, Ratcliffe AA, Morison L, *et al*: Herpes simplex 2 risk among women in a polygynous setting in rural West Africa. *AIDS* 2003, **17**(1):97-103.
17. Allen S, Meinzen-Derr J, Kautzman M, *et al*: Sexual behavior of HIV discordant couples after HIV counseling and testing. *AIDS* 2003, **17**(5):733-40.
18. Kebede Y, Dorigo-Zetsma W, Mengistu Y, *et al*: Transmission of Herpes Simplex Virus Type 2 among factory workers in Ethiopia. *J Infect Dis* 2004, **190**(2):365-72.
19. Mertz GJ, Benedetti J, Ashley R, *et al*: Risk factors for the sexual transmission of genital herpes. *Ann Intern Med* 1992, **116**:197-202.
20. Weiss H: Epidemiology of herpes simplex virus type 2 infection in the developing world. *Herpes* 2004, **11**(Suppl 1):24A-35A.
21. Wald A: Herpes simplex virus type 2 transmission: risk factors and virus shedding. *Herpes* 2004, **11**(Suppl 3):130A-137A.

doi:10.1186/1756-0500-4-64

Cite this article as: Mujugira *et al.*: Risk Factors for HSV-2 Infection among Sexual Partners of HSV-2/HIV-1 Co-Infected Persons. *BMC Research Notes* 2011 **4**:64.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

