


 Cite this: *RSC Adv.*, 2021, 11, 14862

Tuning the sensitivity towards mercury via cooperative binding to D-fructose: dual fluorescent chemosensor based on 1,8-naphthyridine-boronic acid derivative†

 Marina Rajadurai * and E. Ramanjaneya Reddy

A novel fluorescent chemosensor naphthyridine-boronic acid derivative (**1.1**) was synthesized and its ability to act as a selective chemosensor was examined for various metal ions. Compound **1.1** displayed highly selective fluorescence quenching upon interaction with Hg^{2+} , possibly by means of photo induced electron transfer (PET) mechanism. The binding stoichiometry of the naphthyridine-boronic acid- Hg^{2+} complex and the association constant was determined. It was found that in the presence of D-fructose at physiological concentration, the sensitivity of chemosensor **1.1** towards Hg^{2+} improved by at least 7 times, perhaps as a result of the cooperative binding of both D-fructose and mercury ion to the sensor. Till now, the presented dual D-fructose-mercury chemosensor is the first example of utilizing boronic acid-diol complexation for enhancement of the sensor's sensitivity towards a toxic metal ion. The utility of compound **1.1** lays in applications in the food industry, e.g. for detection of mercury contamination of high fructose corn syrup, or in estimation of mercury in polluted biological samples and underground water.

Received 17th March 2021

Accepted 6th April 2021

DOI: 10.1039/d1ra02122b

rsc.li/rsc-advances

1. Introduction

Mercury and mercuric salts are extensively circulated in atmospheric air, water and soil, and considered to be highly toxic and hazardous to humans and the environment.¹ For humans, mercury contamination may cause a wide variety of symptoms, including neuro-disorders, neuromuscular changes, memory loss and carcinogenic diseases.² Despite the danger mercury possesses, especially to unborn children, it is still used in industry. Its contaminants are often detected in various products, including food industry products, for example in high fructose corn syrup, which is cheaper and sweeter than regular sugar. Traces of mercury were found in high fructose corn syrup containing beverages/confectioneries, and the alarming amounts of up to 0.570 micrograms of mercury per gram of high fructose corn syrup were determined.³ Various methods have been employed for the detection of Hg^{2+} , including cold vapor atomic absorption spectroscopy,⁴ high performance liquid chromatography⁵ and inductively coupled plasma atomic emission spectrometry.⁶ The above methods are extremely sensitive and able to detect Hg^{2+} in the nanomolar range. However, their drawbacks are: costly equipment, time-consuming and laborious procedures, and need for the

trained professionals in order to perform the analysis. At the same time, mercury detection based on fluorescence changes allows rapid, convenient and inexpensive detection. Fluorescence chemosensors offer definite advantages, such as high selectivity, high sensitivity, accuracy and possibility to investigate molecule-molecule recognition in both environmental and biological samples.⁷ Till date, numerous fluorescent chemosensors for Hg^{2+} are developed based on rhodamine and BOD-IPY,⁸ various nanoparticles, for example silver,⁹ modified naphthalimide derivatives and other sensors.¹⁰ However, these sensors often suffer from such drawbacks, as laborious, complex or expensive synthesis, low aqueous solubility or limited selectivity as result of interference with other metal ions.

1,8-Naphthyridine and its derivatives are extensively studied for molecular recognitions events, including nucleosides sensing (e.g. for guanine, cytosine and thymidine),¹¹ mono-saccharide sensing¹² and heavy transition metal ions sensing¹³ (e.g. for Zn^{2+} , Hg^{2+} , Cd^{2+}). In order to improve fluorescent properties, affinity, aqueous solubility and stability of 1,8-naphthyridine based chemosensors, we report design and synthesis of novel fluorescent chemosensor **1.1** [(2-(((5-(7-acetamido-1,8-naphthyridin-2-yl)thiophen-2-yl)methyl)(methyl)amino)methyl)phenyl)boronic acid]. In order to strengthen an interaction of mercury ion with the chemosensor, 1,8-naphthyridine was conjugated to thiophene moiety, keeping in mind high affinity of mercury to sulfur. At the same time, boronic acid group, which is known to easily form cyclic ester with

Center for Innovation in Molecular & Pharmaceuticals Sciences (CIMPS), Dr Reddy's Institute of Life Sciences, University of Hyderabad Campus, Gachibowli, Hyderabad 500046, Telangana, India. E-mail: marinasraj@gmail.com

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra02122b



monosaccharides, was additionally introduced in the same scaffold aiming cooperative action of boronic acid and metal chelate, which could possibly enhance sensitivity and selectivity either to metal ion or sugar. There are very few examples are known for using boronic acid containing compounds for direct detection of metal ions.¹⁴

Thus, we report the synthesis of naphthyridine-boronic acid based fluorescent chemosensor **1.1**, its fluorescence properties and investigation of interaction with various metal ions and monosaccharides (*D*-glucose, *D*-fructose, *D*-mannose and *D*-galactose). The ability of chemosensor **1.1** to form reversible covalent bonds with *D*-fructose was employed for the first time to significantly enhance selectivity towards metal ion, namely Hg²⁺.

2. Materials and methods

2.1. General

The materials 2-thiophene carboxaldehyde, 2,6-diaminopyridine, *n*-butyl lithium, tributyltin chloride, bis(triphenylphosphine) palladium(II) dichloride, acetic anhydride, methyl amine (40% aqueous solution), 2-thiophenemethylamine, potassium bis(trimethylsilyl)amide, 2-(bromomethyl)phenylboronic acid and paraformaldehyde were purchased from Sigma Aldrich. *N*-Bromosuccinimide, phosphorus trichloride oxide, azobisisobutyronitrile, di-*tert*-butyl dicarbonate, trifluoroacetic acid, methyl iodide were purchased from Spectro Chem. Commercial reagents (*DL*-malic acid, ethane-1,2-diol, sodium borohydride, ammonium bicarbonate, mercury(I) chloride and HCl) were purchased from Merck and Rankem and were used as received. The solvents THF, DMF, toluene, 1,4-dioxane and DCM were distilled and dried before reactions and for extracting purposes. All reactions were carried out under an inert atmosphere with dry solvents, unless otherwise stated. Syringes and needles for the transfer of reagents were dried at 100 °C and allowed to cool in a desiccator over P₂O₅ before use. Reactions were monitored by thin layer chromatography (TLC) on silica gel plates (60 F254), using UV light detection. Merck silica gel (particle size 100–200 mesh) was used for column chromatography. For UV-vis and fluorescence measurements spectroscopic grade solvents were used.

Buffers and samples preparations for fluorescence studies and synthesis of the compounds of routes 2–4 are given in ESI.†

2.2. Synthesis of target compound **1.1** using route I (Schemes 1 and 2)

2.2.1. *tert*-Butyl(thiophen-2-ylmethyl)carbamate (2.1).¹⁵ Thiophene-2-methyl amine (0.5 g, 4.42 mmol) was dissolved in dry DCM (20 mL), and triethyl amine (0.95 mL, 6.84 mmol) was added to the solution. Boc-anhydride (1.22 mL, 5.31 mmol) was added portion wise to the above reaction mixture. After complete addition of Boc-anhydride the reaction mixture was stirred at room temperature for 4 h. After 4 h, the reaction mixture was washed with water and extracted with DCM. The organic phase was dried with anhydrous sodium sulphate, concentrated under reduced pressure, and the resulting residue **2.1** (0.8 g, 86%) was used without further purification.

Spectroscopic data matches with the one from reported procedure. ¹H NMR (CDCl₃, 400 MHz) δ: 7.22–7.20 (1H, m), 6.94–6.92 (2H, m), 4.87 (1H, brs), 4.47 (2H, d, *J* = 5.2 Hz), 1.46 (9H, s).

2.2.2. *tert*-Butyl((5-(tributylstannyl)thiophen-2-yl)methyl)carbamate (2.2).¹⁵ To a solution of compound **2.1** (0.4 g, 2.62 mmol) in dry THF (30 mL) was added *n*-butyl lithium (1.6 M in hexane, 3.58 mL, 5.73 mmol) at –78 °C. After 1 h at –78 °C, tributyl tin chloride (1.57 mL, 5.81 mmol) was added drop wise to the above reaction mixture. After stirring over 6 h at room temperature, the reaction mixture was quenched with 20 mL of saturated aqueous ammonium chloride solution and extracted with THF. The organic phase was dried with anhydrous sodium sulphate, concentrated under reduced pressure, and the resulting residue **2.2** (0.87 g, 93%) was used without further purification. Spectroscopic data matches with the one from reported procedure. ¹H NMR (CDCl₃, 400 MHz) δ: 7.05 (1H, d, *J* = 3.2 Hz), 6.99 (1H, d, *J* = 2.8 Hz), 4.87 (1H, brs), 4.51 (2H, s), 1.59–1.51 (6H, m), 1.46 (9H, s), 1.37–1.28 (6H, m), 1.07–1.04 (6H, m), 0.91–0.87 (9H, m).

2.2.3. *tert*-Butylmethyl((5-(tributylstannyl)thiophen-2-yl)methyl)carbamate (1.3). To a solution of compound **2.2** (5.54 g, 11.0 mmol) in 30 mL of dry THF was added 1 M KHMDS (22 mL, 22 mmol) dropwise at –10 °C. The reaction was maintained at –10 °C for 1.5 h, and methyl iodide (2.05 mL, 33.0 mmol) was added drop wise. The ice bath was removed and reaction mixture was stirred at room temperature for 4 h. The reaction mixture was washed with brine (4 × 100 mL) and extracted with hexane. The organic layer was collected, dried over anhydrous sodium sulphate, concentrated under reduced pressure to provide compound **1.3** (5.45 g, 96% yield) as a dark brown oil, which was used without further purification. TLC (10% EtOAc in hexane): *R*_f = 0.39; ¹H NMR (CDCl₃, 400 MHz) δ: 7.03 (1H, d, *J* = 3.2 Hz), 7.01 (1H, d, *J* = 3.2 Hz), 4.56 (2H, s), 2.86 (3H, s), 1.67–1.62 (2H, m), 1.59–1.54 (4H, m), 1.49 (9H, s), 1.39–1.31 (6H, m), 1.10–1.06 (6H, m), 0.98–0.93 (9H, m). ¹³C NMR (CDCl₃, 100 MHz) δ: 155.3, 146.1, 136.7, 134.8, 127.1, 79.9, 47.3, 33.7, 28.9, 28.4, 27.2, 13.6, 10.7. MS (EI, 70 eV): *m/z* [M]⁺ calcd for C₂₃H₄₃NO₂SSn: 517.2, found 517.2 [M]⁺. Anal. calcd for C₂₃H₄₃NO₂SSn: C, 53.50; H, 8.39; N, 2.71; O, 6.20; S, 6.21; Sn, 22.99, found: C, 53.46; H, 8.36; N, 2.75.

2.2.4. 7-Amino-1,8-naphthyridin-2-ol (2.4).¹⁶ 2,6-Diaminopyridine **2.3** (1.1 g, 10.07 mmol) and malic acid (1.48 g, 11.07 mmol) were placed in a three necked round bottom flask outfitted with an additional funnel, reflux condenser and mechanical stirrer. The mixture was cooled to 0 °C and conc. H₂SO₄ (5 mL) was added dropwise. After addition of conc. H₂SO₄ was completed, the reaction mixture was slowly heated to 110 °C and stirred for 3 h. The reaction mixture was cooled to 0 °C and the solution was made alkaline (pH = 8) by careful addition of aqueous ammonium hydroxide. The crude solid was collected by vacuum filtration and washed thoroughly with water and methanol to give compound **2.4** as a yellowish solid (1.38 g, 86%). Spectroscopic data matches with the one from reported procedure. ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 11.91 (1H, s), 7.65 (2H, d, *J* = 9.2 Hz), 7.03 (2H, s), 6.34 (1H, d, *J* = 8.4 Hz), 6.12 (1H, d, *J* = 9.2 Hz).

2.2.5. *N*-(7-Hydroxy-1,8-naphthyridin-2-yl)acetamide (2.5).¹⁷ A suspension of compound **2.4** (5.0 g, 15.5 mmol) in 50 mL of acetic anhydride was stirred at 110 °C for 3 h. After 3 h, reaction mixture was cooled to room temperature and the precipitate was collected by vacuum filtration, washed with hexane and dried at reduced pressure to give compound **2.5** as a yellow solid (5.33 g, 89%). Spectroscopic data matches with the one from reported procedure. ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 11.90 (1H, s), 10.50 (1H, s), 8.02 (1H, d, *J* = 8.4 Hz), 7.90 (1H, d, *J* = 8.8 Hz), 7.82 (1H, d, *J* = 9.6 Hz), 6.40 (1H, d, *J* = 9.6 Hz), 2.12 (3H, s).

2.2.6. *N*-(7-Chloro-1,8-naphthyridin-2-yl)acetamide (1.4).¹⁸ A mixture of compound **2.5** (10.0 g, 49.25 mmol) and POCl₃ (175 mL) was heated at 100 °C for 2 h. The reaction mixture was cooled to room temperature, and excess of POCl₃ was removed by distillation. The crude residue was dissolved in ice water, and the solution was made alkaline (pH = 8) by careful addition of concentrated ammonium hydroxide. The crude solid was collected by vacuum filtration, air-dried, and continuously extracted (Soxhlet extraction) with chloroform for 12 h. Chloroform was removed at reduced pressure, and the crude product was purified using column chromatography on silica gel using MeOH/DCM (1 : 9) as an eluent to yield compound **1.4** in form of a golden needles (6.47 g, 60%). ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 8.70 (1H, s), 8.56 (1H, d, *J* = 8.8 Hz), 8.20 (1H, d, *J* = 8.8 Hz), 8.18 (1H, d, *J* = 8.4 Hz), 7.40 (1H, d, *J* = 8.0 Hz), 2.30 (3H, S).

2.2.7. *tert*-Butyl((5-methyl(7-acetamido-1,8-naphthyridin-2-yl)thiophen-2-yl)methyl)carbamate (2.6). To a solution of compound **1.3** (6.2 g, 12.0 mmol) in dry 1,4-dioxane (50 mL) were added compound **1.4** (2.0 g, 9.0 mmol) and PdCl₂(PPh₃)₂ (0.315 g, 0.45 mmol) under nitrogen atmosphere. After 12 h at reflux, the solvent was removed under reduced pressure, and the crude residue was dissolved in EtOAc, passed through celite and solvent was evaporated. Crude product was purified by column chromatography using 50% EtOAc in hexane to give **2.6** as yellowish solid (2.5 g, 67%). Mp: 115–118 °C; TLC (50% EtOAc in hexane): *R*_f = 0.26; ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 11.07 (1H, bs), 8.33 (3H, m), 8.01 (1H, d, *J* = 8.8 Hz), 7.87 (1H, d, *J* = 4.4 Hz), 7.08 (1H, d, *J* = 3.6 Hz), 4.53 (2H, s), 2.80 (3H, s), 2.14 (3H, s), 1.43 (9H, s). ¹³C NMR (CDCl₃, 100 MHz) δ: 169.5, 155.6, 154.6, 153.9, 145.7, 143.5, 139.2, 139.0, 137.0, 126.8, 119.3, 116.9, 114.3, 114.0, 80.2, 48.0, 33.8, 28.4, 24.9. MS (EI, 70 eV): *m/z* [M]⁺ calcd for C₂₁H₂₄N₄O₃S: 412.16, found 413.2 [M + 1]⁺. Anal. calcd for C₂₁H₂₄N₄O₃S: C, 61.14; H, 5.86; N, 13.58; O, 11.64; S, 7.77, found: C, 61.26; H, 5.89; N, 13.45.

2.2.8. *N*-(7-(5-((Methylamino)methyl)thiophen-2-yl)-1,8-naphthyridin-2-yl)acetamide (1.2). Compound **2.6** (0.293 g, 0.71 mmol) was dissolved in TFA : DCM (3 mL : 3 mL) and stirred at RT for 2 h. After 2 h solvents were removed and the reaction mixture was treated with sodium bicarbonate solution and extracted with DCM. The organic solvent was removed under reduced pressure and the crude residue was then purified using column chromatography on silica gel using MeOH/DCM (2 : 8) as an eluent to yield compound **1.2** as a yellowish solid (0.189 g, 85%). Mp: 260–262 °C; TLC (20% MeOH in DCM): *R*_f = 0.21. ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 11.08 (1H, brs), 8.35–8.28 (3H, m),


8.01 (1H, d, *J* = 8.4 Hz), 7.86 (1H, d, *J* = 4.4 Hz), 7.08 (1H, d, *J* = 3.6 Hz), 3.93 (2H, s), 2.35 (3H, s), 2.14 (3H, s). ¹³C NMR (DMSO-*d*₆, 100 MHz) δ: 170.5, 155.2, 154.9, 154.8, 143.8, 139.6, 138.0, 127.9, 127.5, 119.2, 116.8, 114.2, 49.7, 35.2, 24.5. MS (EI, 70 eV): *m/z* [M]⁺ calcd for C₁₆H₁₆N₄O₃S: 312.1, found 313.1 [M + 1]⁺. Anal. calcd for C₁₆H₁₆N₄O₃S: C, 61.52; H, 5.16; N, 17.93; O, 5.12; S, 10.26, found: C, 61.42; H, 5.23; N, 17.85.

2.2.9. 2-(((5-(7-Acetamido-1,8-naphthyridin-2-yl)thiophen-2-yl)methyl)(methylamino)methyl)phenyl)boronic acid (1.1). Anhydrous K₂CO₃ (0.176 g, 1.32 mmol) was added to a stirred suspension of compound **1.2** (0.14 g, 0.44 mmol) in dry DMF (2 mL). After 30 min, a solution of 2-(bromomethyl)phenylboronic acid (0.096 g, 0.44 mmol) in DMF (2 mL) was added, and the reaction mixture was stirred at room temperature for 12 h. DMF was removed under reduced pressure, and the crude residue was washed with water, hexane and toluene to yield compound **1.1** as a white solid (0.14 g, 70%). Mp: 210–212 °C. ¹H NMR (DMSO-*d*₆, 400 MHz) δ: 11.12 (1H, s), 9.03 (2H, brs), 8.34–8.27 (3H, m), 8.0 (1H, d, *J* = 8.4 Hz), 7.87 (1H, d, *J* = 3.6 Hz), 7.72–7.69 (1H, m), 7.35–7.25 (3H, m), 7.10 (1H, d, *J* = 3.6 Hz), 3.75 (2H, s), 3.70 (2H, s), 2.14 (3H, s), 2.13 (3H, s). ¹¹B NMR (CD₃OD, 130 MHz) δ: 26.5. ¹³C NMR (CD₃OD, 100 MHz) δ: 170.5, 155.2, 154.9, 154.8, 143.8, 139.6, 138.0, 127.9, 127.5, 119.2, 116.8, 114.2, 49.7, 35.2, 24.5. MS (EI, 70 eV): *m/z* [M]⁺ calcd for C₂₃H₂₃BN₄O₃S: 446.1, found 447.1 [M + 1]⁺. Anal. calcd for C₂₃H₂₃BN₄O₃S: C, 60.38; H, 4.73; N, 18.78; O, 5.36; S, 10.75, found: C, 61.76; H, 5.12; N, 12.65.

3. Results and discussion

3.1. Synthesis of naphthyridine-boronic acid (1.1)

The key intermediate in synthesis of the chemosensor **1.1** is compound **1.2**, which in its turn could be synthesized through four different routes (Scheme 1). We selected the route I which allowed to successfully synthesizing key intermediate **1.2** and target compound **1.1** as described in Scheme 2. Thiophene methyl amine, which was protected with Boc-anhydride to give compound **2.1** served as the starting material for this synthetic route.¹⁵ Compound **2.1** was subjected to lithiation using *n*-BuLi followed by treatment with tributyl tin chloride resulting in compound **2.2** in excellent yield.¹⁵ Further, compound **2.2** was methylated using methyl iodide and potassium hexamethyldisilazane as a base, to give compound **1.3** in 96% yield. In order to build naphthyridine moiety, 2,6-diamino pyridine (**2.3**), was cyclised using DL-malic acid in the presence of conc. H₂SO₄, according to standard procedure,¹⁶ to give 7-amino-1,8-naphthyridin-2-ol (**2.4**) in 86% yield. Amino group of 7-amino-1,8-naphthyridin-2-ol was protected using acetic anhydride to obtain acetamide protected compound **2.5** in 89% yield.¹⁷ This step allowed subsequent selective conversion of hydroxyl group to the halogen *via* chlorination with phosphorus oxychloride¹⁸ to obtain **1.4** in 60% yield. In the next step, coupling of compound **1.3** with **1.4** in Stille coupling conditions in 1,4-dioxane using palladium(II) catalyst resulted in compound **2.6** in 67% yield. In order to introduce phenyl-boronic acid moiety in this scaffold, Boc-protection of the compound **2.6** was removed using trifluoroacetic acid to give key intermediate **1.2**



Scheme 1 Plausible retrosynthesis of key intermediate **1.2** and target molecule **1.1**.

in 82% yield, followed by *N*-alkylation with commercially available 2-bromomethylphenylboronic acid, using potassium carbonate as a base. Final compound was obtained in 70% yield and its structure was confirmed by different methods, including NMR spectroscopy and mass-spectrometry. Particularly, ^1H NMR spectrum showed an appearance of characteristic $\text{N}-\text{CH}_2$ peak at 3.72 ppm and $\text{B}-(\text{OH})_2$ peak at 9.03 ppm, additionally EI-MS spectrum showed predominant peak with m/z value of 447.1 corresponding to target molecule (ESI Fig. S1 and S2 †).

Apart from the successful route described above, other attempts to synthesize target compound **1.1** were made as sketched in Scheme 1 (see experimental procedures and discussion in ESI, Schemes S1–S3 †), including:


(a) Imine formation followed by reduction using intermediate **1.5**, as presented in route II.

(b) Conversion of bromine to amine, *via* azide formation followed by *N*-alkylation using intermediate **1.6**, as presented in route III.

(c) *N*-Alkylation using intermediate **1.7**, as presented in route IV.

3.2. Photophysical properties of compound **1.1**

To explore utility of the final fluorophore **1.1**, we investigated the photophysical properties of **1.1** in MeOH and MeOH/H₂O solutions. In both solutions, MeOH and MeOH/H₂O



Scheme 2 Route I towards key intermediate **1.2** and target compound **1.1**.




Fig. 1 UV-vis absorption spectra of compound **1.1** in MeOH and MeOH/H₂O, and its emission spectra in MeOH and MeOH/H₂O, $\lambda_{\text{ex}} = 356$ nm (insert).

fluorophore **1.1** displayed identical absorption maxima at 356 nm and a shoulder at 371 nm (Fig. 1 and Table S1 in ESI[†]). Fluorescent emission spectra ($\lambda_{\text{ex}} = 356$ nm) in both solvents exhibited similar fluorescence bands, with a maximum at 401 nm for **1.1** in MeOH and 403 nm in MeOH/H₂O. The solution state quantum yield of **1.1** in MeOH/H₂O ($\Phi_f = 0.26$) was slightly higher than that in MeOH ($\Phi_f = 0.20$), Stokes shift of the new fluorophore **1.1** was 31 nm.

3.3. Examination of binding ability of **1.1** to metal ions

Fluorescence response of compound **1.1** in presence of various metal ions was investigated. The study was carried out in aqueous

methanolic buffer at pH 7.5, and metal ions such as Ag¹⁺, Zn²⁺, Cu¹⁺, Mg²⁺, Pb²⁺, Cd²⁺, Hg²⁺, Li¹⁺ and Fe³⁺ were tested. Among them, addition of Hg²⁺ to **1.1** showed a significant decrease of fluorescence intensity by 75% (Fig. 2A). This can be explained by the fact that after coordination to metal ion fluorophore becomes electron deficient, and as a result the PET from proximal tertiary nitrogen to adjacent chromophore increases and the fluorescence intensity of the fluorophore decreases. Apart from that, there could be possible contribution of strong spin-orbit coupling associated with Hg²⁺ to the luminescence quenching of **1.1**. Fluorescence properties of compound **1.1** (10^{-5} M) were also studied upon increasing the concentration of Hg²⁺ (1–10 eq.), and it was observed that as the concentrations of Hg²⁺ increases, the fluorescent intensity of **1.1** gradually decreases as shown in Fig. 2B and C. In order to estimate the binding stoichiometry of naphthyridine-boronic acid–Hg²⁺ complex, Job plot experiments were employed.¹⁹ In Job method, equal concentrations of Hg²⁺ and **1.1** were prepared, and mixed in different proportions maintaining a total concentration of 10 μM . Using the obtained data, a graph was plotted between the emission intensity (fixed at 408 nm) and a mole fraction of compound **1.1** (Fig. 2D). The minimum emission intensity was reached when the mole fraction was 0.5. This result indicates the formation of naphthyridine-boronic acid–Hg²⁺ complex in a 1:1 ratio. The association constant (K_a) of naphthyridine-boronic acid–Hg²⁺ was calculated from the variation of fluorescence intensity as a function of the concentration of Hg²⁺ by using Benesi–Hildebrand equation²⁰ (eqn (1)) and fitting the emission wavelength at 408 nm. The calculated K_a for a complex of **1.1** with Hg²⁺ was $2.842 \times 10^5 \text{ M}^{-1}$.




Fig. 2 (A) Fluorescence spectra of compound **1.1** in MeOH/H₂O in presence of various metal ions; (B) fluorescence spectra of compound **1.1** in MeOH/H₂O in presence of Hg²⁺ at various concentrations (0–10 eq.); (C) Benesi–Hildebrand plot of the naphthyridine-boronic acid–Hg²⁺ complex in MeOH/H₂O solutions; and (D) Job plot of naphthyridine-boronic acid–Hg²⁺ complex in MeOH/H₂O.

$$1/(F - F_0) = 1/(K_a(F_{\max} - F_0)[\text{Hg}^{2+}]) + 1/(F_{\max} - F_0), \quad (1)$$

where F is the fluorescence intensity at 408 nm at any given concentration of Hg^{2+} , F_0 is the fluorescence intensity at 408 nm in the absence of Hg^{2+} , F_{\max} is the fluorescence intensity at 408 nm in the presence of Hg^{2+} in solution and K_a is association constant.

In order to understand the structure of naphthyridine-boronic acid- Hg^{2+} complex, proton NMR spectroscopy was employed.²¹ Addition of Hg^{2+} to the DMSO solution of **1.1** resulted in a downfield chemical shift of the H_a proton from 7.10 ppm to 7.55 ppm, additionally a moderate down field chemical shifts observed for H_e and $\text{H}_{h,i,j}$ (Fig. 3). These changes indicate that Hg^{2+} interacts mainly with the sulphur atom, which was expected as high affinity of mercury for sulphur atoms is well known, and partly binds to the N_1 and N_8 of the naphthyridine moiety; at the same time there are no interaction with N -methyl nitrogen (no shift observed in NMR spectrum, not shown in Fig. 3). The decrease of fluorescence intensity or down field chemical shifts of H_a , H_e and $\text{H}_{h,i,j}$ upon addition of Hg^{2+} are due to decrease of electron density after coordination of Hg^{2+} to the respective binding site as shown in Fig. 3. Therefore, these both observations support the theory of PET mechanism for presented here chemosensor.

3.4. Examination of binding ability of **1.1** to monosaccharides

Fluorescence titration studies of compound **1.1** (10^{-5} M) in presence of various monosaccharides was carried out in aqueous methanolic phosphate buffer at pH 8.20. As expected, the fluorescence intensity of compound **1.1** increased upon increasing of saccharides concentration (D -glucose, D -mannose, D -fructose and D -galactose). As reported by Shinkai *et al.*²² such fluorescence response explained by the fact that boronic acids (such as compound **1.1**) form cyclic esters with *cis* 1,2-diol or 1,3-diol of mono saccharides, which leads to increase of the

Lewis acidity of the central boron atom. As result B-N (proximal tertiary nitrogen) interactions are strengthening²³ and PET from proximal tertiary nitrogen to adjacent fluorophore is suppressed and the fluorescence of the fluorophore is enhancing.^{22,24} The fluorescence enhancement I/I_0 obtained for D -glucose, D -galactose, D -mannose and D -fructose is in the order D -fructose > D -galactose > D -mannose > D -glucose as shown in Fig. 4.

The binding selectivity of **1.1** towards D -fructose is around two and half fold higher compared to other monosaccharides (D -galactose, D -mannose, D -glucose) at lower concentrations (<10 mM). The association constant (K_a) of naphthyridine-boronic acid based fluorescence chemo sensor **1.1** with D -fructose, D -glucose, D -galactose and D -mannose were calculated from the variation of fluorescence intensity (Fig. 4 and S21–S23 in ESI†) as a function of the concentration of D -monosaccharides, using Benesi–Hildebrand equation²⁰ (eqn (2)) and fitting the emission wavelength at 403, 409, 410 and 408 respectively. The highest affinity of **1.1** was observed towards fructose (the association constants (K_a) = 104.91 M^{-1}), while to other sugars it was significantly lower (K_a = 14.42 – 39.46 M^{-1}). The affinity for **1.1** were in the following order: D -fructose > D -galactose > D -mannose > D -glucose, and the calculated association constant are listed in Tables S2–S5 in ESI.†

$$1/(F - F_0) = 1/(K_a(F_{\max} - F_0)[\text{sugar}]) + 1/(F_{\max} - F_0), \quad (2)$$

where F is the fluorescence intensity at 403/409/410/408 nm at any given concentration of D -fructose/ D -galactose/ D -mannose/ D -glucose respectively, F_0 is the fluorescence intensity at 403/408/409/410 nm in the absence of D -saccharides, F_{\max} is the fluorescence intensity at 403/409/410/408 nm in the presence of D -fructose/ D -galactose/ D -mannose/ D -glucose respectively in solution, and K_a is association constant.

Fluorescence response of the boronic acid based fluorescent chemosensors is usually pH dependent. Therefore, fluorescence response of naphthyridine-boronic acid fluorescent




Fig. 3 (A) Proposed coordination complex structure of Hg^{2+} ions with **1.1**, and (B) ^1H NMR spectra of compound **1.1** along and in the presence of Hg salt (1 : 3 eq.) in DMSO-d_6 .




Fig. 4 (A) Fluorescence intensity of compound **1.1** in MeOH/H₂O (*v/v* = 1 : 1, 5 mM PBS, pH 8.20) in the presence of D-glucose, D-galactose, D-mannose and D-fructose; (B) fluorescence spectra of compound **1.1** (10^{-5} M) in presence of D-fructose at various concentrations (0–10 mM); (C) pH vs. fluorescent intensity of **1.1** (10^{-5} M) in presence and absence of D-fructose in MeOH/H₂O (*v/v* = 1 : 1, 5 mM), at various pH solutions and (D) Benesi–Hildebrand plot of the naphthyridine-boronic acid–fructose complex in MeOH/H₂O (*v/v* = 1 : 1, 5 mM PBS, pH 8.20) solutions, the emission wavelength was monitored at 403 nm.

chemosensor **1.1** was measured at different pH, both in the presence and absence of D-fructose. As Fig. 4C displays, in both cases the fluorescent response was similar. Fluorescent intensity was increasing upon pH change from 2 to 5.5 due to protonation of tertiary amine, and was decreasing as pH raised from 5.5 to 12 due to deprotonation of tertiary amine and OH[−] adduct formation with boronic acid group.²⁵

3.5. Competitive sensing of Hg²⁺ and D-monosaccharide

As compound **1.1** has incorporated phenyl boronic acid moiety which is expected to bind to mono-saccharides, its fluorescence

response was examined in presence of Hg²⁺ and various D-monosaccharides. It was expected, that cooperative binding of both sugar and metal ion will result in increased sensitivity of the chemosensor to either of analytes, decreasing/eliminating at the same time interference by various factors, which are common for chemosensors acting through “turn-off” mechanism.^{25a} The study was carried out in aqueous methanolic buffer at pH 7.5. Addition of Hg²⁺ to a solution containing both chemosensor **1.1** and D-fructose complex resulted in dramatic decrease of fluorescence intensity by seven folds (Fig. 5A). This fluorescent response was ~2.8 times higher compared to one,




Fig. 5 Fluorescence spectra of **1.1** alone (10^{-5} M), **1.1** with Hg²⁺ and **1.1** with Hg²⁺ and D-monosaccharides in MeOH/H₂O (*v/v* = 1 : 1, 5 mM, at various pH) solutions (A) and comparative bar graph of the emission intensity change for the same solutions (B).

when compound **1.1** was interacting with mercury without addition of D-fructose (Fig. 5B). At the same time, no any significant change was observed in the emission intensity of the probe **1.1** in the presence of other hexose sugars.

Till date, there are no reports, where the sensitivity of the chemosensor for metal ions was enhanced as result of interaction with fructose, although there are few investigations utilizing metal–sensor coordination in order to improve sensitivity towards sugar. For example, Seiji Shinkai *et al.* reported^{25a} new saccharide receptor for uronic acids, based on two-point interactions of boronic acid and coordination with zinc(II), or another work¹⁴ employing cooperative binding of fructose and metal ion performed by group of Tony D. James. In contrary, we obtained an interesting result indicating that the sensitivity of the chemosensor **1.1** towards mercury ions increases significantly in presence of D-fructose moiety. Thus, presented here work is the first example of employing boronic acid–diol complexation to achieve significant enhancement of the sensor's sensitivity towards toxic metal ion.

4. Conclusions

In conclusion, we present the design and synthesis of novel fluorescent chemosensor **1.1** [(2-(((5-(7-acetamido-1,8-naphthyridin-2-yl)thiophen-2-yl)methyl)(methyl)amino)methyl)phenyl)boronic acid], its fluorescence properties and interaction with various metal ions and monosaccharides. Chemosensor **1.1** was designed to combine naphthyridine as a fluorescent metal chelating site and phenyl boronic acid as a saccharide recognition unit in one scaffold. The combination of these properties allowed investigation of the sensing abilities of **1.1** over various metal ions and D-mono-saccharides. Fluorescence change, mediated by PET mechanism, was utilized as a reporting property of the chemosensor. Compound **1.1** found to have dramatic decrease of fluorescence intensity selectively upon binding to Hg²⁺, probably due to decrease of electron density after coordination of Hg²⁺ to the respective binding site. As a result, PET from proximal tertiary nitrogen to adjacent fluorophore enhanced and the fluorescence intensity changed by ~2.5 folds. The binding stoichiometry of the naphthyridine-boronic acid–Hg²⁺ complex was found to be 1 : 1, according to Job plot experiments. The association constant for **1.1** with Hg²⁺ was found to be $2.842 \times 10^5 \text{ M}^{-1}$. Interestingly, the sensitivity of chemosensor **1.1** towards mercury ion greatly improved (7 fold) in the presence of D-fructose at physiological concentration, perhaps as result of the cooperative binding of both D-fructose and mercury ion to the sensor. This work is the first example of utilizing boronic acid–diol complexation to manipulate and enhance the sensor's sensitivity towards toxic metal ion. This finding is especially important for applications in food industry, *e.g.* for detection of mercury contamination of high fructose corn syrup, which often contains dangerous amounts of mercury residues,²⁶ or for estimation of mercury in polluted biological samples and underground water.

Author contributions

E. Ramanjaneya Reddy: investigation, writing – original draft. Marina Rajadurai: conceptualization, supervision, writing – review & editing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

ERR acknowledges CSIR-New Delhi for a Senior Research Fellowship. This research was supported by grants from DST (SR/SI/CS-131/2009), CSIR (01(2411)/10/EMR-II) and DBT (BT/PR32386/MED/32/681/2019), New Delhi to MR.

Notes and references

- 1 P. D. Selid, H. Xu, E. M. Collins, M. S. Face-Collins and J. X. Zhao, *Sensors*, 2009, **9**, 5446–5459.
- 2 F. M. M. Morel, A. M. L. Kraepiel and M. Amyot, *Annu. Rev. Ecol. Syst.*, 1998, **29**, 543–566.
- 3 R. Dufault, B. LeBlanc, R. Schnoll, C. Cornett, L. Schweitzer, D. Wallinga, J. Hightower, L. Patrick and W. Lukiw, *J. Environ. Health*, 2009, **8**, 2–8.
- 4 A. F. Danet, M.-C. Bratu, M.-C. Radulescu and A. Bratu, *Sens. Actuators, B*, 2009, **137**, 12–16.
- 5 H. Hashemi-Moghaddam and M. J. Saber-Tehrani, *J. AOAC Int.*, 2008, **91**, 1453–1458.
- 6 O. T. Butler, J. M. Cook, C. F. Harrington, S. J. Hill, J. Rieuwerts and D. L. Miles, *J. Anal. At. Spectrom.*, 2006, **21**, 217–243.
- 7 (a) W. Lin, L. Long and W. Tan, *Chem. Commun.*, 2010, **46**, 1503–1505; (b) D.-H. Wang, Y. Zhang, Z. Gong, R. Sun, D.-Z. Zhao and C.-L. Sun, *RSC Adv.*, 2015, **5**, 50540–50543.
- 8 (a) J. Hu, Z. Hu, Y. Cui, X. Zhang, H.-W. Gao and K. Uvdal, *Sens. Actuators, B*, 2014, **203**, 452–458; (b) M. Vedamalai, D. Kedaria, R. Vasita, S. Mori and I. Gupta, *Dalton Trans.*, 2016, **45**, 2700–2708; (c) W. Sun, R. Chen, X. Cheng and L. Marin, *New J. Chem.*, 2018, **42**, 19224–19231; (d) A. N. Kursunlu, M. Oguz and M. Yilmaz, *IEEE Sens. J.*, 2019, **19**, 2009–2015.
- 9 S. Balasurya, A. Syed, A. M. Thomas, N. Marraiki, A. M. Elgorban, L. L. Raju, A. Das and S. S. Khan, *Spectrochim. Acta, Part A*, 2020, **228**, 117712–117719.
- 10 (a) P. Mahato, S. Saha, P. Das, H. Agarwalla and A. Das, *RSC Adv.*, 2014, **4**, 36140–36174; (b) J. Liu, K. Vellaisamy, G. Yang, C.-H. Leung and D.-L. Ma, *Sci. Rep.*, 2017, **7**, 3620, DOI: 10.1038/s41598-017-03952-x; (c) M. Saleem, M. Rafiq and M. Hanif, *J. Fluoresc.*, 2017, **27**, 31–58.
- 11 (a) S. Djurdjevic, D. A. Leigh, H. McNab, S. Parsons, G. Teobaldi and F. Zerbetto, *J. Am. Chem. Soc.*, 2007, **129**, 476–477; (b) H. Suda, A. Kobori, J. Zhang, G. Hayashi and K. Nakatani, *Bioorg. Med. Chem.*, 2005, **13**, 4507–4512; (c) S.-H. Lu, S. Selvi and J.-M. Fang, *J. Org. Chem.*, 2007, **72**, 117–122.

- 12 J.-M. Fang, S. Selvi, J.-H. Liao, Z. Slanina, C.-T. Chen and P.-T. Chou, *J. Am. Chem. Soc.*, 2004, **126**, 3559–3566.
- 13 (a) M.-M. Yu, Z.-X. Li, L.-H. Wei, D.-H. Wei and M.-S. Tang, *Org. Lett.*, 2008, **10**, 5115–5118; (b) M. K. Chahal and M. Sankar, *Anal. Methods*, 2015, **7**, 4552–4559; (c) X. Liu, M. Chen, Z. Liu, M. Yu, L. Wei and Z. Li, *Tetrahedron*, 2014, **70**, 658–663.
- 14 M. Li, H. Ge, R. L. Arrowsmith, V. Mirabello, S. W. Botchway, W. Zhu, S. I. Pascu and T. D. James, *Chem. Commun.*, 2014, **50**, 11806–11809.
- 15 C. P. Harvey and J. D. Tovar, *J. Polym. Sci., Part A: Polym. Chem.*, 2011, **49**, 4861–4874.
- 16 C. A. Anderson, P. G. Taylor, M. A. Zeller and S. C. Zimmerman, *J. Org. Chem.*, 2010, **75**, 4848–4851.
- 17 P. S. Corbin, S. C. Zimmerman, P. A. Thiessen, N. A. Hawryluk and T. J. Murray, *J. Am. Chem. Soc.*, 2001, **123**, 10475–10488.
- 18 J.-H. Jeon, K. Tanaka and Y. Chujo, *Org. Biomol. Chem.*, 2014, **12**, 6500–6506.
- 19 J.-T. Yeh, W.-C. Chen, S.-R. Liu and S.-P. Wu, *New J. Chem.*, 2014, **38**, 4434–4439.
- 20 (a) H. A. Benesi and J. H. Hildebrand, *J. Am. Chem. Soc.*, 1949, **71**, 2703–2707; (b) R. B. Singh, S. Mahanta and N. Guchhait, *J. Mol. Struct.*, 2010, **963**, 92–97.
- 21 M. Vedamalai and S.-P. Wu, *Org. Biomol. Chem.*, 2012, **10**, 5410–5416.
- 22 (a) T. D. James, K. R. A. S. Sandanayake and S. Shinkai, *J. Chem. Soc., Chem. Commun.*, 1994, **4**, 477–478; (b) T. D. James, K. R. A. S. Sandanayake and S. Shinkai, *Angew. Chem., Int. Ed.*, 1994, **33**, 2207–2209.
- 23 (a) L. I. Bosch, T. M. Fyles and T. D. James, *Tetrahedron*, 2004, **60**, 11175–11190; (b) L. Zhu, S. H. Shabbir, M. Gray, V. M. Lynch, S. Sorey and E. V. Anslyn, *J. Am. Chem. Soc.*, 2006, **128**, 1222–1232.
- 24 A. P. de Silva, H. Q. N. Gunaratne, T. Gunnlaugsson, A. J. M. Huxley, C. P. McCoy, J. T. Rademacher and T. E. Rice, *Chem. Rev.*, 1997, **97**, 1515–1566.
- 25 (a) M. Takeuchi, M. Yamamoto and S. Shinkai, *Chem. Commun.*, 1997, **18**, 1731–1732; (b) W. Yang, J. Yan, G. Springsteen, S. Deeter and B. Wang, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 1019–1022; (c) R. Hosseinzadeh, M. Mohadjerani and M. Pooryousef, *Luminescence*, 2015, **30**, 549–555.
- 26 R. Dufault, B. LeBlanc, R. Schnoll, *et al.*, *Environ. Health*, 2009, **8**, 2.