

 Open Access Full Text Article

LETTER

Economic Analysis of Intravenous Iron in Patients with Iron Deficiency Anemia Due to Inflammatory Bowel Disease: Considerations for Clinicians [Letter]

Richard F Pollock¹, Anjan Dhar², Matthew Johnson³

¹Health Economics and Outcomes Research, Covalence Research Ltd, London, UK; ²Department of Gastroenterology, County Durham & Darlington NHS Foundation Trust, Darlington, Co. Durham, UK; ³Department of Gastroenterology, Luton and Dunstable University Hospital NHS Foundation Trust, Luton, UK

Correspondence: Richard F Pollock, Health Economics and Outcomes Research, Covalence Research Ltd, London, UK, Tel +44 20 8638 6525, Email pollock@covalence-research.com

Dear editor

We read, with interest, the recently published article by Aksan et al (2021),¹ which presents the results of an economic analysis comparing the cost-effectiveness of intravenous (IV) iron products in the treatment of iron deficiency anemia (IDA) associated with inflammatory bowel disease (IBD), in the UK setting. The authors concluded that ferric carboxymaltose (FCM; Vifor Pharma, Glattbrugg, Switzerland) is likely to be the least costly, and most effective, IV iron therapy compared with ferric derisomaltose (FDI; previously known as iron isomaltoside; Pharmacosmos A/S, Holbæk, Denmark) and iron sucrose (IS; Vifor Pharma, Glattbrugg, Switzerland).¹ We would like to share our experience of health economics in the field of IV iron therapy to present some important considerations for clinicians when interpreting these findings, to ensure rational and evidence-based treatment decisions.

Firstly, the clinical effectiveness inputs used in the Aksan et al (2021) analysis were based on the results of an earlier network meta-analysis (NMA) comparing the efficacy and tolerability of different IV iron products (including FDI, FCM, and IS) in patients with IDA due to IBD (published by Aksan et al in 2017).^{1,2} The NMA showed no statistically significant differences in efficacy (treatment response: hemoglobin [Hb] normalization or increase in Hb ≥ 2 g/dL) between FDI and FCM.² The shortcomings of this NMA have been detailed in a previous communication,³ but the Aksan et al (2021) analysis does not seem to incorporate changes to overcome these short-falls.

Secondly, in the Aksan et al (2021) analysis, the authors acknowledge the weakness of using indirect evidence to obtain the data for the clinical effectiveness inputs, justifying this approach by the lack of direct evidence from head-to-head trials conducted in a relevant patient population.¹ Indeed, Aksan et al (2021) did not include the data from the head-to-head PHOSPHARE-IDA trial showing comparable efficacy (iron and anemia parameters, assessed as secondary endpoints) between FCM and FDI, on the basis that the IDA was due to a gynecological condition, not to IBD.^{1,4} Evidence is now available from the head-to-head PHOSPHARE-IBD trial, which was conducted, specifically, in patients with IBD (n=96); equivalent doses of FCM and FDI (initial dose of 1000 mg at baseline followed by 500 mg or 1000 mg after 5 weeks) were compared, with assessment of Hb as a secondary endpoint.⁵ As shown in Table 1, the PHOSPHARE-IBD trial showed no statistically significant difference between FCM and FDI in the post-treatment Hb changes from baseline.⁵

The evidence from PHOSPHARE-IBD provides confirmation of equivalent hematological efficacy of FCM and FDI in an IBD population with IDA.⁵ Therefore, the suggestion that FCM is more effective than FDI (as was concluded for the Aksan et al (2017) NMA),² is not completely borne out by published evidence.

Table I Mean Change from Baseline in Hb Following Treatment with FCM or FDI

	FDI (n=49)	FCM (n=48)	p-value
Day 1	0.13 (0.45) (n=48)	-0.01 (0.50) (n=48)	0.1165
Week 1	0.58 (0.73) (n=45)	0.47 (0.71) (n=44)	0.2293
Week 2	1.16 (0.87) (n=46)	1.18 (0.94) (n=47)	0.9393
Week 5	1.77 (1.01) (n=42)	1.84 (1.06) (n=44)	0.7778
Week 6	2.08 (1.07) (n=43)	2.06 (1.13) (n=41)	0.8295
Week 7	2.36 (1.27) (n=42)	2.38 (1.23) (n=43)	0.6428
Week 10	2.51 (1.41) (n=43)	2.44 (1.49) (n=41)	0.9257

Notes: Data presented are mean (SD). Adapted from EU Clinical Trials Register. Clinical Trial Results: A randomized, double-blinded, comparative trial comparing the incidence of hypophosphatemia in relation to repeated treatment courses of iron isomaltoside and ferric carboxymaltose in subjects with iron deficiency anaemia due to inflammatory bowel disease. EudraCT number: 2017-002452-87. Available from: <https://www.clinicaltrialsregister.eu/ctr-search/trial/2017-002452-87/results>.⁵

Abbreviations: FCM, ferric carboxymaltose; FDI, ferric derisomaltose; Hb, hemoglobin; SD, standard deviation.

Thirdly, the Aksan et al (2021) analysis acknowledged that the impact of safety issues on cost and clinical outcomes was not considered, and that such information should be captured in future cost-effectiveness studies.¹ This is, indeed, an important reflection given that the Pharmacovigilance Risk Assessment Committee of the European Medicines Agency considers there to be a possible causal relationship between hypophosphatemic osteomalacia and FCM, specifically⁶ – it is not a class effect for all IV iron products. In November 2020, the regulatory authorities in the UK published an update to the product information for FCM to include a requirement for serum phosphate monitoring in patients who receive multiple administrations of FCM at higher doses, or long-term FCM treatment (and in those with existing risk factors for hypophosphatemia, including IBD).⁷ The cost of this additional monitoring is particularly relevant to the Aksan et al (2021) analysis, given that it was focused on a population of IBD patients in the UK.¹

We hope that the issues highlighted in this letter will give a broader perspective to the findings of the Aksan et al (2021) economic analysis for prescribing clinicians and decision-makers in the UK. Of particular importance for real-world clinical practice are the recent changes to the FCM product label, which translate into an additional economic impact of FCM treatment in the management of IDA due to IBD.

Funding

This letter was funded by Pharmacosmos A/S. Writing and editorial assistance was provided by 'Cambridge – a Prime Global agency'.

Disclosure

RFP is a director and shareholder in Covalence Research Ltd, which has received consultancy fees from Pharmacosmos A/S (the marketing authorization holder for Monofer) and Pharmacosmos UK Limited, a wholly-owned subsidiary of Pharmacosmos A/S. RFP has also received personal honoraria from Pharmacosmos UK Limited. AD has received

speaker fees from Pharmacosmos, Pfizer, Takeda, Tillotts UK, and Janssen, advisory honoraria from Takeda, Tillotts UK, Dr Falk Pharma, Pfizer, and conference support from Pharmacosmos, Tillotts, Takeda, Janssen, Dr Falk Pharma, and Cook Medical. MJ has received conference support from Pharmacosmos, Janssen, Dr Falk Pharma, Cook Medical, Pfizer, AbbVie, Amgen Ltd, Ferring Pharmaceuticals, Norgine Pharmaceuticals, Allergen, Aquilant Endoscopy, FujiFilm, MSD, Hospira, NAPP, Warner Chilcott, Sucampo Pharmaceuticals, Regenerus Laboratories, Takeda UK, Tillotts Pharma UK, Mylan, Vifor Pharma, Shield Therapeutics, Kyowa Kirin, and Viatris. The authors report no other conflicts of interest in this communication.

References

1. Aksan A, Beales ILP, Baxter G, et al. Evaluation of the cost-effectiveness of iron formulations for the treatment of iron deficiency anaemia in patients with inflammatory bowel disease in the UK. *Clinicoecon Outcomes Res.* 2021;13:541–552. doi:10.2147/CEOR.S306823
2. Aksan A, Işık H, Radeke HH, Dignass A, Stein J. Systematic review with network meta-analysis: comparative efficacy and tolerability of different intravenous iron formulations for the treatment of iron deficiency anaemia in patients with inflammatory bowel disease. *Aliment Pharmacol Ther.* 2017;45(10):1303–1318. doi:10.1111/apt.14043
3. Pollock RF, Muduma G. Response: an economic evaluation of iron isomaltoside 1000 versus ferric carboxymaltose in patients with inflammatory bowel disease and iron deficiency anemia in Denmark. *Adv Ther.* 2019;36(8):1821–1825. doi:10.1007/s12325-019-00993-8
4. Wolf M, Rubin J, Achebe M, et al. Effects of iron isomaltoside vs ferric carboxymaltose on hypophosphatemia in iron-deficiency anemia: two randomized clinical trials. *JAMA.* 2020;323(5):432–443. doi:10.1001/jama.2019.22450
5. EU Clinical Trials Register. Clinical Trial Results: a randomized, double-blinded, comparative trial comparing the incidence of hypophosphatemia in relation to repeated treatment courses of iron isomaltoside and ferric carboxymaltose in subjects with iron deficiency anaemia due to inflammatory bowel disease. EudraCT number: 2017-002452-87. Available from: <https://www.clinicaltrialsregister.eu/ctr-search/trial/2017-002452-87/results>. Accessed September 25, 2021.
6. European Medicines Agency. Scientific conclusions and grounds for the variation to the terms of the Marketing Authorisation(s). Available from: https://www.ema.europa.eu/en/documents/psusa/iron-parenteral-preparations-except-iron-dextran-cmdh/scientific-conclusions-grounds-variation/00010236/202001_en.pdf. Accessed November 3, 2021.
7. GOV.UK. Drug Safety Update. Ferric carboxymaltose (Ferinject▼): risk of symptomatic hypophosphataemia leading to osteomalacia and fractures; 2020. <https://www.gov.uk/drug-safety-update/ferric-carboxymaltose-ferinject-risk-of-symptomatic-hypophosphataemia-leading-to-osteomalacia-and-fractures>. Accessed November 3, 2021.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the ClinicoEconomics and Outcomes Research ‘letters to the editor’ section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the ClinicoEconomics and Outcomes Research editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

ClinicoEconomics and Outcomes Research

Dovepress

Publish your work in this journal

ClinicoEconomics and Outcomes Research is an international, peer-reviewed open-access journal focusing on Health Technology Assessment, Pharmacoconomics and Outcomes Research in the areas of diagnosis, medical devices, and clinical, surgical and pharmacological intervention. The economic impact of health policy and health systems organization also constitute important areas of coverage. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinicoeconomics-and-outcomes-research-journal>