

INVITED REVIEW

Drug discovery: Insights from the invertebrate *Caenorhabditis elegans*

Sebastián Giunti^{1,2} | Natalia Andersen^{1,2} | Diego Rayes^{1,2}  | María José De Rosa^{1,2} 

¹Instituto de Investigaciones Bioquímicas de Bahía Blanca (INIBIBB) CCT UNS-CONICET, Bahía Blanca, Argentina

²Dpto de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, Bahía Blanca, Argentina

Correspondence

Diego Rayes and María José De Rosa, Instituto de Investigaciones Bioquímicas de Bahía Blanca (INIBIBB) CCT UNS-CONICET, Camino La Carrindanga Km 7, Bahía Blanca, Argentina.
Email: drayes@criba.edu.ar (D. R.); mjderosa@criba.edu.ar (M. J. R.)

Funding information

This work was supported by Consejo Nacional de Investigaciones Científicas y Técnicas (PIP 2015–2017 CO 11220150100182CO to DR), Agencia Nacional de Promoción Científica y Tecnológica (PICT 2017 0566 to MJDR), Universidad Nacional del Sur (PGI 2016 24/B261 to MJDR and PGI 2018 24/B291 to DR).

Abstract

Therapeutic drug development is a long, expensive, and complex process that usually takes 12–15 years. In the early phases of drug discovery, in particular, there is a growing need for animal models that ensure the reduction in both cost and time. *Caenorhabditis elegans* has been traditionally used to address fundamental aspects of key biological processes, such as apoptosis, aging, and gene expression regulation. During the last decade, with the advent of large-scale platforms for screenings, this invertebrate has also emerged as an essential tool in the pharmaceutical research industry to identify novel drugs and drug targets. In this review, we discuss the reasons why *C. elegans* has been positioned as an outstanding cost-effective option for drug discovery, highlighting both the advantages and drawbacks of this model. Particular attention is paid to the suitability of this nematode in large-scale genetic and pharmacological screenings. High-throughput screenings in *C. elegans* have indeed contributed to the breakthrough of a wide variety of candidate compounds involved in extensive fields including neurodegeneration, pathogen infections and metabolic disorders. The versatility of this nematode, which enables its instrumentation as a model of human diseases, is another attribute also herein underscored. As illustrative examples, we discuss the utility of *C. elegans* models of both human neurodegenerative diseases and parasitic nematodes in the drug discovery industry. Summing up, this review aims to demonstrate the impact of *C. elegans* models on the drug discovery pipeline.

KEYWORDS

C. elegans, drug discovery, genetic screenings, human disease models, pharmacological screenings

1 | INTRODUCTION: THE NEMATODE *Caenorhabditis elegans*

Since its introduction into research by Sydney Brenner in the early 60s, the nematode *Caenorhabditis elegans* has played a pivotal role in

different areas of biomedical investigation.^{1–9} Studies on *C. elegans* have contributed to fundamental breakthroughs in life science, such as the discovery of genetic regulators of programmed cell death, the use of the green fluorescent protein (GFP) as a protein marker, and the discovery of RNA interference (RNAi).^{10–12}

Abbreviations: AChR, muscle nicotinic receptor; AT, α 1-antitrypsin; A β , β -amyloid; CACNA1A, calcium voltage-gated channel subunit alpha 1A; CGC, *Caenorhabditis Genetics Center*; HTS, high-throughput screening; MOA, Mechanism of action; NGM, Nematode Growth Medium; PBS, phenotypic-based drug screens; TBS, target-based drug screens.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Pharmacology Research & Perspectives* published by John Wiley & Sons Ltd, British Pharmacological Society and American Society for Pharmacology and Experimental Therapeutics.

C. elegans is a tiny (~1 mm in length), soil, free-living, and bacteria-eating nematode^{13,14} which can develop as either of two sexes, self-fertilizing hermaphrodites or males. Males represent a minor population (~0.2% of population).^{14,15} Healthy hermaphrodites produce up to 300 self-progeny although they are capable of giving approximately 1000 offspring if they are mated with males.

This nematode typically lives for about 3 weeks at 20°C in the laboratory. Unlike other members of the Nematoda phylum, *C. elegans* is a nonhazardous and nonpathogenic animal that can be manipulated with standard safety rules. A unique benefit of using *C. elegans* in the laboratory is that it can be frozen and stored in liquid nitrogen or -80°C ultra-freezers until needed.¹⁴ Moreover, its use does not raise the ethical issues associated with the use of vertebrates (Figure 1).

C. elegans life cycle, which is very short, takes approximately 2.5–3 days and is divided into embryogenesis (~16 h), four larval stages (L1–L4, ~28 h) and the adult molt, from L4 to adult (~12 h). Adverse environmental conditions (starvation, high population density, high temperature) in the late L1 larval stage trigger entry into a developmental arrest phase called dauer.^{16,17} Dauer larvae can survive stressful conditions for months until encountering favorable conditions that permit resuming reproductive development. The presence of separate sexes and the short life cycle permit rapid genetic crosses in the laboratory. It has an invariant number of somatic cells: 959 in hermaphrodites and 1031 in males. *C. elegans* cellular genealogies have been traced from embryo to newly hatched larva. Unlike mammals, the pattern of cell lineages is invariant among

individuals, which has been key to understand the roles of developmental control genes.^{18–20}

Despite its apparent simplicity, *C. elegans* contains organs and tissues present in more complex organisms, such as muscles, hypodermis, intestine, reproductive and excretory system, and a well-described nervous system (Figure 1). It is also the only organism where the complete neural wiring diagram has been established.^{21,22} Its transparency allows the easy visualization of specific cells and subcellular structures through Nomarski (differential interference contrast, DIC) optics using whole-live animals. The visualization of protein expression patterns, protein subcellular localization, and activity of specific cells using transgenic reporters and genetically encoded calcium indicators is feasible in living animals at any point in their lives (Figure 1).^{23–25}

The fact that *C. elegans* was the first multicellular organism to have its complete genome sequenced (*C. elegans* Sequencing Consortium 1998) as well as its amenable genetic manipulation helped to turn this worm into a powerful organism for genetic screens.^{26,27} 83% of *C. elegans* proteome is predicted to have human homolog genes²⁸ and it has been estimated that ~50% of *C. elegans* protein-coding genes have a functional ortholog in humans.²⁹ Essential biological molecules and cell signaling pathways, such as the insulin-signaling pathway, innate immunity regulatory pathways, synaptic machinery, and systems involved in protein homeostasis are highly conserved between worms and mammals.^{30–34}

Taken together, all these features make *C. elegans* an excellent model organism that bridges the gap between the simplicity of cultured cells and the complexity of multicellular organisms.




FIGURE 1 *Caenorhabditis elegans* as a versatile platform for drug discovery. Its genetic amenability and ease of transgenesis allows elucidation of MOAs by forward and reverse genetic screens and generation of "humanized worms" to emulate conditions seen in humans. All this combined with the feasibility for HTS and automation of easily scored phenotypes converts *C. elegans* into a powerful model for pharmacological research. For more detailed anatomy and other *C. elegans* resources visit: WormAtlas (www.wormatlas.org), Wormbase (www.wormbase.org), Wormbook (www.wormbook.org)

2 | *C. elegans* IN DRUG DISCOVERY

The advances in molecular biology techniques and genome knowledge allowed drug screening to be carried out directly on target proteins. Therefore, target-based drug screens (TBS) have become the most commonly used approach for drug discovery. In this type of approach, compounds are screened for their capacity to bind or alter the activity of specific target proteins using cell extracts or cell cultures. However, as targets are previously identified and validated, TBS does not permit the discovery of novel targets and the elucidation of mechanisms of action (MOAs) is frequently limited to agonism or antagonism of known receptors/pathways. Moreover, once hits are detected they need to be tested in the context of a whole organism to further analyze its biological efficacy and toxicity. In contrast to TBS, phenotypic-based drug screens (PBS) use cells or animal disease models to identify compounds that rescue or ameliorate the disease phenotype. This animal-based approach includes the effects of cell communications and tissue interactions in the effect of a given compound. This unbiased strategy allows the identification of compounds with novel MOAs. One of the main caveats of PBS is that the identification of targets and MOAs could be complicated, particularly when mammalian animal models are used. This drawback can be significantly mitigated using invertebrates, such as the nematode *C. elegans*, which has lately gained consideration as an excellent platform for the identification of new drug targets and drug discovery.

Automation of worm transfer, image acquisition, and data analysis allows the use of *C. elegans* for high-throughput screening (HTS) assays.^{35–39} Several companies (Celescreen, Sunnybiotech, Nagi Biosciences) offer these types of assays to the pharmaceutical industry for the identification of new potential compounds and validation of pharmacological targets. In the last 10 years, HTS in *C. elegans* has become—in fact—a useful tool to identify candidate compounds as potential treatments for several pathological conditions (Table 1).

The fact that *C. elegans* shares the nematode phylum with several parasitic roundworms makes it a straightforward model to be used for the development of anthelmintic drugs. Its use has allowed the MOA elucidation of several nematocidal drugs^{75–78} (Table 2). However, its use goes far beyond than merely applying *C. elegans* to the discovery of drugs with anthelmintic potential. *C. elegans* is also exploited as an accessible platform to recapitulate distinctive phenotypes of human diseases, such as cancer, diabetes, neurodegenerative disorders, and pathogen infection.^{79–89} The molecular pathways underlying a biological process that, when altered, could lead to physiological diseases are usually conserved throughout the animal kingdom. The alteration of these conserved pathways in *C. elegans* often leads to specific phenotypes and behaviors, such as modifications in lifespan, stress resistance, or locomotion. All these phenotypes are easily measurable using automated protocols and devices, making the screening of drugs that ameliorate or rescue these defects a relatively simple procedure.

Several genetic tools for the manipulation of single genes or groups of genes are readily available in *C. elegans* (chemical mutagenesis, transgenesis, RNAi, and CRISPR/Cas9) (Figure 1). Animals

carrying mutations of highly conserved biochemical pathways or expressing exogenous pathognomonic protein of a specific disease can be engineered in few weeks. Mutant strains and animals expressing human transgenes are available at a very low cost at the *Caenorhabditis* Genetics Center (CGC). Moreover, the rapid life-cycle and the high brood size of *C. elegans* permit the analysis of the effects of compounds not only in the exposed animal but also in the progeny in a very short time-frame.

Cell cultures or cell extracts are typically the initial step in drug development. Using invertebrate animal models in these early steps is ideal to reduce research costs and time as they help to identify compounds that, apart from being effective in interacting with the active target, maintain their efficacy even after absorption, distribution, metabolism, and excretion (ADME) processes. They also provide information about potential systemic toxicity and biocompatibility.⁹⁶

Although ADME processes are important sources of information compared with cell cultures, in *C. elegans* they have certain limitations that should be taken into account at the moment of processing results from drug-screening assays. In particular, the thick cuticle, which forms a strong barrier that limits drug absorption, can usually render false-negative results.⁹⁷

C. elegans' low-cost maintenance, small size, genetic amenability, and conservation of key molecular pathways with mammalian animals convert this worm into an excellent alternative to significantly reduce drug development costs. Although the use of *C. elegans* as a platform for drug screening is recent, there are compounds that have been identified in this nematode and which, once validated on vertebrate models, are currently being evaluated in human patients. For example, hits obtained in a drug screening performed in a *C. elegans* model of amyotrophic lateral sclerosis were validated in zebrafish and mice. One of these hits, the neuroleptic Pimozide, a Ca²⁺ channel blocker that stabilizes neuromuscular transmission in *C. elegans*, has also shown efficiency in a short randomized controlled trial of sporadic ALS subjects.⁴³ Another report evaluated the pearl powder (a Chinese medicine) by combining *C. elegans* studies with a clinical trial. The authors have correlated the lifespan extension found in *C. elegans* with the higher antioxidant capacity in blood samples of patients treated with pearl powder.⁹⁸ These two examples serve as proofs of concept for the translational potential of drug screenings in the invertebrate *C. elegans*. Since the number of drug screenings in this nematode is increasing year after year (Table 1), more clinical trials for hits identified in *C. elegans* are expected in the near future.

2.1 | Genetic screening assays

One of the greatest advantages of using *C. elegans* for biomedical research is undoubtedly its amenability for genetic manipulation. Genetic screens are widely used in *C. elegans* to either discover gene function or find genes involved in relevant biological pathways. Furthermore, the feasibility of rapid genetic crossing permits epistatic analyses to find new players in known genetic pathways.⁹⁹

TABLE 1 *Caenorhabditis elegans* phenotypic-based drug screenings. The table shows relevant drug screenings assays performed during the last decade in *C. elegans* models for human and animal diseases

Biological activity	Initial screening set	Identified hits	References
Anti-proteotoxicity Neuroprotection	10 positive modulators of healthspan	Metformin, lithium, and curcumin	40
	18 compounds	α -methyl- α -phenylsuccinimide	41
	87 flavonoids and 13 neurosteroids	12 flavonoids (e.g., isoquercitrin) and 2 steroids (3 β -Methoxy-Pregnenolone and 17 β -estradiol)	42
	3 850 compounds	13 hits (e.g., pimozone)	43
	983 FDA-approved drugs	4 hits (dronedarone, tofranil, bendrofluzide, buspar)	44
	4 polyphenolic compounds	Ferulic acid	45
	30 FDA-approved drugs	Tannic acid, bacitracin	46
	115 000 compounds	Four tetrahydroquinolinones	47
Anti-aging	1386 FDA-approved drugs	Verapamil	48
	32 compounds	1 hit (chalcone like-compound)	49
	~100 serine hydrolase inhibitors	JZL184	50
	107 FDA-approved drugs	Tiagabine	51
	15 FDA-approved drugs	Captopril	52
	33 000 compounds	57 hits (e.g., nitrophenyl piperazine-containing compounds)	53
	1280 compounds	57 hits (e.g., minocycline)	54
	normal and disease-associated endogenous metabolites	α -ketoglutarate	55
Anti-microbial	69 compounds	5 hits (phenyl triazine compounds)	56
	82 000 compounds	185 hits (e.g., synthetic retinoid CD437)	57
	86 000 compounds	195 hits (e.g., 5-fluorouracil)	58
	21 500 compounds	318 hits (e.g., phenylsulfonyl pyrazinecarbonitrile)	59
	640 FDA-approved drugs	42 hits (e.g., closantel)	60
	1600 compounds	18 hits (e.g., iron-chelator ciclopirox olamine)	61
	1300 extracts (from endophytic fungi)	4 hits	62
	2560	12 hits (e.g., natural saponins)	63
Anthelmintic	12 benzopyrano pyrazol compounds	4 hits	64
	575 compounds	29 hits (e.g., arylidene ketones)	65
	11 imidazole-derivatives	2 hits (e.g., diisopropylphenyl-imidazole)	66
	400 compounds (Pathogen Box library)	18 hits (e.g., isoxazole compounds)	39
	480 compounds	20 hits (e.g., dihydrobenzoxazepinones)	67
	67 012 compounds	30 hits (e.g., ethyl benzamide moiety compounds)	68
Anti-tumoral	4 in vitro anticancer compounds	4 hits	69
	30 plant extracts	Harmine (from the plant <i>Peganum harmala</i>)	70
	~9000 compounds	2 hits (an EGFR inhibitor and a MEK inhibitor)	71
Anti-metabolic disorders (obesity, insulin resistance, type II diabetes)	24 plants and fungal extracts	2 hits (extracts of <i>Inonotus obliquus</i> and <i>Gardenia jasminoides</i>)	72
	350 natural products	1 hit (swertiamarin)	73
	8 natural plant compounds	2 hits (isoquinoline alkaloids)	74

To date, two strategies to perform screenings in *C. elegans* have been developed: forward genetics and reverse genetics. Whereas the former uses mutagens to randomly generate mutations that induce or reverse a given phenotype, the latter analyzes the phenotype obtained after altering or knocking down a specific known gene. Forward genetics therefore goes from phenotype to gene, whereas reverse genetics works the other way around, that is, from gene to phenotype.

2.1.1 | Forward genetics

Mutations are important tools for gene function discovery. Mutagenesis can be accomplished using a variety of chemical agents, such as ethyl methanesulfonate (EMS), N-ethyl-N-nitrosourea (ENU), trimethylpsoralen (TMP), among others. Hermaphrodites (PO) are

TABLE 2 Recent contributions in deciphering drug-targets and mechanism of action. Representative compounds for which the use of *C. elegans* has been useful in elucidating their mechanisms of action

Compound	Field	Identified target/Mechanism of action	References
Resveratrol	Neurodegeneration	Reduces β -amyloid by targeting UBL-5 and XBP-1, proteins implicated in UPR ^{mt} and UPR ^{ER}	90
Minocycline	Aging	Increases lifespan by decreasing mARN translation	91
RPW-24	Microbial infection	Stimulates innate immune response through pmk-1/p38 MAPK pathway, and the transcription factor, <i>atf-7</i>	92
Monepantel	Helminth infection	Targets ACR-20 and ACR-23 from the DEG-3 subgroup of nAChR subunits	93
Metformin	Cancer	Induces tumor growth inhibition and lifespan extension by targeting nuclear pore complex (NPC) and acyl-CoA dehydrogenase family member-10 (ACAD10)	94
Hesperidin	Metabolic disorders	Inhibits lipid accumulation by downregulating lipid metabolism genes (<i>fat-6</i> and <i>fat-7</i>)	95

Abbreviations: ACR, acetylcholine receptor subunit; *atf-7*, Cyclic AMP-dependent transcription factor 7; MAPK, mitogen-activated protein kinase; pmk-1, p38 MAPK; UBL-5, Ubiquitin-like protein 5; UPR^{ER}, unfolded protein response of the endoplasmic reticulum; UPR^{mt}, mitochondrial unfolded protein response; XBP-1, X-Box Binding Protein 1.

randomly mutagenized at the late L4/early adult stages and then distributed in Petri dishes. F1 hermaphrodite progeny that are heterozygous for these mutations can be allowed to self-fertilize. The F2 animals, which bring mutations to homozygosity, are isolated based on the phenotype of interest.²⁷ Several mutations generate clearly altered phenotypes, such as changes in development, lifespan, stress resistance, uncoordinated movement (*unc*), or dumpy-shaped animals (*dpy*). Today, mutated genes underlying a given phenotype can be identified by whole-genome sequencing.^{100,101}

Another approach, which has proven to be very useful for identifying novel gene interactions, consists in performing mutagenesis directly in strains already carrying mutations that induce a strong phenotype.¹⁰²⁻¹⁰⁴ Mutant animals are then mutagenized allowing enhancer or suppressor genes to be detected. For example, a suppressor screening recently performed on a hyperactive mutant in UNC-2, the worm ortholog for calcium voltage-gated channel subunit alpha 1A (CACNA1A) channel, has led to the identification of key modulators of channel function.¹⁰⁴

Chemical genetic screening is another strategy which involves the use of drugs that induce a strong phenotype. After mutagenesis, which may affect the drug target gene or other genes involved in the same pathway, either resistant or hypersensitive mutants can be identified.^{75,105-107}

In *C. elegans*, forward genetic screens have been traditionally used to identify molecular targets and MOAs for anthelmintic agents.^{75-77,108} However, more recently, this approach has also proved to be useful for discovering targets and MOAs for drugs with potential relevance in metabolic, neurological, and oncological disorders.^{68,94,109-112}

2.1.2 | Reverse genetics

In 1998, Fire and Mello found that injection of double-stranded RNA (dsRNA) into worm gonads allows gene knocking down in the progeny.¹² Since then, RNAi knockdown has become a powerful tool in other animals and cultured cells.¹¹³⁻¹¹⁵ However, in

C. elegans it can be easily achieved by feeding worms with bacteria expressing dsRNA or just soaking worms in dsRNA solution. Therefore, this technique has served for functional analyses of a plethora of genes in worms.

Genomic RNAi bacterial feeding libraries that cover most of the *C. elegans* genome have been generated,^{116,117} avoiding the requirement of in vitro synthesis of dsRNA. *C. elegans*, one of the few organisms for which genome-wide RNAi screens are feasible, has been found to be a useful tool to identify genes involved in essential biological processes^{2,118-120} and pathological conditions.^{107,121,122}

A historical limitation of RNAi knock-down in *C. elegans* has been the poor penetrance of RNAi in neurons. However, several strategies have been developed to solve this caveat, namely strains expressing the transmembrane proteins required for RNAi transportation SID-1 pan-neuronally,¹²³ strains deficient in genes coding for ribonuclease enzymes,¹²⁴ neurons transformed to an immature and more permeable stage¹²⁵ or even a combination of them.¹²⁶

In the last decade, several customized endonucleases, such as zinc-finger nucleases (ZFNs) and transcription activator-like effector nucleases (TALENs),^{127,128} have been used in genome editing. Other techniques, such as CRISPR/Cas9 and Mos-1 single copy insertion that allow the introduction of almost any change in any gene, have also been developed, further widening the tool repertoire to perform reverse genetics in the nematode.¹²⁹⁻¹³¹

2.2 | Drug screening assays

Drug ingestion is probably the main way through which xenobiotic molecules can gain access to target tissues in wild-type *C. elegans*.^{132,133} However, the amphids, a pair of anterior sensory structures opened to the outside environment, and the vulva are also involved in drug and nanoparticles intake, respectively.^{134,135} Moreover, the use of animals that have a compromised cuticle can significantly increase drug absorption.⁹⁷ *C. elegans* is typically grown in the laboratory on petri-dishes containing NGM agar and *E. coli* as a food source. In the

past, during the first drug screens, compounds were generally dissolved in the agar.^{14,136} Although this method has contributed to the elucidation of MOAs for several compounds previously proven to be biologically active,^{137,138} the agar-based technique does not permit large scale screenings because it is labor- and time-consuming and needs large quantities of drugs.¹⁴ Even with limitations, these assays are still very valuable for assessing small compound libraries.⁶⁶

The introduction of *C. elegans* liquid cultures in multi-well format together with workflow automation from worm transfer to data analysis enabled the development of HTS assays.^{139,140} Moreover, the breakthrough of microfluidic devices for *C. elegans* have improved the strength of HTS mainly thanks to the small amounts of compound (microscale) that are required and to the possibility of observing worms in parallel or in a serial manner.¹⁴¹

2.3 | Challenges of using *C. elegans* model in drug discovery

In spite of the obvious advantages of using *C. elegans* for drug screening, some caveats need to be considered. First, *C. elegans* cultures require bacterial co-culture as a food source. Many drugs and compounds are metabolized by bacteria and could alter drug efficacy. These limitations can be circumvented using killed bacteria as a food source.^{142,143} However, feeding worms with dead bacteria can also affect some phenotypes (e.g., it slows worm development).¹⁴⁴ Therefore, the chosen food source will depend on the phenotype to be assessed, the compounds to be tested, and the bacteria species to be used for worm feeding.^{142,145}

A second disadvantage is related to the thick *C. elegans* cuticle that usually affects drug uptake. Some studies revealed that internal tested-drug concentration could be less than half of that applied externally.¹⁴⁶ This implies that some drugs are discarded as hits due to their inefficacy to pass through the cuticle barrier and that some compounds need very high drug concentrations to produce a biological effect. To reduce these limitations, mutant strains with compromised cuticles have been used to enhance drug uptake.⁹⁷ The selection of these mutants depends on the trade-off between enhanced cuticle permeability and animal fitness.⁹⁷ Although *C. elegans* cuticle impermeability is indeed a limitation, it increases the potential significance of a positive hit in specific drug screening assays. In line with this, hits in anthelmintic search assays that induce death in *C. elegans* are extremely likely to kill parasitic nematodes (which in general have thinner cuticles) at lower concentrations.⁶⁸

Finally, *C. elegans* lacks a circulatory system and many vital organs present in mammals. It has an innate immune system but lacks an adaptive immune system or a myelination system. This is the reason why it is difficult to model diseases affecting these organs/systems in *C. elegans*. In some cases, it is nonetheless possible to study a disease affecting a particular mammal organ nonexistent in the worm using phenologues. The latter are described as a group of overlapping genes defining a pathway that, when disrupted, cause different phenotypes in different species. For example, osteogenesis imperfecta (known as brittle

bone disease) is caused by mutations in the human collagen COL1A1 and COL1A2 genes.¹⁴⁷ Although *C. elegans* lacks bones, it expresses collagen genes. When worm collagen genes are mutated, they lead to cuticle defects, easily showing recognized dumpy phenotypes.¹⁴⁸ Thus, the dumpy phenotype can be used to perform screenings to study collagen-associated diseases even in the absence of the affected tissue.

Additionally, even if a disease gene ortholog is not present in the worm, a disease model can be artificially generated by expressing human transgenes.^{149,150} Transgenic animals are then used as tools to screen drugs or new genes involved in the disease phenotype with the goal of finding new therapeutic options. For example, mutations in α 1-antitrypsin (AT) lead to lung and liver diseases.^{151,152} Although *C. elegans* has no orthologs for AT, transgenic expression of human AT gene in worms can replicate in the animal intestine the abnormal misfolded protein accumulation observed in patients' livers. A drug screen performed using this model allowed to detect drugs capable of decreasing misfolded protein accumulation.³⁵

3 | RELEVANT *C. elegans* APPLICATIONS FOR DRUG TARGET DISCOVERIES

As illustrative examples of the use of *C. elegans* model in pharmacology research, we will discuss findings that show the advantages of using this nematode in the drug discovery field.

3.1 | Neurodegenerative diseases

As a result of the continuous increase in the proportion of the elderly population, age-related disorders have become a major health concern. Neurodegenerative diseases (NDs), in particular, such as Alzheimer's disease (AD), Parkinson's disease (PD), and Huntington's disease (HD), are considered as top ten lethal illnesses (G7 Academies' Joint Statements 2017). Despite their growing incidence worldwide and decades of intense research, there is still neither a cure nor a set of effect-mitigation strategies for these diseases. Research efforts are currently aimed at precluding or—at least—delaying their progression. The discovery of curative treatments for the majority of these NDs remains elusive due to several factors. NDs are heterogeneous in etiology (while some of them have a genetic or hereditary origin, others have no related cause). They are also heterogeneous in the collection of neurons affected, leading to an assortment of clinical manifestations ranging from cognitive to progressive motor dysfunction.¹⁵³ They nonetheless share some pathological hallmarks. For example, aging is a strong risk factor related to their development.¹⁵⁴ Moreover, the abnormal formation and deposition of misfolded protein aggregates in specific neurons is a trademark of most NDs. Because of this attribute, these disorders are generally referred to as “proteinopathies.” Therefore, current research is focused on the above-mentioned common features by targeting pathological aggregation or by delaying physiological aging.¹⁵⁵

TABLE 3 List of the main *C. elegans* models of ND, employed for the understanding of disease molecular mechanisms and for performing genetic and pharmacological screenings

Model of	Strategy	Expression	Transgene	Phenotype
HD	polyQ expression	Constitutive muscle	<i>Punc-54::polyQ-yfp</i>	Age-dependent aggregation and motility defects ⁸²
		Constitutive pan-neuronal	<i>Pgef-1::polyQ-yfp</i>	Age-dependent aggregation and neuronal dysfunction like trashing and pharyngeal pumping ⁸³
polyQ-HTT expression		Constitutive muscle	<i>Punc-54::htt513Q128::yfp</i>	Age-dependent aggregation and motility defects ¹⁸³
		Chemosensory neurons	<i>Posm-10::gfp+Posm-10::httQ150+dpv-20(+)</i>	Accelerated ASH neuronal degeneration ¹⁸⁴
		Mechanosensory neurons	<i>Pmec-3::htt57Q128::cfp+Pmec3::yfp</i>	Aggregated polyQ, morphological abnormalities and dysfunction of mechanosensory neurons ¹⁸⁵
AD	human A β expression	Constitutive muscle	<i>Punc-54::Aβ₁₋₄₂</i>	Age-dependent aggregation and paralysis ¹⁷⁸
		Temperature-induced muscle	<i>smg-1(cc54);Is[Pmyo-3::Aβ₁₋₄₂::letUTR+rol6(su1006)]</i>	Rapid paralysis by temperature upshift ¹⁸⁶
		Constitutive pan-neuronal	<i>Pmyo-2::yfp+Punc-119::Aβ₁₋₄₂</i>	Impaired locomotion and chemotaxis behavior ¹⁸⁷
		Temperature-induced pan-neuronal	<i>smg-1(cc54);Is[snb-1::Aβ₁₋₄₂+mtl-2::gfp]</i>	Defective chemotaxis, formation of amyloid deposits, and serotonin hypersensitivity ¹⁸⁸
	human Tau expression	Constitutive pan-neuronal	<i>Paex-3::h4R1NTau WT, V337M and P301L</i>	Insoluble tau accumulation, neurodegeneration and uncoordinated movement (Unc) ¹⁸⁹
	Single copy knock-in in mechanosensory neurons		Single copy human Tau mutants (T231E and K274/281Q)	Diminished touch response and neuronal morphological abnormalities ¹⁹⁰
PD	α -synuclein expression	Constitutive muscle	<i>Punc-54::α-syn::yfp+unc-119(+)</i>	Motility deficits and α -syn aggregation ^{5,159}
		Constitutive pan-neuronal	<i>Paex-3::α-syn (WT, A53T)+Paex-3::gfp/Pdat-1::gfp</i>	Motility deficits, significant dopaminergic neuron loss ^{158,191}
		Constitutive dopaminergic neurons	<i>Pdat-1::α-syn (WT, A53T)+Pdat-1::gfp</i>	Dopaminergic neuronal loss and dendritic breaks, α -syn accumulation ^{158,192}
Neurotoxin-induced	Dopaminergic neurons	[<i>Pdat-1::gfp</i>] subject to 6-OHDA or MPTP	Morphological dopaminergic neuronal defects ¹⁹³	
ALS	SOD-1 models	Temperature-induced muscle	<i>Phsp-16.2::sod-1 (WT, A4V, G37R, G93A) + Pmyo-3::sod-1 (WT, A4V)::GFP +rol-6(su1006)</i>	Increased sensitivity to oxidative stress, and oxidative stress induced- SOD-1 aggregation ¹⁹⁴
		Constitutive muscle	<i>Punc-54::sod=1(WT, G85R, G93A, G127insTGGstop)::yfp</i>	Aggregation of mutant SOD-1 and motility defects ¹⁹⁵
		Constitutive pan-neuronal	<i>Psnb-1::sod-1(WT, G85R)::yfp</i>	Aggregation of mutant SOD-1 and motility defects ¹⁹⁶
		Single copy knock-in	Single copy knock-in expression of mutants forms of <i>sod-1</i> (A4V, H71Y, L84V, G85R, G93A)	Aggregation of mutant SOD-1 and oxidative stress induced neurodegeneration ¹⁸⁰
	TPD-43 models	Constitutive pan-neuronal	<i>Psnb-1::tdp-43 (WT, G290A, A315T, M337V)+snb-1::gfp</i>	Uncoordinated movement and GABAergic motor neuron degeneration ^{197,198}
FUS models		Constitutive GABAergic motor neurons	<i>Punc-47::tpd-43 (WT, A315T)</i>	Age-dependent motility defects and neuronal degeneration ¹⁹⁹
		Constitutive pan-neuronal	<i>Pgef-1::fus (WT, R514G, R521G, R522G, R524S, P525L)</i>	Aggregation and paralysis ²⁰⁰
		Single copy knock-in	Single-copy knock-in of R524S and P525L equivalent mutations into <i>fust-1</i> (FUS orthologous gene)	Impaired neuronal and muscular autophagy ¹⁸¹

Abbreviations: ALS, Amyotrophic lateral sclerosis; a-syn, a-synuclein; FUS, RNA-binding protein Fused in Sarcoma; HTT, huntingtin; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; 6-OHDA, 6-hydroxydopamine; polyQ, polyglutamine; SOD-1, superoxide dismutase 1; TPD-43, Transactive response DNA-binding protein-43.

In this respect, *C. elegans* is a powerful *in vivo* model for the development of targeted therapeutics. Most of the advantages that convert this animal into a superb candidate for drug discovery have already been mentioned in this review and are also applicable to the neurodegeneration research field. Still, there are some topics that deserve special attention. Features that confer *C. elegans* an extra value as a model for age-related proteinopathies research include, among others: (i) the remarkable similarities at the molecular and cellular levels between nematode and vertebrate neurons, (ii) the continuous design of mutant and transgenic *C. elegans* models of human NDs,^{82,156–161} (iii) the thoroughly studied whole-animal neuronal connectome which can be individually visualized using fluorescent reporters in the living transparent worm,^{11,21,22,162} (iv) its short and genetically tractable lifespan,^{7,163–166} and (v) the recently developed high-throughput (even automatized) drug screening platforms.^{35,54,167–173} However, the limitations on using a “humanized” worm model for neurodegenerative proteinopathies should be taken into account, particularly its simple nervous system, glia and lack of myelin. Furthermore, genes encoding for voltage-gated sodium channels are absent in the *C. elegans* genome.¹⁷⁴ Unlike vertebrates, the action potentials detected in muscles and neurons of *C. elegans* depend on voltage-gated calcium channels.^{175–177} Despite these caveats, *C. elegans* is an excellent whole-animal platform to identify genes, drug targets, and compounds with neuroprotective roles in human NDs.

3.1.1 | *C. elegans* models for ND

The first *C. elegans* model of a human ND was built 25 years ago,¹⁷⁸ and, since then, an extensive list of animal models for most of the NDs have been generated (Table 3). The methods that are regularly used include knocking-down or -out the homologous gene involved in the human disease if it is present in the worm genome or, if there is no orthologous, the method of choice involves the transgenic expression of the human gene to mimic a disease-related phenotype. Most of the *C. elegans* models of human NDs are transgenic strains developed by overexpression of human wild-type or disease-associated mutant genes. However, overexpressing genes, even as wild-type, sometimes results in an artificial condition that can be pathogenic. CRISP/Cas9 technology has been fortunately established in *C. elegans*¹⁷⁹ and recent single-copy knock-in models of some NDs have been developed (Table 3).^{180–182}

Since *C. elegans* has no orthologous gene for β -amyloid (A β), α -synuclein, or huntingtin, models of NDs have been essentially produced by transgenic overexpression of the disease causing-protein in either body-wall muscle cells, in all neurons or in a specific subset of neurons.^{82,83,150,158,178,195,197,201} In most cases, the protein is tagged with GFP, which permits monitoring protein aggregation in whole live animals. Furthermore, according to the promoter used, that is, either muscle or neuronal promoters, toxic protein aggregation is reflected by specific pronounced phenotypes which

are indispensable for high-throughput screenings. For instance, muscle-expressing strains usually manifest proteotoxicity as age-dependent locomotion defects. In addition, muscle cells are large, allowing the visualization of protein aggregation more easily than in neurons. Moreover, RNAi is more efficient in muscles than in neurons, making these models more amenable for RNAi screens. On the other hand, transgenic strains with neuronal expression of protein aggregates are more accurate models of NDs. In the latter, proteotoxicity can be assessed by means of indirect phenotypes, such as neurodegeneration or specific morphological and/or behavioral neuronal defects. Thus, according to the selected strain, different read-outs have been described, which can be used in HTS with the aim to evaluate genes and drug modifiers as potential therapeutic interventions.

Genetic and pharmacological screenings have been performed in these models by tracking protein aggregation and assessing a simple phenotype (such as locomotion or neuronal behavior defects) in whole-live animals. In fact, large RNAi-mediated reverse screens, performed in *C. elegans* models of NDs, have identified gene modifiers of protein homeostasis, including genes with roles in protein synthesis, folding, degradation, and vesicle trafficking.^{3,5,159,202–205} Forward mutagenesis genetic screens have also been developed in these models. For example, a study using a *C. elegans* model of HD identified a novel gene, called modifier of aggregation 4 (*moag-4*) as an enhancer of protein aggregation.²⁰⁶ Pharmacological screenings have also contributed with hundreds of neuroprotective compounds, classified in a wide range of categories, including natural and synthetic products, herbal medicines, and also FDA-approved drugs (Table 1). These findings highlight the strength of *C. elegans* in finding novel genetic and drug modifiers for human NDs, reinforcing its potential in preclinical drug discovery.

Despite the complexity of the pathology underlying NDs, most of the pathways involved are conserved or they can be mimicked in the worm. The aim of using *C. elegans* as a platform of NDs is to dig inside molecular mechanisms to unravel pathological features and to develop effective strategies to treat these diseases.

3.2 | *C. elegans* as model of parasite nematodes

Parasitic nematodes infect a wide range of species, including humans, companion animals, livestock, and crops producing a devastating impact on human life quality and economy. The World Health Organization (WHO) claims that helminth infections are the most common neglected tropical diseases and estimates that 30% of the human world population is infected with at least one parasite (https://www.who.int/neglected_diseases/en/). This prevalence could be even higher in rural areas and in low-income countries. Although helminth infections are, in general, not lethal, human helminthiasis are associated with morbidity. The consequences are particularly serious in children, impairing growth, nutrition, cognition, and school performance.

Several issues are of concern in relation to helminthiasis management and treatment, namely (i) the loss of effectiveness caused by parasite resistance, (ii) the environmental impact of drugs used for crop protection, and (iii) the lack of interest in this field by the pharmaceutical industry.

The development of novel pharmacological agents for helminthiasis treatment has been delayed for decades and the repertoire of available anthelmintics is limited.^{77,207,208} Since drug-resistant parasitic nematodes have been reported for all classes of currently used anthelmintics,^{209–213} there is an urgent need to advance in pharmacological research to develop new antiparasitic drugs.

Due to their complex life cycles, growing and maintaining parasitic nematodes under standard laboratory conditions are both challenging. The limited molecular genetic tools available for these nematodes hampers the study of molecular mechanisms.²¹⁴ It has been shown that compounds that induce *C. elegans* death likely also kill parasitic nematodes,⁶⁸ highlighting the potential of this worm as an anthelmintic screening platform (Table 1).^{39,215,216}

3.2.1 | Current use of *C. elegans* in anthelmintic drug discovery

In the past, *C. elegans* was used to dissect the target pathways and molecular mechanisms of known anthelmintics, such as levamisole, ivermectin, benzimidazoles, nitazoxanide, and amino-acetonitrile derivatives.^{75,77,78,108,217–219} Nowadays, it is a recognized platform to screen new compounds and drug repurposing²²⁰ that must be subsequently tested in parasites. Many of the currently used anthelmintics are imidazole derivatives.²²¹ In general, the mechanisms underlying the anthelmintic effect can induce worm death or generate paralysis to facilitate parasite expulsion. The anthelmintic mechanisms of imidazole-derivatives are diverse. For example, levamisole causes nematode spastic paralysis through the potent activation of a muscle nicotinic receptor (AChR)^{218,222–225} whereas the anthelmintic action of benzimidazoles (e.g., albendazole and mebendazole) arises from their capacity to block tubulin polymerization in nematode cells.⁷⁸ Thus, while the presence of the imidazole ring appears to be important for bioactivity, it does not restrict the molecular targets where imidazole-containing anthelmintics can act. Recently, taking advantage of *C. elegans* as an established model for parasitic nematodes, we screened the nematicidal potential of novel imidazolium and imidazole derivatives.⁶⁶ We identified a new compound, diisopropylphenyl-imidazole (DII), that is lethal to *C. elegans* through a novel mode of action that includes differential targeting in larvae and adult nematodes. This lethal effect appears to be specific for nematodes because at DII concentrations, proven to be toxic to *C. elegans*, no significant lethality on bacteria, *Drosophila melanogaster* and HEK-293 cells has been detected. Using *C. elegans* mutant and transgenic strains, we found that DII effects on adult nematodes rely on a previously unidentified UNC-29-containing muscle AChR, different from the classical Levamisole-sensitive AChR. Interestingly, DII targets appear to be different between larvae and adults as *unc-29* null mutant

larvae are sensitive to the drug.⁶⁶ Summing up, using the model *C. elegans* we demonstrated that DII fulfills the major criteria necessary for the development of a novel anthelmintic, namely phylogenetic specificity and a novel biochemical mode of action. The next step will be to expose parasite worms to this already characterized drug.

Other researchers performed a screen of a small-molecule library using *C. elegans* to test the anthelmintic activity of FDA-approved drugs with the aim of repurposing their clinical activity.²²⁶ The availability of data on their toxicity and pharmacokinetic characteristics will expedite its potential use for new therapeutic indications. From this screen, they found that the neuromodulatory drugs sertraline, paroxetine, and chlorpromazine kill *C. elegans* at multiple life stages and inhibit worm feeding. *C. elegans* mutants with resistance to known anthelmintic drugs are as sensitive as wild-type worms to these three drugs, suggesting that they may act through novel targets. They also demonstrated that these drugs affect divergent parasitic helminth species, such as *Trichuris muris* and *Ancylostoma caninum* and the trematode *Schistosoma mansoni*. These researchers therefore conclude that these drugs may represent a new class of anthelmintic drugs that could be used in combination with classic anthelmintics to boost effectiveness as well as to avoid parasite drug resistance.

As a result of the limited repertoire of available anthelmintics and the alarming emergence of anthelmintic resistance, there is an urgent need to develop novel anthelmintics. *C. elegans* becomes a useful platform to accelerate the screening process for anthelmintic drug discovery. The use of *C. elegans* can permit the development of new parasite control strategies by identifying new drugs with better and broad-spectrum, repurpose already approved drugs, and combine different therapeutic options.

4 | CONCLUSIONS

Traditional preclinical drug discovery is a complex process that takes more than 10 years. Today, innovative strategies are required to satisfy both the increasing demand for better and more efficient therapies and the need to save resources and time for drug development. Under this scenario, the use of invertebrate animals, such as the nematode *C. elegans*, during the early phases of drug development becomes a very convenient strategy to achieve these two aims. Since the introduction of *C. elegans* into research studies, this nematode has been extensively used to understand fundamental biological processes. During the last years, *C. elegans* has also become an invaluable tool for the screening of compounds with potential therapeutic uses. This nematode is probably the best cost-effective choice for early target validation scheme among animal models. Primary hits could be identified in *C. elegans* as an initial filter. The efficacy and safety information gathered by the use of this nematode, significantly reduce costs, time, and the number of vertebrate animals required.

In this review, we have outlined several attributes that highlight the strength of *C. elegans* as a model for drug screening, drug repurposing, drug target identification, drug combination,

and molecular drug mechanism unraveling. The advances in massive and automated high-throughput drug discovery assays for *C. elegans*, the amenability of genetic manipulation, and the extended information available on gene structures, mutant and RNAi phenotypes, microarray data and protein–protein interactions (Wormbase) should be further exploited to facilitate the development of new drugs. The variety of *C. elegans* disease models now available in the drug discovery pipeline does secure a significant boost to our understanding of human diseases and accelerates the reach of effective disease treatments.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

AUTHOR CONTRIBUTIONS

All authors were involved in manuscript preparation for this project.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Diego Rayes  <https://orcid.org/0000-0002-5448-1431>

María José De Rosa  <https://orcid.org/0000-0002-2149-6127>

REFERENCES

- Dillin A, Hsu AL, Arantes-Oliveira N, et al. Rates of behavior and aging specified by mitochondrial function during development. *Science*. 2002;298(5602):2398-2401. <https://doi.org/10.1126/science.1077780>
- Lee SS, Kennedy S, Tolonen AC, Ruvkun G. DAF-16 target genes that control *C. elegans* life-span and metabolism. *Science*. 2003;300(5619):644-647. <https://doi.org/10.1126/science.1083614>
- Nollen EA, Garcia SM, van Haaften G, et al. Genome-wide RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. *Proc Natl Acad Sci USA*. 2004;101(17):6403-6408. <https://doi.org/10.1073/pnas.0307697101>
- Kim Y, Sun H. Functional genomic approach to identify novel genes involved in the regulation of oxidative stress resistance and animal lifespan. *Aging Cell*. 2007;6(4):489-503. <https://doi.org/10.1111/j.1474-9726.2007.00302.x>
- van Ham TJ, Thijssen KL, Breitling R, Hofstra RM, Plasterk RH, Nollen EA. *C. elegans* model identifies genetic modifiers of alpha-synuclein inclusion formation during aging. *PLoS Genet*. 2008;4(3):e1000027. <https://doi.org/10.1371/journal.pgen.1000027>
- Habchi J, Arosio P, Perni M, et al. An anticancer drug suppresses the primary nucleation reaction that initiates the production of the toxic Aβ42 aggregates linked with Alzheimer's disease. *Sci Adv*. 2016;2(2):e1501244. <https://doi.org/10.1126/sciadv.1501244>
- De Rosa MJ, Veuthey T, Florman J, et al. The flight response impairs cytoprotective mechanisms by activating the insulin pathway. *Nature*. 2019;573(7772):135-138. <https://doi.org/10.1038/s41586-019-1524-5>
- Pirri JK, Rayes D, Alkema MJ. A change in the ion selectivity of ligand-gated ion channels provides a mechanism to switch behavior. *PLOS Biol*. 2015;13(9):15. <https://doi.org/10.1371/journal.pbio.1002238>
- Jones AK, Rayes D, Al-Diwani A, et al. A Cys-loop mutation in the *Caenorhabditis elegans* nicotinic receptor subunit UNC-63 impairs but does not abolish channel function. *J Biol Chem*. 2011;286(4):2550-2558. <https://doi.org/10.1074/jbc.M110.177238>
- Hengartner MO, Horvitz HR. Programmed cell death in *Caenorhabditis elegans*. *Curr Opin Genet Dev*. 1994;4(4):581-586. [https://doi.org/10.1016/0959-437X\(94\)90076-F](https://doi.org/10.1016/0959-437X(94)90076-F)
- Chalfie M, Tu Y, Euskirchen G, Ward WW, Prasher DC. Green fluorescent protein as a marker for gene expression. *Science*. 1994;263(5148):802-805. <https://doi.org/10.1126/science.8303295>
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*. 1998;391(6669):806-811. <https://doi.org/10.1038/35888>
- Samuel BS, Rowedder H, Braendle C, Félix M-A, Ruvkun G. *Caenorhabditis elegans*; responses to bacteria from its natural habitats. *Proc Natl Acad Sci*. 2016;113(27):E3941. <https://doi.org/10.1073/pnas.1607183113>
- Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics*. 1974;77(1):71-94.
- Girard LR, Fiedler TJ, Harris TW, et al. WormBook: The online review of *Caenorhabditis elegans* biology. *Nucleic Acids Res*. 2007;35(Database):D472-D475. <https://doi.org/10.1093/nar/gkl894>
- Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant of the nematode *Caenorhabditis elegans*. *Dev Biol*. 1975;46(2):326-342. [https://doi.org/10.1016/0012-1606\(75\)90109-8](https://doi.org/10.1016/0012-1606(75)90109-8)
- Golden JW, Riddle DL. A pheromone influences larval development in the nematode *Caenorhabditis elegans*. *Science*. 1982;218(4572):578-580. <https://doi.org/10.1126/science.6896933>
- Sulston JE, Horvitz HR. Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev Biol*. 1977;56(1):110-156. [https://doi.org/10.1016/0012-1606\(77\)90158-0](https://doi.org/10.1016/0012-1606(77)90158-0)
- Kimble J, Hirsh D. The postembryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. *Dev Biol*. 1979;70(2):396-417. [https://doi.org/10.1016/0012-1606\(79\)90035-6](https://doi.org/10.1016/0012-1606(79)90035-6)
- Sulston JE, Schierenberg E, White JG, Thomson JN. The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev Biol*. 1983;100(1):64-119. [https://doi.org/10.1016/0012-1606\(83\)90201-4](https://doi.org/10.1016/0012-1606(83)90201-4)
- White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philos Trans R Soc Lond B Biol Sci*. 1986;314(1165):1-340.
- Cook SJ, Jarrell TA, Brittin CA, et al. Whole-animal connectomes of both *Caenorhabditis elegans* sexes. *Nature*. 2019;571(7763):63-71. <https://doi.org/10.1038/s41586-019-1352-7>
- Kerr R, Lev-Ram V, Baird G, Vincent P, Tsien RY, Schafer WR. Optical imaging of calcium transients in neurons and pharyngeal muscle of *C. elegans*. *Neuron*. 2000;26(3):583-594. [https://doi.org/10.1016/S0896-6273\(00\)81196-4](https://doi.org/10.1016/S0896-6273(00)81196-4)
- Nagel G, Brauner M, Liewald JF, Adeishvili N, Bamberg E, Gottschalk A. Light activation of channelrhodopsin-2 in excitable cells of *Caenorhabditis elegans* triggers rapid behavioral responses. *Current Biol*. 2005;15(24):2279-2284. <https://doi.org/10.1016/j.cub.2005.11.032>
- Nguyen JP, Shipley FB, Linder AN, et al. Whole-brain calcium imaging with cellular resolution in freely behaving *Caenorhabditis elegans*. *Proc Natl Acad Sci USA*. 2016;113(8):E1074-E1081. <https://doi.org/10.1073/pnas.1507110112>
- Singh J. Harnessing the power of genetics: Fast forward genetics in *Caenorhabditis elegans* [published online ahead of print, 2020

- Sep 4]. *Mol Genet Genomics*. 2020. <https://doi.org/10.1007/s00438-020-01721-6>
27. Jorgensen EM, Mango SE. The art and design of genetic screens: *Caenorhabditis elegans*. *Nat Rev Genet*. 2002;3(5):356-369. <https://doi.org/10.1038/nrg794>
 28. Lai CH, Chou CY, Ch'ang LY, Liu CS, Lin W. Identification of novel human genes evolutionarily conserved in *Caenorhabditis elegans* by comparative proteomics. *Genome Res*. 2000;10(5):703-713. <https://doi.org/10.1101/gr.10.5.703>
 29. Kim W, Underwood RS, Greenwald I, Shaye DD. OrthoList 2: A new comparative genomic analysis of human and *Caenorhabditis elegans* genes. *Genetics*. 2018;210(2):445-461. <https://doi.org/10.1534/genetics.118.301307>
 30. Bargmann CI. Neurobiology of the *Caenorhabditis elegans* genome. *Science*. 1998;282(5396):2028-2033. <https://doi.org/10.1126/science.282.5396.2028>
 31. Das D, Arur S. Conserved insulin signaling in the regulation of oocyte growth, development, and maturation. *Mol Reprod Dev*. 2017;84(6):444-459. <https://doi.org/10.1002/mrd.22806>
 32. Jia K, Levine B. Autophagy is required for dietary restriction-mediated life span extension in *C. elegans*. *Autophagy*. 2007;3(6):597-599. <https://doi.org/10.4161/auto.4989>
 33. Kim DH, Feinbaum R, Alloing G, et al. A conserved p38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science*. 2002;297(5581):623-626. <https://doi.org/10.1126/science.1073759>
 34. Jia K, Chen D, Riddle DL. The TOR pathway interacts with the insulin signaling pathway to regulate *C. elegans* larval development, metabolism and life span. *Development*. 2004;131(16):3897-3906. <https://doi.org/10.1242/dev.01255>
 35. Gosai SJ, Kwak JH, Luke CJ, et al. Automated high-content live animal drug screening using *C. elegans* expressing the aggregation prone serpin α 1-antitrypsin Z. *PLoS One*. 2010;5(11):e15460. <https://doi.org/10.1371/journal.pone.0015460>
 36. Mathew MD, Mathew ND, Ebert PR. WormScan: A technique for high-throughput phenotypic analysis of *Caenorhabditis elegans*. *PLoS One*. 2012;7(3):e33483. <https://doi.org/10.1371/journal.pone.0033483>
 37. Swierczek NA, Giles AC, Rankin CH, Kerr RA. High-throughput behavioral analysis in *C. elegans*. *Nat Methods*. 2011;8(7):592-598. <https://doi.org/10.1038/nmeth.1625>
 38. Xian B, Shen J, Chen W, et al. WormFarm: A quantitative control and measurement device toward automated *Caenorhabditis elegans* aging analysis. *Aging Cell*. 2013;12(3):398-409. <https://doi.org/10.1111/accel.12063>
 39. Partridge FA, Brown AE, Buckingham SD, et al. An automated high-throughput system for phenotypic screening of chemical libraries on *C. elegans* and parasitic nematodes. *Int J Parasitol Drugs Drug Resist*. 2018;8(1):8-21. <https://doi.org/10.1016/j.ijpddr.2017.11.004>
 40. Teo E, Fong S, Tolwinski N, Gruber J. Drug synergy as a strategy for compression of morbidity in a *Caenorhabditis elegans* model of Alzheimer's disease. *GeroScience*. 2020;42(3):849-856. <https://doi.org/10.1007/s11357-020-00169-1>
 41. Wong SQ, Pontifex MG, Phelan MM, et al. α -Methyl- α -phenylsuccinimide ameliorates neurodegeneration in a *C. elegans* model of TDP-43 proteinopathy. *Neurobiol Dis*. 2018;118:40-54. <https://doi.org/10.1016/j.nbd.2018.06.013>
 42. Farina F, Lambert E, Commeau L, et al. The stress response factor daf-16/FOXO is required for multiple compound families to prolong the function of neurons with Huntington's disease. *Sci Rep*. 2017;7(1):4014. <https://doi.org/10.1038/s41598-017-04256-w>
 43. Patten SA, Aggad D, Martinez J, et al. Neuroleptics as therapeutic compounds stabilizing neuromuscular transmission in amyotrophic lateral sclerosis. *JCI Insight*. 2017;2(22):e97152. <https://doi.org/10.1172/jci.insight.97152>
 44. Mondal S, Hegarty E, Martin C, Gökçe SK, Ghorashian N, Ben-Yakar A. Large-scale microfluidics providing high-resolution and high-throughput screening of *Caenorhabditis elegans* polyglutamine aggregation model. *Nat Commun*. 2016;7:13023. <https://doi.org/10.1038/ncomms13023>
 45. Jagota S, Rajadas J. Effect of phenolic compounds against A β aggregation and A β -induced toxicity in transgenic *C. elegans*. *Neurochem Res*. 2012;37(1):40-48. <https://doi.org/10.1007/s11064-011-0580-5>
 46. Lublin A, Isoda F, Patel H, et al. FDA-approved drugs that protect mammalian neurons from glucose toxicity slow aging dependent on Cbp and protect against proteotoxicity. *PLoS One*. 2011;6(11):e27762. <https://doi.org/10.1371/journal.pone.0027762>
 47. Su LJ, Auluck PK, Outeiro TF, et al. Compounds from an unbiased chemical screen reverse both ER-to-Golgi trafficking defects and mitochondrial dysfunction in Parkinson's disease models. *Dis Models Mech*. 2010;3(3-4):194-208. <https://doi.org/10.1242/dmm.004267>
 48. Liu W, Lin H, Mao Z, et al. Verapamil extends lifespan in *Caenorhabditis elegans* by inhibiting calcineurin activity and promoting autophagy. *Aging*. 2020;12(6):5300-5317. <https://doi.org/10.18632/aging.102951>
 49. Zhang P, Zhai Y, Cregg J, Ang KK, Arkin M, Kenyon C. Stress resistance screen in a human primary cell line identifies small molecules that affect aging pathways and extend *Caenorhabditis elegans*' lifespan. *G3 (Bethesda, MD)*. 2020;10(2):849-862. <https://doi.org/10.1534/g3.119.400618>
 50. Chen AL, Lum KM, Lara-Gonzalez P, et al. Pharmacological convergence reveals a lipid pathway that regulates *C. elegans* lifespan. *Nat Chem Biol*. 2019;15(5):453-462. <https://doi.org/10.1038/s41589-019-0243-4>
 51. Bazopoulou D, Chaudhury AR, Pantazis A, Chronis N. An automated compound screening for anti-aging effects on the function of *C. elegans* sensory neurons. *Sci Rep*. 2017;7(1):9403. <https://doi.org/10.1038/s41598-017-09651-x>
 52. Kumar S, Dietrich N, Kornfeld K. Angiotensin Converting Enzyme (ACE) inhibitor extends *Caenorhabditis elegans* life span. *PLoS Genet*. 2016;12(2):e1005866. <https://doi.org/10.1371/journal.pgen.1005866>
 53. Lucanic M, Garrett T, Yu I, et al. Chemical activation of a food deprivation signal extends lifespan. *Aging Cell*. 2016;15(5):832-841. <https://doi.org/10.1111/accel.12492>
 54. Ye X, Linton JM, Schork NJ, Buck LB, Petrascheck M. A pharmacological network for lifespan extension in *Caenorhabditis elegans*. *Aging Cell*. 2014;13(2):206-215. <https://doi.org/10.1111/accel.12163>
 55. Chin RM, Fu X, Pai MY, et al. The metabolite α -ketoglutarate extends lifespan by inhibiting ATP synthase and TOR. *Nature*. 2014;510(7505):397-401. <https://doi.org/10.1038/nature13264>
 56. Hummell NA, Revtovich AV, Kirienko NV. Novel immune modulators enhance *Caenorhabditis elegans* resistance to multiple pathogens. *mSphere*. 2021;6(1):e00950-20. <https://doi.org/10.1101/2020.10.23.352229>
 57. Kim W, Zhu W, Hendricks GL, et al. A new class of synthetic retinoid antibiotics effective against bacterial persisters. *Nature*. 2018;556(7699):103-107. <https://doi.org/10.1038/nature26157>
 58. Kirienko DR, Revtovich AV, High-Content KNVA. Phenotypic screen identifies fluorouridine as an inhibitor of pyoverdine biosynthesis and *Pseudomonas aeruginosa* virulence. *mSphere*. 2016;1(4):e00217-e316. <https://doi.org/10.1128/mSphere.00217-16>
 59. Rajamuthiah R, Jayamani E, Majed H, et al. Antibacterial properties of 3-(phenylsulfonyl)-2-pyrazinocarboxitrile. *Bioorg Med Chem Lett*. 2015;25(22):5203-5207. <https://doi.org/10.1016/j.bmcl.2015.09.066>

60. Rajamuthiah R, Fuchs BB, Jayamani E, et al. Whole animal automated platform for drug discovery against multi-drug resistant *Staphylococcus aureus*. *PLoS One*. 2014;9(2):e89189. <https://doi.org/10.1371/journal.pone.0089189>
61. Kiriienko NV, Kiriienko DR, Larkins-Ford J, Wählby C, Ruvkun G, Ausubel FM. *Pseudomonas aeruginosa* disrupts *Caenorhabditis elegans* iron homeostasis, causing a hypoxic response and death. *Cell Host Microbe*. 2013;13(4):406-416. <https://doi.org/10.1016/j.chom.2013.03.003>
62. Zhou YM, Shao L, Li JA, et al. An efficient and novel screening model for assessing the bioactivity of extracts against multidrug-resistant *Pseudomonas aeruginosa* using *Caenorhabditis elegans*. *Biosci Biotechnol Biochem*. 2011;75(9):1746-1751. <https://doi.org/10.1271/bbb.110290>
63. Coleman JJ, Okoli I, Tegos GP, et al. Characterization of plant-derived saponin natural products against *Candida albicans*. *ACS Chem Biol*. 2010;5(3):321-332. <https://doi.org/10.1021/cb900243b>
64. Milišiuonaitė V, Kadlecová A, Žukauskaitė A, et al. Synthesis and anthelmintic activity of benzopyrano[2,3-c]pyrazol-4(2H)-one derivatives. *Mol Diversity*. 2020;24(4):1025-1042. <https://doi.org/10.1007/s11030-019-10010-3>
65. Risi G, Aguilera E, Ladós E, et al. *Caenorhabditis elegans* infrared-based motility assay identified new hits for nematocide drug development. *Vet Sci*. 2019;6(1):29. <https://doi.org/10.3390/vetsci6010029>
66. Blanco MG, Vela Gurovic MS, Silbestri GF, et al. Diisopropylphenylimidazole (DII): A new compound that exerts anthelmintic activity through novel molecular mechanisms. *PLoS Negl Trop Dis*. 2018;12(12):e0007021. <https://doi.org/10.1371/journal.pntd.0007021>
67. Partridge FA, Murphy EA, Willis NJ, et al. Dihydrobenz[e][1,4]oxazepin-2(3H)-ones, a new anthelmintic chemotype immobilising whipworm and reducing infectivity in vivo. *PLoS Negl Trop Dis*. 2017;11(2):e0005359. <https://doi.org/10.1371/journal.pntd.0005359>
68. Burns AR, Luciani GM, Musso G, et al. *Caenorhabditis elegans* is a useful model for anthelmintic discovery. *Nat Commun*. 2015;6:7485. <https://doi.org/10.1038/ncomms8485>
69. Medina PM, Ponce JM, Cruz CA. Revealing the anticancer potential of candidate drugs in vivo using *Caenorhabditis elegans* mutant strains. *Transl Oncol*. 2020;14(1):100940. <https://doi.org/10.1016/j.tranon.2020.100940>
70. Ji J, Yuan J, Guo X, et al. Harmine suppresses hyper-activated Ras-MAPK pathway by selectively targeting oncogenic mutated Ras/Raf in *Caenorhabditis elegans*. *Cancer Cell Int*. 2019;19:159. <https://doi.org/10.1186/s12935-019-0880-4>
71. Bae YK, Sung JY, Kim YN, et al. An in vivo *C. elegans* model system for screening EGFR-inhibiting anti-cancer drugs. *PLoS One*. 2012;7(9):e42441. <https://doi.org/10.1371/journal.pone.0042441>
72. Zwirchmayr J, Kirchweiger B, Lehner T, Tahir A, Pretsch D, Rollinger JM. A robust and miniaturized screening platform to study natural products affecting metabolism and survival in *Caenorhabditis elegans*. *Sci Rep*. 2020;10(1):12323. <https://doi.org/10.1038/s41598-020-69186-6>
73. Wang J, He J. Swertiamarin decreases lipid accumulation dependent on 3-ketoacyl-coA thiolase. *Biomed Pharmacother*. 2019;112:108668. <https://doi.org/10.1016/j.biopha.2019.108668>
74. Chow YL, Sato F. Screening of isoquinoline alkaloids for potent lipid metabolism modulation with *Caenorhabditis elegans*. *Biosci Biotechnol Biochem*. 2013;77(12):2405-2412. <https://doi.org/10.1271/bbb.130523>
75. Lewis JA, Wu CH, Berg H, Levine JH. The genetics of levamisole resistance in the nematode *Caenorhabditis elegans*. *Genetics*. 1980;95(4):905-928.
76. Lewis JA, Wu CH, Levine JH, Berg H. Levamisole-resistant mutants of the nematode *Caenorhabditis elegans* appear to lack pharmacological acetylcholine receptors. *Neuroscience*. 1980;5(6):967-989.
77. Kaminsky R. A new class of anthelmintics effective against drug-resistant nematodes. *Nature*. 2008;452(7184):176-180. <https://doi.org/10.1038/nature06722>
78. Driscoll M, Dean E, Reilly E, Bergholz E, Chalfie M. Genetic and molecular analysis of a *Caenorhabditis elegans* beta-tubulin that conveys benzimidazole sensitivity. *J Cell Biol*. 1989;109(6 Pt 1):2993-3003.
79. Apfeld J, Alper S. What can we learn about human disease from the nematode *C. elegans*? *Methods Mol Biol*. 2018;1706:53-75. https://doi.org/10.1007/978-1-4939-7471-9_4
80. Gao AW, Uit de Bos J, Sterken MG, Kammenga JE, Smith RL, Houtkooper RH. Forward and reverse genetics approaches to uncover metabolic aging pathways in *Caenorhabditis elegans*. *Biochim Biophys Acta*. 2018;1864(9 Pt A):2697-2706. <https://doi.org/10.1016/j.bbadis.2017.09.006>
81. Serrat X, Kukhtar D, Cornes E, et al. CRISPR editing of sftb-1/SF3B1 in *Caenorhabditis elegans* allows the identification of synthetic interactions with cancer-related mutations and the chemical inhibition of splicing. *PLoS Genet*. 2019;15(10):e1008464. <https://doi.org/10.1371/journal.pgen.1008464>
82. Morley JF, Brignull HR, Weyers JJ, Morimoto RI. The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA*. 2002;99(16):10417-10422. <https://doi.org/10.1073/pnas.152161099>
83. Brignull HR, Moore FE, Tang SJ, Morimoto RI. Polyglutamine proteins at the pathogenic threshold display neuron-specific aggregation in a pan-neuronal *Caenorhabditis elegans* model. *J Neurosci*. 2006;26(29):7597-7606. <https://doi.org/10.1523/jneurosci.0990-06.2006>
84. Teixeira-Castro A, Ailion M, Jalles A, et al. Neuron-specific proteotoxicity of mutant ataxin-3 in *C. elegans*: Rescue by the DAF-16 and HSF-1 pathways. *Hum Mol Genet*. 2011;20(15):2996-3009. <https://doi.org/10.1093/hmg/ddr203>
85. Styer KL, Singh V, Macosko E, Steele SE, Bargmann CI, Aballay A. Innate immunity in *Caenorhabditis elegans* is regulated by neurons expressing NPR-1/GPCR. *Science*. 2008;322(5900):460-464. <https://doi.org/10.1126/science.1163673>
86. Singh V, Aballay A. Regulation of DAF-16-mediated Innate Immunity in *Caenorhabditis elegans*. *J Biol Chem*. 2009;284(51):35580-35587. <https://doi.org/10.1074/jbc.M109.060905>
87. Singh J, Aballay A. Intestinal infection regulates behavior and learning via neuroendocrine signaling. *Elife*. 2019;8:e50033. <https://doi.org/10.7554/eLife.50033>
88. Schlotterer A, Kukudov G, Bozorgmehr F, et al. *C. elegans* as model for the study of high glucose-mediated life span reduction. *Diabetes*. 2009;58(11):2450-2456. <https://doi.org/10.2337/db09-0567>
89. Zhu G, Yin F, Wang L, Wei W, Jiang L, Qin J. Modeling type 2 diabetes-like hyperglycemia in *C. elegans* on a microdevice. *Integr Biol*. 2016;8(1):30-38. <https://doi.org/10.1039/C5IB00243E>
90. Regitz C, Fitzenberger E, Mahn FL, Dußling LM, Wenzel U. Resveratrol reduces amyloid-beta (Aβ₁₋₄₂)-induced paralysis through targeting proteostasis in an Alzheimer model of *Caenorhabditis elegans*. *Eur J Nutr*. 2016;55(2):741-747. <https://doi.org/10.1007/s00394-015-0894-1>
91. Solis GM, Kardakaris R, Valentine ER, et al. Translation attenuation by minocycline enhances longevity and proteostasis in old post-stress-responsive organisms. *Elife*. 2018;7:e40314. <https://doi.org/10.7554/eLife.40314>
92. Pukkila-Worley R, Ausubel FM. Immune defense mechanisms in the *Caenorhabditis elegans* intestinal epithelium. *Curr Opin Immunol*. 2012;24(1):3-9. <https://doi.org/10.1016/j.coi.2011.10.004>

93. Baur R, Beech R, Sigel E, Rufener L. Monepantel irreversibly binds to and opens *Haemonchus contortus* MPTL-1 and *Caenorhabditis elegans* ACR-20 receptors. *Mol Pharmacol*. 2015;87(1):96-102. <https://doi.org/10.1124/mol.114.095653>
94. Wu L, Zhou B, Oshiro-Rapley N, et al. An ancient, unified mechanism for metformin growth inhibition in *C. elegans* and cancer. *Cell*. 2016;167(7):1705-1718.e13. <https://doi.org/10.1016/j.cell.2016.11.055>
95. Peng H, Wei Z, Luo H, et al. Inhibition of fat accumulation by hesperidin in *Caenorhabditis elegans*. *J Agr Food Chem*. 2016;64(25):5207-5214. <https://doi.org/10.1021/acs.jafc.6b02183>
96. Agotegaray M, Blanco MG, Campelo A, et al. β -cyclodextrin coating: Improving biocompatibility of magnetic nanocomposites for biomedical applications. *J Mater Sci - Mater Med*. 2020;31(2):22. <https://doi.org/10.1007/s10856-020-6361-4>
97. Xiong H, Pears C, Woollard A. An enhanced *C. elegans* based platform for toxicity assessment. *Sci Rep*. 2017;7(1):9839. <https://doi.org/10.1038/s41598-017-10454-3>
98. Chiu HF, Hsiao SC, Lu YY, et al. Efficacy of protein rich pearl powder on antioxidant status in a randomized placebo-controlled trial. *J Food Drug Anal*. 2018;26(1):309-317. <https://doi.org/10.1016/j.jfda.2017.05.010>
99. Wang Z, Sherwood DR. Dissection of genetic pathways in *C. elegans*. *Methods Cell Biol*. 2011;106:113-157. <https://doi.org/10.1016/b978-0-12-544172-8.00005-0>
100. Lehrbach NJ, Ji F, Sadreyev R. Next-generation sequencing for identification of EMS-induced mutations in *Caenorhabditis elegans*. *Curr Protoc Mol Biol*. 2017;117(7):29.1-7.29.12. <https://doi.org/10.1002/cpmb.27>
101. Doitsidou M, Jarriault S, Poole RJ. Next-generation sequencing-based approaches for mutation mapping and identification in *Caenorhabditis elegans*. *Genetics*. 2016;204(2):451-474. <https://doi.org/10.1534/genetics.115.186197>
102. Anderson P. Mutagenesis. *Methods Cell Biol*. 1995;48:31-58.
103. Kutscher LM, Shaham S. Forward and reverse mutagenesis in *C. elegans*. *WormBook*. 2014:1-26. <https://doi.org/10.1895/wormbook.1.167.1>
104. Huang YC, Pirri JK, Rayes D, et al. Gain-of-function mutations in the UNC-2/CaV2alpha channel lead to excitation-dominant synaptic transmission in *Caenorhabditis elegans*. *Elife*. 2019;8:e45905. <https://doi.org/10.7554/eLife.45905>
105. Nass R, Hahn MK, Jessen T, McDonald PW, Carvelli L, Blakely RD. A genetic screen in *Caenorhabditis elegans* for dopamine neuron insensitivity to 6-hydroxydopamine identifies dopamine transporter mutants impacting transporter biosynthesis and trafficking. *J Neurochem*. 2005;94(3):774-785. <https://doi.org/10.1111/j.1471-4159.2005.03205.x>
106. Liu J, Banskota AH, Critchley AT, Hafting J, Prithiviraj B. Neuroprotective effects of the cultivated *Chondrus crispus* in a *C. elegans* model of Parkinson's disease. *Marine Drugs*. 2015;13(4):2250-2266. <https://doi.org/10.3390/md13042250>
107. Mathew MD, Mathew ND, Miller A, et al. Using *C. elegans* forward and reverse genetics to identify new compounds with anthelmintic activity. *PLoS Negl Trop Dis*. 2016;10(10):e0005058. <https://doi.org/10.1371/journal.pntd.0005058>
108. Dent JA, Smith MM, Vassiliadis DK, Avery L. The genetics of ivermectin resistance in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA*. 2000;97(6):2674-2679
109. Kwok TCY, Hui K, Kostecki W, et al. A genetic screen for dihydropyridine (dhp)-resistant worms reveals new residues required for DHP-blockage of mammalian calcium channels. *PLoS Genet*. 2008;4(5):e1000067. <https://doi.org/10.1371/journal.pgen.1000067>
110. Ung H, Hall R, Kikis E. Chemical mutagenesis of *Caenorhabditis elegans* uncovers genetic modifiers of huntingtin protein aggregation. *Micro Publ Biol*. 2020;2020:10.17912/micropub.biology.000202. <https://doi.org/10.17912/micropub.biology.000202>
111. Kawamura K, Maruyama IN. Forward genetic screen for *Caenorhabditis elegans* mutants with a shortened locomotor healthspan. *G3 (Bethesda, Md)*. 2019;9(8):2415-2423. <https://doi.org/10.1534/g3.119.400241>
112. Sheng M, Hosseinzadeh A, Muralidharan SV, Gaur R, Selstam E, Tuck S. Aberrant fat metabolism in *Caenorhabditis elegans* mutants with defects in the defecation motor program. *PLoS One*. 2015;10(4):e0124515. <https://doi.org/10.1371/journal.pone.0124515>
113. Heigwer F, Port F, Boutros M. RNA interference (RNAi) screening in *Drosophila*. *Genetics*. 2018;208(3):853-874. <https://doi.org/10.1534/genetics.117.300077>
114. Yang X, Niu L, Zhang W, et al. Robust RNAi-mediated resistance to infection of seven potyvirids in soybean expressing an intron hairpin Nib RNA. *Transgenic Res*. 2017;26(5):665-676. <https://doi.org/10.1007/s11248-017-0041-2>
115. Rytlewski JA, Beronja S. RNAi in the mouse: rapid and affordable gene function studies in a vertebrate system. *WIREs*. 2015;4(1):45-57. <https://doi.org/10.1002/wdev.164>
116. Kamath RS, Ahringer J. Genome-wide RNAi screening in *Caenorhabditis elegans*. *Methods*. 2003;30(4):313-321. [https://doi.org/10.1016/s1046-2023\(03\)00050-1](https://doi.org/10.1016/s1046-2023(03)00050-1)
117. Rual JF, Ceron J, Koreth J, et al. Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. *Genome Res*. 2004;14(10b):2162-2168. <https://doi.org/10.1101/gr.2505604>
118. Ashrafi K, Chang FY, Watts JL, et al. Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature*. 2003;421(6920):268-272. <https://doi.org/10.1038/nature01279>
119. Gönczy P, Echeverri C, Oegema K, et al. Functional genomic analysis of cell division in *C. elegans* using RNAi of genes on chromosome III. *Nature*. 2000;408(6810):331-336. <https://doi.org/10.1038/35042526>
120. Zugasti O, Thakur N, Belougne J, et al. A quantitative genome-wide RNAi screen in *C. elegans* for antifungal innate immunity genes. *BMC Biol*. 2016;14:35. <https://doi.org/10.1186/s12915-016-0256-3>
121. Mao Y-Q, Han S-F, Zhang S-L, et al. An approach using *Caenorhabditis elegans* screening novel targets to suppress tumour cell proliferation. *Cell Prolif*. 2020;53(6):e12832. <https://doi.org/10.1111/cpr.12832>
122. Liachko NF, Saxton AD, McMillan PJ, et al. Genome wide analysis reveals heparan sulfate epimerase modulates TDP-43 proteinopathy. *PLoS Genet*. 2019;15(12):e1008526. <https://doi.org/10.1371/journal.pgen.1008526>
123. Calixto A, Chelur D, Topalidou I, Chen X, Chalfie M. Enhanced neuronal RNAi in *C. elegans* using SID-1. *Nat Methods*. 2010;7(7):554-559. <https://doi.org/10.1038/nmeth.1463>
124. Kennedy S, Wang D, Ruvkun G. A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*. *Nature*. 2004;427(6975):645-649. <https://doi.org/10.1038/nature02302>
125. Wang D, Kennedy S, Conte D Jr, et al. Somatic misexpression of germline P granules and enhanced RNA interference in retinoblastoma pathway mutants. *Nature*. 2005;436(7050):593-597. <https://doi.org/10.1038/nature04010>
126. Sieburth D, Ch'ng Q, Dybbs M, et al. Systematic analysis of genes required for synapse structure and function. *Nature*. 2005;436(7050):510-517. <https://doi.org/10.1038/nature03809>
127. Hsu PD, Lander ES, Zhang F. Development and applications of CRISPR-Cas9 for genome engineering. *Cell*. 2014;157(6):1262-1278. <https://doi.org/10.1016/j.cell.2014.05.010>

128. Maggio I, Gonçalves MA. Genome editing at the crossroads of delivery, specificity, and fidelity. *Trends Biotechnol.* 2015;33(5):280-291. <https://doi.org/10.1016/j.tibtech.2015.02.011>
129. Frøkjær-Jensen C. Transposon-assisted genetic engineering with Mos1-Mediated Single-Copy Insertion (MosSCI). *Methods Mol Biol.* 2015;1327:49-58. https://doi.org/10.1007/978-1-4939-2842-2_5
130. Dickinson DJ, Goldstein B. CRISPR-based methods for *Caenorhabditis elegans* genome engineering. *Genetics.* 2016;202(3):885-901. <https://doi.org/10.1534/genetics.115.182162>
131. Nance J, Frøkjær-Jensen C. The *Caenorhabditis elegans* transgenic toolbox. *Genetics.* 2019;212(4):959-990. <https://doi.org/10.1534/genetics.119.301506>
132. Baiga TJ, Guo H, Xing Y, et al. Metabolite induction of *Caenorhabditis elegans* dauer larvae arises via transport in the pharynx. *ACS Chem Biol.* 2008;3(5):294-304. <https://doi.org/10.1021/cb700269e>
133. Kamal M, Moshiri H, Magomedova L, et al. The marginal cells of the *Caenorhabditis elegans* pharynx scavenge cholesterol and other hydrophobic small molecules. *Nat Commun.* 2019;10(1):3938. <https://doi.org/10.1038/s41467-019-11908-0>
134. Scharf A, Piechulek A, von Mikecz A. Effect of nanoparticles on the biochemical and behavioral aging phenotype of the nematode *Caenorhabditis elegans*. *ACS Nano.* 2013;7(12):10695-10703. <https://doi.org/10.1021/nn403443r>
135. Page AP. The sensory amphidial structures of *Caenorhabditis elegans* are involved in macrocyclic lactone uptake and anthelmintic resistance. *Int J Parasitol.* 2018;48(13):1035-1042. <https://doi.org/10.1016/j.ijpara.2018.06.003>
136. Kwok TC, Ricker N, Fraser R, et al. A small-molecule screen in *C. elegans* yields a new calcium channel antagonist. *Nature.* 2006;441(7089):91-95. <https://doi.org/10.1038/nature04657>
137. Hu Y, Xiao SH, Aroian RV. The new anthelmintic tribendimidine is an L-type (levamisole and pyrantel) nicotinic acetylcholine receptor agonist. *PLoS Negl Trop Dis.* 2009;3(8):e499. <https://doi.org/10.1371/journal.pntd.0000499>
138. Tam ZY, Gruber J, Ng LF, Halliwell B, Gunawan R. Effects of lithium on age-related decline in mitochondrial turnover and function in *Caenorhabditis elegans*. *J Gerontol: Series A.* 2014;69(7):810-820. <https://doi.org/10.1093/gerona/glt210>
139. van Haaften G, Vastenhouw NL, Nollen EAA, Plasterk RHA, Tijsterman M. Gene interactions in the DNA damage-response pathway identified by genome-wide RNA-interference analysis of synthetic lethality. *Proc Natl Acad Sci USA.* 2004;101(35):12992. <https://doi.org/10.1073/pnas.0403131101>
140. Moy TI, Ball AR, Anklesaria Z, Casadei G, Lewis K, Ausubel FM. Identification of novel antimicrobials using a live-animal infection model. *Proc Natl Acad Sci USA.* 2006;103(27):10414-10419. <https://doi.org/10.1073/pnas.0604055103>
141. Ben-Yakar A, Bourgeois F. Ultrafast laser nanosurgery in microfluidics for genome-wide screenings. *Curr Opin Biotechnol.* 2009;20(1):100-105. <https://doi.org/10.1016/j.copbio.2009.01.008>
142. Zheng SQ, Ding AJ, Li GP, Wu GS, Luo HR. Drug absorption efficiency in *Caenorhabditis elegans* delivered by different methods. *PLoS One.* 2013;8(2):e56877. <https://doi.org/10.1371/journal.pone.0056877>
143. Edwards C, Canfield J, Copes N, Rehan M, Lipps D, Bradshaw PC. D-beta-hydroxybutyrate extends lifespan in *C. elegans*. *Aging.* 2014;6(8):621-644. <https://doi.org/10.18632/aging.100683>
144. Kim DH. Bacteria and the aging and longevity of *Caenorhabditis elegans*. *Annu Rev Genet.* 2013;47:233-246. <https://doi.org/10.1146/annurev-genet-111212-133352>
145. Cabreiro F, Gems D. Worms need microbes too: Microbiota, health and aging in *Caenorhabditis elegans*. *EMBO Mol Med.* 2013;5(9):1300-1310. <https://doi.org/10.1002/emmm.20110.0972>
146. Burns AR, Wallace IM, Wildenhain J, et al. A predictive model for drug bioaccumulation and bioactivity in *Caenorhabditis elegans*. *Nat Chem Biol.* 2010;6(7):549-557. <https://doi.org/10.1038/nchembio.380>
147. van Dijk FS, Cobben JM, Kariminejad A, et al. Osteogenesis imperfecta: A review with clinical examples. *Molecular Syndromology.* 2011;2(1):1-20. <https://doi.org/10.1159/000332228>
148. Johnstone IL, Shafi Y, Barry JD. Molecular analysis of mutations in the *Caenorhabditis elegans* collagen gene dpy-7. *Embo J.* 1992;11(11):3857-3863.
149. McColl G, Roberts BR, Pukala TL, et al. Utility of an improved model of amyloid-beta (A β ₁₋₄₂) toxicity in *Caenorhabditis elegans* for drug screening for Alzheimer's disease. *Mol Neurodegener.* 2012;7:57. <https://doi.org/10.1186/1750-1326-7-57>
150. Vartiainen S, Aarnio V, Lakso M, Wong G. Increased lifespan in transgenic *Caenorhabditis elegans* overexpressing human alpha-synuclein. *Exp Gerontol.* 2006;41(9):871-876. <https://doi.org/10.1016/j.exger.2006.05.005>
151. Perlmutter DH. The role of autophagy in alpha-1-antitrypsin deficiency: A specific cellular response in genetic diseases associated with aggregation-prone proteins. *Autophagy.* 2006;2(4):258-263. <https://doi.org/10.4161/auto.2882>
152. Silverman G, Pak S, Perlmutter D. Disorders of protein misfolding: Alpha-1-antitrypsin deficiency as prototype. *J Pediatr.* 2013;163(2):320-326. <https://doi.org/10.1016/j.jpeds.2013.03.077>
153. Gitler AD, Dhillon P, Shorter J. Neurodegenerative disease: Models, mechanisms, and a new hope. *Dis Models Mech.* 2017;10(5):499. <https://doi.org/10.1242/dmm.030205>
154. Mattson MP, Magnus T. Ageing and neuronal vulnerability. *Nat Rev Neurosci.* 2006;7(4):278-294. <https://doi.org/10.1038/nrn1886>
155. Chen X, Barclay JW, Burgoyne RD, Morgan A, et al. Using *C. elegans* to discover therapeutic compounds for ageing-associated neurodegenerative diseases. *Chemistry Central Journal.* 2015;9:65. <https://doi.org/10.1186/s13065-015-0143-y>
156. Ma L, Zhao Y, Chen Y, Cheng B, Peng A, Huang K. *Caenorhabditis elegans* as a model system for target identification and drug screening against neurodegenerative diseases. *Eur J Pharmacol.* 2018;819:169-180. <https://doi.org/10.1016/j.ejphar.2017.11.051>
157. Brignull HR, Morley JF, Garcia SM, Morimoto RI. Modeling polyglutamine pathogenesis in *C. elegans*. *Methods Enzymol.* 2006;412:256-282. [https://doi.org/10.1016/S0076-6879\(06\)12016-9](https://doi.org/10.1016/S0076-6879(06)12016-9)
158. Lakso M, Vartiainen S, Moilanen AM, et al. Dopaminergic neuronal loss and motor deficits in *Caenorhabditis elegans* overexpressing human alpha-synuclein. *J Neurochem.* 2003;86(1):165-172.
159. Hamamichi S, Rivas RN, Knight AL, Cao S, Caldwell KA, Caldwell GA. Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proc Natl Acad Sci USA.* 2008;105(2):728-733. <https://doi.org/10.1073/pnas.0711018105>
160. Saldi TK, Ash PE, Wilson G, et al. TDP-1, the *Caenorhabditis elegans* ortholog of TDP-43, limits the accumulation of double-stranded RNA. *EMBO J.* 2014;33(24):2947-2966. <https://doi.org/10.15252/embj.201488740>
161. Yao C, Johnson WM, Gao Y, et al. Kinase inhibitors arrest neurodegeneration in cell and *C. elegans* models of LRRK2 toxicity. *Hum Mol Genet.* 2013;22(2):328-344. <https://doi.org/10.1093/hmg/dds431>
162. Corsi AK, Wightman B, Chalfie M. A transparent window into biology: A primer on *Caenorhabditis elegans*. *Genetics.* 2015;200(2):387-407. <https://doi.org/10.1534/genetics.115.176099>
163. Samuelson AV, Carr CE, Ruvkun G. Gene activities that mediate increased life span of *C. elegans* insulin-like signaling mutants. *Genes Dev.* 2007;21(22):2976-2994. <https://doi.org/10.1101/gad.1588907>
164. Uno M, Nishida E. Lifespan-regulating genes in *C. elegans*. *NPJ Aging Mech Dis.* 2016;2:16010. <https://doi.org/10.1038/npjamd.2016.10>
165. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang RA. A *C. elegans* mutant that lives twice as long as wild type. *Nature.* 1993;366(6454):461-464. <https://doi.org/10.1038/366461a0>

166. Apfeld J, Kenyon C. Regulation of lifespan by sensory perception in *Caenorhabditis elegans*. *Nature*. 1999;402(6763):804-809. <https://doi.org/10.1038/45544>
167. Kinser HE, Pincus Z. High-throughput screening in the *C. elegans* nervous system. *Mol Cell Neurosci*. 2017;80:192-197. <https://doi.org/10.1016/j.mcn.2016.06.001>
168. O'Reilly LP, Luke CJ, Perlmutter DH, Silverman GA, Pak SC. *C. elegans* in high-throughput drug discovery. *Adv Drug Del Rev*. 2014;70:247-253. <https://doi.org/10.1016/j.addr.2013.12.001>
169. Conery AL, Larkins-Ford J, Ausubel FM, Kirienko NV. High-throughput screening for novel anti-infectives using a *C. elegans* pathogenesis model. *Curr Protoc Chem Biol*. 2014;6(1):25-37. <https://doi.org/10.1002/9780470559277.ch130160>
170. Leung CK, Wang Y, Malany S, et al. An ultra high-throughput, whole-animal screen for small molecule modulators of a specific genetic pathway in *Caenorhabditis elegans*. *PLoS One*. 2013;8(4):e62166. <https://doi.org/10.1371/journal.pone.0062166>
171. Petrascheck M, Ye X, Buck LB. An antidepressant that extends lifespan in adult *Caenorhabditis elegans*. *Nature*. 2007;450(7169):553-556. <https://doi.org/10.1038/nature05991>
172. Perni M, Casford S, Aprile FA, et al. Automated behavioral analysis of large *C. elegans* populations using a wide field-of-view tracking platform. *J Vis Exp*. 2018;(141):e58643. <https://doi.org/10.3791/58643>
173. Perni M, Challa PK, Kirkegaard JB, et al. Massively parallel *C. elegans* tracking provides multi-dimensional fingerprints for phenotypic discovery. *J Neurosci Methods*. 2018;306:57-67. <https://doi.org/10.1016/j.jneumeth.2018.02.005>
174. Bargmann CI. Neurobiology of the *Caenorhabditis elegans* genome. *Science*. 1998;282(5396):2028-2033. <https://doi.org/10.1126/science.282.5396.2028>
175. Liu Q, Kidd PB, Dobosiewicz M, Bargmann CI. *C. elegans* AWA olfactory neurons fire calcium-mediated all-or-none action potentials. *Cell*. 2018;175(1):57-70 e17. <https://doi.org/10.1016/j.cell.2018.08.018>
176. Liu P, Ge Q, Chen B, Salkoff L, Kotlikoff MI, Wang ZW. Genetic dissection of ion currents underlying all-or-none action potentials in *C. elegans* body-wall muscle cells. *J Physiol*. 2011;589(Pt 1):101-117. <https://doi.org/10.1113/jphysiol.2010.200683>
177. Shtonda B, Avery L. CCA-1, EGL-19 and EXP-2 currents shape action potentials in the *Caenorhabditis elegans* pharynx. *J Exp Biol*. 2005;208(11):2177-2190. <https://doi.org/10.1242/jeb.01615>
178. Link CD. Expression of human beta-amyloid peptide in transgenic *Caenorhabditis elegans*. *Proc Natl Acad Sci USA*. 1995;92(20):9368-9372. <https://doi.org/10.1073/pnas.92.20.9368>
179. Dickinson DJ, Ward JD, Reiner DJ, Goldstein B. Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat Methods*. 2013;10(10):1028-1034. <https://doi.org/10.1038/nmeth.2641>
180. Baskoylu SN, Yersak J, O'Hern P, et al. Single copy/knock-in models of ALS SOD1 in *C. elegans* suggest loss and gain of function have different contributions to cholinergic and glutamatergic neurodegeneration. *PLoS Genet*. 2018;14(10):e1007682. <https://doi.org/10.1371/journal.pgen.1007682>
181. Baskoylu SN, Chapkis N, Unsal B, et al. Disrupted autophagy and neuronal dysfunction in *C. elegans* knock-in models of FUS amyotrophic lateral sclerosis. *bioRxiv*. 2019:799932. <https://doi.org/10.1101/799932>
182. Guha S, Fischer S, Johnson GVW, Nehrke K. Tauopathy-associated tau modifications selectively impact neurodegeneration and mitophagy in a novel *C. elegans* single-copy transgenic model. *Mol Neurodegeneration*. 2020;15:65. <https://doi.org/10.1186/s13024-02000410-7>
183. Lee AL, Ung HM, Sands LP, Kikis EA. A new *Caenorhabditis elegans* model of human huntingtin 513 aggregation and toxicity in body wall muscles. *PLoS One*. 2017;12(3):e0173644. <https://doi.org/10.1371/journal.pone.0173644>
184. Faber PW, Alter JR, MacDonald ME, Hart AC. Polyglutamine-mediated dysfunction and apoptotic death of a *Caenorhabditis elegans* sensory neuron. *Proc Natl Acad Sci USA*. 1999;96(1):179-184. <https://doi.org/10.1073/pnas.96.1.179>
185. Parker JA, Connolly JB, Wellington C, Hayden M, Dausset J, Neri C. Expanded polyglutamines in *Caenorhabditis elegans* cause axonal abnormalities and severe dysfunction of PLM mechanosensory neurons without cell death. *Proc Natl Acad Sci USA*. 2001;98(23):13318-13323. <https://doi.org/10.1073/pnas.231476398>
186. Drake J, Link CD, Butterfield DA. Oxidative stress precedes fibrillar deposition of Alzheimer's disease amyloid beta-peptide (1-42) in a transgenic *Caenorhabditis elegans* model. *Neurobiol Aging*. 2003;24(3):415-420. [https://doi.org/10.1016/s0197-4580\(02\)00225-7](https://doi.org/10.1016/s0197-4580(02)00225-7)
187. Fong S, Teo E, Ng LF, et al. Energy crisis precedes global metabolic failure in a novel *Caenorhabditis elegans* Alzheimer disease model. *Sci Rep*. 2016;6(1):33781. <https://doi.org/10.1038/srep33781>
188. Wu Y, Wu Z, Butko P, et al. Amyloid-beta-induced pathological behaviors are suppressed by Ginkgo biloba extract EGb 761 and ginkgolides in transgenic *Caenorhabditis elegans*. *J Neurosci*. 2006;26(50):13102-13113. <https://doi.org/10.1523/jneurosci.3448-06.2006>
189. Kraemer BC, Zhang B, Leverenz JB, Thomas JH, Trojanowski JQ, Schellenberg GD. Neurodegeneration and defective neurotransmission in a *Caenorhabditis elegans* model of tauopathy. *Proc Natl Acad Sci USA*. 2003;100(17):9980-9985. <https://doi.org/10.1073/pnas.1533448100>
190. Guha S, Fischer S, Johnson GVW, Nehrke K. Tauopathy-associated tau modifications selectively impact neurodegeneration and mitophagy in a novel *C. elegans* single-copy transgenic model. *Mol Neurodegener*. 2020;15(1):65. <https://doi.org/10.1186/s13024-020-00410-7>
191. Kuwahara T, Koyama A, Koyama S, et al. A systematic RNAi screen reveals involvement of endocytic pathway in neuronal dysfunction in alpha-synuclein transgenic *C. elegans*. *Hum Mol Genet*. 2008;17(19):2997-3009. <https://doi.org/10.1093/hmg/ddn198>
192. Kuwahara T, Koyama A, Gengyo-Ando K, et al. Familial Parkinson mutant alpha-synuclein causes dopamine neuron dysfunction in transgenic *Caenorhabditis elegans*. *J Biol Chem*. 2006;281(1):334-340. <https://doi.org/10.1074/jbc.M504860200>
193. Nass R, Hall DH, Miller DM 3rd, Blakely RD. Neurotoxin-induced degeneration of dopamine neurons in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA*. 2002;99(5):3264-3269. <https://doi.org/10.1073/pnas.042497999>
194. Oeda T, Shimohama S, Kitagawa N, et al. Oxidative stress causes abnormal accumulation of familial amyotrophic lateral sclerosis-related mutant SOD1 in transgenic *Caenorhabditis elegans*. *Hum Mol Genet*. 2001;10(19):2013-2023. <https://doi.org/10.1093/hmg/10.19.2013>
195. Gidalevitz T, Krupinski T, Garcia S, Morimoto RI. Destabilizing protein polymorphisms in the genetic background direct phenotypic expression of mutant SOD1 toxicity. *PLoS Genet*. 2009;5(3):e1000399. <https://doi.org/10.1371/journal.pgen.1000399>
196. Wang J, Farr GW, Hall DH, et al. An ALS-linked mutant SOD1 produces a locomotor defect associated with aggregation and synaptic dysfunction when expressed in neurons of *Caenorhabditis elegans*. *PLoS Genet*. 2009;5(1):e1000350. <https://doi.org/10.1371/journal.pgen.1000350>
197. Liachko NF, Guthrie CR, Kraemer BC. Phosphorylation promotes neurotoxicity in a *Caenorhabditis elegans* model of TDP-43 proteinopathy. *J Neurosci*. 2010;30(48):16208-16219. <https://doi.org/10.1523/jneurosci.2911-10.2010>
198. Ash PE, Zhang YJ, Roberts CM, et al. Neurotoxic effects of TDP-43 overexpression in *C. elegans*. *Hum Mol Genet*. 2010;19(16):3206-3218. <https://doi.org/10.1093/hmg/ddq230>
199. Vaccaro A, Tauffenberger A, Ash PE, Carlomagno Y, Petrucelli L, Parker JA. TDP-1/TDP-43 regulates stress signaling and

- age-dependent proteotoxicity in *Caenorhabditis elegans*. *PLoS Genet*. 2012;8(7):e1002806. <https://doi.org/10.1371/journal.pgen.1002806>
200. Murakami T, Yang SP, Xie L, et al. ALS mutations in FUS cause neuronal dysfunction and death in *Caenorhabditis elegans* by a dominant gain-of-function mechanism. *Hum Mol Genet*. 2012;21(1):1-9. <https://doi.org/10.1093/hmg/ddr417>
 201. Witan H, Kern A, Koziollek-Drechsler I, Wade R, Behl C, Clement AM. Heterodimer formation of wild-type and amyotrophic lateral sclerosis-causing mutant Cu/Zn-superoxide dismutase induces toxicity independent of protein aggregation. *Hum Mol Genet*. 2008;17(10):1373-1385. <https://doi.org/10.1093/hmg/ddn025>
 202. van der Goot AT, Zhu W, Vazquez-Manrique RP, et al. Delaying aging and the aging-associated decline in protein homeostasis by inhibition of tryptophan degradation. *Proc Natl Acad Sci USA*. 2012;109(37):14912-14917. <https://doi.org/10.1073/pnas.1203083109>
 203. Jadiya P, Fatima S, Baghel T, Mir SS, Nazir A. A systematic RNAi screen of neuroprotective genes identifies novel modulators of alpha-synuclein-associated effects in transgenic *Caenorhabditis elegans*. *Mol Neurobiol*. 2016;53(9):6288-6300. <https://doi.org/10.1007/s12035-015-9517-3>
 204. Knight Adam L, Yan X, Hamamichi S, et al. The glycolytic enzyme, GPI, is a functionally conserved modifier of dopaminergic neurodegeneration in Parkinson's models. *Cell Metab*. 2014;20(1):145-157. <https://doi.org/10.1016/j.cmet.2014.04.017>
 205. Silva MC, Fox S, Beam M, Thakkar H, Amaral MD, Morimoto RI. A genetic screening strategy identifies novel regulators of the proteostasis network. *PLoS Genet*. 2011;7(12):e1002438. <https://doi.org/10.1371/journal.pgen.1002438>
 206. van Ham TJ, Holmberg MA, van der Goot AT, et al. Identification of MOAG-4/SERF as a regulator of age-related proteotoxicity. *Cell*. 2010;142(4):601-612. <https://doi.org/10.1016/j.cell.2010.07.020>
 207. Holden-Dye L, Walker RJ. Anthelmintic drugs and nematocides: Studies in *Caenorhabditis elegans*. *WormBook*. 2014;1-29. <https://doi.org/10.1895/wormbook.1.143.2>
 208. Weng HB, Chen HX, Wang MW. Innovation in neglected tropical disease drug discovery and development. *Infect Dis Poverty*. 2018;7(1):67. <https://doi.org/10.1186/s40249-018-0444-1>
 209. Wolstenholme AJ, Fairweather I, Prichard R, von Samson-Himmelstjerna G, Sangster NC. Drug resistance in veterinary helminths. *Trends Parasitol*. 2004;20(10):469-476. <https://doi.org/10.1016/j.pt.2004.07.010>
 210. Osei-Atweneboana MY, Eng JK, Boakye DA, Gyapong JO, Prichard RK. Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of ivermectin in endemic communities in Ghana: A two-phase epidemiological study. *Lancet*. 2007;369(9578):2021-2029. [https://doi.org/10.1016/S0140-6736\(07\)60942-8](https://doi.org/10.1016/S0140-6736(07)60942-8)
 211. Kaminsky R, Bapst B, Stein PA, et al. Differences in efficacy of monepantel, derquantel and abamectin against multi-resistant nematodes of sheep. *Parasitol Res*. 2011;109(1):19-23. <https://doi.org/10.1007/s00436-010-2216-0>
 212. Geurden T, Hoste H, Jacquiet P, et al. Anthelmintic resistance and multidrug resistance in sheep gastro-intestinal nematodes in France, Greece and Italy. *Vet Parasitol*. 2014;201(1-2):59-66. <https://doi.org/10.1016/j.vetpar.2014.01.016>
 213. Krucken J, Fraundorfer K, Mugisha JC, et al. Reduced efficacy of albendazole against *Ascaris lumbricoides* in Rwandan schoolchildren. *Int J Parasitol Drugs Drug Resist*. 2017;7(3):262-271. <https://doi.org/10.1016/j.ijpddr.2017.06.001>
 214. Ward JD. Rendering the intractable more tractable: tools from *caenorhabditis elegans* ripe for import into parasitic nematodes. *Genetics*. 2015;201(4):1279-1294. <https://doi.org/10.1534/genetics.115.182717>
 215. Katiki LM, Ferreira JF, Zajac AM, et al. *Caenorhabditis elegans* as a model to screen plant extracts and compounds as natural anthelmintics for veterinary use. *Vet Parasitol*. 2011;182(2-4):264-268. <https://doi.org/10.1016/j.vetpar.2011.05.020>
 216. Kumarasingha R, Palombo EA, Bhave M, et al. Enhancing a search for traditional medicinal plants with anthelmintic action by using wild type and stress reporter *Caenorhabditis elegans* strains as screening tools. *Int J Parasitol*. 2014;44(5):291-298. <https://doi.org/10.1016/j.ijpara.2014.01.008>
 217. Richmond JE, Jorgensen EM. One GABA and two acetylcholine receptors function at the *C. elegans* neuromuscular junction. *Nat Neurosci*. 1999;2(9):791-797. <https://doi.org/10.1038/12160>
 218. Rayes D, Flamini M, Hernando G, Bouzat C. Activation of single nicotinic receptor channels from *Caenorhabditis elegans* muscle. *Mol Pharmacol*. 2007;71(5):1407-1415. <https://doi.org/10.1124/mol.106.033514>
 219. Fonseca-Salamanca F, Martinez-Grueiro MM, Martinez-Fernandez AR. Nematocidal activity of nitazoxanide in laboratory models. *Parasitol Res*. 2003;91(4):321-324. <https://doi.org/10.1007/s00436-003-0974-7>
 220. Weeks JC, Robinson KJ, Lockery SR, Roberts WM. Anthelmintic drug actions in resistant and susceptible *C. elegans* revealed by electrophysiological recordings in a multichannel microfluidic device. *Int J Parasitol Drugs Drug Resist*. 2018;8(3):607-628. <https://doi.org/10.1016/j.ijpddr.2018.10.003>
 221. Geary TG, Woo K, McCarthy JS, et al. Unresolved issues in anthelmintic pharmacology for helminthiasis of humans. *Int J Parasitol*. 2010;40(1):1-13. <https://doi.org/10.1016/j.ijpara.2009.11.001>
 222. Atchison WD, Geary TG, Manning B, VandeWaa EA, Thompson DP. Comparative neuromuscular blocking actions of levamisole and pyrantel-type anthelmintics on rat and gastrointestinal nematode somatic muscle. *Toxicol Appl Pharmacol*. 1992;112(1):133-143.
 223. Martin RJ, Verma S, Levandoski M, et al. Drug resistance and neurotransmitter receptors of nematodes: Recent studies on the mode of action of levamisole. *Parasitology*. 2005;131(Suppl):S71-S84. <https://doi.org/10.1017/S0031182005008668>
 224. Qian H, Martin RJ, Robertson AP. Pharmacology of N-, L-, and B-subtypes of nematode nAChR resolved at the single-channel level in *Ascaris suum*. *FASEB J*. 2006;20(14):2606-2608. <https://doi.org/10.1096/fj.06-6264fje>
 225. Sloan MA, Reaves BJ, Maclean MJ, Storey BE, Wolstenholme AJ. Expression of nicotinic acetylcholine receptor subunits from parasitic nematodes in *Caenorhabditis elegans*. *Mol Biochem Parasitol*. 2015;204(1):44-50. <https://doi.org/10.1016/j.molbiopara.2015.12.006>
 226. Weeks JC, Roberts WM, Leasure C, et al. Sertraline, paroxetine, and chlorpromazine are rapidly acting anthelmintic drugs capable of clinical repurposing. *Sci Rep*. 2018;8(1):975. <https://doi.org/10.1038/s41598-017-18457-w>

How to cite this article: Giunti S, Andersen N, Rayes D, De Rosa MJ. Drug discovery: Insights from the invertebrate *Caenorhabditis elegans*. *Pharmacol Res Perspect*. 2021;9:e00721. <https://doi.org/10.1002/prp2.721>