

Contents lists available at ScienceDirect

### Preventive Medicine Reports



journal homepage: www.elsevier.com/locate/pmedr

Review article

# Social determinants of health and vaccine uptake during the first wave of the COVID-19 pandemic: A systematic review

Constantine Vardavas <sup>a,b,\*</sup>, Katerina Nikitara <sup>a</sup>, Katerina Aslanoglou <sup>a</sup>, Ioanna Lagou <sup>a</sup>, Valia Marou <sup>a</sup>, Revati Phalkey <sup>c</sup>, Jo Leonardi-Bee <sup>c</sup>, Esteve Fernandez <sup>d,e,f,g</sup>, Victoria Vivilaki <sup>h</sup>, Apostolos Kamekis <sup>a</sup>, Emmanouil Symvoulakis <sup>a</sup>, Teymur Noori <sup>i</sup>, Andrea Wuerz <sup>i</sup>, Jonathan E. Suk <sup>i</sup>, Charlotte Deogan <sup>i</sup>

<sup>a</sup> School of Medicine, University of Crete, Heraklion, Crete, Greece

<sup>b</sup> Department of Oral Health Policy and Epidemiology, Harvard School of Dental Medicine, Harvard University, Boston, MA, USA

<sup>c</sup> Health Centre for Evidence Based Healthcare, School of Medicine, University of Nottingham, Nottingham, UK

<sup>d</sup> Tobacco Control Unit, WHO Collaborating Centre for Tobacco Control, Institut Català d'Oncologia-ICO, L'Hospitalet de Llobregat (Barcelona), Spain

<sup>e</sup> Tobacco Control Research Group, Institut d'Investigació Biomèdica de. Ellvitge-IDIBELL, L'Hospitalet de Llobregat (Barcelona), Spain

<sup>f</sup> School of Medicine and Health Sciences, Campus of Bellvitge, Universitat de Barcelona, Spain

g Centre of Biomedical Research Network on Respiratory Diseases (CIBERES de Enfermedaes Respiratorias), Insituto de Salud Carlos III, Madrid, Spain

h Department of Midwifery, University of West Attica, Athens, Greece

<sup>i</sup> Emergency Preparedness and Response Support, European Centre for Disease Prevention and Control, Solna, Sweden

ARTICLE INFO

Keywords.

COVID-19

Disparities

Pandemic

Vaccination

Public health

Health determinants

#### ABSTRACT

Social determinants of health significantly impact population health status. The aim of this systematic review was to examine which social vulnerability factors or determinants of health at the individual or county level affected vaccine uptake within the first phase of the vaccination program. We performed a systematic review of peer-reviewed literature published from January 2020 until September 2021 in Medline and Embase (Bagaria et al., 2022) and complemented the review with an assessment of pre-print literature within the same period. We restricted our criteria to studies performed in the EU/UK/EEA/US that report vaccine uptake in the general population as the primary outcome and included various social determinants of health as explanatory variables. This review provides evidence of significant associations between the early phases of vaccination uptake for SARS-CoV-2 and multiple socioeconomic factors including income, poverty, deprivation, race/ethnicity, education and health insurance. The identified associations should be taken into account to increase vaccine uptake in socially vulnerable groups, and to reduce disparities in uptake, in particular within the context of public health preparedness for future pandemics. While further corroboration is needed to explore the generalizability of these findings across the European setting, these results confirm the need to consider vulnerable groups and social determinants of health in the planning and roll-out of SARS-CoV-2 vaccination programs and within the context of future respiratory pandemics.

#### 1. Introduction

The COVID-19 pandemic has had a vast and unprecedented impact on population health in terms of morbidity and mortality also in social and economic terms (Vardavas, 2022). Although age has been observed as the strongest factor related to adverse COVID-19 outcomes (Vardavas et al., 2022; Romero Starke et al., 2021), socially and economically disadvantaged groups have been severely affected by the pandemic, particularly during the first waves of the Alpha and Delta variants (Control, 2020). A large body of evidence describes how social determinants of health are associated with general health status and even play a more prominent role than lifestyle choices and medical care in affecting individual health (Braveman and Gottlieb, 1974). Furthermore, health disparities can be worsened by multiple factors, among which are infectious diseases that can disproportionately affect vulnerable groups (Semenza et al., 2010).

https://doi.org/10.1016/j.pmedr.2023.102319

Received 29 November 2022; Received in revised form 7 July 2023; Accepted 10 July 2023 Available online 16 July 2023 2211-3355/@ 2023 Published by Elsevier Inc. This is an open access article under the CC BV-NC-ND license (htt

2211-3355/© 2023 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

<sup>\*</sup> Corresponding author at: School of Medicine, University of Crete, Heraklion, Greece. *E-mail address:* vardavasc@uoc.gr (C. Vardavas).

Although the implementation of non-pharmaceutical interventions (NPI), applied across the European Union, European Economic Area, United Kingdom (EU/EEA/UK) and the United States (US), played an important role in mitigating the pandemic (Mendez-Brito et al., 2021), pharmaceutical measures, such as vaccines, are essential in reducing the risk of adverse outcomes and death (Xing, 2021). Previous research has identified variations in vaccine coverage between ethnic groups for other vaccinations (Ward et al., 2017; Loiacono, 2020; Jain et al., 2018) and in the intention to accept the SARS-CoV-2 vaccine (Omaduvie et al., 2021). Initial vaccination strategies focused on older adults, healthcare workers and those of high clinical vulnerability, and then expanded to the general population. Evidence shows that sociodemographic factors (Martin et al., 2021; Mena et al., 2021), which is consistent with the findings of studies examining immunization programs outside the context of global pandemics (Glatman-Freedman and Nichols, 2012; Yeung et al., 2016). Moreover, as described in the European Centre for Disease Prevention and Control (ECDC) series of reports on the SARS-CoV-2 vaccine deployment, EU/EEA countries face challenges in increasing uptake in different population groups, such as socially vulnerable populations (Control, 2021). Comprehending which factors are associated with vaccine uptake could be of importance for policymakers when preparing a public health emergency plan for future respiratory pandemics. By exploring the first phase one can derive valuable information on what triggers or impedes vaccination uptake during the first phase of public health emergency preparedness, i.e. in the critical phase of the COVID-19 pandemic. With the above in mind, this systematic review aimed to examine which social determinants of health affected vaccine uptake during the first phase of the COVID-19 vaccination rollout in the EU/EEA, UK and the US.

#### 2. Materials and methods

#### 2.1. Search strategy and inclusion criteria

A systematic review was conducted from the 1st of January 2020 through the 22nd of September 2021 using the PRISMA framework (Preferred Reporting Items for Systematic Reviews and Meta-analyses) (Moher et al., 2009) within Medline and Embase (Bagaria et al., 2022) and the MedRxiv pre-print server. Subject headings relating to COVID-19, social determinants of health, and epidemiological study design terms were used to develop a comprehensive search strategy presented in the Online Supplementary Appendix A. Studies of all relevant analytical epidemiological designs were considered eligible provided: (i) they evaluated humans of all ages irrespective of health status, excluding healthcare workers given that this group was expected to have been included in the initial vaccination emergency response; (ii) they presented data on the association between a determinant of health and SARS-CoV-2 vaccine uptake as the outcome; (iii) they referred to actual vaccine uptake, not intention/willingness for vaccination or vaccine hesitancy; (iv) the full text was written in English. Reference lists of all included studies and relevant reviews were also screened to identify additional eligible studies.

#### 2.2. Appraisal of methodological quality

The methodological quality of each included study was evaluated independently by two reviewers using the Joanna Briggs Institute (JBI) standardised critical appraisal tool for each appropriate study design (Moola, 2020). Disagreements were resolved through discussion and, when necessary, by a senior third reviewer. The results from the quality appraisal are presented in Online Supplementary Appendix B.

#### 2.3. Data - extraction and synthesis

Studies identified from the searches were uploaded into Covidence, and duplicates were deleted. A two-stage screening process was used to identify eligible studies. Initially, a pilot training screening process was used where a random sample of 100 titles and abstracts was screened separately for eligibility by two reviewers to enable consistency in screening and identify areas for amendments in the inclusion criteria. Since a high measure of inter-rater agreement was achieved (percentage of agreement = 88.7%), the remaining titles and abstracts were shared between two reviewers. Full-text papers were screened independently with 89.3% inter-rater agreement. Disagreements or uncertainties in the screening stages were resolved through discussion and consensus with a third reviewer.

Data were extracted independently by two reviewers using a predefined data extraction sheet that included study characteristics (first author's name, year of publication), geographical context (country/ area), study design, study population, recruitment date, vaccination data source, follow-up duration, and related quantitative or descriptive findings with regard to the social determinants of health and the primary outcome of interest (vaccine uptake, which was noted as percentages, prevalence ratios or odds ratios). The data were inversed where odds ratios were provided as the likelihood to decline vaccination. Where available, adjusted odds ratios (aOR) were extracted in preference to crude estimates.

#### 3. Results

#### 3.1. Study overview

Through the initial search, 2,932 peer-reviewed studies were identified, of which 2,485 passed onto the title/abstract review process after the removal of duplicates. Of the 118 studies that were found to meet the inclusion criteria through the title and abstract screening process, 16 met all inclusion criteria while 102 studies were excluded due to study design (n = 6), outcomes (n = 67 assessed the intention for vaccination, not actual vaccination), language (n = 1), irrelevancy to COVID-19 (n = 18), irrelevant region (n = 1), and irrelevant population (n = 9). An additional five studies were identified using the MedRxiv pre-print server. The flowchart of study selection and exclusion is presented in Fig. 1.

All 21 studies included in this systematic review (Agarwal et al., 2021; Barry et al., 2021; Brown et al., 2021; Dolby, 2021; Dryden-Peterson, et al., 2021; Gharpure et al., 2021; Glampson et al., 2021; Khubchandani et al., 2021; Kim, 2021; Lindemer, 2021; MacKenna, et al., 2021; Murthy et al., 2021; Nafilyan et al., 2021; Nguyen, 2021; Perry et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Sun and Monnat, 2021; Tram, 2021; Wang et al., 2021; Whiteman et al., 2021) investigated the relationship between vaccination status and at least one social determinant of health (Table 1). The identified determinants included age and sex, ethnicity/minorities, deprivation and income, household composition and marital status, education, employment status, area of residence, household composition, housing type/tenure, health insurance, and disability. Indexes for measuring social vulnerability, such as the Social Vulnerability Index (SVI), the COVID-19 Community Vulnerability Index and the Index of Multiple Deprivation (IMD), were also extracted. Where available, information on demographics including age and gender was also extracted.

Concerning the geographical distribution, 15 were conducted in the US (Agarwal et al., 2021; Barry et al., 2021; Brown et al., 2021; Dryden-Peterson, et al., 2021; Gharpure et al., 2021; Khubchandani et al., 2021; Kim, 2021; Lindemer, 2021; Murthy et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Sun and Monnat, 2021; Tram, 2021; Wang et al., 2021; Whiteman et al., 2021), five were performed in the UK (Dolby, 2021; Glampson et al., 2021; MacKenna, et al., 2021; Nafilyan et al., 2021; Perry et al., 2021) and one study combined data from the US and the UK (Nguyen, 2021). The overall recruitment timeframe of the included studies covered the period from December 2020 to August 2021, reflecting hence the initial rollout of vaccinations.




Fig. 1. PRISMA flowchart of the study selection on the social determinants of health and vaccine uptake during the first wave of the COVID-19 pandemic.

#### 3.2. Vaccination coverage by age and gender

Among the determinants identified in the included studies of this review, age was reported in 13 studies (Barry et al., 2021; Dryden-Peterson, et al., 2021; Glampson et al., 2021; Khubchandani et al., 2021; Kim, 2021; Lindemer, 2021; Murthy et al., 2021; Nafilyan et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Sun and Monnat, 2021; Tram, 2021; Whiteman et al., 2021), presented in Table 2. In all mentioned cases, vaccination coverage increased with age at either the participant or community level, reflecting the age-staggered prioritization of the vaccination. Gender was assessed as a possible contributing factor in vaccine uptake in most of the included studies (n = 11, Table 2) (Glampson et al., 2021; Khubchandani et al., 2021; Kim, 2021; MacKenna, et al., 2021; Murthy et al., 2021; Nafilyan et al., 2021; Perry et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Tram, 2021; Whiteman et al., 2021). Among the seven studies conducted in the US (Khubchandani et al., 2021; Kim, 2021; Murthy et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Tram, 2021; Whiteman et al., 2021), four studies estimated odds ratios (ORs) and found from self-reported data that women were more likely compared to men to have been vaccinated (Kim, 2021; Nafilyan et al., 2021; Perry et al., 2021; Singh et al., 2021), while six other studies with only descriptive results indicated higher vaccination proportions in men, without reporting the statistical significance of these differences (Glampson et al., 2021; Khubchandani et al., 2021; MacKenna, et al., 2021; Murthy et al., 2021; Ryerson et al., 2021; Tram, 2021; Whiteman et al., 2021). Two out of the four included UK studies agreed that women were slightly more likely to receive vaccination than men (Glampson et al., 2021; Nafilyan et al., 2021; Perry et al., 2021). With regard to older participants, vaccination rates were slightly higher among men within populations of UK adults 80 + years of age (Women: 94.5%, men: 94.9%) (MacKenna, et al., 2021) and among US adults 65 + years of age (women: 77.5%, men: 79.6%) (Whiteman et al., 2021).

#### Table 1

Descriptive characteristics of the studies included within this systematic review on the social determinants of health and vaccine uptake during the first wave of the COVID-19 pandemic.

Lead author, Year	Country	Population (n)	Study design	Timeframe	Vaccination data source	Outcome
(Agarwal et al., 2021)	US	General population, $(n = 756 \text{ counties})$	Cross-sectional	as of 19/4/ 2021	Official websites of Departments of Health in each state	Difference in rates with at least 1 dose
(Barry et al., 2021)	US	General population $(n = 3,142 \text{ counties})$	Cross-sectional	14/12/ 2020–1/5/ 2021	CDC COVID-19 data tracker	At least 1 dose
(Brown et al., 2021)	US	General population, $(n = 2,415 \text{ counties})$	Cross-sectional	as of 25/05/ 2021	CDC COVID-19 data tracker	Fully vaccinated <sup>2</sup>
(Dolby, 2021)	UK	General population (n = $35,223,466$ adults – UK registrar)	Not mentioned	as of 15 September 2021	NHS Improvement's National Immunisation Management System	At least 1 dose
(Dryden-Peterson, et al., 2021)	US	General population, MA (n = $6,795,000, n = 291$ communities)	Not mentioned	29/1/2020–9/ 4/2021	Massachusetts Department of Public Health and the Boston Public Health Commission	Vaccination-to- infection risk (VIR) <sup>3</sup>
(Gharpure et al., 2021)	US	Long term care residents and staff $(n = 35,854 \text{ communities})$	Cross-sectional	18/12/ 2020–21/04/ 2021	Partner pharmacies which reported to CDC	At least 1 dose
(Glampson et al., 2021)	UK	General population, London (n $=$ 2,183,939)	Retrospective cohort	08/12/ 2020–15/2/ 2021	GP primary care records	Declining vaccination <sup>4</sup>
(Khubchandani et al., 2021)	US	General population, $(n = 1,602)$	Cross-sectional	as of 6/2021	Self-reported: Online survey via mTurk	At least 1 dose
(Kim, 2021)	US	General population, Household Pulse Survey (two waves, $n = 68,348$ and $n = 80,567$ )	Cross-sectional	6–18/1/2021 17–29/3/2021	Self-reported: US Census Bureau's Household Pulse Survey	At least 1 dose
(Lindemer, 2021) (now published as Donadio)	US	General population ( $n = 1510$ counties)	Not mentioned	as of 1/3/2021	CDC COVID Data Tracker	At least 1 dose
(MacKenna, et al., 2021)	UK	Primary care patients > 80 years old not resident in care homes (2,422,476)	Retrospective cohort	8/12/ 2020–25/3/ 2021	OpenSAFELY-TPP platform	At least 1 dose
(Murthy et al., 2021)	US	General population (n $=$ 113,554,259)	Cross-sectional	14/12/ 2020–10/04/ 2021	CDC COVID-19 data tracker	At least 1 dose
(Nafilyan et al., 2021)	UK	Elderly adults > 70, (n = $6,829,643$ )	Cohort	as of 15/3/ 2021	National Immunisation Management System	Unvaccinated <sup>5</sup>
(Nguyen, 2021)	US & UK	General population (N = 1,341,682; US n = 87,388; UK n = 1254294)	Cohort study	24/3/ 2020–16/2/ 2021	Self-reported: COVID Symptom Study (CSS)	At least 1 dose
(Perry et al., 2021)	UK	General population (n = 1256412)	Cross-sectional	as of 25/4/ 2021	Wales Immunisation System (Wise, 2021)	At least 1 dose
(Ryerson et al., 2021)	US	General population ( $n = 56,749$ )	Cross-sectional	30/5/ 2021–26/6/ 2021	Self-reported: NIS-ACM Survey	At least 1 dose
(Singh et al., 2021)	US	General population ( $n = 224,458$ )	Cross-sectional	6/1-15/2/ 2021	Self-reported: US Census Bureau's Household Pulse Survey	At least 1 dose
(Sun and Monnat, 2021)	US	General population, $(n = 2,869 counties)$	Cross-sectional	as of 11/8/ 2021	CDC COVID-19 data tracker	At least 1 dose
(Tram, 2021)	US	General population (N = 459 235)	Cross-sectional	6/1–29/3/ 2021	Self-reported: US Census Bureau's Household Pulse Survey	At least 1 dose
(Wang et al., 2021)	US	General population, CT (n = 168 counties)	Cross-sectional	as of 8/3/2021	Connecticut Department of Public Health (CTDPH)	At least 1 dose
(Whiteman et al., 2021)	US	General population (n = 42,736,710)	Cross-sectional	14/12/ 2020–10/4/ 2021	CDC COVID-19 data tracker	At least 1 dose

1: Difference in rates with at least 1 dose1.

2 Fully vaccinated reported as having received both doses of a two dose vaccination series or a single dose of a one-dose series.

3: Vaccination-to-infection risk (VIR) VIR ratio was calculated for each community as the quotient of the number of fully vaccinated individuals divided by the cumulative number of confirmed SARS-CoV-2 infections.

4: declining a vaccine if they indicated that they did not want a vaccine to their GP and did not then receive a vaccine.

5: aOR have been reversed to indicate the likelihood of vaccine uptake.

#### 3.3. Race/ethnicity/minorities

Sixteen studies focused on race and ethnicity (Table 2) (Agarwal et al., 2021; Barry et al., 2021; Brown et al., 2021; Dolby, 2021; Dryden-Peterson, et al., 2021; Gharpure et al., 2021; Glampson et al., 2021; Khubchandani et al., 2021; Kim, 2021; Lindemer, 2021; MacKenna, et al., 2021; Nafilyan et al., 2021; Nguyen, 2021; Perry et al., 2021; Ryerson et al., 2021; Singh et al., 2021; Sun and Monnat, 2021; Tram, 2021; Wang et al., 2021; Whiteman et al., 2021), of which 11 were set in the US (Agarwal et al., 2021; Barry et al., 2021; Brown et al., 2021; Dryden-Peterson, et al., 2021; Gharpure et al., 2021; Khubchandani

et al., 2021; Kim, 2021; Lindemer, 2021; Nguyen, 2021; Ryerson et al., 2021; Singh et al., 2021; Sun and Monnat, 2021; Tram, 2021; Wang et al., 2021; Whiteman et al., 2021). In principle, the results showed an increased vaccination coverage in Asians and White, while Black people had lower odds of vaccination. Tram et al. (Tram, 2021), noted that Asians were found to have the highest vaccination proportion, followed by non-Hispanic and White individuals. The lowest proportions were noted in those declared  $\geq 2$  ethnic descents, Hispanic and Black. At the county level, US counties with a greater percentage of Hispanic and Black individuals were found to have lower vaccination coverage when compared to counties with higher shares of Asians, White, and

Author, Year

(Agarwal et al., 2021)

Age

#### Table 2

Effect of Age, Sex and Ethnicity/minorities status on vaccine uptake during the first wave of the COVID-19 pandemic.

Sex

0.93 (0.74–1.17)

(continued on next page)

(1.22–1.39)

n vaccine uptake during	Author, Year	Age	Sex	Race/Ethnicity/ minorities
Race/Ethnicity/	(Dryden-	vaccination		vaccination relative
minorities	Peterson,	relative to infection		to infection risk>209
Race	et al., 2021)	<i>risk</i> per 5% higher		of the community
White: 35.9%		community		Black and/or Latinx
Black: 19.9%		proportion of		(aRR 0.67, 95% CI
Percentage point		residents aged 65+		0.56 to 0.81, p $<$
change in vaccinatio	n	(aRR		0.001).
disparity associated		1.23,1.15–1.31, p		
with one SD increase		< 0.001)		
in the predictor	(Gharpure			Social Vulnerability
Segregation index	et al., 2021)			Index
Coeff. = 1.43 (0.69)				Racial/Ethnic
Percentage of racial				Minority and
and ethnic minority				language: Residents i
residents:				assisted living
Counties below				Q1: 72% vaccine
median: 48.5%				uptake per 100beds
vaccinated				Q2: 75% vaccine
Counties at or above				uptake per 100beds
median: 55.1%				Q3: 75% vaccine
vaccinated				uptake per 100beds
Percentage of person				Decial /Ethal-
speaking English less				Racial/Ethnic
than well:				Minority and
Counties below				language: Residents i residential care
median: 45.8%				Q1: 8/10 vaccine
vaccinated				uptake per ten beds
Counties above				Q2: 9/10 vaccine
median: 55.2%				uptake per ten beds
vaccinated				Q3: 10/10 vaccine
COVID-19				uptake per ten beds
Community	I (Glampson	Male: 5.83%	Age:18-64	Black/Black British:
Vulnerability Index	et al., 2021)	declining	Male: 3.16%	16.14% declining
Overall Community	ct u., 2021)	vaccination	declining	vaccination
vulnerability index: Effect of a 10point		Female: 5.92%	vaccination	Asian/Asian British:
increase = -1.2, p <		declining	Female:	3.21% declining
0.001		vaccination	2.17%	vaccination
Age: 18–65			declining	White: 4.92%
Minority status and			vaccination	declining vaccination
language: Effect of a			Age: $\geq 65$	Mixed ethnicity
10point increase =			Male: 7.08%	groups: 10.39%
1.1, p < 0.001			declining	declining vaccination
Q1: 40.9% vaccinate	1		vaccination	Other ethnic groups:
Q2: 44.4% vaccinate			Female:	9.95% declining
Q3: 44.4% vaccinate			0.94%	vaccination
Q4: 45.6% vaccinate			declining	Unrecorded ethnic
Q5: 50.3% vaccinate			vaccination	groups: 8.52%
Age: >65				declining vaccination
Minority status and	(Khubchandani	18–25: 162/199	Male: 830/	White: 1040/1293
language: Effect of a	et al., 2021)	vaccinated (81%)	1034	vaccinated (80.4%)
10point increase =		26-35: 558/704	vaccinated	African-American:
0.2, p = 0.166		vaccinated (79%)	(80.3%)	121/161 vaccinated
Community		36-45: 282/343	Female: 435/	(75.2%)Asian: 69/86
vulnerability index:		vaccinated (82%)	552	vaccinated
Effect of a 10point		46-59: 185/243	vaccinated	(80.2%)Other: 47/62
increase = -1.2, $p <$		vaccinated	(78.8%)	vaccinated
0.001		(76%)>=60: 77/95		(75.8%)
Q1: 67.4% vaccinate	1	vaccinated		EthnicityHispanic:
Q2: 70.5% vaccinate	1	(81%)		415/459 vaccinated
Q3: 71.1% vaccinate				(90.4%)Non-Hispani
Q4: 71.3% vaccinate				850/1127 vaccinated
Q5: 72.0% vaccinate				(75.4%)
Vaccination rates we	e (Kim, 2021)	6-18/1/2021	6-18/1/2021	6-18/1/2021
highest among White		18–29: Ref	Male: Ref	Non-Hispanic White:
British and Indian		30–39: aOR = 1.21	Female: aOR	Ref
ethnic groups, and		(1.00–1.46)	= 1.69	Non-Hispanic Black:
lowest among Black		40–49: aOR = 1.34	(1.53–1.88)	aOR = 0.92
Caribbean, Black		(1.10–1.64)		(0.76–1.10)
African, Mixed, and		50–64: aOR = 1.47	17-29/3/	Non-Hispanic Asian:
Pakistani ethnic		(1.20 - 1.80)	2021	aOR = 1.79
groups.		65–74: aOR = 0.95	Male: Ref	(1.48–2.18)
- I		(0.75–1.19)	Female: aOR	Non-Hispanic
		75+: aOR = <b>1.55</b>	= 1.30	multiracial: aOR =
		(1.13 - 2.12)	(1.22 - 1.39)	0.93(0.74 - 1.17)

(1.13–2.12)

		change in vaccination
		disparity associated
		with one SD increase
		in the predictor
		Segregation index
		Coeff. = 1.43 (0.69)
(Barry et al.,	Percentage of the	Percentage of racial
2021)	population $\geq$ 65	and ethnic minority
	Counties below median: 54.9%	<i>residents:</i> Counties below
	vaccinated	median: 48.5%
	Counties above	vaccinated
	median: 49.4%	Counties at or above
	vaccinated	median: 55.1%
	Percentage of the	vaccinated
	population < 18	Percentage of persons
	Counties below	speaking English less
	median: 56.8%	than well:
	vaccinated	Counties below
	Counties above	median: 45.8%
	median: 51.3%	vaccinated
	vaccinated	Counties above
		median: 55.2%
		vaccinated
(Brown et al.,		COVID-19
2021)		Community
		Vulnerability Index <sup>1</sup>
		Overall Community vulnerability index:
		Effect of a 10point
		increase = -1.2, $p <$
		0.001
		Age: 18–65
		Minority status and
		language: Effect of a
		10point increase =
		1.1,  p < 0.001
		Q1: 40.9% vaccinated
		Q2: 44.4% vaccinated
		Q3: 44.4% vaccinated
		Q4: 45.6% vaccinated
		Q5: 50.3% vaccinated Age: >65
		Minority status and
		language: Effect of a
		10point increase =
		0.2, p = 0.166
		Community
		vulnerability index:
		Effect of a 10point
		increase = -1.2, p $<$
		0.001
		Q1: 67.4% vaccinated
		Q2: 70.5% vaccinated
		Q3: 71.1% vaccinated Q4: 71.3% vaccinated
		Q5: 72.0% vaccinated
(Dolby, 2021)		Vaccination rates were
(2010), 2021)		highest among White
		British and Indian
		ethnic groups, and
		lowest among Black
		Caribbean, Black
		African, Mixed, and
		Pakistani ethnic
		groups.

5

orts 35 (2023) 102319

Author, Year	Age	Sex	Race/Ethnicity/ minorities	А
	17-29/3/2021		Hispanic: aOR = 1.06	
	18–29: Ref		(0.88–1.26)	
	30-39: aOR = 1.42			
	(1.24–1.63)		17-29/3/2021	
	40-49: aOR = 1.81		Non-Hispanic White:	
	(1.57-2.07)		Ref	
	50-64: aOR = <b>3.04</b> (2.65-3.49)		Non-Hispanic Black: aOR = 1.07	
	(2.03-3.49) 65–74: aOR =		(0.95-1.20)	
	12.76		Non-Hispanic Asian:	
	(10.81-15.05)		aOR = 1.28	
	75+: aOR = <b>25.73</b>		(1.10 - 1.50)	
	(20.29–32.67)		Non-Hispanic	
			multiracial: $aOR =$	
			0.98 (0.81–1.18) Hispanic: aOR = 1.21	
			(1.08-1.36)	
(Lindemer,	Difference in rate,		Difference in rate, by	
2021)	by Quartile of		Quartile of vaccinated	
	vaccinated counties		counties (Q1 vs. Q4)	
	(Q1 vs. Q4)		%Black: RR = 0.635	(]
	<18: RR = 0.87		(0.634–0.635)	
	(0.869, 0.87) $\geq 65: RR = 1.235$		%Hispanic: RR = 0.642 (0.641-0.642)	
	$\geq 0.5$ . RR $= 1.233$ (1.234, 1.236)		%American Indian:	
	()		RR = 0.9	
			(0.896–0.904)	
			%Hawaiian/Pacific	
			Islander: RR = 1.045	
			(1.035–1.055)	
			%Asian: RR = 1.198 (1.196–1.201)	(]
			%White: $RR = 1.228$	(
			(1.228–1.229)	
(MacKenna,		Female:	Among the 80 $+$	
et al., 2021)		94.4%	population not	
		(1,400,532/	resident in care	
		1,481,970)	homesWhite: 96.2%	
		Male: 94.9% (1,021,944/	(1,458,548/ 1,515,535)Mixed:	
		1,076,936)	77.7%	
		_,,	(6,650/8,554)South	
			Asian: 84.6%	
			(55,846/65,975)	
			Black: 68.3%	
			(23,590/34,517)	
			Other: 78.3%	(
			(8,561/10,934) Unknown: 94.1%	(
			(869,260/923,363)	
(Murthy et al.,	18-64: 36.6%	Female:	,	
2021)	≥65:74.7%	47.4%		
GY (1)	<b>FO F</b> 4 <b>F</b> 4	Male: 41.0%	11111, 10, 1-1-1, 10, 1	
(Nafilyan et al.,	70–74: Ref	Female:	White British: Ref	
2021)	75-79: aOR = 1.51 (1.51-1.53)	Reference Male: OR =	Bangladeshi: aOR = 0.39 (0.37–0.41)	
	(1.51-1.55) 80–84: aOR = 2.22	1.03	Black African: $aOR =$	
	(2.17-2.22)	(1.02–1.03)	0.19 (0.19–0.20)	
	85-89: aOR = 1.96		Black Caribbean: aOR	
	(1.92—1.96)		= 0.20 (0.20-0.21)	
	90-94: aOR = 1.56		Chinese: $aOR = 0.37$	(
	(1.53-1.58)		(0.36–0.39)	
	95-99: aOR = 1.22		Indian: $aOR = 0.74$	
	(1.19-1.25) $\geq 100: aOR = 0.62$		(0.72-0.75) Mixed: aOR = 0.45	
	$\geq 100. \text{ aOK} = 0.02$ (0.59-0.65)		(0.43-0.46)	
			Other: $aOR = 0.40$	
			(0.4–0.41)	
			(011 011-)	
			Pakistani: $aOR = 0.27$	
			Pakistani: aOR = 0.27 (0.27–0.28)	
			Pakistani: $aOR = 0.27$ (0.27–0.28) White other: $aOR =$	
(Names 2001)			Pakistani: aOR = 0.27 (0.27–0.28) White other: aOR = 0.51 (0.51–0.52)	
(Nguyen, 2021)			Pakistani: $aOR = 0.27$ (0.27–0.28) White other: $aOR =$	

Author, Year	Age	Sex	Race/Ethnicity/ minorities
			(0.64 – 0.79) Hispanic: aOR = 0.93
			(0.84-1.02)
			Asian: $aOR = 1.00$
			(0.93–1.09)
			More than one/other:
			aOR = 0.94
			(0.81–1.08)
			UK White: Ref
			Black: $aOR = 0.98$
			(0.92–1.04)
			South Asian: aOR =
			1.18 (1.13–1.23)
			Middle East/East
			Asian: $aOR = 1.01$
			(0.94–1.08) More than one/other:
			aOR = 0.99
			(0.93–1.04)
(Perry et al.,		Male: Ref	White: Ref
2021)		Female: aOR	Black: $aOR = 0.22$
		= 1.18	(0.21–0.24)
		(1.16–1.19)	Asian: $aOR = 0.41$
			(0.39–0.43) Minodu aOD = 0.26
			Mixed: $aOR = 0.36$ (0.34-0.38)
			Other: $aOR = 0.24$
			(0.22–0.27)
			Unknown: aOR = 0.20
			(0.19-0.20)
(Ryerson et al.,	Age (No disability	Sex (No	Ethnicity (No
2021)	vs. disability)	disability vs.	disability vs.
	18–24: 46.4% vs.	disability)	disability)White (Ref)
	33.5% 25–29: 49.8% vs.	Male: 61.9% vs. 66.4%	: 66.6% vs. 69.0% Hispanic: 61.8%
	35.5%	Female: 67%	vs.67.2%
	30–39: 52.9% vs.	vs. 67.3%	Black: 56.3% vs 60.1%
	48.8%		AI/AN: 39.2% vs.
	40-49: 60.8% vs.		38.2%
	54.4%		Asian: 85.5% vs.
	50–64: 71.9% vs.		74.7%
	62.8%) 65–75: 88.6% vs.		NHPI: 59.1% vs. 71.1%
	82.7%		Multiple race/Other:
	75+: 90.0% vs.		49.2
(Singh et al.,	88.2% >= 75: Ref	Male: Ref	Non-Hispanic White:
2021)	64-74: aOR = 0.48	Female: aOR	Ref
-	(0.44–0.53)	= 1.54	Non-Hispanic Black:
	55-64:aOR = 0.25	(1.47–1.62)	aOR = 0.89
	(0.22–0.27)		(0.82–0.98)
	45-54: aOR = 0.22		Asian: $aOR = 1.5$
	(0.20-0.24) 35-44: aOR = 0.20		(1.36–1.67) Other & multiple race
	(0.18-0.22)		aOR = $1.02$
	(0.13-0.22) 25-34: aOR = 0.19		(0.89-1.17)
	(0.17–0.21)		Hispanic: $aOR = 1.03$
	18–24: aOR = 0.14		(0.95–1.12)
	(0.11–0.17)		
(Sun and	% age 65+:		%Non-Hispanic Black
Monnat,	Coefficient = $1.36$ ,		(1st quartile) Ref%
2021)	p < 0.001		Non-Hispanic Black (2nd quartile)
			: Coeff. = $-0.4$ , p =
			0.425%Non-Hispanic
			Black (3rd quartile)
			: Coeff. = -0.5, p =
			0.395%Non-Hispanic
			Black (4th quartile)
			: Coeff. = -4.18,
			<0.001%Hispanic (1s
			quartile) Ref%Hispanic (2nd
			quartile)
			(continued on next page

#### Table 2 (continued)

Author, Year	Age	Sex	Race/Ethnicity/ minorities
(Tram, 2021)	$\begin{array}{l} 18-24 = 9.7\% \\ (9.0-10.4) \\ 25-39 = 17.1\% \\ (16.8-17.4) \\ 40-54 = 20.2\% \\ (19.9-20.6) \\ 55-64 = 24.3\% \\ (23.8-24.8) \\ \geq 65 = 45.4\% \\ (44.9-45.9) \end{array}$	Female = 27.1% (26.8–27.3) male = 22.0% (21.7–22.3)	: Coeff. = 0.05, p = 0.927%Hispanic (3rd quartile) : Coeff. = 0.59, p = 0.276%Hispanic (4th quartile) : Coeff. = 1.96, p = 0.002 Black = 20.6% (20.0-21.1) White = 25.4% (25.2-25.6) Asian = 29.2% (28.3-30.1) $\ge 2$ Races + other = 18.5% (17.7-19.3) Hispanic/Latino = 18.9% (18.3-19.5)
(Wang et al., 2021)			Not Hispanic/Latino = 25.8% (25.6–26.0) Minority aspect of the SVI Age: 65–74Q1: 78.81
			(11.05)Q4: 70.18 (11.49)Diff: 8.63 (3.74–13.53) Age: >75Q1: 95.80 (17.39)Q4: 74.80 (10.88)Diff: 21.00 (14.70–27.30)
(Whiteman et al., 2021)	65–74: 79.6% vaccinated >=75: 78.3% vaccinated	Male: 79.6% vaccinated Female: 77.5% vaccinated	Average percentage of older adults indicating race/ethnicity other than White<50% vaccination rates
			(8.0%; 95%  CI = $4.9\%-11.1\%) \ge 75\%$ vaccination rates (9.3%; 95%  CI = 6.4%-12.1%)

\* The number of fully vaccinated individuals divided by the cumulative number of confirmed SARS-CoV-2 infections.

\* The difference in the COVID-19 vaccination rate between White and Black residents in a county, where vaccination rates are based on the total population of a given race in a county.

\*\* The number of fully vaccinated individuals divided by the cumulative number of confirmed SARS-CoV-2 infections.

1: CCVI is computed using 40 measures within 7 themes: socioeconomic status (SES); minority status and language; housing type, transportation, household composition, and disability ("housing type/composition" hereafter); epidemiological factors; healthcare system factors; high risk environments; and population density (Mendez-Brito et al., 2021).

Hawaiian/Pacific Islanders (Lindemer, 2021). Furthermore, significantly lower vaccination rates were noted in US counties with larger percentages of Black people (Sun and Monnat, 2021), findings that were corroborated by other studies (Agarwal et al., 2021; Nguyen, 2021; Singh et al., 2021), with the latter estimating that non-Hispanic Black people had lower odds of vaccination compared to non-Hispanic Whites, while Asians were more likely to be vaccinated. The minority status and language theme of the COVID-19 Community vulnerability index also indicated the US counties with lower levels of vulnerability had higher vaccination rates compared to those with the highest level of vulnerability (Brown et al., 2021), while the minority aspect of the SVI also was associated with significant differences in vaccine uptake (Wang et al., 2021).

In line with the findings of the preceding US studies, the included studies of this review which took place in the UK during the first four months of 2021 suggested that White people and Asians were more likely to have been vaccinated against COVID-19 when compared to Black and Black British (MacKenna, et al., 2021; Nguyen, 2021) while multivariable analyses also confirmed the above findings (Nafilyan et al., 2021; Perry et al., 2021). In the first study, White British at the age of  $\geq$  70 years old were found to have lower odds of being unvaccinated compared to White, Bangladeshi, and Pakistani, Black African/Caribbean, Chinese, Indian, mixed, and other ethnicity groups at the same age range (Nafilyan et al., 2021). In the second study, White individuals at 50 years of age or older were more likely to have been partially vaccinated than Black, Asian, mixed, other, and unknown ethnicity groups (Perry et al., 2021). Finally, in the Dolby et al. study (Dolby, 2021), vaccination rates were highest among White British and Indian ethnic groups and lowest among Black Caribbean, Black African, Mixed, and Pakistani ethnic groups.

#### 3.4. Deprivation and, income

A key social determinant of health identified in this review is poverty/deprivation and overall socioeconomic status for which all studies noted significant associations and are presented in Table 3. In all studies that used the IMD as an indicator of deprivation, vaccination was lower for those within the most deprived communities compared to those within the most affluent (Government, 2015). Perry et al. (Perry et al., 2021), noted the odds of being vaccinated were significantly lower for deprived people compared to those least deprived (most deprived: OR = 0.59, 95% C.I: 0.57–0.60). These results are similar to those of Nafilyan et al. (Nafilyan et al., 2021) (most deprived: OR = 0.63; 95 %C. I: 0.62–0.64). Comparing vaccination rates between quintiles of the IMD, MacKenna et al. (MacKenna, et al., 2021) also indicated that the most deprived had lower rates of being vaccinated compared to the least deprived. Using the COVID-19 Community Vulnerability Index, Brown et al. (Brown et al., 2021), also noted that vaccination rates were lower among those more vulnerable while Wang et al., noted that the differences in vaccination rates were statistically significant between quartiles of the poverty SVIs (Wang et al., 2021).

Income was studied as a possible predictor and lower personal income, either at the individual or the county level, was associated with lower vaccination rates in all identified studies. Notably, two US studies reporting odds ratios of respondent level data, noted that participants with an income of <25,000\$ had significantly lower odds of having been vaccinated with at least one vaccine dose (OR = 0.67, p < 0.05), (Kim, 2021; Singh et al., 2021). Lower odds were also noted for those within the 25,000–34,999\$ (OR = 0.81, p < 0.05) and 35,000–49,999\$ (OR = 0.87, p < 0.05) income range, while Tram et al. (Tram, 2021) found that as reported income increased, so did the proportion of people who received vaccination, ranging from 15.4% (14.6-16.2) among those with an income of < 25,000 to 33.0% (32.2–33.8) for those with an income of > 200,000\$. Additionally, using county level data, rural and urban US counties with a higher median household income had significantly higher vaccination rates (Coeff. = 2.78, p < 0.001), while in the study by Barry et al. (Barry et al., 2021), those with a median income of <26,245\$ had lower vaccination rates (42.7%) compared to those above the median income (56.7%). The study by Nguyen et al. (Nguyen, 2021), with data from the US and the UK, presented slightly higher vaccination rates in higher-income participants, while the median per capita income was also identified to be strongly associated with vaccine roll-out. According to Kim (Kim, 2021), the odds of being vaccinated were significantly lower for those experiencing very difficult financial hardship [OR = 0.76 (0.63–0.90), p = 0.002] and also for those experiencing somewhat difficult financial hardship [OR = 0.85 (0.73-0.99), p = 0.03].

#### 3.5. Education

Ten studies aimed to examine the relationship between education level and vaccine uptake (Table 3), nine of which were based in the US, while one study included data both from the US and the UK. All studies indicated that increased vaccination rates were noted amongst those

#### Table 3

Preventive Medicine Reports 35 (2023) 102319

Table 3 (continued)

Study	e first wave of the CC Education	Income/ Deprivation/ Poverty	Occupation/ Occupational setting/	(Gharpure		Deprivation/ Poverty Social	Occupational setting/ Employment
		5	Employment	et al., 2021)		Social Vulnerability Index	
Agarwal et al., 2021)	Percentage point change in vaccination	Percentage point change in vaccination		ct all, 2021)		Socioeconomical status Residents in assisted living Q1: 75% vaccine	
	disparity associated with one SD increase in the predictor County High school	disparity associated with one SD increase in the predictor County Median				uptake per 100beds Q2: 74% vaccine uptake per 100beds Q3: 71% vaccine uptake per 100beds	
	graduation rate: Coeff. = 1.22, p > 0.05 County High school disparity: Coeff. =	income: Coeff. = -2.2, p < 0.05 County Median income disparity: Coeff. = 0.89, p >				Socioeconomical status: Residents in residential care	
Barry et al., 2021)	2.01, p < 0.001 No high school diploma Counties below	0.05 <b>Income per capita</b> Counties below median: 42.7%	<b>Unemployment:</b> Counties below median: 56.6%			Q1: 10/10 vaccine uptake per ten beds Q2: 9/10 vaccine uptake per ten beds	
	median: 56.5% vaccinated Counties above median: 50.40% vaccinated	vaccinated Counties at or above median: 56.7% vaccinated	vaccinated Counties at or above median: 51.9% vaccinated	(Glampson et al., 2021)		Q3: 8/10 vaccine uptake per ten beds Deprivation and rate of declining vaccination	
		Percentage of people living in poverty Counties below				Overall association: r = -0.94; P = 0.002) <b>By Postcodes</b> Most deprived 13.5%	
		median: 57.4% vaccinated Counties at or above median: 49.8% vaccinated				(1980/14,571) declining vaccinationLeast deprived 0.98%	
(Brown et al., 2021)		COVID-19 Community Vulnerability		(Khubchandani	High school or less:	(869/9609) declining vaccination	Full-time: 1121
		Index <sup>2</sup> Age: 18–65 Socioeconomical status: Effect of a 10point increase = -1.2, $p < 0.001Q1: 53.8%vaccinated$		et al., 2021)	A4/90 (49%) College: 93/149 (62%)Bachelor's degree: 822/992 (83%)>=Master's degree: 306/355 (86%)		1393 vaccinated (80.5%)Part- time: 91/116 vaccinated (78.4%)Not employed: 53/7 vaccinated
		Q2: 53.0% vaccinated Q3: 48.3%		(Kim, 2021)	<b>6–18/1/2021</b> ≥College: Ref	Income 6–18/1/2021	(68.8%)
		vaccinated Q4: 44.6% vaccinated Q5: 45.2%			<high aor<br="" school:="">= <b>0.41</b> (0.27–0.63) High school: aOR = <b>0.40</b> (0.34–0.48) Some college: aOR</high>	<pre>&gt;\$200,000: Ref \$150,000- \$199,999: aOR = 1.06 (0.83, 1.34) \$100,000-</pre>	
		vaccinated Age: >65 Socioeconomical status: Effect of a			= <b>0.76</b> (0.69–0.85) <b>17–29/3/2021</b> ≥College: Reference	\$149,999: aOR = 0.96 (0.78, 1.17) \$75,000-\$99,999: aOR = 0.79 (0.63,	
		10point increase = -1.4, p < 0.001 $Q1: 76.7%$ $vaccinated$			<high aor<br="" school:="">= <b>0.29</b> (0.24–0.36) High school: aOR = <b>0.41</b> (0.37–0.45)</high>	0.997) \$50,000-\$74,999: aOR = 0.78 (0.62, 0.98)	
		Q2: 76.0% vaccinated Q3: 72.7% vaccinated Q4: 69.0%			Some college: aOR = <b>0.58</b> (0.54–0.62)	\$35,000-\$49,999: aOR = 0.68 (0.51, 0.91) \$25,000-\$34,999: aOR = 0.63 (0.46,	
(Dryden- Peterson,		vaccinated Q5: 66.9% vaccinated Decreased VIR per Quartile increase				0.87) <\$25,000: aOR = 0.53 (0.36, 0.77) <b>17-29/3/2021</b> ≥\$200,000: Ref	
et al., 2021)		aRR 0.82; 95 %C.I: (0.77–0.87)				\$150,000- \$199,999: aOR = 0.98 (0.82, 1.18)	

## Tab St

(continued on next page)

able 3 (continue	-	<b>T</b> = = = = = (	0	Table 3 (continu		To a second d	0
Study	Education	Income/ Deprivation/ Poverty	Occupation/ Occupational setting/ Employment	Study	Education	Income/ Deprivation/ Poverty	Occupation/ Occupational setting/ Employment
(Lindemer, 2021)	Difference in rate, by Quartile of vaccinated counties (Q1 vs. Q4) High school graduation: $RR =$ <b>1.019</b> (1.017-1.021)	\$100,000- \$149,999: $aOR =$ 1.00 (0.87, 1.16) \$75,000-\$99,999: aOR = 0.91 (0.77, 1.08) \$50,000-\$74,999: aOR = 0.91 (0.78, 1.07) \$35,000-\$49,999: aOR = 0.87 (0.73, 1.04) \$25,000-\$49,999: aOR = 0.86 (0.71, 1.04) <\$25,000: $aOR =0.66 (0.54, 0.81)Financialhardship:6-18/1/2021Not at all difficult:RefA little difficult:aOR = 0.89(0.78-1.00)Somewhat difficult:aOR = 0.69(0.59-0.80)Very difficult: aOR= 0.56$ (0.44-0.70) 17-29/3/2021 Not at all difficult: Ref A little difficult: aOR = 0.95 (0.87-1.03) Somewhat difficult: aOR = 0.85 (0.73-0.99) Very difficult: $aOR$ = 0.76 (0.63-0.90)	Difference in rate, by Quartile of vaccinated counties (Q1 vs. Q4) Unemployment: RR = <b>0.87</b> (0.867–0.872)	(Nguyen, 2021)	USLower education (Quartile 1) : $3991/17936$ vaccinatedWhite: 1.0 (ref.) Black: $aOR = 0.65$ (0.54 to 0.78) Hispanic: $aOR = 1.06$ (0.90 to 1.26) Asian: $aOR = 1.10$ (0.92 to 1.31) Other: $aOR = 0.86$ (0.64 to 1.14) Higher education (Quartile 4) : $4153/17851$ vaccinatedWhite: 1.0 (ref.) Black: $aOR = 0.93$ (0.70 to 1.24) Hispanic: $aOR = 0.93$ (0.70 to 1.24) Hispanic: $aOR = 1.09$ (0.05 to 1.42) Other: $aOR = 1.09$ (0.80 to 1.50) UKLower education (Quartile 1) : $40794/284004$ vaccinatedWhite: 1.0 (ref.) Black: $aOR = 1.08$ (0.97 to 1.20) Hispanic: $aOR = 1.09$ (0.94 to 1.27) Other: $aOR = 1.09$	$\begin{array}{c} (0.91-0.93) \\ \text{IMD 3: OR} = 0.85 \\ (0.85-0.86) \\ \text{IMD 2: OR} = 0.75 \\ (0.74-0.75) \text{IMD 1} \\ (\text{most deprived}) \\ : OR = 0.63 \\ (0.62-0.64) \\ \text{USLower income} \\ (Quartile 1) \\ : 4094/17865 \\ \text{vaccinatedWhite:} \\ 1.0 \\ (ref.) \\ \text{Black: aOR} = 0.73 \\ (0.61 to 0.86) \\ \text{Hispanic: aOR} = 0.73 \\ (0.61 to 0.86) \\ \text{Hispanic: aOR} = 0.94 \\ (0.79 to 1.14) \\ \text{Asian: aOR} = 0.90 \\ (0.70 to 1.16) \\ \text{Other: aOR} = 1.08 \\ (0.81 to 1.44) \\ \text{Higher income} \\ (Quartile 4) \\ : 4111/17845 \\ \text{vaccinatedWhite:} \\ 1.0 \\ (ref.) \\ \text{Black: aOR} = 0.83 \\ (0.62 to 1.11) \\ \text{Hispanic: aOR} = 0.97 \\ (0.84 to 1.22) \\ \text{Other: aOR} = 1.09 \\ (0.81 to 1.45) \\ \end{array}$	
(MacKenna, et al., 2021)	Some college: RR = <b>1.143</b> (1.143–1.144)	Index of Multiple DeprivationIMD: 5 (Least deprived): 612,731/634,340 (96,6%)IMD: 4: 564,410/588,546 (95.9%)IMD: 3: 514,682/541,737 (95%)IMD: 2: 402,836/433,622 (92.9%)IMD: 1 (Most deprived): 309,190/340,928		(Perry et al.,	(0.83 to 1.05) Higher education (Quartile 4) : 37191/226383 vaccinatedWhite: 1.0 (ref.) Black: aOR = 0.96 (0.81 to 1.12) Hispanic: aOR = 0.93 (0.81 to 1.23) Asian: aOR = 0.93 (0.81 to 1.06) Other: aOR = 0.96 (0.86 to 1.07)	Higher income (Quartile 4) : 30809/198904 vaccinatedWhite: 1.0 (ref.) Black: aOR = 1.08 (0.90 to 1.31) Hispanic: aOR = 1.08 (0.90 to 1.33) Asian: aOR = 0.91 (0.77 to 1.08) Other: aOR = 1.01 (0.88 to 1.15) IMD: 5 (least	
(Nafilyan et al., 2021)		(90,7%)Unknown: 18,613/19,691 (94.5%) IMD: 5 (least deprived): Ref IMD 4: OR = 0.92		(Perry et al., 2021)		MD: 5 (feast deprived): Ref IMD 4: aOR = <b>0.81</b> (0.79 - 0.83) IMD 3: aOR = <b>0.78</b> (0.76 - 0.79) IMD 2: aOR = <b>0.71</b>	

9

(18.4-19.2)

= 23.1%

(22.9-23.4)

(33.4-33.9)

Some college or

associate's degree

Bachelor's degree

or higher = 33.6%

(14.6-16.2)

(19.6 - 21.2)

(22.0-23.2)

(25.4-26.8) 75,000-99,999\$

20.3

22.6%

25,000-34,999\$ =

35,000-49,999\$ =

50,000-74,999\$ =26.1%

#### Table

Occupation/ Occupational

setting/ Employment

Study	Education	Income/ Deprivation/ Poverty	Occupation/ Occupational setting/ Employment	Study	Education
		(0.70 – 0.73),IMD 1 (most deprived) : aOR = <b>0.59</b> (0.57–0.60)			
(Ryerson et al., 2021)		SVI (No disability vs. disability) Low: 69.9% vs. 68% Moderate: 65.1% vs. 68.6%			
		High: 60.4% vs. 64.8% Poverty status and household income (No disability vs. disability) Above poverty > 75 K: 72.5% vs.		(Wang et al., 2021)	Without highschoo diploma Age 65–74:Q1: 79.77 (8.19)Q4: 66.86 (10.14)Diff: 12.91 (8.91–16.92)
		78.0% Above poverty < 75 K: 61.1% vs. 68.9% Below poverty: 48.6% vs. 55.5% Unknown: 64.3%			Age > 75:Q1: 98.67 (13.53)Q4: 74.47 (14.91)Diff: 24.15 (17.97–30.33)
(Singh et al., 2021)	Graduate degree or higher (>=17 y.o.): RefCollege degree, (16 y.o.) : aOR = 0.71 (0.68-0.75)Some college (13-15 y.o.): aOR = 0.59 (0.56-0.62) High school (12 y. o.) : aOR = 0.39 (0.36-0.42)High school	vs. 66.6% >= 200,000 (in 2019): Ref 150,000-199,999 (in 2019): aOR = 1.07 (0.97-1.17) 100,000-149,999 (in 2019): aOR = 0.97 (0.90-1.04) 75,000-99,999 (in 2019): aOR = 0.95 (0.87-1.03) 50,000-74,999 (in 2019): aOR = 0.92 (0.84-1.00)	Employed: Ref Unemployed: aOR = <b>0.5</b> ( <b>0.47–0.54</b> )	(Whiteman et al., 2021)	
	(<12 y.o.): aOR = 0.36 (0.30–0.43)	35,000-49,999 (in 2019): aOR = 0.87 (0.78-0.97) 25,000-34,999 (in 2019): aOR = 0.81 (0.72-0.91)< 25,000 (in 2019) : aOR = 0.67 (0.59-0.77) Unknown: aOR = 0.98 (0.88-1.10)		county-level as An increase was noted in U et al. (Khubcha 2021) estimate bachelor's deg degree or thos	lucation both wi sociations. in the vaccination JS counties in the ndani et al., 2021 ed a higher vaccin ree compared to e who had comple d Monnat, 2021)
(Sun and Monnat, 2021)	Rural & urban % with Bachelor or higher: Coeff. = 0.13, p = 0.713 Urban % with Bachelor or higher: Coeff. = 2.33, p < 0.001	County Median household income: Coeff. = 2.78, p < 0.001		percentage of likely to have uation status v (Agarwal et al. in US counties diploma (Barr higher levels	residents with Ba better vaccine upt vas also associate , 2021), while at with a higher perce y et al., 2021). A of education were , 2021; Lindemer
(Tram, 2021)	High school or less = $18.8\%$	Income<25,000\$ =15.4%		amongst those	65 + years old trelated with vacc

(Wang et al., 2021)	Without highschool diploma Age 65–74:Q1: 79.77 (8.19)Q4: 66.86 (10.14)Diff: 12.91 (8.91–16.92) Age > 75:Q1: 98.62	=27.7% (26.9-28.5) 100,000-149,999\$ = 29.4% (28.8-30.0) 150,000-199,999\$ $=32.0\%$ (31.0-32.9) $\geq$ 200,000\$ = 33.0% (32.2-33.8) Below poverty Age: 65-74Q1: 79.58 (9.99)Q4: 69.12 (12.37)Diff: 10.46 (5.59-15.35) Age: >75Q1: 98.01 (14.92)Q4: 73.87	Unemployed Age: 65–74Q1: 75.60 (9.39)Q4: 70.99 (12.43)Diff: 4.61 (0.14–9.36) Age: >75Q1: 88.75
	(13.53)Q4: 74.47 (14.91)Diff: 24.15 (17.97–30.33)	(13.70)Diff: 24.14 (17.92–30.35)	(15.64)Q4: 83.72 (19.24)Diff: 5.03 (2.54–12.58)
(Whiteman et al., 2021)		Counties with $<$ 50% vaccination initiation rates:10.3% (95% CI = 9.2% – 11.4%) in poverty Counties with >=75% vaccination initiation rates:7.6% (95% CI = 7.0%– 8.2%) in poverty	

Income/

-27 7%

Deprivation/ Poverty

ithin participant-level associations and

on rate as the educational level increased e self-reported survey by Khubchandani 1). Similarly in the US, Tram et al. (Tram, ine uptake among those with at least a individuals with a college or associate leted high school or less. Likewise, Sun noted that US counties with a greater achelor's degrees or higher were more take. The disparity in high school graded with increased vaccination disparity the county-level vaccination was lower rcentage of residents with no high school Additional studies also indicated that ere positively associated with vaccine er, 2021; Singh et al., 2021). Similarly, the lack of a high school diploma was ccination coverage (Wang et al., 2021). Finally, Nguyen et al. (Nguyen, 2021), who presented data both from the US and the UK for adults at the age of 18 and older, also confirmed that vaccination rates were higher in participants with higher education.

#### 3.6. Employment status

The role of employment status in vaccine uptake was explored in five studies (Table 3), all of which took place in the US and which, in principle, indicated that unemployed individuals had a lower likelihood or lower percentages of vaccination either at the population level (Khubchandani et al., 2021; Lindemer, 2021; Singh et al., 2021) or through county-level unemployment rates (Barry et al., 2021).

#### 3.7. Marital status

Four studies investigated marital status as a possible factor for vaccine uptake in the general population (n = 4). In three studies, married participants were found to have the highest vaccination rates (Khubchandani et al., 2021); (Tram, 2021); (Kim, 2021), while one (Singh et al., 2021) indicated that widowed people were more likely to have received vaccination than unmarried. With regard to household composition, findings differed based on the definition of composition across studies and on the associations between participant and countylevel data.

#### 3.8. Housing

Among the social determinants of health identified in this review, housing tenure was studied in three studies, two from the US adults (Lindemer, 2021; Singh et al., 2021) and one among adults > 70 years of age in the UK (Nafilvan et al., 2021). In all included studies, the odds of being vaccinated were higher for owners than for renters. With regard to housing type, we identified three studies (Barry et al., 2021; Gharpure et al., 2021; Wang et al., 2021); two of them reported data from the general US population and one from long-term care residents and staff in the US. Barry et al. (Barry et al., 2021), using the housing component of the SVI, indicated higher vaccination coverage in US counties with a higher percentage of housing structures with  $\geq 10$  units and lower coverage in counties with higher percentages of mobile homes. On the contrary, Gharpure et al. (Gharpure et al., 2021) reported increased vaccine uptake with decreasing SVI quartiles related to housing type among long-term care residents and Wang et al. (Wang et al., 2021) estimated higher rates of vaccination in mobile homes with residents aged 75 and older.

#### 3.9. Urbanicity

The association between urbanicity and vaccination coverage in principle differed depending on whether the study was performed in the US or the UK. Data from the CDC COVID-19 data tracker suggested a better vaccination status in urban areas, with a lower likelihood of vaccination for the residents of rural US counties (RR = 0.96, 95 %CI: 0.96–0.97) (Lindemer, 2021), a direction of association similar to that reported by Murthy et al., also in the US (Murthy et al., 2021). On the contrary, two large studies with data from the UK reported higher vaccination rates in rural areas. According to Perry et al. (Perry et al., 2021), the odds of being vaccinated were lower for individuals over 50 years of age residing in urban areas compared to rural (OR = 0.86, 95% CI: 0.84–0.87), which is similar to the results of Nafilyan et al., 2021).

#### 3.10. Disability

Disability was identified as a factor in five studies of this review with ambiguous results (Barry et al., 2021; MacKenna, et al., 2021; Nafilyan et al., 2021; Ryerson et al., 2021; Wang et al., 2021). The largest sample populations were from the UK, where people aged  $\geq$  70 years old were more likely to be vaccinated if they reported to be a little limited due to disability (OR = 1.08 (1.07–1.08) or a lot limited due to disability (OR = 1.29 (1.28–1.30) (Nafilyan et al., 2021), while among those > 80 years old, 93.6% of the participants with a learning disability were vaccinated (MacKenna, et al., 2021). On the contrary, a population-based study in the US (Ryerson et al., 2021) indicated that participants with disabilities were less likely to be vaccinated (PR = 0.88, p=<0.05), while studies assessing disability at the county level did not identify significant

associations (Barry et al., 2021; Wang et al., 2021).

#### 3.11. Health insurance

Among the social determinants of health identified in this review, health insurance was analysed in three US studies (Lindemer, 2021; Singh et al., 2021; Sun and Monnat, 2021) (Online Supplementary Appendix C). Based on the study of Singh et al. (Singh et al., 2021), the odds of being vaccinated were significantly lower for participants that didn't have insurance (OR = 0.59, p=<0.05). Similarly, Lindemer et al. (Lindemer, 2021) showed that the odds of being vaccinated were significantly lower for uninsured adults [OR = 0.74 (0.74–0.74), p=<0.001)] and children [OR = 0.79 (0.79–0.79), p=<0.001]. Finally, Sun et al. (Sun and Monnat, 2021) using county-level data did not identify a significant difference in vaccination status between insured and uninsured adults.

#### 4. Discussion

This systematic review aimed to examine the relationship between social determinants of health and vaccine uptake for SARS-CoV-2 vaccines in the first phase of the vaccination programs. Our findings indicated significant differences in vaccine uptake by almost all factors assessed within the context of this review, including but not limited to age, ethnicity/minority status, income, indexes of social deprivation, marital status, health insurance, housing, employment, and disability. Notably, many of the groups with the lowest rates of COVID-19 vaccine uptake, are also the groups that have been disproportionately affected by the pandemic, with increased morbidity and mortality – further highlighting the significance of vaccine acceptance and uptake among these population subgroups (Mena et al., 2021; Elliott et al., 2021; Gross et al., 2020; Mackey et al., 2021).

With regard to age, most findings showed a high vaccination coverage in individuals at the age of  $\geq$  65 years compared to younger age groups, which reflects the age-based prioritization in the deployment strategies. The studies included in this current review mainly reflect the first 3-4 months after the initiation of the vaccination campaign in late December 2020, when most of the vaccination strategies initially targeted the rapid reduction of severe outcomes, hospitalizations, and deaths from SARS-COV-2 that were predominantly age dependent (Cilloniz et al.,). Hence, the initial phase focused on high-risk individuals, including older people, care home residents and staff, and front-line healthcare workers (Org 2020). With regard to gender, our results were ambiguous and indicated higher vaccine uptake by males in some studies and by females in others. Previous research on other diseases has indicated that among the general population women had higher odds of being vaccinated compared to men, which could be explained by the fact that women have an increased likelihood of seeking medical assistance, using preventive healthcare services, including other vaccinations, and having better health behaviours (Vaidya et al., 2012; Kopsidas et al.,). Studies also have shown that men are less compliant with personal protective measures for COVID-19 (Mahalik et al., 2022; Vardavas et al., 2020; Sultana et al., 2022; Boutsikari et al.,).

Consistent with our findings, prior analyses of COVID-19 vaccination coverage among the general population by county of residence identified lower vaccination coverage in counties with higher social vulnerability related to socioeconomic status (including income, employment status, and education level) and household composition/disability (including age, single-parent household status, and disability status) but higher vaccination coverage in counties with higher social vulnerability related to racial or ethnic minority status and limited English language fluency (Barry et al., 2021; Gharpure et al., 2021). In our review, indexes of social or community deprivation indicated a strong association between increased community deprivation and reduced vaccine uptake. As for income, lower income was associated with a lower likelihood of vaccination in all the included studies that assessed these parameters.

We also identified that ethnicity was strongly associated with disparities in vaccine uptake. According to the findings of the included studies in the current review, which were predominantly from the US, white individuals were more likely to have been vaccinated compared to other racial and ethnic groups. However, in addition to concerns regarding access to vaccination, an important barrier to vaccine uptake in different racial and ethnic groups was vaccine hesitancy (Hamel, 2020). The lack of diverse ethnical representation in clinical trials (Loomba et al., 2021) may have played a role in the hesitancy among these groups. Racial and ethnic disparities in COVID-19 vaccination shown here are mainly based on US and UK context, and are consistent with similar disparities in flu vaccine uptake showing significantly lower rates of flu vaccination among Hispanics, Black individuals, and American Indians/Alaska Natives compared to Asians and non-Hispanic Whites (Control, 2021).

It is important to note that, within the included studies, migrant status was not reported as a variable and hence ethnicity should be interpreted as the ethnicity of origin that does not necessarily reflect recent migratory status. Specifically for migrant populations in the EU/ EEA, evidence shows that some migrant communities may be at increased risk of exposure to SARS-CoV-2, as well as infection with it, and are disproportionately represented in cases, hospitalisations, and deaths (Control, 2021). Also in Europe, some migrant sub-populations have lower coverage of routine vaccination, and higher rates of hesitancy towards vaccination compared with the general population (Control, 2021) (Crawshaw, et al., 2021). Region of birth has also shown to be significantly associated with vaccination coverage for COVID-19 (Martin et al., 2021; Folkhalsomyndigheten, 2022).

While the studies included in this systematic review report for vaccine uptake of at least one dose, it is interesting to note that these differences also persist with uptake of two doses of the vaccine. Notably, the COVID-19 health inequalities monitoring for England (CHIME) tool, including data up to March 2022, indicate a large gap uptake of two doses of the vaccine related to ethnic group, housing status and deprivation quintile (most deprived 78.9% vs. 92.7% for least deprived) (COVID-19 Health Inequalities Monitoring for England (CHIME) tool, 2022). Moreover, currently, there is a continued sub-optimal uptake of the second and subsequent booster doses in the European setting related to social determinants of health, as observed in the UK among elementary trade workers and manual workers (Office for National Statistics -Public Health Data Asset and Coronavirus, 2022) and to country of birth, as observed in Sweden. In Sweden, the uptake for non-Swedish-born citizens of all age categories increased more rapidly than for Swedishborn individuals in the later phases of the rollout, indicating a catchup effect. However, the uptake of the second dose and of subsequent booster doses continued to be lower among non-Swedish born individuals of all ages indicating the importance of social determinants in all phases of the COVID-19 vaccine deployment (Folkhalsomyndigheten, 2022).

#### 5. Strengths and limitations

Our systematic review has a number of strengths, including the systematic approach to study identification, data extraction, and quality appraisal. Some limitations should, however, be acknowledged. This systematic review mainly identified studies from the US, and hence the results here may not reflect the associations identified in other areas of the globe or be relevant to the European context. Moreover, included studies adjusted for important confounders such as age, yet there is a probability that not all factors related to priority groups for vaccination could be accounted for; hence associations require further assessment. Also, significant heterogeneity was noted with regard to population characteristics, sample sizes, methodology and outcomes which did not allow for a meaningful *meta*-analysis. As most of the included studies reflect the first months after the release of the COVID-19 vaccine, they

reflect the prioritization strategies used at that time point that targeted specific population groups and thus impacted vaccine uptake (Control, 2022). Furthermore, due to the time lag between data extraction and manuscript publication, it is likely that additional studies have been published that may possibly alter the association of some of the identified analyses, although the addition of pre-print literature may have mitigated to some extent this time lag. Finally, as some of the studies used county-level estimates it is possible that these factors may vary within large counties and hence only ecological associations can be drawn. These limitations could have an impact on the generalisability of the findings. Future research could also focus on vaccination uptake within the later stages of the COVID-19 pandemic, so as to assess if the socioeconomical determinants that were associated with vaccination uptake differ between initial vaccination and vaccination later on within the COVID-19 pandemic. However, this review provides an overview of the factors associated with the initial vaccination uptake within the context of an emergency public health response and can be used by policymakers when designing roll out strategies in future respiratory pandemics.

#### 6. Conclusion

This review provides evidence of the association between vaccine uptake for SARS-CoV-2 and multiple socioeconomic factors including income, poverty, deprivation, ethnicity, education, and health insurance. This evidence needs to be taken into account to protect socially and medically vulnerable groups and reduce disparities in vaccine uptake within the context of possible emerging variants of concern and future respiratory pandemics. The results of this analysis could provide healthcare professionals with guidance for addressing vaccination inequity among individuals as the significance of social variables in ensuring equitable healthcare delivery is increasingly recognised. While further corroboration is needed to explore the generalisability of these findings across the European setting, these results confirmed the need for an integrated approach to understand the drivers of sub-optimal vaccine acceptance and uptake in a given population and to develop targeted interventions to address these early on.

#### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Data availability

No data was used for the research described in the article.

#### Acknowledgments

We would like to thank Chrysa Chatzopoulou, Katerina Papathanasaki and Konstantinos Skouloudakis for contributing to data archiving and data management.

#### Contributions

CV, JLB, CD, and JES designed the study. KN and KA undertook the literature review and extracted the data with help from CV. JLB and RP developed the search code and performed the search. CV, KN, IL, VV, and KA analysed and interpreted the data. EF, CD, TN, and JES participated in data interpretation along with CV, JLB, RP, KN, KZ, KA, AK, ES, VV, VM and IL. CV and KN prepared the first draft of the manuscript with input from all authors with further revisions by VM. All authors reviewed and revised subsequent drafts and approved the final text.

#### Funding

This report was commissioned by the European Centre for Disease Prevention and Control (ECDC), to the PREP-EU Consortium, coordinated by Dr. Vardavas under specific contract No. 13 ECD.11980 within Framework contract ECDC/2019/001.

#### Data sharing statement

Data sharing is not applicable to this article as no new data were created or analysed in this study.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.pmedr.2023.102319.

#### References

- Agarwal, R., et al., 2021. Socioeconomic privilege and political ideology are associated with racial disparity in COVID-19 vaccination. Proc. Natl. Acad. Sci. USA 118 (33).
- Bagaria, J., et al., 2022. Rapidly adapting primary care sentinel surveillance across seven countries in Europe for COVID-19 in the first half of 2020: strengths, challenges, and lessons learned. Eurosurveillance 27 (26).
- Barry, V., et al., 2021. Patterns in COVID-19 vaccination coverage, by social vulnerability and urbanicity—United States, December 14, 2020–May 1, 2021. Morbid. Mortal. Weekly Rep. 70 (22), 818–824.
- Boutsikari, E.C., et al., 2021. Greek population's perceptions of nonpharmacological interventions towards the first wave of COVID-19 pandemic mitigation: A regressionbased association analysis. Pneumon 34 (3), 1–11.
- Braveman, P., Gottlieb, L., 2014. The social determinants of health: it's time to consider the causes of the causes. Public Health Rep. (Washington, D.C.: 1974) 129 (Suppl 2), 19–31.
- Brown, C.C., Young, S.G., Pro, G.C., 2021. COVID-19 vaccination rates vary by community vulnerability: A county-level analysis. Vaccine 39 (31), 4245–4249.
- Cilloniz, C., et al., 2021. COVID-19 in patients aged 80 years and over during the peaks of the first three pandemic waves at a Spanish tertiary hospital. Pneumon 34 (4), 1–6.
- Control, E.C.f.D.P.a., Guidance on the provision of support for medically and socially vulnerable populations in EU/EEA countries and the United Kingdom during the COVID-19 pandemic. 2020: Stockholm.
- Control, E.C.f.D.a. Flu Disparities Among Racial and Ethnic Minority Groups. 2021; Available from: https://www.cdc.gov/flu/highrisk/disparities-racial-ethnicminority-groups.html.
- Control, E.C.f.D.P.a., COVID-19 situation update for the EU/EEA, as of 29 October 2021. 2021: Stokholm.
- Control, E.C.f.D.P.a., Reducing COVID 19 transmission and strengthening vaccine uptake among migrant populations in the EU/EEA– 3 June 2021. 2021: Stockholm.
- Control, E.C.f.D.P.a., Overview of the implementation of COVID-19 vaccination strategies and deployment plans in the EU/EEA. 21 April 2022. 2022: Stockholm.
- COVID-19 Health Inequalities Monitoring for England (CHIME) tool. 2022 [cited 2022 2 June ]; Available from: https://analytics.phe.gov.uk/apps/chime/.
- Crawshaw, A.F., et al., Defining the determinants of under-vaccination in migrant populations in Europe to improve routine and COVID-19 vaccine uptake: a systematic review. medRxiv, 2021: p. 2021.11.08.21266058.
- Dolby, T., et al., Monitoring sociodemographic inequality in COVID-19 vaccination coverage in England: a national linked data study. medRxiv, 2021: p. 2021.10.07.21264681.
- Dryden-Peterson, S., et al., Disparities in SARS-CoV-2 Vaccination-to-Infection Risk, Massachusetts 2020-2021. medRxiv, 2021: p. 2021.04.14.21255467.
- Elliott, J., Bodinier, B., Whitaker, M., Delpierre, C., Vermeulen, R., Tzoulaki, I., Elliott, P., Chadeau-Hyam, M., 2021. COVID-19 mortality in the UK Biobank cohort: revisiting and evaluating risk factors. Eur. J. Epidemiol. 36 (3), 299–309.
- Folkhalsomyndigheten. Vaccination coverage by country of birth, income and degree of education - week 21. 2022; Available from: https://www.folkhalsomyndigheten.se/ folkhalsorapportering-statistik/statistikdatabaser-och-visualisering/ vaccinationstatistik/statistik-for-vaccination-mot-covid-19/uppfoljning-avvaccination/vaccinationstackning-i-undergrupper/.
- Gharpure, R., et al., 2021. COVID-19 vaccine uptake among residents and staff members of assisted living and residential care communities-pharmacy partnership for long-term care program, December 2020-April 2021. J. Am. Med. Dir. Assoc. 22 (10), 2016–2020 e2.
- Glampson, B., et al., 2021. Assessing COVID-19 vaccine uptake and effectiveness through the north west London vaccination program: retrospective cohort study. JMIR Public Health Surveill. 7 (9) e30010.
- Glatman-Freedman, A., Nichols, K., 2012. The effect of social determinants on immunization programs. Human Vacc. Immunotherap. 8 (7), 916–920.
- Government, M.o.H.C.L., English indices of deprivation 2015: research report. 2015. Gross, C.P., Essien, U.R., Pasha, S., Gross, J.R., Wang, S.-y., Nunez-Smith, M., 2020. Racial and ethnic disparities in population-level Covid-19 mortality. J. Gen. Intern. Med. 35 (10), 3097–3099.

- Preventive Medicine Reports 35 (2023) 102319
- Hamel L., L.L., Munana C., Artiga S., Brodie M. KFF/The Undefeated Survey on Race and Health. 2020; Available from: https://www.kff.org/racial-equity-and-health-policy/ report/kff-the-undefeated-survey-on-race-and-health/.
- Jain, A., et al., 2018. Zoster vaccination inequalities: A population based cohort study using linked data from the UK Clinical Practice Research Datalink. PLoS One 13 (11).
- Khubchandani, J., Sharma, S., Price, J.H., Wiblishauser, M.J., Webb, F.J., 2021. COVID-19 morbidity and mortality in social networks: does it influence vaccine hesitancy? Internat. J. Environ. Res. Public Health 18 (18), 9448.
- Kim, D., Associations of Race/Ethnicity and Other Demographic and Socioeconomic Factors with Vaccine Initiation and Intention During the COVID-19 Pandemic in the United States. medRxiv, 2021: p. 2021.02.16.21251769.
- Kopsidas, I., et al., 2021. COVID-19 pandemic impact on seasonal flu vaccination: A cross-sectional study. Pneumon 34 (2), 1–9.
- Lindemer, E., et al., Counties with lower insurance coverage are associated with both slower vaccine rollout and higher COVID-19 incidence across the United States. medRxiv, 2021: p. 2021.03.24.21254270.
- Loiacono, M.M., et al., 2020. Patient and practice level factors associated with seasonal influenza vaccine uptake among at-risk adults in England, 2011 to 2016: An agestratified retrospective cohort study. Vaccine X 4, 100054.
- Loomba, S., de Figueiredo, A., Piatek, S.J., de Graaf, K., Larson, H.J., 2021. Measuring the impact of COVID-19 vaccine misinformation on vaccination intent in the UK and USA. Nature Human Behav. 5 (3), 337–348.
- MacKenna, B., et al., Trends, regional variation, and clinical characteristics of COVID-19 vaccine recipients: a retrospective cohort study in 23.4 million patients using OpenSAFELY. medRxiv, 2021: p. 2021.01.25.21250356.
- Mackey, K., Ayers, C.K., Kondo, K.K., Saha, S., Advani, S.M., Young, S., Spencer, H., Rusek, M., Anderson, J., Veazie, S., Smith, M., Kansagara, D., 2021. Racial and ethnic disparities in COVID-19-related infections, hospitalizations, and deaths: A systematic review. Ann. Intern. Med. 174 (3), 362–373.
- Mahalik, J.R., Bianca, M.D., Harris, M.P., 2022. Men's attitudes toward mask-wearing during COVID-19: Understanding the complexities of mask-ulinity. J. Health Psychol. 27 (5), 1187–1204.
- Martin, C.A., et al., 2021. SARS-CoV-2 vaccine uptake in a multi-ethnic UK healthcare workforce: A cross-sectional study. PLOS Med. 18 (11).
- Mena, G.E., Martinez, P.P., Mahmud, A.S., Marquet, P.A., Buckee, C.O., Santillana, M., 2021. Socioeconomic status determines COVID-19 incidence and related mortality in Santiago. *Chile.* Science 372 (6545).
- Mendez-Brito, A., El Bcheraoui, C., Pozo-Martin, F., 2021. Systematic review of empirical studies comparing the effectiveness of non-pharmaceutical interventions against COVID-19. J. Infect. 83 (3), 281–293.
- Moher, D., et al., 2009. Preferred reporting items for systematic reviews and metaanalyses: the PRISMA statement. PLoS Med. 6 (7).
- Moola, S., Tufanaru, C., Aromataris, E., Sears, K., Sfetcu, R., Currie, M., Lisy, K., Qureshi, R., Mattis, P., Mu, P., 2020. Chapter 7: Systematic reviews of etiology and risk. JBI Manual Evid. Synthesis.
- Murthy, B.P., et al., 2021. Disparities in COVID-19 vaccination coverage between urban and rural counties—United States, December 14, 2020–April 10, 2021. Morbidity Mortal. Weekly Rep. 70 (20), 759–764.
- Nafilyan, V., et al., 2021. Sociodemographic inequality in COVID-19 vaccination coverage among elderly adults in England: a national linked data study. BMJ Open 11 (7).
- Nguyen, L.H., et al., Racial and ethnic differences in COVID-19 vaccine hesitancy and uptake. medRxiv, 2021.
- Office for National Statistics Public Health Data Asset, N.E.-N.I.M.S. Coronavirus and vaccination rates in people aged 18 to 64 years by occupation and industry, England: 28 February 2022. 2022; Available from: https://www.ons.gov.uk/ peoplepopulationandcommunity/healthandsocialcare/healthinequalities/bulletins/ coronavirusandvaccinationratesinpeopleaged18to64yearsbyoccupationengland/ 28february2022.
- Omaduvie, U., Bachuwa, G., Ayo-Yusuf, O., Vardavas, C., Agaku, I., 2021. State specific estimates of vaccine hesitancy among US adults. Populat. Med. 3 (December), 1–9.
- Organization, W.H., Evidence to recommendations for COVID-19 vaccines: Evidence framework. 2020.
- Perry, M., Akbari, A., Cottrell, S., Gravenor, M.B., Roberts, R., Lyons, R.A., Bedston, S., Torabi, F., Griffiths, L., 2021. Inequalities in coverage of COVID-19 vaccination: A population register based cross-sectional study in Wales, UK. Vaccine 39 (42), 6256–6261.
- Ryerson, A.B., Rice, C.E., Hung, M.-C., Patel, S.A., Weeks, J.D., Kriss, J.L., Peacock, G., Lu, P.-J., Asif, A.F., Jackson, H.L., Singleton, J.A., 2021. Disparities in covid-19 vaccination status, intent, and perceived access for noninstitutionalized adults, by disability status—National Immunization Survey Adult Covid Module, United States, May 30–June 26, 2021. Morbidity Mortality Weekly Rep. 70 (39), 1365–1371.
- Semenza, J.C., Suk, J.E., Tsolova, S., 2010. Social determinants of infectious diseases: a public health priority. Eurosurveillance 15 (27), 19608.
- Singh, G.K., Lee, H., Azuine, R.E., 2021. Marked inequalities in COVID-19 vaccination by racial/ethnic, socioeconomic, geographic, and health characteristics, United States, January 6 – February 15, 2021. Internat. J. Transl. Med. Res. Public Health 5 (2), 103–112.
- Starke, K.R., et al., 2021. The isolated effect of age on the risk of COVID-19 severe outcomes: a systematic review with meta-analysis. BMJ Global Health 6 (12).
- Sultana, M.S., Khan, A.H., Islam, M.R., Hossain, S., Tasdik Hasan, M., Sikder, M.T., 2022. Gender differences in knowledge, attitudes and preparedness to respond to COVID-19 among adults in Bangladesh: A cross-sectional study. Populat. Med. 4 (January), 1–11.
- Sun, Y., Monnat, S.M., 2021. Rural-urban and within-rural differences in COVID-19 vaccination rates. J. Rural Health.

Tram, K.H., et al., 2021. Deliberation, Dissent, and Distrust: Understanding distinct drivers of COVID-19 vaccine hesitancy in the United States. Clin. Infect. Dis.

- Vaidya, V., Partha, G., Karmakar, M., 2012. Gender differences in utilization of
- preventive care services in the United States. J. Womens Health (Larchmt) 21 (2), 140–145.
- Vardavas, C.I., et al., 2022. Prognostic factors for mortality, intensive care unit and hospital admission due to SARS-CoV-2: a systematic review and meta-analysis of cohort studies in Europe. Eur. Respir. Rev. 31 (166), 220098.
- Vardavas, C., Odani, S., Nikitara, K., El Banhawi, H., Kyriakos, C., Taylor, L., Lown, G., Becuwe, N., 2020. Perceptions and practice of personal protective behaviors to prevent COVID-19 transmission in the G7 nations. Populat. Med. 2 (June).
- Vardavas, C., et al., The cost of the COVID-19 pandemic vs the cost-effectiveness of mitigation strategies in the EU/UK/EEA and OECD countries: a systematic review. medRxiv, 2022: p. 2022.05.31.22275813.
- Wang, H., Xu, R., Qu, S., Schwartz, M., Adams, A., Chen, X., 2021. Health inequities in COVID-19 vaccination among the elderly: Case of Connecticut. J. Infect. Public Health 14 (10), 1563–1565.
- Ward, C., Byrne, L., White, J.M., Amirthalingam, G., Tiley, K., Edelstein, M., 2017. Sociodemographic predictors of variation in coverage of the national shingles vaccination programme in England, 2014/15. Vaccine 35 (18), 2372–2378.
- Whiteman, A., et al., 2021. Demographic and social factors associated with COVID-19 vaccination initiation among adults aged≥ 65 years—United States, December 14, 2020–April 10, 2021. Morbidity Mortality Week. Rep. 70 (19), 725–730.
- Wise, J., 2021. Covid-19: EU has lessons to learn from early response to pandemic, say auditors. BMJ 372, n173.
- Xing, K., et al., 2021. Efficacy and safety of COVID-19 vaccines: a systematic review. Zhongguo Dang Dai Er Ke Za Zhi 23 (3), 221–228.
- Yeung, M.P.S., Lam, F.L.Y., Coker, R., 2016. Factors associated with the uptake of seasonal influenza vaccination in adults: a systematic review. J. Public Health 38 (4), 746–753.